

cambridge.org/ags

Crops and Soils Research Paper

Cite this article: Groeneveld M, Grunwald D, Piepho H-P, Koch H-J (2024). Crop rotation and sowing date effects on yield of winter wheat. *The Journal of Agricultural Science* **162**, 139–149. <https://doi.org/10.1017/S0021859624000261>

Received: 20 August 2023

Revised: 8 March 2024

Accepted: 29 April 2024



First published online: 8 May 2024

Keywords:

preceding crop; pre-crop; silage maize; sugar beet; winter oilseed rape

Corresponding author:

Dennis Grunwald;

Email: grunwald@ifz-goettingen.deMarten Groeneveld¹, Dennis Grunwald¹ , Hans-Peter Piepho²  and Heinz-Josef Koch¹

¹Department of Agronomy, Institute of Sugar Beet Research, Holtenser Landstraße 77, 37079 Göttingen, Germany and ²Biostatistics Unit, Institute of Crop Science, University of Hohenheim, Fruwirthstrasse 23, 70599 Stuttgart, Germany

Abstract

Simplified cereal-based crop rotations are widely grown due to economic reasons, leading to the cultivation of wheat after wheat and associated yield losses. In this study, a crop rotation trial was conducted in Northern Germany on a Stagnic Luvisol from 2006 to 2018 with winter wheat after the four most widely used preceding crops in the region (sugar beet, winter wheat, silage maize and winter oilseed rape) in different crop rotations to evaluate potential benefits of different preceding crops. Additionally, the effects of two different sowing dates (2016–2018) and higher crop residue input (whole period) were investigated.

While the pre-preceding crop had no effect, preceding crops winter oilseed rape and sugar beet led to a significantly higher yield of about 1.00 and 0.43 t/ha, respectively, compared to wheat after wheat. This was not modified by crop rotational diversity, including wheat monoculture. Wheat yield tended to be higher for the late sowing date after sugar beet, maize and wheat, while there was no effect of sowing date after oilseed rape. Higher crop residue input led to a significantly higher yield (0.30 t/ha) in wheat after wheat (after pre-preceding crop sugar beet). Overall, sugar beet and winter oilseed rape were found to be favourable preceding crops for winter wheat under the given site conditions. The effect of sowing date on yield and potential modifications of the preceding crop effect by sowing date needs further research in appropriate long-term trials.

Introduction

Since the 1960s crop rotations in Northern Germany comprising winter wheat (*Triticum aestivum* L.) were simplified, i.e. reduced in crop diversity partially through to monoculture, for economic reasons (Christen *et al.*, 1992; Sieling *et al.*, 2005). In 2010, almost 50% of the arable land in the North German state of Lower Saxony was cultivated with simplified crop rotations (Steinmann and Dobers, 2013). Monoculture of winter wheat is simpler to manage than more diverse crop rotations (Angus *et al.*, 2015) but can lead to high yield losses in the first years due to occurrence of root diseases caused by the pathogens *Gaeumannomyces tritici* (Jenkyn *et al.*, 2014) and *Pseudocercospora herpotrichoides*. Generally, yield losses in wheat grown after wheat can range between 8 and 57% compared to wheat grown after more favourable preceding crops (Sieling and Christen, 2015). Alternatively, a sequence of continuous wheat may be broken with the cultivation of an unrelated species, a so-called ‘break crop’ (Angus *et al.*, 2015), by which life cycles of crop-specific pathogens are interrupted (Kirkegaard *et al.*, 2008).

Generally, within a crop rotation, preceding crops were found to have the largest effect on yield of the subsequent crop, while the effect of pre-preceding crops and the overall crop rotation was lower (Sieling and Christen, 2015). Preceding crops including their residues remaining in the field may influence root pathogens, other soil organisms, soil water and soil nitrogen, whereby they can affect the yield of the following crop (Angus *et al.*, 2015). In the case of winter wheat, the incidence and severity of *G. tritici* is crucial for the yield response after different preceding crops (Christen *et al.*, 1992). Besides preceding crops, sowing date influences infestation by *G. tritici* or *P. herpotrichoides*, too: Early sown winter wheat is infested earlier and stronger than late sown (Bateman, 1986; Colbach *et al.*, 1997; Sieling *et al.*, 2007). Sowing date of wheat may also be affected by the preceding crop through different harvest dates. Moreover, sowing of wheat after *Brassica* break crops can be earlier than after wheat as weed populations after these break crops are often low; however, this benefit probably goes unnoticed in many experiments as different sowing dates for the same preceding crop are necessary in field studies to detect it (Angus *et al.*, 2015).

Besides winter wheat, winter oilseed rape (*Brassica napus* L.), sugar beet (*Beta vulgaris* L. var. *altissima* Döll) and maize (*Zea mays* L.) were the most frequently grown preceding crops of winter wheat in Lower Saxony, Germany, in 2009/2010 (Steinmann and Dobers, 2013). Concerning oilseed rape as preceding crop, Sieling *et al.* (2007) found 1 t/ha higher

wheat yield than after winter wheat. Angus *et al.* (2015) determined 0.8 t/ha mean increase of wheat yield after preceding crop oilseed rape compared to wheat after wheat considering different experimental locations in Australia, Europe and North America. After cultivation of oilseed rape, soils in southeastern Australia were more porous, had lower shear strength and more stable aggregates than soils after cultivation of field pea (*Pisum sativum* L.) or barley (*Hordeum vulgare* L.) (Chan and Heenan, 1996). Weed populations can be relatively low in wheat following preceding crop oilseed rape, but it is unclear whether allelopathy or the use of persistent herbicides on oilseed rape is the cause for this (Angus *et al.*, 2015).

Studies with preceding crops sugar beet and (silage) maize before winter wheat are scarce, despite the economic importance and thus widespread cultivation of these crops in rotations with winter wheat. Claupein and Zoschke (1987) found a higher yield of winter wheat in a rotation with preceding crop sugar beet than in continuous winter wheat cultivation by about 0.65 t/ha at experimental station Rauischholzhausen from 1969 to 1985. However, conditions regarding residue treatment in this trial were far from current common practice. Macholdt and Honermeier (2019) noticed a high yield stability in a crop rotation with preceding crop sugar beet, similar to crop rotations with preceding crops winter oilseed rape and field bean, at the same site in a different trial from 1993 to 2017. Berzsenyi *et al.* (2000) found a higher wheat yield in a maize – maize – wheat – wheat rotation compared to wheat monoculture by about 0.38 t/ha in an unfertilized control, while the difference increased to about 0.86 t/ha when averaged over differently fertilized treatments. However, the authors do not mention if maize was grown as silage or grain maize which may severely affect the outcome due to the large difference in aboveground crop residues remaining on the field. Thus, potential benefits of sugar beet and (silage) maize as preceding crops before wheat compared to wheat and other preceding crops require further research.

Beyond the direct impact of preceding crops, diversity of crop rotations was found to affect processing of newly added residues, microbial dynamics and nutrient cycling (McDaniel *et al.*, 2014) and may, thus, ultimately also affect yield of the single crops. However, when winter wheat rotations with different numbers of species (crop diversity) and a wheat monoculture were compared by Smith *et al.* (2008), there was no effect of crop diversity on wheat yield beside a lower yield in monoculture (by around a third) compared to all other treatments. In contrast, Berzsenyi *et al.* (2000) found an effect of the share of wheat in crop rotations on yield: the lower the share, the higher the yield. Further, crop rotations can affect the soil organic matter (Ellerbrock and Gerke, 2016). Although soil organic matter can generally be expected to improve soil fertility, no significant effect on yield could be found in a meta-analysis for Europe (Hijbeek *et al.*, 2017). Macholdt *et al.* (2020) suppose that a higher soil organic carbon content leads to lower yield variability, but this hypothesis could not be confirmed with the collected data in their study. Thus, the overall crop rotation might have an impact on wheat yield beyond the direct preceding crop.

In this study, a crop rotation trial conducted from 2006 to 2018 in Lower Saxony, Germany, was analysed to answer the following research questions: (1) How do preceding crops sugar beet, winter wheat, silage maize and winter oilseed rape affect grain yield of winter wheat and is this modified by the overall crop rotation? (2) Is there a pre-preceding crop effect on grain yield of winter wheat? (3) Does an early (end of September) or late (end of

October) sowing date in combination with different preceding crops influence grain yield and yield components of winter wheat? (4) Is there an effect of the amount of soil organic matter, modified by an increased crop residue input, on grain yield of winter wheat after different preceding crops?

Materials and methods

Site

The study site Harste is located near Göttingen in Lower Saxony, Germany (51°36'23.5"N, 9°51'55.5"E, 155 m a.s.l.). Long-term (1992–2021) mean annual air temperature is 9.5°C and long-term mean sum of annual precipitation is 629 mm (DWD 2022). In 2007–2018, mean air temperature was 9.7°C (range: 8.0–10.6°C) and mean sum of annual precipitation 626 mm (range: 430–892 mm). Soil type at the study site is classified as Stagnic Luvisol (IUSS Working Group WRB, 2014) and soil texture in the upper 30 cm as silty loam (100 g clay/kg soil, 760 g silt/kg soil, 140 g sand/kg soil).

Crop rotations and experimental design

The crop rotation trial Harste was established in 2006 with seven different crop rotations:

1. sugar beet – winter wheat – winter wheat
2. winter wheat monoculture
3. silage maize – winter wheat – winter wheat
4. winter oilseed rape – winter wheat – winter wheat
5. sugar beet – winter wheat – silage maize
6. sugar beet – winter wheat – winter oilseed rape – winter wheat – winter wheat – grain pea
7. sugar beet – winter wheat – winter wheat

Rotation 5 included grain maize instead of silage maize until 2009. As catch crops, mustard (*Sinapis alba* L.) was grown before sugar beet in all crop rotations except for rotation 5 and before silage maize, while phacelia (*Phacelia tanacetifolia* L.) was grown before grain pea. Both catch crops were fertilized with 40–50 kg N/ha. Each crop rotation element was cultivated every year in an α -design, i.e. with complete replications (three in our case) each containing incomplete blocks (in our case six blocks of four plots (16.2 m × 14.0 m) each). Plots of rotation 7 were split into subplots (8.1 m × 14.0 m) for the whole study period and residues of sugar beet were moved from one subplot to the adjacent subplot after each sugar beet harvest, resulting in subplots with double the amount of sugar beet residues or none.

Soil tillage was usually done with a cultivator before sowing at a depth of 15–20 cm. Winter wheat cultivar was Cubus throughout the experimental period. Cubus is rated with a medium ear density, medium to high ear number, medium thousand seed weight (TSW) and a medium grain yield (Bundessortenamt, 2022).

Sowing date was chosen according to preceding crop until 2015, similarly to agricultural practice: early (end of September) after winter wheat and winter oilseed rape and late (end of October) after sugar beet and silage maize. Since this way preceding crop and sowing date were confounded, plots of rotation 2 (wheat monoculture) and plots of the second phase in rotations 1, 3 and 4 (first wheat after sugar beet, silage maize and oilseed rape, respectively) were split into subplots (8.1 m × 14.0 m) with early and late sowing date for the growing seasons of 2016 to

2018 to analyse the possible modification of the preceding crop effect by sowing date differences. For the early sowing date, winter wheat seeding rate was 250–270 seeds/m² and for the late sowing date 350 seeds/m².

Plant protection was conducted according to the specific growing conditions and occurrence of pests and diseases in each year. Generally, except for rotation 7 as described above, crop residues were left on the field. Mineral nitrogen (N) fertilization was adapted to preceding crop (expected N mineralization) and soil mineral N content in 0–90 cm depth (N_{min}) measured in spring each year. Target value as the sum of spring N_{min} and mineral fertilizer N was 240 kg N/ha for preceding crops sugar beet and winter oilseed rape and 260 kg N/ha for preceding crops winter wheat and silage maize. The resulting amount was divided into three to five applications with urea ammonium nitrate solution (except for first application in 2018: ammonium sulphate solution). Total amount of nitrogen applied varied strongly among the years with between 80 and 230 kg N/ha, yet, when keeping in mind the different target values of total available N after the different crops, the overall mean in the study years 2008–2018 was similar for the four preceding crops with 181 (oilseed rape), 187 (sugar beet), 197 (silage maize) and 204 kg N/ha (winter wheat).

Grain yield was determined by machine harvesting the grain fresh matter on a subplot of 1.5 m × 14.0 m of which a sample was taken and dried at 105°C to analyse the dry matter content. Additionally, for the study years 2016–2018, the TSW was determined by weighing while ears were counted by hand after harvest in eight rows at a length of 1 m and scaled up to ears/m². Finally, grains per ear could be calculated from the three aforementioned parameters.

Statistical analyses

All statistical analyses were performed in SAS Version 9.4 (SAS Institute Inc., 2016). Graphs were created in RStudio Version 2022.02.3 (Rstudio, 2022) with R Version 4.2.1 (The R Foundation for Statistical Computing, 2022) and the packages ggplot2 (Wickham, 2016) and ggpubr (Kassambara, 2020).

For the analysis of rotation, preceding crop and pre-preceding crop effects on grain yield in 2008–2018, rotations 1–6 were considered. In cases where plots were split into subplots with different sowing dates (2016–2018), subplots with early sowing date were selected for preceding crops winter oilseed rape and winter wheat, and subplots with late sowing date for preceding crops sugar beet and silage maize in order to be consistent with sowing dates until 2015.

A linear mixed-effects model was fitted with the GLIMMIX procedure of SAS using residual (restricted) maximum likelihood (REML) for estimation of variance components as follows:

$$\begin{aligned} & \text{TIME} + A + A \cdot B + \text{TIME} \cdot A + \text{TIME} \cdot A \cdot B : \text{YEAR} \\ & + \text{YEAR} \cdot A + \text{YEAR} \cdot A \cdot B + \text{REP} + \text{YEAR} \cdot \text{REP} \\ & + \text{BLOCK} \cdot \text{REP} + \text{YEAR} \cdot \text{BLOCK} \cdot \text{REP} + \text{PLOT} \end{aligned} \quad (1)$$

Fixed effects are listed first and separated from random effects by a colon (Piepho *et al.*, 2003). The year effect was partitioned in a fixed part (TIME) and a remaining random part (YEAR) (Loughin *et al.*, 2007; Onofri *et al.*, 2016). TIME is a continuous covariate and describes a linear time trend and YEAR models the year-by-year variation around this trend line. Factors A (rotation)

and B (phase) are fixed effects, while REP (replication), BLOCK·REP (block within replication), and all interactions with YEAR are random effects. Different covariance structures were fitted to account for autocorrelation on the same plot across years. The Akaike information criterion (AIC) value was lowest with compound symmetry structure (Table 1).

Variance components BLOCK·REP and YEAR·A were removed from the model due to values of zero, YEAR·BLOCK·REP was removed after likelihood-ratio test. Fixed effects TIME·A·B and TIME·A were removed from the model after fitting models using maximum likelihood (ML) for estimation of variance components and carrying out likelihood-ratio tests. Custom contrasts were defined with the statement ESTIMATE. P values were adjusted by a simulation-based approach using the multivariate t distribution (option ADJUST = SIMULATE).

To analyse the rotation effect, all phases within rotations with preceding crop sugar beet were compared with each other, both phases within rotations with preceding crop winter oilseed rape were compared and all phases within rotations with preceding crop winter wheat were compared pairwise in a first contrast. This contrast includes the analysis of pre-preceding crop effect. Preceding crop effect was investigated by comparing all phases within rotations with a specific preceding crop with phases within rotations with a specific different preceding crop in a second contrast.

For the analysis of preceding crop and sowing date effects on grain yield in 2016–2018, rotations 1–4 were considered. A linear mixed-effects model was fitted with the MIXED procedure using REML for estimation of variance components as follows:

$$\begin{aligned} & A + B + A \cdot B + \text{REP} : \text{YEAR} + \text{YEAR} \cdot A + \text{YEAR} \\ & \cdot B + \text{YEAR} \cdot A \cdot B + \text{YEAR} \cdot \text{REP} + \text{YEAR} \cdot A \cdot \text{REP} \\ & + \text{SUBPLOT} \end{aligned} \quad (2)$$

where factors A (preceding crop), B (sowing date) and REP (replication) are fixed effects, while YEAR, SUBPLOT, and all interactions with YEAR are random effects. Different covariance structures were fitted to account for autocorrelation on the same main- and subplot across years. AIC was lowest with autoregressive structure of order 1 with heteroscedastic errors by year (Table 2).

Variance components YEAR·A·B and YEAR·REP were removed from the model due to values of zero. Model (2) was

Table 1. Models regarding grain yield 2008–2018 with different within-plot covariance structures

Model	df	–2 Log-likelihood	AIC
ID	30	612.30	626.30
CS	31	611.12	625.12
AR(1)	31	611.64	627.64
Toeplitz	40	602.94	634.94
CSH	41	600.05	636.05
ARH(1)	41	600.39	636.39

df, degrees of freedom (fixed and random effects); AIC, Akaike information criterion; ID, independent (no within-plot correlation); CS, compound symmetry; AR(1), autoregressive of order 1; CSH, compound symmetry with heteroscedastic errors by year; ARH(1), autoregressive of order 1 with heteroscedastic errors by year.

Table 2. Models regarding grain yield 2016–2018 with different within-main- and subplot covariance structures

Model	df	-2 Log-likelihood	AIC
ID	17	80.8	92.8
CS	19	76.2	92.2
AR(1)	19	75.3	91.3
Toeplitz	21	71.3	91.3
CSH	23	66.6	88.6
ARH(1)	23	65.6	85.6

df, degrees of freedom (fixed and random effects); AIC, Akaike information criterion; ID, independent (no within-plot correlation); CS, compound symmetry; AR(1), autoregressive of order 1; CSH, compound symmetry with heteroscedastic errors by year; ARH(1), autoregressive of order 1 with heteroscedastic errors by year.

fitted for the dependent variables ear density, grains per ear and TSW, too, with independent uncorrelated plots (lowest AIC).

The effect of removing or adding aboveground sugar beet residues, representing differences in soil organic matter content, was analysed considering rotation 7 in the MIXED procedure using REML for estimation of variance components with the following linear mixed-effects model:

$$\begin{aligned} & \text{TIME} + A + B + \text{TIME} \cdot A + \text{TIME} \cdot B + A \cdot B \\ & + \text{TIME} \cdot A \cdot B : \text{YEAR} + \text{YEAR} \cdot A + \text{YEAR} \cdot B \\ & + \text{YEAR} \cdot A \cdot B + \text{REP} + \text{YEAR} \cdot \text{REP} + \text{YEAR} \cdot A \\ & \cdot \text{REP} + \text{SUBPLOT} \end{aligned} \quad (3)$$

The year effect was partitioned in a fixed and a random part as in model (1). TIME is a continuous covariate and describes a linear time trend and YEAR models the year-by-year variation around this trend line. Factors A (preceding crop) and B (residues) are fixed effects, while REP, SUBPLOT and all interactions with YEAR are random effects. AIC was lowest for autoregressive structure of order 1 for main- and subplots. Variance components REP, YEAR·REP and YEAR·B were removed from the model due to values of zero. Fixed effects TIME·A·B and TIME·A were removed from the model after fitting models using ML for estimation of variance components and carrying out likelihood-ratio tests.

Denominator degrees of freedom in Wald-type *F* tests were approximated after Kenward and Roger (1997). Mean values of model (2) were compared with the statement LSMEANS (option ADJUST = TUKEY) for significant main effects. If interactions of model (2) or (3) were significant ($P < 0.05$), custom contrasts were defined with the statement LSMESTIMATE and mean values were compared separately for each preceding crop. *P* values were adjusted by a simulation-based approach using the multivariate *t* distribution (option ADJUST = SIMULATE). For all models, normality of residuals and homogeneity of variance were checked using residual plots.

Results

Crop rotation, preceding crop and pre-preceding crop effects

The overall crop rotation did not significantly modify the effect of the preceding crop on wheat yield (Fig. 1). Wheat yield after sugar

beet ranged between 8.7 ± 0.28 t/ha (standard error) and 8.8 ± 0.28 t/ha across three different crop rotations, while wheat yield after wheat ranged between 8.2 ± 0.28 t/ha and 8.5 ± 0.28 t/ha in five different crop rotations, and wheat yield after oilseed rape was 9.3 ± 0.28 t/ha in both studied crop rotations. In any case, differences between the wheat yield after a given preceding crop across different crop rotations were not significant. Thus, wheat yields after the same preceding crop were averaged across crop rotations for the following analyses.

Averaged over all crop rotations considered and for the typical wheat sowing dates after the single preceding crops, yield of winter wheat was significantly ($P < 0.01$) higher after preceding crop winter oilseed rape than after all other preceding crops (Fig. 2). Yield was significantly higher than after sugar beet by about 0.6 ± 0.13 t/ha ($P < 0.001$), than after winter wheat by about 1.0 ± 0.12 t/ha ($P < 0.001$), and than after silage maize by about 0.6 ± 0.17 t/ha ($P < 0.01$) on average. After sugar beet, yield was significantly higher than after winter wheat by about 0.4 ± 0.10 t/ha ($P < 0.001$) on average, while wheat yield after maize was on tendency ($P = 0.0567$) higher than after wheat.

If sown at the same sowing date, differences in wheat yield after the single preceding crops varied for the two sowing dates. For an early sowing (end of September), wheat yield was higher after oilseed rape than after the other preceding crops, while among the other preceding crops no differences were found (Fig. 3). For a late sowing (end of October), wheat yield after oilseed rape was higher than after wheat, while yields after sugar beet and silage maize were not statistically different than yields after either wheat or oilseed rape.

Sowing date and crop residue effects

The effect of sowing date on yield was dependent on preceding crop (Table 3). After sugar beet, wheat yield was significantly ($P < 0.01$) higher for the late sowing date than for the early sowing date by about 0.6 ± 0.17 t/ha on average (Fig. 4a). After the other preceding crops, sowing date did not significantly affect wheat yields, although yields were numerically higher for the late sowing date after silage maize and winter wheat, while there was no difference at all after winter oilseed rape.

Interaction between preceding crop and sowing date was not significant for yield components ear density, grains per ear and TSW (Table 3). On average across sowing dates, grain number was significantly ($P < 0.05$) higher after winter wheat than after silage maize by about 4 grains/ear (Table 4). TSW after preceding crops sugar beet and silage maize was about 3.6 g and 3.8 g higher than after winter wheat, respectively ($P < 0.05$ and < 0.01), while TSW after winter oilseed rape was intermediate (Table 4). The difference of TSW between early sowing date and late sowing date was 1.5 g ($P < 0.05$) on average across preceding crops (Table 4), while ear density and grains per ear were not affected by sowing date.

In the sugar beet – winter wheat – winter wheat rotation with artificially created differences in sugar beet residues, yield of winter wheat after winter wheat was significantly ($P < 0.01$) higher on the subplot with doubled sugar beet residues by about 0.3 t/ha on average (Table 5, Fig. 5). Wheat yield after preceding crop sugar beet was not affected by the residue treatment. Also, yield decrease over time for wheat after wheat was significantly ($P < 0.05$) lower with doubled residues compared to no residues on average (Table 5).




Figure 1. Crop rotation effect on wheat grain yield (cv. Cubus) in long-term crop rotation trial Harste, 2008–2018, with preceding crops (a) sugar beet, (b) winter oilseed rape and (c) winter wheat. •: Least squares means. $n = 33$, preceding crop sugar beet in rotation 1: $n = 32$. Means with the same preceding crop marked with a common letter are not significantly different ($P > 0.05$, multiple t tests with adjusted P values). 1: SB – WW – WW, 2: WW continuous, 3: SM – WW – WW, 4: WR – WW – WW, 5: SB – WW – SM, 6: SB – WW – WR – WW – WW – GP, SB: sugar beet, WW: winter wheat, SM: silage maize, WR: winter oilseed rape, GP: grain pea.




Figure 2. Preceding crop effect in 2008–2018 on yield of winter wheat (cv. Cubus) in the long-term crop rotation trial in Harste with wheat sown early after oilseed rape and wheat and late after silage maize and sugar beet. •: Least squares means, also shown by numbers next to boxes. Means with no letter in common are significantly different ($P < 0.05$, multiple t tests with adjusted P values).

Discussion

Preceding crop and sowing date effects

Yield of winter wheat after preceding crop winter oilseed rape was significantly ($P < 0.001$) higher than after preceding crop winter wheat by about 1.0 t/ha or 10.7% on average, when considering the usual sowing dates of wheat after these crops, which would be rather early. This positive effect of winter oilseed rape is in line with results of Sieling *et al.* (2007) and Sieling and Christen (2015) from German study sites but higher than the mean increase of wheat yield (0.8 t/ha) after preceding crop oilseed rape determined in the review by Angus *et al.* (2015). However, Angus *et al.* (2015) considered different experimental locations in Australia, Europe and North America in their review, thus site conditions were probably diverse and not necessarily comparable to the study site Harste. For Germany, Weiser *et al.* (2018)

found a break crop benefit of winter oilseed rape of 0.56 t/ha (7.3%) in a dataset containing randomly selected farms. However, Weiser *et al.* (2018) compared yield after preceding crop winter oilseed rape to yield after cereals in general (mainly (82%) wheat, but also winter and spring barley, rye, triticale and further cereals other than wheat, without maize and oats), in contrast to yield after wheat in this study. After other cereals than wheat, severity of *G. tritici* and *P. herpotrichoides* can be lower than in continuous cultivation of wheat and therefore the negative impact on yield may not be as high (Schönhammer and Fischbeck, 1987a).

Another possible difference to the other mentioned studies might be the sowing date. While a sowing date difference of around four weeks did not affect the wheat yield after oilseed rape in the present study, wheat grown after wheat showed a (non-significantly) higher yield for the later sowing date, resulting




Figure 3. Preceding crop effect in 2016–2018 on yield of winter wheat (cv. Cubus) in the long-term crop rotation trial in Harste for (a) an early sowing date (end of September) and (b) a late sowing date (end of October). Least squares means with standard errors from a linear mixed-effects model. Means with no letter in common are significantly different ($P < 0.05$, multiple t tests with adjusted P values). SB: sugar beet, WW: winter wheat, SM: silage maize, WR: winter oilseed rape.

Table 3. P values from Wald-type F tests for preceding crop and sowing date effect on yield and yield components ear density, grains per ear and thousand seed weight (TSW) from 2016 to 2018

Effect	df	Ear density	Grains/ear	TSW	Yield
Replication	2	0.2772	0.2590	0.4250	0.0239
Preceding crop	3	0.1375	0.0057	0.0074	0.0466
Sowing date	1	0.9873	0.1942	0.0161	0.1024
Preceding crop \times sowing date	3	0.8341	0.7819	0.7342	0.0294

df, numerator degrees of freedom.

in a lower relative yield benefit for wheat after oilseed rape for later sowing dates. Thus, even if wheat after oilseed rape and after wheat were sown on the same date in the other studies, differences might have been lower if wheat sowing took place later than in this study.

Regarding potential mechanisms leading to the observed benefits, no continuous measurement programme was in place throughout the study years in our trial. However, another study on the same trial for the study year 2018 found a positive effect of oilseed rape as preceding crop on the soil microbial biomass of subsequent wheat when compared to the other preceding crops considered here (Hamer *et al.*, 2021). For later years of the trial beyond the temporal scope of this study, Arnhold *et al.* (2023a) further found a higher subsoil (30–120 cm) root length density of wheat grown after oilseed rape than after wheat which was also independent of take-all occurrence. However, this positive effect was not connected to soil structural benefits of oilseed rape as preceding crop as might have been expected, at least not in the upper 45 cm of soil by spring, as found in a separate study on the same plots (Arnhold *et al.*, 2023b). Also, N benefits by oilseed rape as preceding crop might have been low since soil mineral N levels in spring were taken into account for

the calculation of N fertilization as was also an expected higher N mineralization, resulting in a lower N target value after oilseed rape than after wheat. Thus, N benefits after oilseed rape might primarily occur in the growth period before fertilization.

After silage maize, yield was on tendency higher than after winter wheat by about 0.38 t/ha when considering the usual sowing dates of wheat after these crops (early after wheat, late after maize). This yield advantage corresponds well to the 0.38 t/ha higher yield of wheat after maize compared to wheat after wheat found by Berzsenyi *et al.* (2000) without N fertilization. However, Berzsenyi *et al.* (2000) did not specify whether grain or silage maize was cultivated in their study. As the maize benefit in their study was even higher for regularly fertilized treatments (0.86 t/ha) and Berzsenyi *et al.* (2000) speculate that the reason for the benefit might be related to soil physical and biological improvements following supplemental C sources from crop residues, the higher benefit of regularly fertilized maize as preceding crop to winter wheat compared to the present study seems to be connected to the cultivation as grain maize. In the case of silage maize cultivation, as in this trial, aboveground maize residue input is reduced to low amounts which can be expected to lead to a decrease in soil organic C, as also found in the studied trial (Grunwald *et al.*, 2021), which may ultimately lead to negative effects on crop yields. In agricultural practice, this is often mitigated by organic fertilization; in the studied trial, however, no organic fertilizer was used. The benefit of silage maize as preceding crop before winter wheat under practical conditions with organic fertilization may thus be higher than found in this study.

Nonetheless, when sown at the same date, wheat after wheat and after silage maize showed lower and non-significant yield differences, as shown for the subplots with different sowing dates from 2016 to 2018 in this study. This suggests that at least part of the benefit of silage maize as a preceding crop in contrast to wheat is owed to the later harvest of the former and the resulting later sowing of wheat which would be a rather indirect effect in contrast to an effect based directly on traits of the contrasting crops. In future studies this needs to be addressed, either by




Figure 4. Sowing date effect as dependent on preceding crop in 2016–2018 on yield of winter wheat (cv. Cubus) in the long-term crop rotation trial in Harste. Least squares means with standard errors from a linear mixed-effects model. *P* values from *t* tests for sowing date effect separated by preceding crop (adjusted by a simulation-based approach using the multivariate *t* distribution). early: end of September, late: end of October.

Table 4. Preceding crop and sowing date effects in 2016–2018 on yield structure components of winter wheat (cv. Cubus) in the long-term crop rotation trial Harste

	Ear density (Ears/m ²)	Grains/ear	TSW (g)
<i>Preceding crop</i>			
Sugar beet	478 ± 17.2	38 ± 2.1	41.7 ± 1.04
Winter wheat	482 ± 17.2	40 ± 2.1	38.1 ± 1.04
Silage maize	497 ± 17.2	36 ± 2.1	42.0 ± 1.04
Winter oilseed rape	526 ± 17.2	38 ± 2.1	40.7 ± 1.04
<i>Sowing date</i>			
Early (end of September)	496 ± 15.2	37 ± 2.2	41.4 ± 0.97
Late (end of October)	496 ± 15.2	39 ± 2.2	39.9 ± 0.97

TSW, thousand seed weight. Least squares means with standard errors from a linear mixed-effects model.

Table 5. Wald-type *F* tests for time trend, preceding crop effect, and residues effect (type III sums of squares) for crop rotation 7 in 2008–2018

Effect	df ₁	df ₂	<i>F</i>	<i>P</i>
Time	1	9.12	13.29	0.0053
Preceding crop	1	9.74	3.27	0.1034
Residues	1	20.3	6.09	0.0214
Preceding crop × Residues	1	13.3	6.69	0.0237
Time × Residues	1	21.5	6.97	0.0151

df₁, numerator degrees of freedom; df₂, denominator degrees of freedom. Degrees of freedom with Kenward-Roger approximation.

including several sowing dates for all preceding crops or by setting one sowing date for all treatments in order to understand the mechanisms leading to differences in wheat yield.

Preceding crop sugar beet led to a significantly ($P < 0.001$) higher yield of winter wheat compared to wheat after wheat of about 0.43 t/ha or 4.9%. Although sugar beet is known as beneficial preceding crop, published studies reporting yield of wheat after sugar beet in comparison to other preceding crops are scarce. In a long-term trial at experimental station Rauischholzhausen near Gießen, Claupein and Zoschke (1987) found a higher wheat yield of about 0.65 t/ha or 13% after sugar beet compared to a winter wheat monoculture over 17 years. However, in that study the straw management also differed between the two rotations with considerable benefits for the crop rotation with sugar beet which might have superimposed the pure preceding crop effect. Concerning yield stability, Macholdt and Honermeier (2019) found benefits for a crop rotation with sugar beet before winter wheat in a long-term trial which were comparable to rotations with oilseed rape and field bean before wheat. Although no yield values were included in that study, this underlines the apparently beneficial effect of sugar beet before wheat. However, similar to maize, the positive yield effect of wheat grown after sugar beet compared to wheat grown after wheat is considerably lower and non-significant when the same sowing date for wheat is chosen after both preceding crops. Thus, future studies need to clear up if there are advantages of sugar beet related to certain crop traits or if the positive effect found here is rather an indirect one connected mainly to the choice of wheat sowing date.

As mentioned above, the split-up of the plots in 2016–2018 to investigate the effect of an early (end of September) and a late (end of October) sowing date after each preceding crop led to contrasting results. Concerning sugar beet, winter wheat and silage maize, the late sowing date led to a higher yield of subsequent winter wheat, but the difference was only significant after sugar beet (Fig. 4). After preceding crop winter oilseed rape, sowing date did not affect yield of wheat. Winter oilseed rape may improve the soil structure (Schönhammer and Fischbeck, 1987b; Chan and Heenan, 1996; Sieling *et al.*, 2007; Kirkegaard *et al.*, 2008) and is harvested earlier than silage maize and sugar beet. Therefore, time for cultivation and seedbed preparation is longer




Figure 5. Effect of sugar beet residue management in 2008–2018 after preceding crops (a) sugar beet and (b) winter wheat on yield of winter wheat (cv. Cubus) in a sugar beet – winter wheat – winter wheat rotation in the long-term crop rotation trial in Harste. •: Least squares means, also shown by numbers next to boxes. $n = 33$ (preceding crop winter wheat, residues doubled: $n = 31$). Means with the same preceding crop with no letter in common are significantly different ($P < 0.05$, multiple t tests with adjusted P values).

after oilseed rape than after the latter. Additionally, weed populations can be low after winter oilseed rape, whereby an earlier sowing of wheat without detrimental effects on yield, such as for the other preceding crops in this study, is possible (Angus *et al.*, 2015). A later sowing date, therefore, possibly had no advantages regarding root development and competition with weeds in wheat following winter oilseed rape. The early sowing date after preceding crops winter wheat, silage maize and sugar beet, however, led to a shorter time for cultivation and seedbed preparation. In combination with non-inversion tillage and possibly a generally poorer soil structure compared to oilseed rape as preceding crop, this may have caused comparatively poorly developed wheat roots, which can increase vulnerability to stress and root diseases, and consequently cause yield losses. This seems to be most clear for the early wheat sowing date after sugar beet which took place immediately after the typically rather late harvest of sugar beet. However, these assumed mechanisms need to be confirmed by accordingly planned studies.

Regarding the yield components, TSW in 2016–2018 was significantly ($P < 0.05$) higher after preceding crops sugar beet and silage maize than after winter wheat but not after winter oilseed rape. TSW of wheat was significantly ($P < 0.01$) higher after an early sowing date than after a late sowing date after all preceding crops (Table 4). An earlier sowing generally prolongs the grain filling period and leads to a higher TSW (Ortiz-Monasterio *et al.*, 1994; Ozturk *et al.*, 2006; Koppensteiner *et al.*, 2022). Also, the slightly (non-significantly) lower grain number per ear after an early sowing date seems to be balanced out by a the higher TSW later in the season. Despite this, early sowing is one of the biggest risk factors for an infection with *G. tritici* in wheat after wheat (Jenkyn *et al.*, 2014), which can reduce TSW (Sieling *et al.*, 2005). The higher TSW after early sowing in this study possibly indicates that no considerable infection with *G. tritici* occurred in the observed years, including wheat grown after wheat (see also below).

Crop rotation and crop residue effects

The diversity of different crop rotations had no significant effect on grain yield of winter wheat in this trial. There was neither a difference of wheat yield between wheat after wheat within crop

rotations and wheat monoculture nor between wheat after sugar beet or winter oilseed rape in simpler and in more diverse crop rotations. Contrastingly, Berzsenyi *et al.* (2000) found that yield of wheat decreased with an increasing share of wheat (25% to 100%) in the crop rotation. However, this effect on yield could possibly rather be attributed to the preceding crop than to the share of wheat in the rotation. This view would confirm the observation of Sieling and Christen (2015) that the preceding crop effect superimposes a possible crop rotation effect. Nonetheless, Sieling and Christen (2015) also found a lower yield of wheat in short rotations with a high share of cereals. The most diverse rotation in Harste, rotation 6, consisted of four different crops (sugar beet, winter wheat, winter oilseed rape and field pea) and two catch crops (mustard and phacelia). Share of winter wheat was 50% in this rotation. Rotation 5 consisted of three different crops (sugar beet, winter wheat and silage maize) and one catch crop (mustard). This rotation had the lowest share of winter wheat (33.3%), while continuous cultivation of winter wheat had the highest (100%). The lack of an effect of these large differences in wheat share in the total rotation might be connected to the relatively high use of pesticides in all rotations in Harste. Andert *et al.* (2016) assume that more diverse crop rotations can reduce use of herbicides and fungicides. Although reduction of pesticides was not investigated in this study as all rotations were treated in the same way, the cultivation of spring crops in a crop rotation, like sugar beet, silage maize or field pea, could potentially be beneficial in the control of annual grass weeds and thus offer possibilities regarding wheat yield maintenance at lower pesticide inputs.

In contrast to previous studies from Kirkegaard and Ryan (2014) and Sieling and Christen (2015), no pre-preceding crop effect could be found in this study. Angus *et al.* (2015) compared data from experiments in Australia, Canada, Sweden and the United Kingdom and found that the yield effect of break crops like oilseed rape, oats or grain legumes on the second wheat after a break-crop was less pronounced than the preceding crop effect on the first wheat (Angus *et al.*, 2015). After oilseed break-crops, the preceding crop effect was less persistent in Sweden than in Australia, leading to the assumption that the persistence of the preceding crop effect (which is similar to pre-preceding crop

effect) is higher under arid conditions and more inconsistent in humid and semi-arid conditions. This could be due to longer suppression of pathogens after a single break-crop under dry conditions, or the longer pathogen persistence after wheat (Angus *et al.*, 2015). In a humid climate, Sieling and Christen (2015) found an effect of the pre-preceding crop in wheat after winter oilseed rape but not in wheat after wheat. This indicates that under central European conditions the preceding crop effect is clearly more important than the pre-preceding crop effect (Sieling and Christen, 2015).

The take-all disease caused by *G. tritici* can lead to high yield losses in the first years of continuous winter wheat cultivation (Jenkyn *et al.*, 2014) but yield losses can decrease subsequently through the take-all decline phenomenon (Gerlagh, 1968). Yield data from the studied trial does not suggest that take-all decline occurred in continuous winter wheat cultivation, although *G. tritici* was not rated or analysed in this study. Typically, yield decreases until the third to fourth year after a high incidence of *G. tritici* and increases slightly in following years due to a decrease of *G. tritici* incidence without reaching the yield level of the first year again (Gerlagh, 1968; Sieling and Hanus, 1990; Jenkyn *et al.*, 2014). However, in the studied trial, yield of the wheat monoculture decreased and increased over the years without a clear trend hinting at take-all decline (data not shown). One requirement for the occurrence of take-all decline phenomenon is a heavy infection with *G. tritici* (Jenkyn *et al.*, 2014). Hence, if there had been no heavy infection with *G. tritici* in the continuous cultivation of winter wheat, a take-all decline might in turn have not occurred. This might explain the lack of a difference in wheat yield between the long-term wheat monoculture and the second wheat grown after a pre-preceding break crop.

Interestingly, however, for later years in the same trial (2021 and 2022), Arnhold *et al.* (2023a) rated the occurrence of *G. tritici* and correlated these data with grain yield for the first and second wheat grown after oilseed rape as well as wheat monoculture. They found a strong disease severity in rotational wheat grown after wheat and a substantially lower severity in wheat monoculture in a wet year (2021), hinting at the occurrence of a take-all decline, at least in later years of the trial. For the study year in question, take-all severity correlated well with grain yield which was significantly lower for the more strongly infested rotational wheat after wheat than for the wheat monoculture. In a dry study year (2022), without the occurrence of take-all, no difference in yield was found. This may further highlight the strong differentiation between single study years in terms of weather conditions for pathogens and the dependence of a beneficial effect of monoculture cultivation on weather conditions of a single year.

In rotation 7 (sugar beet – winter wheat – winter wheat), sugar beet residues were moved from one subplot to the adjacent subplot after harvest. This higher amount of organic input (doubled sugar beet residues) led to a significantly ($P < 0.05$) higher yield in winter wheat after winter wheat by about 0.30 t/ha on average over eleven years compared to a lower organic input (sugar beet residues removed). For these plots, Grunwald *et al.* (2021) found that doubled sugar beet residues caused significantly higher soil organic carbon stocks in 0–10 cm soil depth by 2018/19. Soil organic carbon is an important indicator of soil quality (Reeves, 1997). However, despite the general improvement of numerous soil properties by an increased soil organic matter, Hijbeek *et al.* (2017) found no yield increment of winter wheat in Europe through organic input in a meta-analysis. In contrast, wheat yield responded positively to soil organic matter or soil

organic carbon increases in China (Pan *et al.*, 2009), Uruguay (Rubio *et al.*, 2021) and in a global meta-analysis (Oldfield *et al.*, 2019). The higher soil organic carbon stocks in the rotation with doubled sugar beet residues in Harste found by Grunwald *et al.* (2021) and the higher yield found in this study in this rotation potentially indicate a yield benefit of soil organic carbon on yield of winter wheat. However, this benefit seems to occur solely after an unfavourable preceding crop such as winter wheat and not after sugar beet. Thus, the rather positive preceding crop effect of sugar beet possibly superimposed the benefit of a higher soil organic matter content. Still, an even stronger effect for wheat after wheat and possibly also a positive effect on wheat after sugar beet might be expected after a longer period than considered here, as soil organic matter levels in the soil need to build up over time. Thus, in the first few years of this treatment, there might have been no effect at all while the benefit found here might be caused by advantages in the latter half of the sampling period.

Practical considerations

In Lower Saxony in 2009/10, 30.0% of winter wheat was cultivated after winter wheat, 19.3% was grown after winter oilseed rape, 15.4% after sugar beet and 12.5% after maize on average (Steinmann and Dobers, 2013). As cultivation of wheat after wheat will be restricted in the European Union (EU 2021/2115), wheat after wheat must be substituted in crop rotations if the farmer wants to receive direct payments from the European Union. European farmers generally prefer changes in crop rotation to other management practices like growing cover crops or wildflower strips to contribute to biodiversity enhancement (Kleijn *et al.*, 2019), indicating a certain potential to an increased cultivation of wheat after beneficial break crops. Beyond the studied preceding crops in this study, other possible alternatives to wheat grown after wheat in crop rotations like other cereals (barley, oat or rye) or grain legumes (pea or bean) might be considered in future studies. Nevertheless, in this study, sugar beet and winter oilseed rape were found to be favourable preceding crops for winter wheat under current site conditions in southern Lower Saxony and can be seen as reasonable choices and beneficial alternatives to wheat in crop rotations as preceding crops before winter wheat.

Conclusion

In this study on a long-term trial, preceding crop was the crucial factor influencing grain yield of winter wheat, while crop rotation diversity, different shares of winter wheat and pre-preceding crop were less important. Sugar beet and winter oilseed rape were beneficial preceding crops for winter wheat and should be cultivated in crop rotations containing winter wheat. However, this may be influenced by the chosen wheat sowing date which should be taken into account in future studies. In general, under favourable conditions for arable cropping in Northern Germany, a late sowing date in the second half of October seems to be beneficial, except after oilseed rape for which no such effect was found. A further positive effect on wheat yield by possibly manageable factors was found to be a higher soil organic matter content, in particular for wheat after wheat. However, the reasons for the effects of preceding crop, sowing date and soil organic matter on wheat grain yields were not investigated in this study and need to be clarified in further studies. For the studied trial, an investigation

into the mechanisms involved in the preceding crop effects is currently ongoing, focusing on above- and particularly belowground crop residues and how different preceding crops may affect the growing conditions of subsequent wheat, especially in the rhizosphere.

Acknowledgements. We sincerely thank Ines Wiese, Fritz Evers, Wilfried Hübener, Michael Gerhardy, Marten Steinke, Kristina Winkelbach and Christina Jung for excellent technical assistance.

Author contributions. HJK designed and supervised the field experiment including data gathering. MG conducted the statistical data analysis which was supervised by HPP and wrote the original draft which was supervised by DG. All authors contributed to the discussion and interpretation of the results, reviewed the manuscript and approved the final version.

Funding statement. This research received no specific grant from any funding agency, commercial or not-for-profit sectors.

Competing interests. None.

Ethical standards. Not applicable.

References

- Andert S, Bürger J, Stein S and Gerowitt B (2016) The influence of crop sequence on fungicide and herbicide use intensities in North German arable farming. *European Journal of Agronomy* **77**, 81–89.
- Angus JF, Kirkegaard JA, Hunt JR, Ryan MH, Ohlander L and Peoples MB (2015) Break crops and rotations for wheat. *Crop and Pasture Science* **66**, 523–552.
- Arnhold J, Grunwald D, Braun-Kiewnick A and Koch HJ (2023a) Effect of crop rotational position and nitrogen supply on root development and yield formation of winter wheat. *Frontiers in Plant Science* **14**, 1265994.
- Arnhold J, Grunwald D, Kage H and Koch HJ (2023b) No differences in soil structure under winter wheat grown in different crop rotational positions. *Canadian Journal of Soil Science* **103**, 642–649.
- Bateman GL (1986) Effects of triadimenol-containing seed treatment on winter wheat infected with take-all. *Zeitschrift für Pflanzenkrankheiten und Pflanzenschutz* **93**, 404–414.
- Berzsenyi Z, Györfly B and Lap D (2000) Effect of crop rotation and fertilisation on maize and wheat yields and yield stability in a long-term experiment. *European Journal of Agronomy* **13**, 225–244.
- Bundessortenamt (2022) Beschreibende Sortenliste. Bundessortenamt. Available at https://www.bundessortenamt.de/bsa/media/Files/BSL/bsl_getreide_2022.pdf (accessed on 2 September 2022).
- Chan KY and Heenan DP (1996) The influence of crop rotation on soil structure and soil physical properties under conventional tillage. *Soil and Tillage Research* **37**, 113–125.
- Christen O, Sieling K and Hanus H (1992) The effect of different preceding crops on the development, growth and yield of winter wheat. *European Journal of Agronomy* **1**, 21–28.
- Claupen W and Zoschke M (1987) Einfluß langjähriger Winterweizen-Monokultur auf Ertragsbildung, Krankheitsbefall und Nematodenbesatz im Vergleich zum Winterweizenanbau in der Fruchtfolge. *Journal of Agronomy and Crop Science* **158**, 227–235.
- Colbach N, Lucas P and Meynard JM (1997) Influence of crop management on take-all development and disease cycles on winter wheat. *Phytopathology* **87**, 26–32.
- DWD (2022) Climate Data Center (CDC). Deutscher Wetterdienst (DWD). Available at <https://cdc.dwd.de/portal/> (accessed on 7 April 2022).
- Ellerbrock RH and Gerke HH (2016) Analyzing management-induced dynamics of soluble organic matter composition in soils from long-term field experiments. *Vadose Zone Journal* **15**, 1–10.
- EU 2021/2115. Regulation (EU) 2021/2115 of the European Parliament and of the Council. Official Journal of the European Union.
- Gerlagh M (1968) Introduction of *Ophiobolus graminis* into new polders and its decline. *Netherlands Journal of Plant Pathology* **74**, 1–97.
- Grunwald D, Götz P and Koch H-J (2021) Soil organic carbon stocks in sugar beet rotations differing in residue management and associated rotational crop species. *Journal of Plant Nutrition and Soil Science* **184**, 556–561.
- Hamer U, Meyer MUT, Meyer UN, Radermacher A, Götz P, Koch HJ and Scherber C (2021) Soil microbial biomass and enzyme kinetics for the assessment of temporal diversification in agroecosystems. *Basic and Applied Ecology* **53**, 143–153.
- Hijbeek R, van Ittersum MK, ten Berge H, Gort G, Spiegel H and Whitmore AP (2017) Do organic inputs matter – a meta-analysis of additional yield effects for arable crops in Europe. *Plant and Soil* **411**, 293–303.
- IUSS Working Group WRB (2014) World Reference Base for Soil Resources 2014. International soil classification system for naming soils and creating legends for soil maps. World Soil Resources Reports No. 106. FAO, Rome.
- Jenkyn JF, Gutteridge RJ and White RP (2014) Effects of break crops, and of wheat volunteers growing in break crops or in set-aside or conservation covers, all following crops of winter wheat, on the development of take-all (*Gaeumannomyces graminis* var. *tritici*) in succeeding crops of winter wheat. *Annals of Applied Biology* **165**, 340–363.
- Kassambara A (2020) ggpubr: ‘ggplot2’ Based Publication Ready Plots. Available at <https://CRAN.R-project.org/package=ggpubr>
- Kenward MG and Roger JH (1997) Small sample inference for fixed effects from restricted maximum likelihood. *Biometrics* **53**, 983–997.
- Kirkegaard JA and Ryan MH (2014) Magnitude and mechanisms of persistent crop sequence effects on wheat. *Field Crops Research* **164**, 154–165.
- Kirkegaard J, Christen O, Krupinsky J and Layzell D (2008) Break crop benefits in temperate wheat production. *Field Crops Research* **107**, 185–195.
- Kleijn D, Bommarco R, Fijen TPM, Garibaldi LA, Potts SG and van der Putten WH (2019) Ecological intensification: bridging the gap between science and practice. *Trends in Ecology & Evolution* **34**, 154–166.
- Koppensteiner LJ, Kaul H-P, Piepho H-P, Barta N, Euteneuer P, Bernas J, Klimek-Kopyra A, Gronauer A and Neugschwandner RW (2022) Yield and yield components of facultative wheat are affected by sowing time, nitrogen fertilization and environment. *European Journal of Agronomy* **140**, 126591.
- Loughin TM, Roediger MP, Milliken GA and Schmidt JP (2007). On the analysis of long-term experiments. *Journal of the Royal Statistical Society: Series A (Statistics in Society)* **170**, 29–42.
- Macholdt J and Honermeier B (2019) Stability analysis for grain yield of winter wheat in a long-term field experiment. *Archives of Agronomy and Soil Science* **65**, 686–699.
- Macholdt J, Piepho H-P, Honermeier B, Perryman S, Macdonald A and Poulton P (2020) The effects of cropping sequence, fertilization and straw management on the yield stability of winter wheat (1986–2017) in the Broadbalk Wheat Experiment, Rothamsted, UK. *The Journal of Agricultural Science* **158**, 65–79.
- McDaniel MD, Grandy AS, Tiemann LK and Weintraub MN (2014) Crop rotation complexity regulates the decomposition of high and low quality residues. *Soil Biology and Biochemistry* **78**, 243–254.
- Oldfield EE, Bradford MA and Wood SA (2019) Global meta-analysis of the relationship between soil organic matter and crop yields. *SOIL* **5**, 15–32.
- Onofri A, Seddaiu G and Piepho H-P (2016) Long-term experiments with cropping systems: case studies on data analysis. *European Journal of Agronomy* **77**, 223–235.
- Ortiz-Monasterio JI, Dhillon SS and Fischer RA (1994) Date of sowing effects on grain yield and yield components of irrigated spring wheat cultivars and relationships with radiation and temperature in Ludhiana, India. *Field Crops Research* **37**, 169–184.
- Ozturk A, Caglar O and Bulut S (2006) Growth and yield response of facultative wheat to winter sowing, freezing sowing and spring sowing at different seeding rates. *Journal of Agronomy and Crop Science* **192**, 10–16.
- Pan G, Smith P and Pan W (2009) The role of soil organic matter in maintaining the productivity and yield stability of cereals in China. *Agriculture, Ecosystems & Environment* **129**, 344–348.
- Piepho H-P, Büchse A and Emrich K (2003) A Hitchhiker’s guide to mixed models for randomized experiments. *Journal of Agronomy and Crop Science* **189**, 310–322.
- Reeves DW (1997) The role of soil organic matter in maintaining soil quality in continuous cropping systems. *Soil and Tillage Research* **43**, 131–167.

- RStudio** (2022) RStudio Version 2022.02.3. Available at <https://rstudio.com>
- Rubio V, Diaz-Rossello R, Quincke JA and van Es HM** (2021) Quantifying soil organic carbon's critical role in cereal productivity losses under annualized crop rotations. *Agriculture, Ecosystems & Environment* **321**, 107607.
- SAS Institute Inc.** (2016) SAS Version 9.4.
- Schönhammer A and Fischbeck G** (1987a) Investigation at cereal dominated crop rotations and cereal monocultures. I. The differentiation of yield performance and yield components during a 15 year cultivation period. *Bayerisches landwirtschaftliches Jahrbuch* **64**, 175–191.
- Schönhammer A and Fischbeck G** (1987b) Investigation at cereal dominated crop rotations and cereal monocultures. III. Changes in soil properties. *Bayerisches landwirtschaftliches Jahrbuch* **64**, 681–694.
- Sieling K and Christen O** (2015) Crop rotation effects on yield of oilseed rape, wheat and barley and residual effects on the subsequent wheat. *Archives of Agronomy and Soil Science* **61**, 1–19.
- Sieling K and Hanus H** (1990) Yield reaction of winter wheat in monoculture in dependence upon weather and soil. *Journal of Agronomy and Crop Science* **165**, 151–158.
- Sieling K, Stahl C, Winkelmann C and Christen O** (2005) Growth and yield of winter wheat in the first 3 years of a monoculture under varying N fertilization in NW Germany. *European Journal of Agronomy* **22**, 71–84.
- Sieling K, Ubben K and Christen O** (2007) Effects of preceding crop, sowing date, N fertilization and fluquinconazole seed treatment on wheat growth, grain yield and take-all. *Journal of Plant Diseases and Protection* **114**, 213–220.
- Smith RG, Gross KL and Robertson GP** (2008) Effects of crop diversity on agroecosystem function: crop yield response. *Ecosystems* **11**, 355–366.
- Steinmann H-H and Dobers ES** (2013) Spatio-temporal analysis of crop rotations and crop sequence patterns in Northern Germany: potential implications on plant health and crop protection. *Journal of Plant Diseases and Protection* **120**, 85–94.
- The R Foundation for Statistical Computing** (2022) R Version 4.2.1. Available at <https://r-project.org>
- Weiser C, Fuß R, Kage H and Flessa H** (2018) Do farmers in Germany exploit the potential yield and nitrogen benefits from preceding oilseed rape in winter wheat cultivation? *Archives of Agronomy and Soil Science* **64**, 25–37.
- Wickham H** (2016) *ggplot2: Elegant Graphics for Data Analysis*. New York: Springer, 260 S.