

## THE WEIGHTED $g$ -DRAZIN INVERSE FOR OPERATORS

A. DAJIĆ and J. J. KOLIHA 

(Received 10 August 2004; revised 14 February 2005)

Communicated by A. Pryde

### Abstract

The paper introduces and studies the weighted  $g$ -Drazin inverse for bounded linear operators between Banach spaces, extending the concept of the weighted Drazin inverse of Rakočević and Wei (*Linear Algebra Appl.* **350** (2002), 25–39) and of Cline and Greville (*Linear Algebra Appl.* **29** (1980), 53–62). We use the Mbekhta decomposition to study the structure of an operator possessing the weighted  $g$ -Drazin inverse, give an operator matrix representation for the inverse, and study its continuity. An open problem of Rakočević and Wei is solved.

2000 *Mathematics subject classification*: primary 47A60, 46H30, 47A05, 47A10.

*Keywords and phrases*: Banach algebra without unit,  $g$ -Drazin inverse, bounded linear operator, weighted  $g$ -Drazin inverse, the Mbekhta decomposition, ascent and descent.

### 1. Introduction

In recent papers [13, 14], Rakočević and Wei defined and investigated the weighted Drazin inverse for bounded linear operators between Banach and Hilbert spaces, extending the concept of a weighted Drazin inverse for rectangular matrices introduced by Cline and Greville [5]. The weighted Drazin inverse for operators was previously introduced and studied by Qiao in [12], and further investigated by Wang in [16, 17]. The main purpose of this paper is to introduce and study the weighted  $g$ -Drazin inverse for bounded linear operators between Banach spaces  $X$  and  $Y$ , thus further extending the above mentioned works.

Let  $\mathcal{B}(X, Y)$  denote the set of all bounded linear operators between  $X$  and  $Y$ , and let  $W$  be a nonzero operator in  $\mathcal{B}(Y, X)$ . The  $W$ -weighted  $g$ -Drazin inverse (the  $Wg$ -Drazin inverse for short) can be studied in the framework of Banach algebras when we introduce on the space  $\mathcal{B}(X, Y)$  the  $W$ -product  $A \ B = AWB$ , and the

$W$ -norm  $\|A\|_W = \|A\|\|W\|$ . This elegant approach which turns  $\mathcal{B}(X, Y)$  into a Banach algebra was suggested to the authors of [13, 14] by an anonymous referee. Unless  $W$  is invertible (and this would require the spaces  $X$  and  $Y$  to be isomorphic and homeomorphic), the resulting algebra is without unit.

In our work we remove the restriction of finite polarity of the operator  $WA$  (and  $AW$ ) adopted by Rakočević and Wei [13]. In addition, we solve an open problem posed in [13], and complete and extend the results of Buoni and Faires [3] on the ascent and descent of  $AB$  and  $BA$ .

In Section 2 we gather relevant results on the  $g$ -Drazin inverse in Banach algebras without unit in order to study the  $Wg$ -Drazin inverse within the space  $\mathcal{B}(X, Y)$ , without having to adjoin a unit. Section 3 introduces and studies the weighted  $g$ -Drazin inverse between two different Banach spaces. In Section 4 we explore some properties of the weighted  $g$ -Drazin inverse, including the core decomposition and an integral representation for the weighted inverse. The ascent and descent for  $WA$  and  $AW$  is studied in Section 5, and a solution to an open problem posed by Rakočević and Wei in [13] is given there. In Section 6 we compare the Mbekhta decomposition for the operators  $WA$  and  $AW$  and recover and sharpen a result of Yukhno [19] on rectangular matrices. In the remaining sections we give an operator matrix representation for the  $Wg$ -Drazin inverse, compare it with the Moore–Penrose inverse in Hilbert spaces, and give necessary and sufficient conditions for its continuity.

## 2. The $g$ -Drazin inverse in Banach algebras without unit

Let  $\mathcal{A}$  be a Banach algebra. We write  $\mathcal{A}^{\text{qnil}}$  for the set of all quasinilpotent elements in  $\mathcal{A}$ , that is, elements  $a$  satisfying  $\lim_{n \rightarrow \infty} \|a^n\|^{1/n} = 0$ ; the set of all nilpotent elements is denoted by  $\mathcal{A}^{\text{nil}}$ . If  $\mathcal{A}$  is unital, we denote by  $\mathcal{A}^{\text{inv}}$  the group of all invertible elements in  $\mathcal{A}$ . An element  $a \in \mathcal{A}$  is *quasipolar* if 0 is not an accumulation point of the spectrum of  $a$ . In an algebra without unit, this is equivalent to 0 being an isolated spectral point of  $a$ . The set of all quasipolar elements of  $\mathcal{A}$  will be denoted by  $\mathcal{A}^{\text{qpol}}$ . An element  $a \in \mathcal{A}$  is *polar* if it is quasipolar and 0 is at most a pole of the resolvent of  $a$ . The set of all polar elements is denoted by  $\mathcal{A}^{\text{pol}}$ .

The following holds [7, Theorems 4.2 and 5.1]:

LEMMA 2.1. *Let  $\mathcal{A}$  be a unital Banach algebra. Then  $a \in \mathcal{A}$  is quasipolar (polar) in  $\mathcal{A}$  if and only there exists  $p \in \mathcal{A}$  such that*

$$(2.1) \quad p^2 = p, \quad ap = pa \in \mathcal{A}^{\text{qnil}} \quad (ap = pa \in \mathcal{A}^{\text{nil}}), \quad a + p \in \mathcal{A}^{\text{inv}}.$$

*The resolvent  $R(\lambda; a) = (\lambda 1 - a)^{-1}$  has a Laurent expansion in some punctured*

neighbourhood  $0 < |\lambda| < r$  of 0 given by

$$(2.2) \quad R(\lambda; a) = \sum_{n=0}^{\infty} \lambda^{-n-1} a^n p - \sum_{n=1}^{\infty} \lambda^{n-1} b^n,$$

where  $b = (a + p)^{-1}(1 - p)$ .

The element  $p$  is uniquely determined by the conditions of the theorem; it is called the *spectral idempotent* of  $a$ , and it double commutes with  $a$ . The element  $q = 1 - p$  is the *support idempotent* of  $a$ . The support idempotent of a quasipolar element exists in an algebra without a unit, but not the spectral idempotent. The element  $b = (a + p)^{-1}(1 - p)$  defines the  $g$ -Drazin inverse of  $a$  in the case of a unital algebra;  $b$  also double commutes with  $a$ . We write  $a^\pi$  and  $a^\sigma$  for the spectral idempotent and the support idempotent of a quasipolar element  $a$ , respectively.

From now on we assume that  $\mathcal{A}$  is a complex Banach algebra without unit.

The *unitisation* of  $\mathcal{A}$  is the unital Banach algebra  $\mathcal{A}_1 = \mathcal{A} \oplus \mathbb{C}$  containing  $\mathcal{A}$  as a two sided ideal of codimension 1 [2, page 15]. Given  $a \in \mathcal{A}$ , we define the *spectrum*  $\text{Sp}(a)$  of  $a$  in  $\mathcal{A}$  as the spectrum of  $a$  considered as an element of the unital Banach algebra  $\mathcal{A}_1$ , that is, the set of all  $\lambda \in \mathbb{C}$  such that  $\lambda 1 - a \notin \mathcal{A}_1^{\text{inv}}$ . Observe that 0 is always in the spectrum of any element of a Banach algebra without unit.

**PROPOSITION 2.2.** *Let  $\mathcal{A}$  be a Banach algebra without unit. Then  $a \in \mathcal{A}^{\text{qpol}}$  ( $a \in \mathcal{A}^{\text{pol}}$ ) if and only if there exists  $b \in \mathcal{A}$  such that*

$$(2.3) \quad ab = ba, \quad bab = b, \quad a - aba \in \mathcal{A}^{\text{qnil}} \quad (a - aba \in \mathcal{A}^{\text{nil}}).$$

The element  $b$ , if it exists, is unique.

**PROOF.** We embed  $\mathcal{A}$  into its unitisation  $\mathcal{A}_1$ .

If  $a$  is quasipolar in  $\mathcal{A}$ , then it is also quasipolar in  $\mathcal{A}_1$ . Let  $p$  be the spectral idempotent of  $a$  in  $\mathcal{A}_1$ , and  $b = (a + p)^{-1}(1 - p)$  the Drazin inverse of  $a$  in  $\mathcal{A}_1$ . Since  $1 - p$  is in  $\mathcal{A}$ , so is  $b$  ( $\mathcal{A}$  is an ideal). The equations (2.3) are then easily verified.

Conversely, let equations (2.3) hold. Then  $p = 1 - ab$  is the spectral idempotent of  $a$  in  $\mathcal{A}_1$  [7, Theorem 4.2], and  $a$  is quasipolar, both in  $\mathcal{A}_1$  and  $\mathcal{A}$ . From

$$(a + p)b = (a + 1 - ab)b = ab + b - bab = ab = 1 - p$$

and the invertibility of  $a + p$  in  $\mathcal{A}_1$ , we get  $b = (a + p)^{-1}(1 - p)$  in  $\mathcal{A}_1$  (and in  $\mathcal{A}$ ). This proves the uniqueness of  $b$  satisfying (2.3). □

**DEFINITION 2.3.** Let  $\mathcal{A}$  be a Banach algebra without unit and let  $a \in \mathcal{A}^{\text{qpol}}$ . We define the  $g$ -Drazin inverse  $a^D$  of  $a$  to be the unique element  $b$  satisfying (2.3). The *Drazin index* of a quasipolar element  $a$  is defined by

$$i(a) = \inf \{k \in \mathbb{N} : (a - a^2 a^D)^k = 0\}$$

( $\inf \emptyset = \infty$ ). The  $g$ -Drazin inverse of a polar element is called the *Drazin inverse*.

We observe that  $a \in \mathcal{A}$  is polar if and only if it is quasipolar and has a finite Drazin index.

As in the unital case, any  $g$ -Drazin invertible element  $a$  of  $\mathcal{A}$  has the ‘core’ decomposition.

**PROPOSITION 2.4.** *Let  $\mathcal{A}$  be a Banach algebra without unit. Then  $a \in \mathcal{A}^{\text{qpol}}$  if and only if  $a = c + u$ , where  $c$  is simply polar,  $u$  quasinilpotent, and  $cu = 0 = uc$ . Such a decomposition is unique. In addition,*

$$(2.4) \quad a^{\text{D}} = c^{\text{D}}, \quad a^{\sigma} = c^{\sigma}, \quad \text{Sp}(c) = \text{Sp}(a).$$

We can show that  $ua^{\sigma} = 0$  and that the element  $c$ , called the *core* of  $a$ , satisfies

$$c = aa^{\sigma} = (a^{\text{D}})^{\text{D}} = a^2 a^{\text{D}}.$$

**PROPOSITION 2.5.** *Let  $\mathcal{A}$  be a Banach algebra without unit and let  $a \in \mathcal{A}^{\text{qpol}}$ . Then  $a^{\text{D}} = a$  if and only if  $a^3 = a$ .*

**PROOF.** Suppose that  $a^3 = a$  and let  $a = c + u$  be the core decomposition of  $a$ . We observe that  $a^3 = c^3 + u^3$  is the core decomposition for  $a^3 = a$ . From the uniqueness,  $c^3 = c$  and  $u^3 = u$ . Since  $u^3 = u \in \mathcal{A}^{\text{qnil}}$ , we conclude that  $u = 0$ :

$$\lim_{n \rightarrow \infty} \|u\|^{1/3^n} = \lim_{n \rightarrow \infty} \|u^{3^n}\|^{1/3^n} = r(u) = 0.$$

Thus  $a = c = aa^{\sigma}$  is simply polar, and

$$a^{\text{D}} = (a^{\text{D}})^2 a = (a^{\text{D}})^2 a^3 = (a^{\text{D}} a^2)(a^{\text{D}} a) = aa^{\sigma} = a.$$

Conversely, if  $a^{\text{D}} = a$ , then  $a = (a^{\text{D}})^2 a = a^3$ . □

As an example of further properties of the  $g$ -Drazin inverse in Banach algebras without unit we prove the following result, which for matrices reduces to Theorem 7.8.4 of Campbell and Meyer [4].

**PROPOSITION 2.6.** *Let  $\mathcal{A}$  be a Banach algebra without unit, and let  $a, b \in \mathcal{A}$  be such that  $(ba)^2 \in \mathcal{A}^{\text{qpol}}$ . Then both  $ab$  and  $ba$  are  $g$ -Drazin invertible, and*

$$(2.5) \quad (ab)^{\text{D}} = a((ba)^2)^{\text{D}} b.$$

PROOF. If  $(ba)^2 \in \mathcal{A}^{qpol}$ , then also  $(ab)^2$ ,  $ab$  and  $ba$  are quasipolar, and  $w = ((ba)^2)^D = ((ba)^D)^2$  commutes with  $ba$ . Set  $c = a((ba)^2)^D b = awb$ . It is not difficult to show that  $(ab)c = c(ab)$  and  $(ab)c^2 = c$ . The element  $ab - (ab)^2c = (a - a(ba)^2w)b$  is quasinilpotent if and only if  $x = b(a - a(ba)^2w) = ba - (ba)^3w$  is quasinilpotent. Imbedding  $\mathcal{A}$  into its unitisation  $\mathcal{A}_1$ , we recall that  $p = 1 - (ba)^2w$  is idempotent; hence  $x = (ba)p \in \mathcal{A}^{qnil}$  if and only if  $x^2 = (ba)^2p \in \mathcal{A}^{qnil}$  if and only if  $(ba)^2 - (ba)^4w \in \mathcal{A}^{qnil}$ . This completes the proof.  $\square$

### 3. The weighted $g$ -Drazin inverse for operators

Throughout this section we assume that  $X, Y$  are nonzero complex Banach spaces and  $W$  is a fixed nonzero operator in  $\mathcal{B}(Y, X)$ , the set of all bounded linear operators on  $Y$  to  $X$ . First we turn  $\mathcal{B}(X, Y)$  into a Banach algebra  $\mathcal{B}_w(X, Y)$  (in general without a unit) by introducing a multiplication of elements of  $\mathcal{B}(X, Y)$  facilitated by the operator  $W$ , and imposing a suitable norm on  $\mathcal{B}(X, Y)$ .

LEMMA 3.1. *Let  $\mathcal{B}_w(X, Y)$  be the space  $\mathcal{B}(X, Y)$  equipped with the multiplication*

$$(3.1) \quad A \ B = AWB,$$

*and norm  $\|A\|_w = \|A\|\|W\|$ . Then  $\mathcal{B}_w(X, Y)$  becomes a complex Banach algebra;  $\mathcal{B}_w(X, Y)$  has a unit if and only if  $W$  is invertible, in which case  $W^{-1}$  is that unit.*

PROOF. The verification of most Banach algebra axioms is straightforward. The positive definiteness of the norm is ensured by the fact that  $W \neq 0$ . We check the submultiplicativity of the norm. If  $A, B \in \mathcal{B}(X, Y)$ , then

$$(3.2) \quad \|A \ B\|_w = \|AWB\|\|W\| \leq \|A\|\|W\|\|B\|\|W\| = \|A\|_w\|B\|_w.$$

If  $W$  is invertible, then  $W^{-1} \in \mathcal{B}(X, Y)$  is the unit in  $\mathcal{B}_w(X, Y)$ . Conversely, assume that  $P \in \mathcal{B}(X, Y)$  is the unit for  $\mathcal{B}_w(X, Y)$ . Then

$$(3.3) \quad AWP = A = PWA \quad \text{for all } A \in \mathcal{B}(X, Y).$$

For each  $y \in Y$  and  $f \in X^*$  define  $f \otimes y : X \rightarrow Y$  by  $(f \otimes y)x = f(x)y$  for all  $x \in X$ ; then  $f \otimes y \in \mathcal{B}(X, Y)$ . From  $PW(f \otimes y)x = (f \otimes y)x$  we get  $f(x)PWy = f(x)y$ . Selecting  $x$  and  $f$  so that  $f(x) \neq 0$ , we obtain  $PWy = y$  for any  $y \in Y$ . From  $(f \otimes y)WPx = (f \otimes y)x$  we get  $f(WPx)y = f(x)y$ . Selecting  $y \neq 0$ , yields  $f(WPx) = f(x)$  for all  $f \in X^*$ , which implies  $WPx = x$  for any  $x \in X$ . Then  $W$  is invertible; setting  $A = W^{-1}$  in (3.3), we get  $W^{-1} = PWW^{-1} = P$ .  $\square$

We observe that if  $\mathcal{B}_W(X, Y)$  has the unit  $W^{-1}$ , the spaces  $X$  and  $Y$  are isomorphic and homeomorphic; in particular,  $X$  and  $Y$  are of the same dimension. Moreover, the norm of the unit in  $\mathcal{B}_W(X, Y)$  is equal to  $\|W^{-1}\|_W = \|W^{-1}\| \|W\| = \kappa(W)$ , known as the condition number of  $W$ .

For any  $n \in \mathbb{N}$  we write  $A^n = A \cdots A$  ( $n$  factors). Observe that

$$(3.4) \quad A^n = (AW)^{n-1}A = A(WA)^{n-1}.$$

We write  $r_W(\cdot)$  for the spectral radius of elements of  $\mathcal{B}_W(X, Y)$ . We show that

$$(3.5) \quad r_W(A) = r(AW) = r(WA),$$

where  $r(\cdot)$  is the spectral radius in  $\mathcal{B}(Y)$  or  $\mathcal{B}(X)$ . Indeed,

$$\begin{aligned} r(AW) &= \lim_{n \rightarrow \infty} \|(AW)^n\|^{1/n} \leq \lim_{n \rightarrow \infty} (\|(AW)^{n-1}A\| \|W\|)^{1/n} \\ &= \lim_{n \rightarrow \infty} \|A^n\|_W^{1/n} = r_W(A). \end{aligned}$$

Conversely,

$$\begin{aligned} r_W(A) &= \lim_{n \rightarrow \infty} \|A^n\|_W^{1/n} = \lim_{n \rightarrow \infty} \|A^n\|^{1/n} \|W\|^{1/n} \\ &= \lim_{n \rightarrow \infty} \|(AW)^{n-1}A\|^{1/n} \|W\|^{1/n} \\ &\leq \lim_{n \rightarrow \infty} \|(AW)^{n-1}\|^{1/n} \lim_{n \rightarrow \infty} (\|A\| \|W\|)^{1/n} \\ &= \lim_{n \rightarrow \infty} \|(AW)^{n-1}\|^{1/n} = r(AW) \end{aligned}$$

as  $\lim_{n \rightarrow \infty} \|(AW)^{n-1}\|^{1/n} = \lim_{n \rightarrow \infty} \|(AW)^n\|^{1/n}$ . The second equality in (3.5) follows by symmetry.

**DEFINITION 3.2.** Let  $W$  be a fixed nonzero operator in  $\mathcal{B}(Y, X)$ . An operator  $A \in \mathcal{B}(X, Y)$  is called *Wg-Drazin invertible* if  $A$  is quasipolar in the Banach algebra  $\mathcal{B}_W(X, Y)$ . The *Wg-Drazin inverse*  $A^{D,W}$  of  $A$  (or *W-weighted g-Drazin inverse*) is then defined as the *g-Drazin inverse*  $B$  of  $A$  in the Banach algebra  $\mathcal{B}_W(X, Y)$ ;  $i_W(A)$  is the Drazin index of  $A$  in  $\mathcal{B}_W(X, Y)$ . A polar element of  $\mathcal{B}_W(X, Y)$  is called *W-Drazin invertible*, with the *W-Drazin inverse*  $A^{D,W} = B$ .

The *Wg-Drazin inverse* is unique if it exists (Proposition 2.2), and is characterised by the following theorem.

**THEOREM 3.3.** *Let  $W$  be a fixed nonzero operator in  $\mathcal{B}(Y, X)$ . Then  $A \in \mathcal{B}(X, Y)$  is *Wg-Drazin invertible* with the *Wg-Drazin inverse*  $A^{D,W} = B \in \mathcal{B}(X, Y)$  if and only if one of the following equivalent conditions holds:*

- (i) *AW is quasipolar in  $\mathcal{B}(Y)$  with  $(AW)^D = BW$ ;*

- (ii)  $WA$  is quasipolar in  $\mathcal{B}(X)$  with  $(WA)^D = WB$ ;
- (iii) There exists  $B \in \mathcal{B}(X, Y)$  satisfying

$$(AW)B = (BW)A, \quad (BW)^2A = B, \quad (AW)^2BW - AW \in \mathcal{B}(Y)^{qnil};$$

- (iv) There exists  $B \in \mathcal{B}(X, Y)$  satisfying

$$A(WB) = B(WA), \quad A(WB)^2 = B, \quad WB(WA)^2 - WA \in \mathcal{B}(X)^{qnil}.$$

The  $Wg$ -Drazin inverse  $A^{D,W}$  of  $A$  then satisfies

$$(3.6) \quad A^{D,W} = ((AW)^D)^2A = A((WA)^D)^2.$$

PROOF. Suppose that  $A$  has the  $Wg$ -Drazin inverse  $B$ .

The conditions

$$A B = B A, \quad B A B = B, \quad A B A - A \in \mathcal{B}_w(X, Y)^{qnil},$$

translate to

$$(3.7) \quad AWB = BWA, \quad (BW)^2A = B, \quad T = (AW)^2B - A \in \mathcal{B}_w(X, Y)^{qnil}.$$

Let  $C = BW$ . Then  $(AW)C = C(AW)$  and  $C^2(AW) = C$  by (3.7). Finally, by (3.5),  $r(TW) = r_w(T) = 0$ . Hence  $(AW)^2C - AW = TW$  is quasinilpotent in  $\mathcal{B}(Y)$ , and (i) is proved.

Condition (ii) follows from a symmetrical argument. Conditions (i) and (iii) (respectively (ii) and (iv)) are equivalent by the characterisation of the  $g$ -Drazin inverse given in Proposition 2.2.

Conversely, suppose that  $AW \in \mathcal{B}(Y)$  has the  $g$ -Drazin inverse  $C$ . Let  $B = C^2A$ . The equations  $(AW)C = C(AW)$  and  $C^2(AW) = C$  imply

$$A B = AWC^2A = C^2AWA = B A, \quad \text{and} \\ B A B = (C^2AW)(AWC^2)A = C^2A = B.$$

Write  $A B A - A = (AWC^2)AWA - A = CAWA - A = S$ . Since  $SW = C(AW)^2 - AW$  is quasinilpotent in  $\mathcal{B}(Y)$ ,  $r_w(S) = r(SW) = 0$ , and  $S$  is quasinilpotent in  $\mathcal{B}_w(X, Y)$ . This proves that condition (i) implies that  $A$  is  $Wg$ -Drazin invertible with  $A^{D,W} = C^2A$ . The rest follows from Proposition 2.2 by symmetry.  $\square$

From (3.6) we find an expression for the support idempotent  $A^{\sigma,W}$  of  $A$  in  $\mathcal{B}_w(X, Y)$ :  $A^{\sigma,W} = A A^{D,W} = AW((AW)^D)^2A = (AW)^DA$ . By symmetry,

$$(3.8) \quad A^{\sigma,W} = (AW)^DA = A(WA)^D.$$

PROPOSITION 3.4. *If  $A \in \mathcal{B}(X, Y)$  is Wg-Drazin invertible, then the Drazin indices  $i_w(A)$ ,  $i(WA)$ , and  $i(AW)$  are all finite or all infinite, and satisfy the inequalities*

$$(3.9) \quad \max \{i(AW), i(WA)\} \leq i_w(A) \leq \min \{i(AW), i(WA)\} + 1.$$

PROOF. Let  $A^{D,W} = B$  be the Wg-Drazin inverse of  $A$  and let  $T = (AW)^2B - A$ . If  $i_w(A) = k < \infty$ , then  $T^k = 0$ . Consequently  $(TW)^k = (TW)^{k-1}TW = T^k W = 0$  and hence  $i(AW) \leq i_w(A)$ .

Let  $AW$  have the  $g$ -Drazin inverse  $C$  and let  $S = CAWA - A$ . If  $i(AW) = k < \infty$ , then  $(SW)^k = 0$ , and  $S^{(k+1)} = (SW)^k S = 0$ , that is,  $i_w(A) \leq k + 1$ . This proves the inequality for  $i(AW)$  in (3.9).

It is known that for any  $A \in \mathcal{B}(X, Y)$  and  $W \in \mathcal{B}(Y, X)$ ,

$$(3.10) \quad \text{Sp}(AW) \setminus \{0\} = \text{Sp}(WA) \setminus \{0\}.$$

Hence  $AW$  is  $g$ -Drazin invertible in  $\mathcal{B}(Y)$  if and only if  $WA$  is  $g$ -Drazin invertible in  $\mathcal{B}(X)$ . The inequality for  $i(WA)$  in (3.9) is obtained by symmetry. □

EXAMPLE 3.5. The inequality  $i(AW) \leq i_w(A)$  (respectively  $i(WA) \leq i_w(A)$ ) in (3.9) can be strict. Let

$$W = \begin{bmatrix} 0 & 0 \\ 1 & 1 \\ 0 & 0 \end{bmatrix},$$

and let  $\mathcal{B}_W$  be the space  $\mathcal{M}_{2,3}(\mathbb{C})$  of all complex  $2 \times 3$  matrices with the multiplication (3.1). By the preceding theorem, every element  $A \in \mathcal{M}_{2,3}(\mathbb{C})$  has a  $g$ -Drazin inverse of finite Drazin index in  $\mathcal{B}_W$  since the matrix  $AW$  has the conventional Drazin inverse  $(AW)^D$  in  $\mathcal{M}_{2,2}(\mathbb{C})$ . Let

$$A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}.$$

Then  $AW = 0 = B$ , where  $B$  is the  $g$ -Drazin inverse of  $A$  in  $\mathcal{B}_W$ , and  $T = (AW)^2B - A = -A$ . Since  $T \neq 0$  and  $T^2 = AWA = 0$ , we have  $i_w(A) = 2$ . On the other hand,  $i(AW) = i(0) = 1$ . An example of a strict inequality between  $i(WA)$  and  $i_w(A)$  can be obtained from the present example and the following proposition involving the dual spaces  $X^*$  and  $Y^*$  of  $X$  and  $Y$ .

PROPOSITION 3.6.  *$A \in \mathcal{B}(X, Y)$  is Wg-Drazin invertible if and only if the adjoint  $A^* \in \mathcal{B}(Y^*, X^*)$  of  $A$  is  $W^*g$ -Drazin invertible. In this case*

$$(3.11) \quad (A^*)^{D,W^*} = (A^{D,W})^*.$$

PROOF. Since  $\text{Sp}((AW)^*) = \text{Sp}(AW)$ ,  $(AW)^*$  is quasipolar if and only if  $AW$  is quasipolar. Hence  $W^*A^* = (AW)^*$  is  $g$ -Drazin invertible if and only if  $AW$  is. By Theorem 3.3,  $A^*$  is  $W^*g$ -Drazin invertible if and only if  $A$  is  $Wg$ -Drazin invertible. Equation (3.11) follows on application of Proposition 2.2.  $\square$

EXAMPLE 3.7 (Rakočević and Wei [13]). If  $A \in \mathcal{B}(X, Y)$  is a finite rank operator, then  $A$  has a finite index  $Wg$ -Drazin inverse for any nonzero  $W \in \mathcal{B}(Y, X)$ . If  $W \in \mathcal{B}(Y, X)$  is a nonzero operator of finite rank, then any  $A \in \mathcal{B}(X, Y)$  has a finite index  $Wg$ -Drazin inverse.

### 4. Further properties of the $Wg$ -Drazin inverse

First we briefly explore a duality between  $A^{D,W}$  and  $W^{D,A}$  provided  $A \in \mathcal{B}(X, Y)$  and  $W \in \mathcal{B}(Y, X)$ . From Theorem 3.3 we see that the weighted  $g$ -Drazin inverse  $W^{D,A}$  exists if and only if  $A^{D,W}$  exists. Equation (3.6) gives rise to the following relations:

$$\begin{aligned} W^{D,A}A &= (WA)^D = WA^{D,W}, \\ AW^{D,A} &= (AW)^D = A^{D,W}W. \end{aligned}$$

We can then express  $W^{D,A}$  in terms of  $A^{D,W}$  and vice versa:

$$\begin{aligned} W^{D,A} &= WA^{D,W}WA^{D,W}W, \\ (4.1) \quad A^{D,W} &= AW^{D,A}AW^{D,A}A. \end{aligned}$$

We observe that in (4.1), the operators  $AW^{D,A}$  and  $W^{D,A}A$  are simply polar (that is, of index 1 or 0): for example,  $AW^{D,A} = AW((AW)^D)^2 = (AW)^D$ . The simple polarity of the  $g$ -Drazin inverse of  $AW$  is well known (see [7]). Specialised to matrices, this proves the necessary part of Theorem 3 in [5].

PROPOSITION 4.1. *Let  $A \in \mathcal{B}(X, Y)$  be  $Wg$ -Drazin invertible. Then the following are true:*

- (i)  $A = A^{D,W}$  if and only if  $A = A^3 = AWAWA$ .
- (ii)  $(A^{D,W})^{D,W} = (AW)^\sigma A = A(WA)^\sigma$ .
- (iii)  $(A^{D,W})^{\sigma,W} = A^{\sigma,W}$ .
- (iv) For any  $n \in \mathbb{N}$ ,  $(A^{D,W})^n = ((AW)^D)^{n+1}A = A((WA)^D)^{n+1} = (A^n)^{D,W}$ .

PROOF. (i) This follows from Proposition 2.5 applied to  $\mathcal{B}_w(X, Y)$ .

(ii) Applying the results of [7] while working in the Banach algebra  $\mathcal{B}_w(X, Y)$ , we have  $(A^{D,W})^{D,W} = A^{-1}A^{\sigma,W} = AW(AW)^DA = (AW)^\sigma A$ .

(iii) In the proof of [7, Theorem 5.2] it is shown that a quasipolar element and its  $g$ -Drazin inverse have the same support idempotent.

(iv) This is shown via induction on  $n$ .  $\square$

Part (ii) of the preceding theorem implies that  $(A^{D,W})^{D,W} = A$  if and only if  $(AW)^\sigma A = A$  ( $A(WA)^\sigma = A$ ). This is equivalent to  $A$  being simply polar in  $\mathcal{B}_W(X, Y)$ .

From [7, Theorem 5.5] we can deduce the following result.

**PROPOSITION 4.2.** *Let  $A, B \in \mathcal{B}(X, Y)$  be Wg-Drazin invertible. If  $AWB = BWA$ , then  $AWB$  is Wg-Drazin invertible with  $(AWB)^{D,W} = A^{D,W}WB^{D,W}$ .*

We now turn our attention to an analogue of the core decomposition for the weighted  $g$ -Drazin inverse.

**THEOREM 4.3.** *An operator  $A \in \mathcal{B}(X, Y)$  is Wg-Drazin invertible if and only if there exist operators  $C, U \in \mathcal{B}(X, Y)$  such that*

$$(4.2) \quad A = C + U, \quad CWU = 0, \quad UWC = 0,$$

$$(4.3) \quad (CW)^\sigma C = C, \quad UW \in \mathcal{B}(Y)^{qnil}.$$

*Such operators are uniquely determined, and  $C = (A^{D,W})^{D,W} = (AW)^\sigma A$ . Further,*

$$(4.4) \quad (AW)^D = (CW)^D, \quad (AW)^\sigma = (CW)^\sigma, \quad \text{Sp}(AW) \cup \{0\} = \text{Sp}(CW).$$

**PROOF.** We apply Theorem 2.4 to  $\mathcal{B}_W(X, Y)$ .  $A$  is Wg-Drazin invertible if and only if there exist  $C, U \in \mathcal{B}(X, Y)$  such that  $A = C + U$ ,  $C \ U = CWU = 0$ ,  $U \ C = UWC = 0$ ,  $C$  is simply polar in  $\mathcal{B}_W(X, Y)$ , and  $U$  is quasinilpotent in  $\mathcal{B}_W(X, Y)$ . The element  $C \in \mathcal{B}_W(X, Y)$  is simply polar if and only if  $C \ C^{\sigma,W} = C$ . From the equation

$$C \ C^{\sigma,W} = CW(CW)^D C = (CW)^\sigma C$$

we conclude that the simple polarity of  $C \in \mathcal{B}_W(X, Y)$  is equivalent to  $(CW)^\sigma C = C$ . Finally,  $r_W(U) = r(UW)$ , and  $UW$  is quasinilpotent in  $\mathcal{B}(X, Y)$  if and only if  $U$  is quasinilpotent in  $\mathcal{B}_W(X, Y)$ . This proves the equivalence of (4.2) and (4.3) to the Wg-Drazin invertibility of  $A$ . Explicitly,  $C = A \ A^{\sigma,W} = (AW)^\sigma A$ .

Towards (4.4) in view of Theorem 2.4,

$$(AW)^D = ((AW)^D)^2 AW = A^{D,W}W = C^{D,W}W = ((CW)^D)^2 CW = (CW)^D.$$

Therefore

$$(CW)^\sigma = (CW)^D C = (AW)^D C = (AW)^D (AW)^\sigma A = (AW)^D A = (AW)^\sigma.$$

If  $\text{Sp}_W(A)$  denotes the spectrum of  $A$  as an element of the Banach algebra  $\mathcal{B}_W(X, Y)$  without unit, then it can be shown that  $\text{Sp}_W(A) = \text{Sp}(AW) \cup \{0\}$ . Hence

$$\text{Sp}(AW) \cup \{0\} = \text{Sp}_W(A) = \text{Sp}_W(C) = \text{Sp}(CW) \quad (\text{as } 0 \in \text{Sp}(CW)).$$

This completes the proof. □

The statement of the theorem remains true when (4.3) is replaced by  $C(WC)^\sigma = C$ ,  $WU \in \mathcal{B}(X)^{\text{qnil}}$ , and  $AW, CW$  in (4.4) are replaced by  $WA, WC$ , respectively.

We close the section with an integral representation of the  $Wg$ -Drazin inverse. The representation of the  $g$ -Drazin inverse given by Castro *et al.* [6, Theorem 2.2] is valid also for Banach algebras without unit. Applying this result to  $\mathcal{B}_W(X, Y)$ , we get the integral representation

$$A^{D,W} = - \int_0^\infty \exp(tA) A^{\sigma,W} dt$$

provided  $A$  is  $Wg$ -Drazin invertible and the nonzero spectrum  $\text{Sp}_W(A) \setminus \{0\}$  lies in the open left half-plane. We express  $\exp(tA) A^{\sigma,W}$  in terms of the usual multiplication of operators:

$$A^n A^{\sigma,W} = (AW)^{n-1}AWA^{\sigma,W} = (AW)^n A^{\sigma,W}.$$

Hence

$$\exp(tA) A^{\sigma,W} = \sum_{n=0}^\infty \frac{t^n}{n!} (AW)^n A^{\sigma,W} = \exp(tAW)A^{\sigma,W}.$$

Note that in general  $\exp(tAW)$  belongs to the unitisation of  $\mathcal{B}_W(X, Y)$  but not to  $\mathcal{B}_W(X, Y)$ , while  $\exp(tAW)A^{\sigma,W}$  is in  $\mathcal{B}_W(X, Y)$ . We summarise our findings.

PROPOSITION 4.4. *Let  $A \in \mathcal{B}(X, Y)$  be  $Wg$ -Drazin invertible such that  $\text{Sp}(WA) \setminus \{0\}$  lies in the left open half-plane. Then*

$$(4.5) \quad A^{D,W} = - \int_0^\infty \exp(tAW)A^{\sigma,W} dt.$$

*If the Drazin index  $i(AW)$  is finite and the set  $\text{Sp}((AW)^{m+1}) \setminus \{0\}$  lies in the left open half-plane for some  $m \geq \min \{i(AW), i(WA)\} + 1$ , then*

$$(4.6) \quad A^{D,W} = - \int_0^\infty \exp(t(AW)^{m+1})(AW)^{m-1}A dt.$$

PROOF. Equation (4.5) follows from our calculations preceding the theorem. For (4.6) we find

$$(A^{(m+1)})^n A^m = A^{(m+1)n+m} = (AW)^{(m+1)n}(AW)^{m-1}A,$$

and

$$\exp(tA^{(m+1)}) A^m = \exp(t(AW)^{m+1})(AW)^{m-1}A.$$

Equation (4.6) then follows from [6, Theorem 2.4]. □

As expected from symmetry, there is also a  $WA$  version of the preceding theorem. If we specialise Equation (4.6) to matrices, we recover [18, Theorem 1]. The inequality  $m \geq \min \{i(AW), i(WA)\} + 1$  in the preceding theorem can be relaxed to  $m \geq i_w(A)$ .

Using the core decomposition of a  $Wg$ -Drazin invertible operator  $A \in \mathcal{B}(X, Y)$ , we obtain yet another integral representation for  $A^{D,W}$ .

**COROLLARY 4.5.** *Let  $A \in \mathcal{B}(X, Y)$  be  $Wg$ -Drazin invertible such that  $\text{Sp}((WA)^2) \setminus \{0\}$  lies in the open left half-plane, and let  $A = C + U$  be the core decomposition of  $A$ . Then*

$$A^{D,W} = C^{D,W} = - \int_0^\infty \exp(t(CW)^2)C dt.$$

**PROOF.** This follows from (4.6) when we note that  $i_w(C) = 1$ . □

### 5. Ascent and descent

We recall that the ascent and descent of an operator  $T \in \mathcal{B}(X)$  are defined by

$$\begin{aligned} \text{asc}(T) &= \inf \{k \in \mathbb{N} \cup \{0\} : N(T^{k+1}) = N(T^k)\}, \\ \text{des}(T) &= \inf \{k \in \mathbb{N} \cup \{0\} : R(T^{k+1}) = R(T^k)\} \end{aligned}$$

( $\inf \emptyset = \infty$ ). Rakočević and Wei [13] ask whether the finiteness of  $\text{asc}(AW)$  and  $\text{des}(WA)$  is sufficient for  $A$  to have the  $W$ -weighted Drazin inverse. An equivalent question is whether  $\text{asc}(AW)$  and  $\text{asc}(WA)$  are always both finite or both infinite.

In this connection it is interesting to recall that Buoni and Faires [3] studied the ascent and descent for the operators  $\lambda I - BA$  and  $\lambda I - AB$ , where  $A, B \in \mathcal{B}(X)$ , and proved, *inter alia*, that for any  $\lambda \neq 0$ ,

$$(5.1) \quad \text{asc}(AB - \lambda I) = \text{asc}(BA - \lambda I), \quad \text{des}(AB - \lambda I) = \text{des}(BA - \lambda I);$$

however, the case  $\lambda = 0$  was left open. Later, Barnes [1] proved by different methods that the ascents of  $I - RS$  and  $I - SR$  are equal for  $R \in \mathcal{B}(X, Y)$  and  $S \in \mathcal{B}(Y, X)$ . It can be shown that the arguments in [3] concerning descent are valid also when  $A \in \mathcal{B}(X, Y)$  and  $B \in \mathcal{B}(Y, X)$ . Thus (5.1) is valid for operators between different spaces. The following theorem, dealing with the ascent and descent in general, completes the results of Buoni and Faires in the case  $\lambda = 0$ .

**THEOREM 5.1.** *Let  $A \in \mathcal{B}(X, Y)$  and  $B \in \mathcal{B}(Y, X)$ . Then the ascents (descents) of  $AB$  and  $BA$  are both finite or both infinite, and satisfy the inequalities*

$$(5.2) \quad \begin{aligned} \text{asc}(AB) - 1 &\leq \text{asc}(BA) \leq \text{asc}(AB) + 1, \\ \text{des}(AB) - 1 &\leq \text{des}(BA) \leq \text{des}(AB) + 1. \end{aligned}$$

PROOF. Suppose that  $\text{asc}(AB) = p < \infty$ . If there existed

$$x \in N((BA)^{p+2}) \setminus N((BA)^{p+1}),$$

we would have  $(AB)^{p+2}Ax = A(BA)^{p+2}x = 0$ , and  $B(AB)^pAx = (BA)^{p+1}x \neq 0$ , that is,  $(AB)^pAx \neq 0$ . Then  $Ax$  would belong to  $N((AB)^{p+2}) \setminus N((AB)^p)$ , which is empty by assumption. This contradiction proves  $N((BA)^{p+1}) = N((BA)^{p+2})$ , which shows that  $\text{asc}(BA) \leq p + 1 = \text{asc}(AB) + 1$ . A symmetrical argument gives  $\text{asc}(AB) \leq \text{asc}(BA) + 1$ . This proves the first inequality in (5.2).

Let  $\text{des}(AB) = p < \infty$ . Suppose

$$(5.3) \quad x \in R((BA)^{p+1}) \setminus R((BA)^{p+2}).$$

Then there exists  $x' \in X$  such that

$$x = (BA)^{p+1}x' = B(AB)^pAx' = By,$$

where  $y = (AB)^pAx' \in R((AB)^p) = R((AB)^{p+2})$ . Hence  $y = (AB)^{p+2}y'$  for some  $y' \in Y$ , and  $(BA)^{p+2}By' = B(AB)^{p+2}y' = By = x$  contrary to (5.3). This proves that  $R((BA)^{p+1}) = R((BA)^{p+2})$ , so that  $\text{des}(BA) \leq p + 1$ .  $\square$

The inequalities in (5.2) can be strict; this follows from Example 3.5 since for matrices  $i(AB) = \text{asc}(AB) = \text{des}(AB)$ .

The following theorem gives a solution to the open problem of Rakočević and Wei [13, page 28].

**THEOREM 5.2.** *Let  $A \in \mathcal{B}(X, Y)$  and  $W \in \mathcal{B}(Y, X) \setminus \{0\}$ . Then  $A$  is  $W$ -Drazin invertible if and only if one of the following equivalent conditions hold:*

- (i)  $AW$  is polar in  $\mathcal{B}(Y)$ ;
- (ii)  $WA$  is polar in  $\mathcal{B}(X)$ ;
- (iii)  $\text{asc}(AW)$  and  $\text{des}(WA)$  are both finite;
- (iv)  $\text{asc}(WA)$  and  $\text{des}(AW)$  are both finite.

PROOF. Suppose that  $A$  is  $W$ -Drazin invertible. By Theorem 3.3,  $AW$  is quasipolar, and by (3.9) we have  $i(AW) \leq i_w(A)$ , which proves that  $AW$  is polar. Conversely, if  $AW$  is polar, then  $i_w(A) \leq i(AW) + 1$ , and  $A$  is  $W$ -Drazin invertible.

(i) implies (ii): Since  $AW$  is quasipolar, so is  $WA$  by (3.10). By (3.9) again,  $i(WA) \leq i(AW) + 1$ , and  $WA$  is polar.

(ii) implies (iii): It is well known that if  $WA$  is polar, then  $\text{asc}(WA)$  and  $\text{des}(WA)$  are both finite. However,  $\text{asc}(AW) \leq \text{asc}(WA) + 1$  by Theorem 5.1 and (iii) follows.

(iii) implies (iv): This follows from Theorem 5.1 as  $\text{asc}(WA) \leq \text{asc}(AW) + 1$  and  $\text{des}(AW) \leq \text{des}(WA) + 1$ .

(iv) implies (i): Since  $\text{asc}(AW) \leq \text{asc}(WA) + 1$ , both  $\text{asc}(AW)$  and  $\text{des}(AW)$  are finite; this implies that  $AW$  is polar.  $\square$

### 6. The Mbekhta decomposition for WA and AW

As before,  $X, Y$  are Banach spaces and  $W$  a nonzero operator in  $\mathcal{B}(Y, X)$ . In order to obtain an operator matrix representation for the weighted  $g$ -Drazin inverse of an operator  $A \in \mathcal{B}(X, Y)$ , we first recall the Mbekhta decomposition for a quasipolar operator. For any operator  $T \in \mathcal{B}(X)$  we define spaces  $H_0(T)$  and  $K(T)$  as follows:

$$H_0(T) = \left\{ x \in X : \lim_{n \rightarrow \infty} \|T^n x\|^{1/n} = 0 \right\},$$

$$K(T) = \left\{ x \in X : \exists x_n \in X, x_n = T x_{n+1}, x_0 = x, \sup_{n \in \mathbb{N}} \|x_n\|^{1/n} < \infty \right\}.$$

Both spaces are hyperinvariant under  $T$ ,  $H_0(T) \supset N(T^n)$ , and  $K(T) \subset R(T^n)$  for all  $n \in \mathbb{N}$ . Further,  $T K(T) = K(T)$  and  $T^{-1} H_0(T) = H_0(T)$ .

PROPOSITION 6.1 (See [8, 11]). *The following conditions on  $T \in \mathcal{B}(X)$  are equivalent:*

- (i)  $T$  is quasipolar;
- (ii)  $X$  is the topological direct sum  $X = K(T) \oplus H_0(T)$ ;
- (iii)  $T = T_1 \oplus T_2$ , where  $T_1$  is invertible and  $T_2$  quasinilpotent.

Condition (ii) can be weakened to  $X = K(T) \oplus H_0(T)$  being only an algebraic sum with at least one of the spaces closed (see [10] and [15]).

THEOREM 6.2. *Let  $A \in \mathcal{B}(X, Y)$  and  $W \in \mathcal{B}(Y, X)$ . If  $WA$  is quasipolar, then so is  $AW$ ,*

$$(6.1) \quad \begin{aligned} A(K(WA)) &= K(AW), & A^{-1}(H_0(AW)) &= H_0(WA), \\ W(K(AW)) &= K(WA), & W^{-1}(H_0(WA)) &= H_0(AW), \end{aligned}$$

*and the spaces  $K(WA), K(AW)$  are isomorphic and homeomorphic.*

PROOF. The result on quasipolarity follows from (3.10). We introduce the following notation

$$(6.2) \quad X_1 = K(WA), \quad X_2 = H_0(WA), \quad Y_1 = K(AW), \quad Y_2 = H_0(AW).$$

Then  $X$  and  $Y$  are decomposed into the topological direct sums  $X = X_1 \oplus X_2$  and  $Y = Y_1 \oplus Y_2$ . The operator matrices

$$(6.3) \quad T = \begin{bmatrix} WA & 0 \\ 0 & AW \end{bmatrix}, \quad S = \begin{bmatrix} 0 & 0 \\ A & 0 \end{bmatrix}$$

represent commuting operators in  $\mathcal{B}(X \oplus Y)$  with  $T$  quasipolar. The support projection  $T^\sigma$  of  $T$  double commutes with  $T$ , that is, the matrix

$$T^\sigma = \begin{bmatrix} (WA)^\sigma & 0 \\ 0 & (AW)^\sigma \end{bmatrix}$$

commutes with the matrix  $S$ . This gives  $A(WA)^\sigma = (AW)^\sigma A$ . Since  $(WA)^\sigma$  is the projection of  $X$  onto  $X_1$  along  $X_2$ , and  $(AW)^\sigma$  is the projection of  $Y$  onto  $Y_1$  along  $Y_2$ , we have  $A(X_i) \subset Y_i$  ( $i = 1, 2$ ). The inclusions  $W(Y_i) \subset X_i$  ( $i = 1, 2$ ) are obtained by symmetry.

Note that  $A(X_2) \subset Y_2$  is equivalent to  $X_2 \subset A^{-1}(Y_2)$ . In order to prove  $A^{-1}(Y_2) \subset X_2$ , assume that  $Ax \in Y_2$ . Then  $x = k + h$  with  $k \in X_1$  and  $h \in X_2$ , and  $Ax = Ak + Ah \in Y_2$  implies that  $Ak = 0$ . From  $k \in N(A) \subset N(WA) \subset X_2$ , we obtain  $k \in X_1 \cap X_2 = \{0\}$ . Hence  $x = h \in X_2$ .

Let  $A_0 : X_1 \rightarrow Y_1$  be the restriction of  $A$ . If  $x \in X_1$  and  $Ax = 0$ , then  $x = 0$  by the argument of the preceding paragraph. Hence  $A_0$  is injective. Suppose that  $y \in Y_1$ . Since  $AWY_1 = Y_1$ , there exists  $u \in Y_1$  such that  $y = AWu$ . But  $WY_1 \subset X_1$ , and so  $Wu \in X_1$ . This proves that  $A_0$  is surjective. Therefore  $A_0$  is a bounded linear bijection from  $X_1$  to  $Y_1$ , and (6.1) is proved. □

In particular, if  $AW$  is quasipolar, then the spaces  $K(AW)$  and  $K(WA)$  have the same dimension being isomorphic.

If  $A$  and  $W$  are rectangular matrices of orders  $m \times n$  and  $n \times m$  respectively, we recover the result of Yukhno [19, Theorem]. For this the operator  $T : \mathbb{C}^m \rightarrow \mathbb{C}^m$  with the matrix  $WA$  is polar, and  $T = T_1 \oplus T_2$ , where  $T_1$  is invertible and  $T_2$  nilpotent;  $T_1$  operates on  $X_1 = K(T) = R(T^p)$ , where  $p$  is the index of  $T$ . The eigenvalues of  $T_1$  are the nonzero eigenvalues of  $WA$ . Let  $\lambda$  be a nonzero eigenvalue of  $WA$ , and  $x_1, \dots, x_k$  a chain of generalised eigenvectors of  $WA$  corresponding to  $\lambda$ , that is,

$$WAx_1 = \lambda x_1 + x_2, \quad \dots, \quad WAx_{k-1} = \lambda x_{k-1} + x_k, \quad WAx_k = \lambda x_k.$$

In view of the decomposition of  $T$  as  $T = T_1 \oplus T_2$ , where  $T_1$  operates on  $X_1$ , we can take  $x_i \in X_1$  for all  $i$ . If  $y_i = Ax_i$  ( $i = 1, \dots, k$ ), then  $y_1, \dots, y_k$  is a chain of generalised eigenvectors of  $AW$  corresponding to  $\lambda$  (this follows from the bijectivity of the operator  $x \mapsto Ax$  restricted from  $X_1$  to  $Y_1$ ). All chains corresponding to nonzero eigenvalues of  $WA$  are matched in this way. This leads to the following structure theorem for  $WA$  and  $AW$ .

**PROPOSITION 6.3.** *Let  $A$  and  $W$  be rectangular matrices of orders  $m \times n$  and  $n \times m$ , respectively. The matrices  $WA$  and  $AW$  (of orders  $n \times n$  and  $m \times m$ , respectively) have Jordan forms*

$$\begin{bmatrix} U & 0 \\ 0 & N_1 \end{bmatrix} \quad \text{and} \quad \begin{bmatrix} U & 0 \\ 0 & N_2 \end{bmatrix},$$

where  $U$  is a matrix in Jordan form corresponding to the nonzero eigenvalues of  $WA$  (and  $AW$ ), while  $N_1$  and  $N_2$  are nilpotent matrices in Jordan form, of different orders in general.

Recall that the entries of  $N_1$  and  $N_2$  are zero except for the superdiagonals, which consist of 0s and 1s.

### 7. An operator matrix representation of the $Wg$ -Drazin inverse

From the Mbekhta decomposition theorem (Proposition 6.1), it follows that an operator  $T \in \mathcal{B}(X)$  is quasipolar if and only if it can be expressed as the direct sum  $T = T_1 \oplus T_2$ , where  $T_1$  is invertible and  $T_2$  quasinilpotent; the  $g$ -Drazin inverse of  $T$  is given by

$$T^D = T_1^{-1} \oplus 0.$$

Our aim is to derive an analogous formula for the  $Wg$ -Drazin inverse using the results of the preceding section.

**THEOREM 7.1.** *Let  $A \in \mathcal{B}(X, Y)$  and  $W \in \mathcal{B}(Y, X) \setminus \{0\}$ . Then  $A$  is  $Wg$ -Drazin invertible if and only if there exist topological direct sums  $X = X_1 \oplus X_2, Y = Y_1 \oplus Y_2$  such that  $A = A_1 \oplus A_2$  and  $W = W_1 \oplus W_2$ , where  $A_i \in \mathcal{B}(X_i, Y_i), W_i \in \mathcal{B}(Y_i, X_i)$ , with  $A_1, W_1$  invertible, and  $W_2A_2$  and  $A_2W_2$  quasinilpotent in  $\mathcal{B}(X_2)$  and  $\mathcal{B}(Y_2)$ , respectively. The  $Wg$ -Drazin inverse of  $A$  is given by  $A^{D,W} = (W_1A_1W_1)^{-1} \oplus 0$  with  $(W_1A_1W_1)^{-1} \in \mathcal{B}(X_1, Y_1)$  and  $0 \in \mathcal{B}(X_2, Y_2)$ .*

**PROOF.** If  $WA$  is quasipolar, the decomposition exists with  $X_i$  and  $Y_i$  given by (6.2). By Theorem 6.2,  $A$  maps  $X_1$  into  $Y_1$ , and  $X_2$  into  $Y_2$ , that is,  $A = A_1 \oplus A_2$ , with  $A_i \in \mathcal{B}(X_i, Y_i), i = 1, 2$ . Similarly, since  $W$  maps  $Y_1$  into  $X_1$  and  $Y_2$  into  $X_2$ ,  $W = W_1 \oplus W_2$ , where  $W_i \in \mathcal{B}(Y_i, X_i), i = 1, 2$ . Hence

$$WA = W_1A_1 \oplus W_2A_2, \quad AW = A_1W_1 \oplus A_2W_2$$

relative to  $X = X_1 \oplus X_2$  and  $Y = Y_1 \oplus Y_2$ . Since  $WA$  and  $AW$  are quasipolar,  $W_1A_1$  and  $A_1W_1$  are invertible, and  $W_2A_2$  and  $A_2W_2$  are quasinilpotent. Hence  $A_1$  and  $W_1$  are invertible.

The  $Wg$ -Drazin inverse of  $A$  is equal to

$$A((WA)^D)^2 = (A_1 \oplus A_2)((W_1A_1)^{-2} \oplus 0) = (W_1A_1W_1)^{-1} \oplus 0.$$

Conversely, if the decompositions with the specified properties exist, then  $AW = (A_1W_1) \oplus (A_2W_2)$  is quasipolar as  $A_1W_1$  is invertible and  $A_2W_2$  quasinilpotent. Then  $A$  is  $Wg$ -Drazin invertible. □

From the necessary part of the preceding theorem we recover [18, Theorem 2] when we specialise the operators to finite matrices. From Theorem 6.2 applied to finite matrices we deduce that the ranks of  $(AW)^m$  and  $(WA)^m$  are equal for any  $m \geq \max \{\text{ind}(AW), \text{ind}(WA)\}$ . (This is used, but not proved, in the derivation of [18, Theorem 2]).

From the commutativity of the operator matrices given in (6.3) and the double commutativity of the  $g$ -Drazin inverse we deduce that  $(AW)^D A = A(WA)^D$ , which leads to the new equality for  $A^{D,W}$  derived from (3.6),

$$A^{D,W} = (AW)^D A(WA)^D.$$

### 8. Relation to the Moore–Penrose inverse

We briefly address the relation of the  $Wg$ -Drazin inverse to the Moore–Penrose inverse in Hilbert spaces (see [13, page 28]). Let  $H, K$  be Hilbert spaces and let  $A \in \mathcal{B}(H, K)$ . It is well known that  $R(A)$  is closed if and only if  $R(A^*)$  is closed,  $R(A^*)$  is closed if and only if  $A^*A$  is simply polar, and  $A^*A$  is simply polar if and only if  $AA^*$  is simply polar. This means that  $A \in \mathcal{B}(H, K)$  is  $A^*g$ -Drazin invertible if and only if the range of  $A$  is closed. We note that

$$(8.1) \quad (A^{D,A^*})^* = (A^*)^{D,A}.$$

We can then prove that the operator  $A^\dagger = (A^*)^{\sigma,A} = A^* A^{D,A^*} A^*$  is the Moore–Penrose inverse characterised by the equations

$$(8.2) \quad A^\dagger AA^\dagger = A^\dagger, \quad AA^\dagger A = A, \quad (A^\dagger A)^* = A^\dagger A, \quad (AA^\dagger)^* = AA^\dagger.$$

We offer a sample of such proof

$$A^\dagger AA^\dagger = (A^*)^{\sigma,A} A (A^*)^{\sigma,A} = (A^*)^{\sigma,A} \circ (A^*)^{\sigma,A} = (A^*)^{\sigma,A} = A^\dagger,$$

where  $T \circ S = T A S$ , and

$$AA^\dagger A = AA^* A^{D,A^*} A^* A = A \quad A^{D,A^*} \quad A = A,$$

where  $T \quad S = T A^* S$ . Other equations in (8.2) can be proved similarly.

### 9. Continuity of the $Wg$ -Drazin inverse

**THEOREM 9.1.** *Let  $A_n \rightarrow A_0$  in  $\mathcal{B}(X, Y)$  and  $W_n \rightarrow W_0 \neq 0$  in  $\mathcal{B}(Y, X)$ , where each  $A_n$  is  $W_n g$ -Drazin invertible,  $n = 0, 1, 2, \dots$ . Then the following conditions are equivalent:*

- (i)  $A_n^{D, W_n} \rightarrow A_0^{D, W_0}$ ;
- (ii)  $\sup_n \|A_n^{D, W_n}\| < \infty$ ;
- (iii)  $(A_n W_n)^D \rightarrow (A_0 W_0)^D$ ;
- (iv)  $A_n^{\sigma, W_n} \rightarrow A_0^{\sigma, W_0}$ .

PROOF. We rely on continuity results for the  $g$ -Drazin inverse obtained in [9]. Condition (i) clearly implies (ii). Suppose that (ii) holds. Since

$$(A_n W_n)^D = ((A_n W_n)^D)^2 (A_n W_n) = A_n^{D, W_n} W_n,$$

we have  $\sup_n \|(A_n W_n)^D\| < \infty$ . By [9, Theorem 2.4],  $(A_n W_n)^D \rightarrow (A_0 W_0)^D$ .

If (iii) holds, then  $A_n^{\sigma, W_n} = (A_n W_n)^D A_n \rightarrow (A_0 W_0)^D A_0 = A_0^{\sigma, W_0}$ .

Let (iv) hold. From the equation

$$(A_n W_n)^\sigma = (A_n W_n)^D A_n W_n = A_n^{\sigma, W_n} W_n,$$

we deduce that  $(A_n W_n)^\sigma \rightarrow (A_0 W_0)^\sigma$ . Using [9, Theorem 2.4] again, we obtain  $(A_n W_n)^D \rightarrow (A_0 W_0)^D$ . Hence  $A_n^{D, W_n} = ((A_n W_n)^D)^2 A_n \rightarrow ((A_0 W_0)^D)^2 A_0 = A_0^{D, W_0}$  and the theorem is proved.  $\square$

From the preceding theorem we recover [13, Theorem 5.1] when we specialise the result to a finite index weighted Drazin inverse.

## References

- [1] B. A. Barnes, 'Common operator properties of the linear operators  $RS$  and  $SR$ ', *Proc. Amer. Math. Soc.* **126** (1998), 1055–1061.
- [2] F. F. Bonsall and J. Duncan, *Complete normed algebras* (Springer, Berlin, 1973).
- [3] J. J. Buoni and J. D. Faires, 'Ascent, descent, nullity and defect of products of operators', *Indiana Univ. Math. J.* **25** (1976), 703–707.
- [4] S. L. Campbell and C. D. Meyer, *Generalized inverses of linear transformations* (Pitman, London, 1979).
- [5] R. E. Cline and T. N. E. Greville, 'A Drazin inverse for rectangular matrices', *Linear Algebra Appl.* **29** (1980), 53–62.
- [6] N. Castro González, J. J. Koliha and Y. Wei, 'On integral representation of the Drazin inverse in Banach algebras', *Proc. Edinburgh Math. Soc.* **45** (2002), 327–331.
- [7] J. J. Koliha, 'A generalized Drazin inverse', *Glasgow Math. J.* **38** (1996), 367–381.
- [8] ———, 'Isolated spectral points', *Proc. Amer. Math. Soc.* **124** (1996), 3417–3424.
- [9] J. J. Koliha and V. Rakočević, 'Continuity of the Drazin inverse II', *Studia Math.* **131** (1998), 167–177.
- [10] J. J. Koliha and P. W. Poon, 'Spectral sets II', *Rend. Circ. Mat. Palermo (2)* **47** (1998), 293–310.
- [11] M. Mbekhta, 'Généralisation de la décomposition de Kato aux opérateurs paranormaux et spectraux', *Glasgow Math. J.* **29** (1987), 159–175.

- [12] S. Z. Qiao, 'The weighted Drazin inverse of a linear operator on a Banach space and its approximation', *Numer. Math. J. Chinese Univ.* **3** (1981), 296–305 (Chinese).
- [13] V. Rakočević and Y. Wei, 'A weighted Drazin inverse and applications', *Linear Algebra Appl.* **350** (2002), 25–39.
- [14] ———, 'The representation and approximation of the  $W$ -weighted Drazin inverse of linear operators in Hilbert spaces', *Appl. Math. Comput.* **141** (2003), 455–470.
- [15] Ch. Schmoeger, 'On isolated points of the spectrum of a bounded linear operator', *Proc. Amer. Math. Soc.* **117** (1993), 715–719.
- [16] G. R. Wang, 'Approximation methods for the  $W$ -weighted Drazin inverse of linear operators in Banach spaces', *Numer. Math. J. Chinese Univ.* **10** (1988), 74–81 (Chinese).
- [17] ———, 'Iterative methods for computing the Drazin inverse and the  $W$ -weighted Drazin inverse of linear operators based on functional interpolation', *Numer. Math. J. Chinese Univ.* **11** (1989), 269–280.
- [18] Y. Wei, 'Integral representation of the  $W$ -weighted Drazin inverse', *Appl. Math. Comput.* **144** (2003), 3–10.
- [19] L. F. Yukhno, 'An eigenvalue property of the product of two rectangular matrices', *Comput. Math. Math. Phys.* **36** (1996), 555–557.

Department of Mathematics and Statistics

The University of Melbourne

VIC 3010

Australia

e-mail: a.dajic@ms.unimelb.edu.au, j.koliha@ms.unimelb.edu.au