

Ergodic theory and dynamical systems


EDITORS

Michel Herman Anatole Katok Anthony Manning William Parry

EDITORIAL BOARD

A Connes (IHES)	D S Ornstein (Stanford University)
J Feldman (University of California)	J Palis (IMPA)
J Franks (Northwestern University)	D J Rudolph (University of Maryland)
H Furstenberg (Hebrew University)	D Ruelle (IHES)
M Gromov (IHES)	K Schmidt (University of Warwick)
M Keane (Technische Hogeschool Delft)	Ya G Sinai (L D Landau Inst for Theoretical Physics, Moscow)
H Kesten (Cornell University)	J P Thouvenot (University of Paris 6)
U Krengel (University of Gottingen)	W A Veech (Rice University)
G A Margulis (USSR Acad of Sciences)	P Walters (University of Warwick)
J N Mather (Princeton University)	B Weiss (Hebrew University)
M Misiurewicz (Warsaw University)	R F Williams (Northwestern University)
S E Newhouse (University of Nth Carolina)	R Zimmer (University of Chicago)

VOLUME 7
1987



CAMBRIDGE UNIVERSITY PRESS

CAMBRIDGE

LONDON NEW YORK NEW ROCHELLE
MELBOURNE SYDNEY

PUBLISHED BY
THE PRESS SYNDICATE OF THE UNIVERSITY OF CAMBRIDGE

The Pitt Building, Trumpington Street, Cambridge CB2 1RP
32 East 57th Street, New York, NY 10022, USA
10 Stamford Road, Oakleigh, Melbourne 3166, Australia

© Cambridge University Press 1987

Printed in Great Britain by J W Arrowsmith Ltd, Bristol

Contents

PART 1 MARCH 1987

The algebraic-geometric AKNS potentials	<i>C de Concini and R A Johnson</i>	1
Topological dynamics and ergodic theory	<i>R Ellis</i>	25
Semi-rigidity of horocycle flows over compact surfaces of variable negative curvature	<i>J Feldman and D Ornstein</i>	49
An extension of Ratner's rigidity theorem to n -dimensional hyperbolic space	<i>L Flaminio</i>	73
A refinement of the Conley index and an application to the stability of hyperbolic invariant sets	<i>A Floer</i>	93
Classes of linear automata	<i>R H Gilman</i>	105
On Gibbs and equilibrium states	<i>N T A Haydn</i>	119
The $C^{1,1}$ conclusions in Gromov's theory	<i>C C Pugh</i>	133
Boshernitzan's criterion for unique ergodicity of an interval exchange transformation	<i>W A Veech</i>	149

PART 2 JUNE 1987

The ordering on permutations induced by continuous maps of the real line	<i>C Bernhardt</i>	155
Orbit types for maps of the interval	<i>L Block and D Hart</i>	161
Periodic orbits for dissipative twist maps	<i>M Casdagli</i>	165
Time-like geodesic flows on Lorentz manifolds	<i>G G Emch and S Hong</i>	175
Eventually positive matrices with rational eigenvectors	<i>D Handelman</i>	193
A characterization of flat metrics on tori by ergodicity	<i>N Innami</i>	197
Necessary and sufficient conditions for a maximal ergodic theorem along subsequences	<i>R L Jones</i>	203
On minimal self-joinings in topological dynamics	<i>A del Junco</i>	211
A rank-one, rigid, simple, prime map	<i>A del Junco and D J Rudolph</i>	229
Expansive dynamics on zero-dimensional groups	<i>B P Kitchens</i>	249
Loose Bernoullicity is preserved under exponentiation by integrable functions	<i>M H Rahe and D J Rudolph</i>	263
The rate of mixing for geodesic and horocycle flows	<i>M Ratner</i>	267

An example of an amenable action from geometry	<i>R J. Spatzier</i>	289
A modulus of 3-dimensional vector fields	<i>Y Togawa</i>	295
Markov measures determine the zeta function	<i>S Tuncel</i>	303
Book Reviews		313

PART 3 SEPTEMBER 1987

Periodic attractors as a result of diffusion	<i>J Barkmejer</i>	319
Weakly mixing PET	<i>V Bergelson</i>	337
Dynamics of Markov chains and stable manifolds for random diffeomorphisms	<i>M Brin and Yu Kifer</i>	351
Generic 3-parameter families of vector fields on the plane, unfolding a singularity with nilpotent linear part The cusp case of codimension 3	<i>F Dumortier, R Roussarie and J Sotomayor</i>	375
Diffeomorphisms on surfaces with a finite number of moduli	<i>W de Melo and S J van Strien</i>	415
A bound for the fixed-point index of area-preserving homeomorphisms on two-manifolds	<i>S Pelikan and E E Slaminka</i>	463

PART 4 DECEMBER 1987

Ergodic theorem for iterated maps	<i>J H Elton</i>	481
Finitely presented dynamical systems	<i>D Fried</i>	489
A counterexample to a C^2 closing lemma	<i>C Gutierrez</i>	509
On ergodic actions whose self-joinings are graphs	<i>A del Junco and D Rudolph</i>	531
Recurrence without uniform recurrence	<i>T W Korner</i>	559
On the generic existence of homoclinic points	<i>F Oliveira</i>	567
Generic properties of periodic reflecting rays	<i>L Stojanov</i>	597
Entropy of expansive flows	<i>R F Thomas</i>	611
Invariant subsets of expanding mappings of the circle	<i>M Urbański</i>	627
Index to Volume 7		647

INSTRUCTIONS TO AUTHORS

1 *Submission of typescripts*

Two copies of the manuscript should be submitted to one of the four Executive Editors (addresses on outside front cover) The editor will acknowledge receipt of the manuscripts **It is important that authors inform the editor of any changes of address** whilst their paper is under consideration

2 *Typescript*

Papers should be typed, double-spaced, on one side only and with generous margins The pages must be numbered

The first page should give the title, the author's name and institution, and a short abstract intelligible to mathematicians

The title, while brief, must be informative (e.g. *A new proof of the ergodic theorem*, whereas *Some applications of a theorem of Birkhoff* would be useless)

3 *Notation*

It is important that mathematical expressions are clear to a printer (who is not a mathematician) For instance, n_k (n sub k) is common usage, but avoid if possible using c sub n sub k . Fractions are generally best expressed by a solidus Complicated exponentials like

$$\exp \{z^2 \sin \theta / (1 + y^2)\}$$

should be shown in this and no other way

In the typescript, italics, small capitals and capitals are specified by single, double and triple underlining Bold-faced type is shown by wavy underlining

It helps if displayed equations or statements which will be quoted later are numbered in order on the right of their line They can then be referred to by, for example, 'from (7)'

The author must enable the printer (if necessary by pencilled notes in the margin) to distinguish between similar symbols such as o , O , o , O , 0 , x , X , \times , ϕ , Φ , \emptyset , l , 1 , ε , ϵ , κ , k .

There is no need to underline Greek or script letters provided these are clearly typed Any special symbols should be explained on a separate sheet of directions for the printer

If an author wishes to mark the end of the proof of a theorem, the sign \square may be used

Footnotes should be avoided

4 *Diagrams*

Figures and drawings should be on separate sheets in black ink Photocopies are acceptable only if

they are as clear as the originals *Symbols, legends and captions should be given on a transparent overlay* Each text figure must be numbered as Figure 1, Figure 2, and its intended position clearly indicated in the typescript The author's name in pencil must be on all separate sheets of diagrams

A figure is expensive to reproduce and should be included only when the subject matter demands it, or when it greatly clarifies the exposition

The publisher recognizes that some authors do not have the facilities for producing drawings of a sufficiently high standard to be reproduced directly and is therefore willing to have such diagrams re-drawn, provided that they are clear

5 *Tables*

Tables should be numbered (above the table) and set out on separate sheets Indicate the position of each in the text as for figures

6 *References*

References should be collected at the end of the paper numbered in alphabetical order of the authors' names A reference to a book should give the title, in italics, and then in roman type the publisher's name and the place and year of publication,

[4] N Dunford & J T Schwartz *Linear Operators* Part I Wiley New York, 1958

A reference to a paper should give in italics the title of the periodical, the number of the volume and year, and the beginning and end pages of the paper Titles should be abbreviated as in *Mathematical Reviews*

[6] J E Littlewood The 'pits effect' for functions in the unit circle *J Analyse Math* **23** (1970), 236-268

7 *Proofs*

Authors receive one set of proofs for correction If excessive alterations to the original manuscript are requested after the paper has been typeset, the author will be charged the cost of resetting For papers with more than one author the proofs are sent to the first named author unless the editor receives other instructions **It is important that proofs are corrected and returned promptly.**

8 *Reprints*

There are 100 reprints, free of charge, for each paper For papers with several authors these reprints are divided between the authors There are no page charges

Ergodic theory and dynamical systems

VOLUME 7 PART 4 DECEMBER 1987

CONTENTS

<i>Elton, J. H.</i> Ergodic theorem for iterated maps	481
<i>Fried, D.</i> Finitely presented dynamical systems	489
<i>Gutierrez, C.</i> A counter-example to a C^2 closing lemma	509
<i>del Junco, A. and Rudolph, D.</i> On ergodic actions whose self-joinings are graphs	531
<i>Körner, T. W.</i> Recurrence without uniform recurrence	559
<i>Oliveira, F.</i> On the generic existence of homoclinic points	567
<i>Stojanov, L.</i> Generic properties of periodic reflecting rays	597
<i>Thomas, R. F.</i> Entropy of expansive flows	611
<i>Urbański, M.</i> Invariant subsets of expanding mappings of the circle	627
Index to Volume 7	647

© Cambridge University Press 1987

CAMBRIDGE UNIVERSITY PRESS

The Pitt Building, Trumpington Street, Cambridge CB2 1RP

32 East 57th Street, New York, NY 10022, USA

10 Stamford Road, Oakleigh, Melbourne 3166, Australia

Printed in Great Britain by J. W. Arrowsmith Ltd, Bristol