

**Disclosure of interest** The authors have not supplied their declaration of competing interest.

<http://dx.doi.org/10.1016/j.eurpsy.2017.02.135>

#### EW0522

### Burnout in medical residents: Prevalence and risk factors

I. Feki<sup>1</sup>, S. Hentati<sup>1</sup>, R. Sallemi<sup>1</sup>, N. Bahloul<sup>2</sup>, D. Zalila<sup>2</sup>, S. Kammoun<sup>2</sup>, J. Masmoudi<sup>1,\*</sup>

<sup>1</sup> University Hospital Hédi Chaker, psychiatry, Sfax, Tunisia

<sup>2</sup> University Hospital Hédi Chaker, pneumology, Sfax, Tunisia

\* Corresponding author.

**Background** Burnout is a state of mental and physical exhaustion related to work. It is manifested in a variety of professions and is prevalent in contexts in which health professionals are required to interact directly with the public.

**Objectives** To estimate the burnout syndrome (BOS) prevalence and to identify its associated factors among medical residents.

**Methods** A descriptive and analytic study conducted during a period from 2012 until 2014, including 184 residents exerting at the two university hospitals of Sfax, Tunisia. Data collection was conducted using an anonymous self-questionnaire, including demographic items and working conditions items. Maslach Burnout Inventory (MBI) was used to assess BOS.

**Results** The average age was  $25.14 \pm 1.47$  years. The sex-ratio was 0.65. The majority (154 residents) was single. Over than half (58.2%) of medical residents endure a high emotional exhaustion, 62.5% a high depersonalisation and 12.5% a low personal accomplishment. About eighty percent (79.3%) of them were in burn out with 37% at intermediate and 11.3% at a severe degree. No statistical association was found between the personal characteristics and BOS. Whereas, several factors related to employment status were correlated with the presence of BOS such us: number of working hours > 30 h/week, a number of free weekends < 2/month and a number of days off  $\leq 30$  days/year. In addition, we found that psychotropic and alcohol consumption were significantly associated to BOS with respectively  $P=0.050$  and  $P=0.002$ .

**Conclusion** Burnout is a worrying reality among young doctors. Individual as well as organizational interventions should be targeted to prevention.

**Disclosure of interest** The authors have not supplied their declaration of competing interest.

<http://dx.doi.org/10.1016/j.eurpsy.2017.02.136>

#### EW0523

### Eating disorders in schizophrenia

R. Sallemi, S. Hentati, I. Feki, J. Masmoudi\*, M. Moala

University hospital Hédi Chaker, psychiatry, Sfax, Tunisia

\* Corresponding author.

**Background** Due to their frequency and negative impact on quality of life, eating disorders in schizophrenia need to be considered and highlighting.

**Objective** To identify the risk of eating disorders (ED) and its correlates among mental patients.

**Methods** It was a descriptive and analytic study. It included 53 inpatients with DSM-5 diagnoses of schizophrenia or schizoaffective disorder, followed in the department of Psychiatry at the Hedi Chaker University Hospital of Sfax in Tunisia, during the three months of August, September and October 2016. Data collections were conducted using questionnaire exploring sociodemographic and medical data. The SCOFF (sick, control, one, fat, food) Questionnaire was used to screen ED. A total score of  $\geq 2$  was used as a cutoff point to select persons at risk of ED.

**Results** The average age of our patients was  $30.47 \pm 9.5$  years old. The majority of our patients was male (71.7%) and single (71%). The

mean of extra Body mass was 27.9. The mean duration of disease was  $9.9 \pm 8.1$  years and patients were mostly (54%) in atypical neuroleptics. According to the SCOFF Questionnaire, 35.8% had a risk of ED. Female gender and treatment with atypical neuroleptics were significantly associated to ED risk with respectively  $P=0.02$  and  $P=0.038$ .

**Conclusion** Eating disorders remain underestimated among patients suffering from schizophrenia. Yet, its screening prevention and management are crucial and must be multidisciplinary for optimal care.

**Disclosure of interest** The authors have not supplied their declaration of competing interest.

<http://dx.doi.org/10.1016/j.eurpsy.2017.02.137>

#### EW0524

### Palatable food access during adolescence increased BDNF expression in the nucleus accumbens and anxiety-/depression-like behaviors in males, but not in females

J.W. Jahng\*, J.Y. Kim, J.H. Lee

School of Dentistry, Seoul National University, Dental Research Institute, Seoul, Republic of Korea

\* Corresponding author.

**Objectives** This study was conducted to examine sexual dimorphic effects of highly palatable food access during adolescence and youth on psychoemotional behaviors of rats and its underlying neural mechanism.

**Methods** Male and female Sprague Dawley pups had free access to chocolate cookie rich in fat (highly palatable food) from post-natal day 28 in addition to ad libitum chow, and the control groups received chow only. The food conditions were continued though out the entire experimental period, and the neurochemical and behavioral measurements were performed during young adulthood. Corticosterone levels during 2 h of restraint stress were analyzed with radioimmunoassay, and  $\Delta$ FosB and brain-derived neurotrophic factor (BDNF) expression in the nucleus accumbens (NAc) with western blot analysis.

**Results** Cookie access did not affect body weight gain and total caloric intake in both sexes; however, it increased retroperitoneal fat depot only in males. The time spent in open arms during elevated plus maze test was decreased and immobility during forced swim test was increased in cookie-fed males, but not in cookie-fed females. Main effect of food condition on the stress-induced corticosterone increase was observed in males, but not in females, and cookie access increased BDNF expression in the NAc only in males.

**Conclusions** Increased BDNF expression in the NAc and fat depot, in addition to the HPA axis dysfunction, may play roles in the pathophysiology of depression- and/or anxiety-like behaviors induced by cookie access.

**Disclosure of interest** The authors have not supplied their declaration of competing interest.

<http://dx.doi.org/10.1016/j.eurpsy.2017.02.138>

#### EW0525

### Investigation of food addiction and impulsivity relations biological rhythms differences and insomnia in university students

A. Kandeger\*, Y. Selvi

Selcuk University, Department of Psychiatry, Konya, Turkey

\* Corresponding author.

**Objective** Increasing prevalence of obesity in the world and increasing role of processed foods in daily life has led to become the focal point of food addiction. This study aims to investigation of food