British Journal of Nutrition

cambridge.org/bjn

Policy Relevant Paper

Cite this article: Zhang M, Alsokhn C, Jakob R, Li N, and Wang Y (2024). Advancing nutritional disorder classification: unleashing the impact of ICD-11 on clinical practice and public health. *British Journal of Nutrition*, page 1 of 2. doi: 10.1017/S0007114524002502

Received: 27 September 2024 Accepted: 13 October 2024

Keywords

ICD-11; Nutritional disorders; International Classification of Diseases; Health data standard

Abbreviations:

ICD, International Classification of Diseases

Corresponding author:

Naishi Li; Email: Ins@medmail.com.cn

[†]The authors contributed equally to this work

© The Author(s), 2024. Published by Cambridge University Press on behalf of The Nutrition Society. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (https://creative.commons.org/licenses/by/4.0/), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.





Advancing nutritional disorder classification: unleashing the impact of ICD-11 on clinical practice and public health

Meng Zhang^{1,2,†}, Carine Alsokhn^{3,†}, Robert Jakob³, Naishi Li^{1,2,4} and Yi Wang^{1,2}

¹Collaborating Center for the WHO Family of International Classifications in China, Beijing, People's Republic of China; ²Department of Medical Records, Peking Union Medical College Hospital, Chinese Academy of Medical Sciences & Peking Union Medical College, Beijing, People's Republic of China; ³Department of Data and Analytics, World Health Organization, Geneva, Switzerland and ⁴Department of Endocrinology, Key Laboratory of Endocrinology of National Health Commission, Peking Union Medical College Hospital, Chinese Academy of Medical Sciences & Peking Union Medical College, Beijing, People's Republic of China

Nutritional disorders represent a major global public health issue, with significant and enduring impacts on society and healthcare systems. The International Classification of Diseases (ICD) is fundamental for diagnosing and classifying diseases and health conditions worldwide, supporting health statistics, monitoring, planning and resource allocation. It is extensively used for recording, reporting, researching and comparing nutritional statuses. The prevalence, mortality and disease burden of nutritional disorders are reported based on ICD-coded data^(1–3). Studies have demonstrated that nutritional assessment and ICD coding can lead to increased case mix indices, patient complexity and co-morbidity levels, and health insurance reimbursements^(4–6).

However, some concepts related to nutritional disorders in ICD-10, which have been in use for over 30 years, are now outdated and do not fully align with current clinical practice and research⁽⁷⁾. ICD-11, approved at the 72nd World Health Assembly and effective from 1 January 2022, introduces a range of substantial advancements. These include a semantic network known as the Foundation, updated classifications and terminologies, enhancements in the coding framework, full digital support and interoperability^(8,9).

Advised by the Nutrition Guidance Expert Advisory Group, the Department of Nutrition for Health and Development of WHO has actively engaged in updating the nutritional disorders in ICD-11 to align with WHO's guidelines for nutrition surveillance and management. As a result, significant updates have been made in the classification of undernutrition from ICD-10 to ICD-11. In ICD-11, undernutrition is now categorised based on anthropometric or clinical standards and deficiencies in specific nutrients. For example, ICD-11 provides distinct codes for underweight, wasting, acute malnutrition and stunting in infants, children and adolescents. ICD-11 is currently available in ten languages, with translations into twenty-five additional languages underway, and more expected in the near future. This language-independent classification standardises the understanding and recording of nutritional disorders globally, facilitating national and international comparisons and research, and enabling the tracking of indicators for the Sustainable Development Goals and the Global Nutrition Monitoring Framework^(10,11).

ICD-11 provides greater granularity than ICD-10, offering more detailed and specific categories and subcategories for nutritional disorders. For instance, while ICD-10 does not differentiate between overweight and obesity, ICD-11 assigns distinct codes for overweight and for obesity due to energy imbalance across different age groups. Additionally, while ICD-10 groups all forms of beriberi under a single code, ICD-11 subdivides beriberi into dry and wet forms. Furthermore, ICD-11 introduces Wernicke–Korsakoff syndrome as a distinct entity with separate codes for Wernicke encephalopathy and Korsakoff syndrome, reflecting the disease's continuum. All entities in ICD-11 can be uniquely identified using a uniform resource identifier (URI) in the Foundation, even those without a statistical code. These updates enhance the accuracy of prevalence data and improve the epidemiological monitoring of these conditions.

ICD-11 enhances clinical reporting with its code combination approach known as cluster coding, to capture detailed clinical information through postcoordination⁽¹²⁾. For instance, a patient with polyneuropathy due to vitamin B_{12} deficiency would be coded as 8C03.3/5B5F, where 8C03.3 indicates polyneuropathy in nutritional deficiency and 5B5F identifies the vitamin B_{12} deficiency. This approach allows for the reporting of both the manifestation and the aetiology, thereby improving statistical precision and contributing to public health strategies and resource allocation. Meanwhile, ICD-11 offers an array of extension codes for use in cluster coding to capture more details, such as severity, temporality, anatomy and medication. For example, obesity induced by Olanzapine could be captured using the cluster 5B81.1&XM6GK7,



2 M. Zhang et al.

where 5B81.1 represents drug-induced obesity and XM6GK7 indicates Olanzapine as the causative agent. This level of detail is invaluable for refining patient management and devising targeted interventions.

The content model used for ICD-11 allows each entity to include a textual definition and diagnostic criteria (13,14). The textual definition provides the meaning and descriptive features of each entity, while the diagnostic criteria outline the necessary diagnostic information. Definitions from WHO sources have been attached to some nutritional disorders in ICD-11, with more anticipated as international consensus is achieved. ICD-11 has significant potential for standardising diagnoses and improving the quality of data reporting. Continuous efforts are required to fully realise these potential benefits.

ICD-11 continues to evolve with medical and scientific advancements, incorporating input from global users. The WHO has established a maintenance mechanism for ICD-11, supported by an open platform and a transparent, rigorous review process. As of August 2024, users from over seventy countries have submitted proposals to the platform, contributing to the ongoing updates of ICD-11. A total of 125 proposals related to nutritional disorders have been received on the maintenance platform, of which 105 (84%) have been evaluated and decided upon. The remaining proposals are currently undergoing further review or awaiting triage, including those submitted by the author to enrich the Foundation with nutrition-related concepts. Some of these suggested concepts were reported missing in ICD in academic publications but had not been communicated to the ICD-11 maintenance process^(9,15). After 2 years following the official coming into effect of ICD-11, 132 Member States and areas are at various phases of implementation (16). As this global implementation progresses, more suggestions and feedback on ICD-11 are anticipated. The nutrition community can contribute to the ICD-11 by directly engaging with the WHO maintenance platform (https://icd.who.int/dev11) when the need for enhancements is identified.

In summary, ICD-11 represents a significant advancement in the classification of nutritional disorders, introducing innovations that enhance clinical practices and public health outcomes. The increased granularity and standardisation of nutritional disorder classifications in ICD-11 support more targeted management, enhance epidemiological monitoring, facilitate international health data comparisons and bolster global health initiatives aimed at addressing these conditions. As ICD-11 continues to evolve with progress in nutrition science, it will unlock its full potential through active engagement of the nutrition community. The transition from ICD-10 to ICD-11 involved collaboration among all relevant stakeholders. Training within nutrition communities and piloting ICD-11 in practice are important steps for a smooth transition.

Acknowledgements. This work was supported by the National High Level Hospital Clinical Research Funding (no. 2022-PUMCH-B-091 and no. 2022-PUMCH-A-223).

The authors declare none.

References

- World Health Organization (1999) WHO Mortality Database List of Causes and Corresponding ICD-10 Codes. https://platform.who.int/ mortality/about/list-of-causes-and-corresponding-icd-10-codes (accessed September 2024).
- GBD 2021 Causes of Death Collaborators (2024) Global burden of 288
 causes of death and life expectancy decomposition in 204 countries and
 territories and 811 subnational locations, 1990–2021: a systematic analysis
 for the Global Burden of Disease Study 2021. Lancet 403, 2100–2132.
- 3. Saengnipanthkul S, Phosuwattanakul J, Thepsuthammarat K, *et al.* (2022) Epidemiological data on nutritional disorders and outcomes in hospitalized Thai children: an analysis of data from the National Health Database 2015–2019. *Epidemiol Health* **44**, e2022047.
- Ockenga J, Freudenreich M, Zakonsky R, et al. (2005) Nutritional assessment and management in hospitalised patients: implication for DRG-based reimbursement and health care quality. Clin Nutr 24, 913–919.
- Li X, Yuan L, Gao B, et al. (2023) Comparison of diagnosis-related group based reimbursement and case-mix index within hospitalized patients before and after modified malnutrition diagnosis. Asia Pac J Clin Nutr 32, 356–361.
- Stollhof LE, Braun JM, Ihle C, et al. (2019) The continuous downgrading of malnutrition in the German DRG system: possible effects on the treatment of patients at risk for malnutrition. Excli J 18, 370–381.
- Cederholm T, Barazzoni R, Austin P, et al. (2017) ESPEN guidelines on definitions and terminology of clinical nutrition. Clin Nutr 36, 49–64.
- 8. Chute CG & Celik C (2022) Overview of ICD-11 architecture and structure. BMC Med Inform Decis Making 21, 378.
- de Onis M, Zeitlhuber J & Martínez-Costa C (2016) Nutritional disorders in the proposed 11th revision of the International Classification of Diseases: feedback from a survey of stakeholders. *Public Health Nutr* 19, 3135–3141.
- United Nations (2015) Sustainable Development Goals. https://sdgs.un. org/goals (accessed May 2024).
- 11. World Health Organization (2017) Global Nutrition Monitoring Framework: Operational Guidance for Tracking Progress in Meeting Targets for 2025. Geneva: World Health Organization. https://www.who.int/publications/i/item/9789241513609 (accessed September 2024).
- Mabon K, Steinum O & Chute CG (2022) Postcoordination of codes in ICD-11. BMC Med Inf Decis Making 21, 379.
- 13. World Health Organization (2021) WHO-FIC Content Model Reference Guide. https://icd.who.int (accessed September 2024).
- Jiang G, Solbrig HR & Chute CG (2013) Using Semantic Web technology to support ICD-11 textual definitions authoring. J Biomed Semant 4, 11.
- Cederholm T, Bosaeus I, Barazzoni R, et al. (2015) Diagnostic criteria for malnutrition - an ESPEN Consensus Statement. Clin Nutr 34, 335–340.
- World Health Organization (2024) WHO Advances Implementation and Integration of ICD-11 and Related Medical Classifications and Terminologies. https://www.who.int/standards/classifications/classificationof-diseases (accessed September 2024).