




Li coefficients and the quadrilateral zeta function

Kajtaz H. Bllaca, Kamel Mazhouda, and Takashi Nakamura 

Abstract. In this note, we study the Li coefficients $\lambda_{n,a}$ for the quadrilateral zeta function. Furthermore, we give an arithmetic and asymptotic formula for these coefficients. Especially, we show that for any fixed $n \in \mathbb{N}$, there exists $a > 0$ such that $\lambda_{2n-1,a} > 0$ and $\lambda_{2n,a} < 0$.

1 Introduction and statement of main results

1.1 Li coefficients

The Riemann hypothesis (RH) is a critical question in analytic number theory. As such, it is interesting to examine different ways to frame it, which may shed more light on its resolution. In 1997, Xian-Jin Li has discovered a new positivity criterion for the RH. In [10], he defined the Li coefficients for the Riemann zeta function as

$$\lambda_n = \frac{1}{(n-1)!} \frac{d^n}{ds^n} \left[s^{n-1} \log \xi(s) \right]_{s=1},$$

where ξ is the completed Riemann zeta function defined by

$$\xi(s) = s(s-1)\pi^{-s/2}\Gamma(s/2)\zeta(s),$$

which satisfies $\xi(s) = \xi(1-s)$ and gave a simple equivalence criterion for the RH: RH is true if and only if these coefficients are nonnegative for every positive integer n . The Li coefficients λ_n can be written as follows:

$$\lambda_n = \sum_{\rho}^* \left[1 - \left(1 - \frac{1}{\rho} \right)^n \right] = \lim_{T \rightarrow \infty} \sum_{\rho: |\operatorname{Im}(\rho)| \leq T} \left[1 - \left(1 - \frac{1}{\rho} \right)^n \right],$$

where the sum runs over the nontrivial zeros of the Riemann zeta function counted with multiplicity. This criterion is generalized by Bombieri and Lagarias [4] for any arbitrarily multiset of numbers assuming certain convergence conditions. Voros [19, Section 3.3] has proved that the RH true is equivalent to the growth of λ_n as $\frac{1}{2}n \log n$ determined by its archimedean part, while the RH false is equivalent to the oscillations of λ_n with exponentially growing amplitude, determined by its finite part. The Li

Received by the editors May 21, 2023; revised March 25, 2024; accepted March 26, 2024.

Published online on Cambridge Core April 8, 2024.

The third author was partially supported by JSPS grant 22K03276.

AMS subject classification: 11M26, 11M35.

Keywords: Li's coefficients, the quadrilateral zeta function, Riemann Hypothesis.



coefficients were generalized in two ways: by generalizing these coefficients to various sets of functions (the Selberg class, the class of automorphic L -functions, zeta function on function fields, ... [8, 11, 17]) and by introducing new parameter in its definition (see [12]). The Li coefficients (and its generalizations) have generated a lot of research interest due to its applicability and simplicity.

1.2 Quadrilateral zeta function

Recall the definitions of Hurwitz and periodic zeta functions. The Hurwitz zeta function $\zeta(s, a)$ is defined by the series

$$\zeta(s, a) := \sum_{n=0}^{\infty} \frac{1}{(n+a)^s}, \quad \sigma > 1, \quad 0 < a \leq 1.$$

The function $\zeta(s, a)$ is a meromorphic function with a simple pole at $s = 1$ whose residue is 1 (see, for example, [1, Section 12]). The periodic zeta function $\text{Li}_s(e^{2\pi ia})$ is defined by

$$\text{Li}_s(e^{2\pi ia}) := \sum_{n=1}^{\infty} \frac{e^{2\pi ina}}{n^s}, \quad \sigma > 1, \quad 0 < a \leq 1$$

(see, for instance, [1, Exercise 12.2]). Note that the function $\text{Li}_s(e^{2\pi ia})$ with $0 < a < 1$ is analytically continuable to the whole complex plane since $\text{Li}_s(e^{2\pi ia})$ does not have any pole, that is shown by the fact that the Dirichlet series of $\text{Li}_s(e^{2\pi ia})$ converges uniformly in each compact subset of the half-plane $\sigma > 0$ when $0 < a < 1$ (see, for example, [9, p. 20]). For $0 < a \leq 1/2$, we define zeta functions

$$\begin{aligned} Z(s, a) &:= \zeta(s, a) + \zeta(s, 1-a), & P(s, a) &:= \text{Li}_s(e^{2\pi ia}) + \text{Li}_s(e^{2\pi i(1-a)}), \\ 2Q(s, a) &:= Z(s, a) + P(s, a), & \xi_Q(s, a) &:= s(s-1)\pi^{-s/2}\Gamma(s/2)Q(s, a). \end{aligned}$$

We can see that $Q(s, a)$ is meromorphic functions with a simple pole at $s = 1$. In addition, we have $Q(0, a) = -1/2 = \zeta(0)$ and $\xi_Q(s, a) = \xi_Q(1-s, a)$, which is proved by

$$(1.1) \quad Q(1-s, a) = \Gamma_{\cos}(s)Q(s, a), \quad \Gamma_{\cos}(s) := \frac{2\Gamma(s)}{(2\pi)^s} \cos\left(\frac{\pi s}{2}\right)$$

(see [13, Theorem 1.1]). Moreover, the function $Q(s, a)$ has the following properties. When $a = 1/6, 1/4, 1/3$, and $1/2$, the RH holds true if and only if all nonreal zeros of $Q(s, a)$ are on the line $\text{Re}(s) = 1/2$ (see [14, Proposition 1.3]). Let $N_Q^{\text{CL}}(T)$ be the number of the zeros of $Q(s, a)$ on the line segment from $1/2$ to $1/2 + iT$. In [13, Theorem 1.2], the third author proved that for any $0 < a \leq 1/2$, there exist positive constants $A(a)$ and $T_0(a)$ such that

$$N_Q^{\text{CL}}(T) \geq A(a)T \quad \text{whenever} \quad T \geq T_0(a).$$

Next, let $N_F(T)$ count the number of nonreal zeros of a function $F(s)$ having $|\text{Im}(s)| < T$. Then, for any $0 < a \leq 1/2$,

$$N_{\zeta}(T) - N_Q(T) = O_a(T),$$

and the third author [14, Proposition 1.8] proved that

$$N_Q(T) = \frac{T}{\pi} \log T - \frac{T}{\pi} \log(2e\pi a^2) + O_a(\log T).$$

Furthermore, he [14, Theorem 1.1] proved that there is a unique absolute $a_0 \in (0, 1/2)$ such that

$$Q(1/2, a) > 0 \iff 0 < a < a_0.$$

In addition, it is proved in [14, Corollary 1.2] that all real zeros of $Q(s, a)$ are simple and are located only at the negative even integers just like $\zeta(s)$ if and only if $a_0 < a \leq 1/2$. Let us note by Z_Q the set of all nontrivial zeros ρ_a of $\xi_Q(s, a)$. Since it is an entire function of order 1, one has

$$(1.2) \quad \xi_Q(s, a) = e^{A+Bs} \prod_{\rho_a \in Z_Q} \left(1 - \frac{s}{\rho_a}\right) e^{\frac{s}{\rho_a}} = \xi_Q(0, a) \prod_{\rho_a \in Z_Q} \left(1 - \frac{s}{\rho_a}\right),$$

where $e^A = 1/2$, $B = \frac{Q'}{Q}(0, a) - 1 - \frac{\gamma + \log \pi}{2}$, and γ denotes the Euler constant. Note that $Q'(0, a)$ is given explicitly in [14, Theorem 1.5].

1.3 Main results

Recall that $\zeta(1-s) = \Gamma_{\cos}(s)\zeta(s)$ and $Q(1-s, a) = \Gamma_{\cos}(s)Q(s, a)$ by (1.1). However, the function $Q(s, a)$ does not have an Euler product except for $a = 1/6, 1/4, 1/3$, and $1/2$. Hence, the function $Q(s, a)$ is a suitable object to consider the influence of not Riemann’s functional equation but an Euler product to zeros of zeta functions. We show a criterion for nonvanishing of $Q(s, a)$ in terms of the positivity of the Li coefficients, an arithmetic and asymptotic formula for these coefficients in Theorems 1.1, 1.2, and 1.4, respectively. It should be emphasized that $\lambda_{n,a}$ defined in (1.3) are the first Li coefficients that we can explicitly give $n \in \mathbb{N}$ such that $\lambda_{n,a} < 0$. There is a possibility that this fact would give an idea to find negative Li coefficients for $\zeta(s)$ if they would exist.

For $n \neq 0$, the Li coefficients attached to $Q(s, a)$ nonvanishing at zero are defined by the sum

$$\lambda_{n,a} := \sum_{\rho_a \in Z_Q}^* \left(1 - \left(1 - \frac{1}{\rho_a}\right)^n\right) = \lim_{T \rightarrow \infty} \sum_{|\text{Im}(\rho_a)| \leq T}^* \left(1 - \left(1 - \frac{1}{\rho_a}\right)^n\right).$$

The symmetry $\rho_a \mapsto 1 - \rho_a$ in the set Z_Q of nontrivial zeros of $Q(s, a)$ implies that $\lambda_{-n,a} = \overline{\lambda_{n,a}} = \lambda_{n,a}$ for all $n \in \mathbb{N}$. So, $\lambda_{n,a}$ are real. We have also

$$(1.3) \quad \lambda_{n,a} := \frac{1}{(n-1)!} \frac{d^n}{ds^n} [s^{n-1} \log \xi_Q(s, a)]_{s=1}.$$

Moreover, from (1.2), we have (see [4, Equations (2.3) and (2.4)] or [17, Appendix A])

$$\sum_{n=0}^{\infty} \lambda_{n+1,a} s^n = \frac{d}{ds} \log \left[\xi_Q \left(\frac{1}{1-s}, a \right) \right].$$

As an analogue of Li coefficients for the Riemann zeta function, we have the following.

Theorem 1.1 *The function $Q(s, a)$ does not vanish when $\text{Re}(s) > 1/2$ if and only if $\lambda_{n,a} \geq 0$ for all $n \in \mathbb{N}$.*

An arithmetic formula for $\lambda_{n,a}$ is stated in the following theorems.

Theorem 1.2 *We have*

$$\lambda_{n,a} = 1 - \frac{n}{2}(\log(4\pi) + \gamma) + \sum_{k=2}^n (-1)^k \binom{n}{k} (1 - 2^{-k}) \zeta(k) + \sum_{k=1}^n \binom{n}{k} \gamma_Q(k-1),$$

where $\gamma_Q(n)$ are defined as follows:

$$\frac{Q'}{Q}(s+1, a) + \frac{1}{s} = \sum_{n=0}^{\infty} \gamma_Q(n) s^n.$$

Theorem 1.3 *For $a = 1/2, 1/3, 1/4, 1/6$, under the RH, we have*

$$\lambda_{n,a} = \frac{n}{2} \log n + \frac{n}{2} (\gamma - 1 - \log 2\pi) + O(\sqrt{n} \log n).$$

For a fixed $l \in \mathbb{N}$, we have the following asymptotic formula of $\lambda_{l,a}$ when $a \rightarrow +0$. We can see that there exists $n \in \mathbb{N}$ such that $\lambda_{n,a} < 0$ by Theorem 1.1 and the fact that $Q(s, a)$ does not satisfy an analogue of the RH when $a \in \mathbb{Q} \cap (0, 1/2) \setminus \{1/6, 1/4, 1/3\}$ (see [14, Proposition 1.4]). Clearly, this argument gives no information on the frequency of $n \in \mathbb{N}$, the smallest $n \in \mathbb{N}$ such that $\lambda_{n,a} < 0$ and so on. However, the next theorem implies that $\lambda_{2n,a} < 0$ if we fix any $n \in \mathbb{N}$ and then we take $a > 0$ sufficiently small.

Theorem 1.4 *Fix $l \in \mathbb{N}$. Then it holds that*

$$\lambda_{l,a} = \frac{(-1)^{l+1}}{(2a)^l} + O_l(a^{1-l} |\log a|), \quad a \rightarrow +0.$$

Especially, for any fixed $n \in \mathbb{N}$, there are $a > 0$ such that

$$\lambda_{2n-1,a} > 0 \quad \text{and} \quad \lambda_{2n,a} < 0.$$

2 Proofs

2.1 Proof of Theorem 1.1

Since $\lambda_{-n,a} = \overline{\lambda_{n,a}} = \lambda_{n,a}$ for all $n \in \mathbb{N}$, then $\text{Re}(\lambda_{-n,a}) = \text{Re}(\lambda_{n,a}) = \lambda_{n,a}$. Using that $\xi_Q(s, a)$ is an entire function of order 1, and its zeros lie in the critical strip $0 < \text{Re}(s) < 1$, we obtain that the series $\sum_{\rho \in Z_Q} \frac{1+|\text{Re}(\rho)|}{(1+|\rho|)^2}$ is convergent. Application of [4, Theorem 1] to the multiset Z_Q of zeros of $Q(s, a)$ gives that $\text{Re}(\rho) \leq 1/2$ if and only if $\lambda_{n,a} \geq 0$ for all $n \in \mathbb{N}$. Now, the application of the same theorem to the multiset $1 - Z_Q = Z_Q$ gives $\text{Re}(\rho) \geq 1/2$ if and only if $\lambda_{n,a} \geq 0$. This completes the proof.

Theorem 1.1 can also be proved by the same argument used in [5, Theorem 1], which is due to Oesterlé.

2.2 Proof of Theorem 1.2

From the expression of $\xi_Q(s, a)$, one has

$$\frac{\xi'_Q}{\xi_Q}(s, a) = \frac{1}{s} + \frac{1}{s-1} - \frac{1}{2} \log \pi + \frac{1}{2} \frac{\Gamma'}{\Gamma}(s/2) + \frac{Q'}{Q}(s, a),$$

which is rewritten as

$$(2.1) \quad \frac{\xi'_Q}{\xi_Q}(s+1, a) = \frac{1}{s+1} + \frac{1}{s} - \frac{1}{2} \log \pi + \frac{1}{2} \frac{\Gamma'}{\Gamma}((s+1)/2) + \frac{Q'}{Q}(s+1, a).$$

Note that $Q(s, a)$ is a meromorphic function on the whole complex plane, which is holomorphic everywhere except for a simple pole at $s = 1$ with residue 1 (see [13, Section 2.1]). Let us define the coefficients $\gamma_Q(n)$ and $\tau_Q(n)$ as follows:

$$(2.2) \quad \frac{Q'}{Q}(s+1, a) + \frac{1}{s} = \sum_{n=0}^{\infty} \gamma_Q(n) s^n$$

and

$$(2.3) \quad -\frac{1}{2} \log \pi + \frac{1}{2} \frac{\Gamma'}{\Gamma}((s+1)/2) = \sum_{n=0}^{\infty} \tau_Q(n) s^n.$$

By Equation (1.2), one has

$$\log \xi_Q(s, a) = \log \xi_Q(0, a) - \sum_{\rho_a \in Z_Q} \sum_{m=1}^{\infty} \frac{1}{m \rho^m} s^m.$$

From the functional equation for the function $\xi_Q(s, a)$, in the neighborhood of $s = 0$, we have

$$(2.4) \quad \frac{\xi'_Q}{\xi_Q}(s+1, a) = -\frac{\xi'_Q}{\xi_Q}(-s, a) = \sum_{m=0}^{\infty} (-1)^m \sum_{\rho_a \in Z_Q} \frac{1}{\rho^{m+1}} s^m.$$

Comparing Equations (2.1)-(2.4), we get

$$(-1)^m \sum_{\rho_a \in Z_Q} \frac{1}{\rho^{m+1}} = (-1)^m + \gamma_Q(m) + \tau_Q(m),$$

for $m \geq 0$. Hence, the definition of $\lambda_{n,a}$ yields

$$\lambda_{n,a} = \sum_{k=1}^n (-1)^{k-1} \binom{n}{k} \sum_{\rho_a \in Z_Q} \frac{1}{\rho^k} = 1 + \sum_{k=1}^n \binom{n}{k} \gamma_Q(k-1) + \sum_{k=1}^n \binom{n}{k} \tau_Q(k-1),$$

where

$$\tau_Q(0) = -\frac{1}{2} \log \pi + \frac{1}{2} \psi(1/2) \text{ and } \tau_Q(k-1) = (-1)^k \sum_{m=0}^{\infty} \frac{1}{(2m+1)^k}$$

using that $\psi(z) = -\gamma - \frac{1}{z} + \sum_{k=1}^{\infty} \frac{z}{k(k+z)}$. Here, $\psi(s) = \frac{\Gamma'(s)}{\Gamma(s)}$ is the logarithmic derivative of the Gamma function. Since $\psi(1/2) = -\gamma - 2 \log 2$, we obtain

$$\begin{aligned} \lambda_{n,a} &= 1 - \frac{n}{2}(\log(4\pi) + \gamma) + \sum_{k=2}^n (-1)^k \binom{n}{k} \sum_{m=0}^{\infty} \frac{1}{(2m+1)^k} + \sum_{k=1}^n \binom{n}{k} \gamma_Q(k-1) \\ &= 1 - \frac{n}{2}(\log(4\pi) + \gamma) + \sum_{k=2}^n (-1)^k \binom{n}{k} (1-2^{-k}) \zeta(k) + \sum_{k=1}^n \binom{n}{k} \gamma_Q(k-1). \end{aligned}$$

The equality above implies Theorem 1.2.

2.3 Proof of Theorem 1.3

Let us note that

$$\sum_{k=2}^n (-1)^k \binom{n}{k} (1-2^{-k}) \zeta(k) = \sum_{k=2}^n (-1)^k \binom{n}{k} \frac{\zeta(k, 1/2)}{2^k},$$

where $\zeta(s, a)$ is the Hurwitz zeta function defined in Section 1.2. With the notation of Flajolet and Vespas [7, Lines 2–4, p. 70], this is $A_n(1, 2)$ and which is equal to

$$\frac{n}{2} \psi(n) + n \left(\gamma - \frac{1}{2} + \frac{1}{2} \log 2 \right) + o(1),$$

where the $o(1)$ error term above is exponentially small and oscillating and equal to

$$\frac{1}{2} \left(\frac{n}{\pi} \right)^{1/4} \exp(-\sqrt{2\pi n}) \cos \left(\sqrt{2\pi n} - \frac{5\pi}{8} \right) + O \left(n^{-1/4} e^{-\sqrt{2\pi n}} \right).$$

Then we have

$$\lambda_{n,a} = \frac{n}{2} \log n + \frac{n}{2} (\gamma - 1 - \log 2\pi) + \sum_{k=1}^n \binom{n}{k} \gamma_Q(k-1) + O(1).$$

It remains to prove that

$$(2.5) \quad \sum_{k=1}^n \binom{n}{k} \gamma_Q(k-1) = O(\sqrt{n} \log n).$$

To do so, we follow very closely the lines of the proof of the corresponding result in [8, Theorem 6.1] or [16, Lemma 3.3] and it will be shortened. We use the following kernel function:

$$k_n(s) := \left(1 + \frac{1}{s} \right)^n - 1 = \sum_{k=1}^n \binom{n}{k} \frac{1}{s^k}.$$

The residue theorem gives

$$\sum_{k=1}^n \binom{n}{k} \gamma_Q(k-1) = \frac{1}{2i\pi} \int_C k_n(s) \left(-\frac{Q'}{Q}(s+1, a) \right) ds,$$

where C is a contour enclosing the point $s = 0$ counterclockwise on a circle of small enough positive radius. The residue comes entirely from the singularity at $s = 0$, as no other singularities lie inside the contour. Let $T = \sqrt{n} + \varepsilon_n$, for some $0 < \varepsilon_n < 1$.

Now we follow very closely the lines in [16, pp. 1106–1107] using that the function $\frac{Q'}{Q}(s, a)$ satisfies the properties¹

$$\frac{Q'}{Q}(s, a) = \sum_{\rho_a; |\text{Im}(\rho_a - s)| < 1} \frac{1}{s - \rho_a} + O(\log(1 + |s|)),$$

for $-2 < \text{Re}(s) < 2$ and

$$\left| \frac{Q'}{Q}(s + 1, a) \right| = O(\log^2 T),$$

for $-2 \leq \text{Re}(s) \leq 2$, and we get

$$\sum_{k=1}^n \binom{n}{k} \gamma_Q(k - 1) = \lambda_{-n, a, T} + O(\sqrt{n} \log n),$$

where

$$\lambda_{-n, a, T} = \sum_{\rho_a \in \mathbb{Z}_Q; |\text{Im}(\rho_a)| \leq T}^* \left(1 - \left(1 - \frac{1}{\rho_a} \right)^n \right),$$

with $T = \sqrt{n} + \varepsilon_n$. For $a = 1/2, 1/3, 1/4, 1/6$, under the RH, since $\left| 1 - \frac{1}{\rho_a} \right| = 1$ and using formula of $N_Q(T)$ given in Section 1.2, we obtain $\lambda_{n, a, T} = O(T \log T + 1)$. Therefore, Equation (2.5) follows from that $\lambda_{-n, a, \sqrt{n}} = \lambda_{-n, a, \sqrt{n}} = O(\sqrt{n} \log n)$.

Remark Since $2Q(s, a) := Z(s, a) + P(s, a)$, from Corollary 2.3 below and [6, Equation (1.18)], we obtain

$$\gamma_Q(n) = \frac{1}{2} \left(\delta_n(a) + \frac{(-1)^n}{n!} (l_n(a) + l_n(1 - a)) \right),$$

where $\delta_n(a) = \frac{|\log a|^n}{an!} + O(1)$ and $l_n(a)$ are the coefficients in the expansion of $\text{Li}_s(e^{2\pi ia})$ at $s = 1$; for $a \notin \mathbb{Z}$, one has

$$\text{Li}_s(e^{2\pi ia}) = \sum_{n=0}^{\infty} \frac{(-1)^n}{n!} l_n(a) (s - 1)^n.$$

2.4 Proof of Theorem 1.4

To show Theorem 1.4, we quote the following lemmas from [2, 3].

Lemma 2.1 [3, Theorem 1] *We set*

$$(s - 1)\zeta(s, a) = 1 + \sum_{n=0}^{\infty} \gamma_n(a)(s - 1)^{n+1}, \quad 0 < a \leq 1.$$

¹These properties are well known for the Riemann zeta function. The proof for the function $Q(s, a)$ is exactly the same since the Riemann–von Mangoldt formula holds for $Q(s, a)$ (see [14, Proposition 2.5] or [18, p. 217]).

Then it holds that

$$\gamma_n(a) = \frac{(-1)^n}{n!} \lim_{m \rightarrow \infty} \left(\sum_{k=0}^m \frac{\log^n(k+a)}{k+a} - \frac{\log^{n+1}(m+a)}{n+1} \right).$$

Lemma 2.2 [2, Equation (26)] *Let $0 < a \leq 1$, and let n be a nonnegative integer. Then one has*

$$\begin{aligned} \zeta^{(n)}(0, a) &= \left(\frac{1}{2} - a \right) |\log a|^n - n! + n! a \sum_{m=n}^{\infty} \frac{|\log a|^m}{m!} \\ &\quad + (-1)^n n \int_0^{\infty} \frac{\varphi(x) \log^{n-1}(x+a)}{(x+a)^2} dx \\ &\quad - (-1)^n n(n-1) \int_0^{\infty} \frac{\varphi(x) \log^{n-2}(x+a)}{(x+a)^2} dx, \end{aligned}$$

where $\varphi(x) = \int_0^x (y - [y] - 1/2) dy$ is periodic with period 1 and satisfies $2\varphi(x) = x(x-1)$ if $0 \leq x \leq 1$.

By using the lemmas above, we immediately obtain the following.

Corollary 2.3 *When $a > 0$ is sufficiently small,*

$$\begin{aligned} (s-1)Z(s, a) &= 2 + \sum_{n=0}^{\infty} \delta_n(a)(s-1)^{n+1}, \quad \delta_n(a) = \frac{|\log a|^n}{an!} + O(1), \\ Z(s, a) &= \sum_{n=1}^{\infty} \varepsilon_n(a)s^n, \quad \varepsilon_n(a) = O(|\log a|^n). \end{aligned}$$

Proof The first formula and estimation are easily proved by Lemma 2.1 (see also [3, Theorem 2]). For the first integral in Lemma 2.2, one has

$$\begin{aligned} \int_0^1 \frac{\varphi(x) \log^{n-1}(x+a)}{(x+a)^2} dx &\ll \int_0^1 \frac{\log^{n-1}(x+a)}{x+a} dx = O(|\log a|^n), \\ \int_1^{\infty} \frac{\varphi(x) \log^{n-1}(x+a)}{(x+a)^2} dx &\ll \int_1^{\infty} \frac{\log^{n-1}(x+a)}{(x+a)^2} dx = O(1) \end{aligned}$$

from $x < x+a$ when $x, a > 0$. In addition, we have

$$a \sum_{m=n}^{\infty} \frac{|\log a|^m}{m!} \leq a \sum_{m=0}^{\infty} \frac{|\log a|^m}{m!} = ae^{|\log a|} = ae^{-\log a} = 1, \quad 0 < a < 1/2.$$

Hence, we obtain

$$\zeta(s, a) = \sum_{n=0}^{\infty} \frac{\zeta^{(n)}(0, a)}{n!} s^n, \quad \zeta^{(n)}(0, a) = O(|\log a|^n).$$

Therefore, we have $\varepsilon_n(a) = O(|\log a|^n)$ and the second formula in this corollary by the definition of $Z(s, a)$ and $Z(0, a) = \zeta(0, a) + \zeta(0, 1-a) = 0$ (see [15, Equation (4.11)]). ■

Proof of Theorem 1.4 Recall the functional equation

$$Z(1 - s, a) = \Gamma_{\cos}(s)P(s, a), \quad \Gamma_{\cos}(s) := \frac{2\Gamma(s)}{(2\pi)^s} \cos\left(\frac{\pi s}{2}\right)$$

(see [15, Lemma 4.11]). By using $\Gamma_{\cos}(s)\Gamma_{\cos}(1 - s) = 1$, we have

$$2Q(s, a) = Z(s, a) + P(s, a) = Z(s, a) + \Gamma_{\cos}(1 - s)Z(1 - s, a).$$

Let $|s - 1|$ be sufficiently small. Then, by $\lim_{s \rightarrow 1}(s - 1)Q(s, a) = 1$, the equation above, and the definitions of $Q(s, a)$ and $\xi_Q(s, a)$, we have

$$\begin{aligned} \frac{d^l}{ds^l} [s^{l-1} \log \xi_Q(s, a)]_{s=1} &= \frac{d^l}{ds^l} [s^{l-1} \log((s - 1)Q(s, a)) + s^{l-1} \log(sp^{-s/2}\Gamma(s/2))]_{s=1} \\ &= \frac{d^l}{ds^l} \left[s^{l-1} \log\left(\frac{s-1}{2} (Z(s, a) + \Gamma_{\cos}(1-s)Z(1-s, a))\right) \right]_{s=1} + O_l(1) \\ &= \frac{d^l}{ds^l} \left[s^{l-1} \log\left(1 + \sum_{n=0}^{\infty} (\delta'_n(a) + \varepsilon'_n(a))(s-1)^{n+1}\right) \right]_{s=1} + O_l(1), \end{aligned}$$

where $\delta'_n(a)$ and $\varepsilon'_n(a)$ are defined by

$$\delta'_n(a) := \frac{\delta_n(a)}{2}, \quad (s - 1)\Gamma_{\cos}(1 - s)Z(1 - s, a) = 2 \sum_{n=0}^{\infty} \varepsilon'_n(a)(s - 1)^{n+1}.$$

Clearly, the second estimation in Corollary 2.3 implies

$$Z(1 - s, a) = \sum_{n=1}^{\infty} \varepsilon_n(a)(1 - s)^n, \quad \varepsilon_n(a) = O(|\log a|^n).$$

Thus, we can see that $\varepsilon'_n(a) = O(|\log a|^{n+1})$ from $\lim_{s \rightarrow 1}(s - 1)\Gamma_{\cos}(1 - s) = -2$ and the fact that the function $(s - 1)\Gamma_{\cos}(1 - s)$ does not depend on a . Put $\eta_n(a) := \delta'_n(a) + \varepsilon'_n(a)$. Then, for $n \geq 0$, we have

$$(2.6) \quad \eta_n(a) = \frac{1}{n!} \frac{|\log a|^n}{2a} + O(|\log a|^{n+1}), \quad a \rightarrow +0$$

by Corollary 2.3. By virtue of

$$\begin{aligned} (a_0x + a_1x^2 + a_2x^3 + \dots)^m &= a_0^m x^m + \binom{m}{1} a_0^{m-1} a_1 x^{m+1} + \dots \\ (a_0x + a_1x^2 + a_2x^3 + \dots)^{m-1} &= a_0^{m-1} x^{m-1} + \binom{m-1}{1} a_0^{m-2} a_1 x^m + \dots \\ &\vdots \\ (a_0x + a_1x^2 + a_2x^3 + \dots)^1 &= \dots + a_m x^m + \dots, \end{aligned}$$

where $m \in \mathbb{N}$ and $a_m, x \in \mathbb{C}$, the coefficient of $(s - 1)^l$ in the function

$$f(s, a) := \sum_{m=1}^{\infty} \frac{(-1)^{m+1}}{m} \left(\sum_{n=0}^{\infty} \eta_n(a)(s - 1)^{n+1} \right)^m$$

is expressed as

$$(2.7) \quad \frac{(-1)^{l+1}}{l}(\eta_0(a))^l + \frac{(-1)^l}{l-1} \binom{l-1}{1} \eta_0(a)^{l-2} \eta_1(a) + \dots + \frac{(-1)^{l+1}}{1} \eta_{l-1}(a).$$

Note that the function above is estimated by

$$(2.8) \quad \frac{(-1)^{l+1}}{l}(\eta_0(a))^l + O_l(\eta_0(a)^{l-2} \eta_1(a)) = \frac{(-1)^{l+1}}{l} (2a)^{-l} + O_l(a^{1-l} |\log a|)$$

from (2.6) when $a \rightarrow +0$. We can find that

$$(s-1) \left(Z(s, a) + \Gamma_{\cos}(1-s) Z(1-s, a) \right) = 1 + \sum_{n=0}^{\infty} \eta_n(a) (s-1)^{n+1}$$

is analytic when $|s-1| < 1$ form the poles of $Z(s, a)$ and $\Gamma_{\cos}(1-s)$. So we can choose $|s-1| > 0$ such that

$$\sum_{n=0}^{\infty} |\eta_n(a)| |s-1|^{n+1} < \frac{1}{2}.$$

Then, from (2.7), the Leibniz product rule, the definition of $\eta_n(a)$, and the Taylor expansion of $\log(1+x)$ with $|x| < 1$, one has

$$\begin{aligned} & \frac{d^l}{ds^l} \left[s^{l-1} \log \xi_Q(s, a) \right]_{s=1} = \frac{d^l}{ds^l} \left[s^{l-1} f(s, a) \right]_{s=1} + O_l(1) \\ (b) \quad & = \binom{l}{l} \frac{(-1)^{l+1}}{l} l! (\eta_0(a))^l + O_l(\eta_0(a)^{l-2} \eta_1(a)) \\ (h) \quad & + \binom{l}{l-1} (l-1) \frac{(-1)^l}{l-1} (l-1)! (\eta_0(a))^{l-1} + O_l(\eta_0(a)^{l-3} \eta_1(a)) \\ (\sharp) \quad & + \dots + \binom{l}{1} (l-1)! \frac{(-1)^{l+1}}{1} (\eta_0(a))^1 + O_l(1). \end{aligned}$$

Note that (b) comes from $f^{(l)}(s, a)$, (h) is deduced by $f^{(l-1)}(s, a)$, and (sharp) derives from $f^{(1)}(s, a)$, $f^{(0)}(s, a)$, and $O_l(1)$ in the left-hand side of the formula above. Therefore, by (2.8), we obtain

$$\begin{aligned} \frac{d^l}{ds^l} \left[s^{l-1} \log \xi_Q(s, a) \right]_{s=1} & = (-1)^{l+1} (l-1)! (\eta_0(a))^l + O_l(\eta_0(a)^{l-2} \eta_1(a)) \\ & = (-1)^{l+1} \frac{(l-1)!}{(2a)^l} + O_l(a^{1-l} |\log a|), \end{aligned}$$

which implies Theorem 1.4. ■

At the end of the paper, we give numerical computation for $\lambda_{n,a}$ by Mathematica 13.0. Let

$$\lambda_{n,a}^{[k]} := \frac{1}{(n-1)!} \frac{d^n}{ds^n} \left[s^{n-1} \log \xi_Q(s, a) \right]_{s=1-10^{-k}}, \quad \lambda_{n,a}^* := \frac{(-1)^{n+1}}{(2a)^n}.$$

Then, we have the following.

For $n = 1$, we have

$$\begin{array}{lll}
 a := 2^{-17} & \lambda_{1,a}^{[10]} = 65, 537\dots & \lambda_{1,a}^{[10]}/\lambda_{1,a}^* = 1.00001\dots \\
 a := 2^{-18} & \lambda_{1,a}^{[10]} = 131, 074\dots & \lambda_{1,a}^{[10]}/\lambda_{1,a}^* = 1.00002\dots \\
 a := 2^{-19} & \lambda_{1,a}^{[10]} = 262, 151\dots & \lambda_{1,a}^{[10]}/\lambda_{1,a}^* = 1.00003\dots \\
 a := 2^{-17} & \lambda_{1,a}^{[11]} = 65, 536.6\dots & \lambda_{1,a}^{[11]}/\lambda_{1,a}^* = 1.00001\dots \\
 a := 2^{-18} & \lambda_{1,a}^{[11]} = 131, 073\dots & \lambda_{1,a}^{[11]}/\lambda_{1,a}^* = 1.00001\dots \\
 a := 2^{-19} & \lambda_{1,a}^{[11]} = 262, 145\dots & \lambda_{1,a}^{[11]}/\lambda_{1,a}^* = 1.00000\dots \\
 a := 2^{-17} & \lambda_{1,a}^{[12]} = 655, 365\dots & \lambda_{1,a}^{[12]}/\lambda_{1,a}^* = 1.00001\dots \\
 a := 2^{-18} & \lambda_{1,a}^{[12]} = 131, 073\dots & \lambda_{1,a}^{[12]}/\lambda_{1,a}^* = 1.00000\dots \\
 a := 2^{-19} & \lambda_{1,a}^{[12]} = 262, 145\dots & \lambda_{1,a}^{[12]}/\lambda_{1,a}^* = 1.00000\dots
 \end{array}$$

For $n = 2$, we have

$$\begin{array}{lll}
 a := 2^{-17} & \lambda_{2,a}^{[10]} = -4.29352\dots \times 10^9 & \lambda_{2,a}^{[10]}/\lambda_{2,a}^* = 0.999663\dots \\
 a := 2^{-18} & \lambda_{2,a}^{[10]} = -1.7177\dots \times 10^{10} & \lambda_{2,a}^{[10]}/\lambda_{2,a}^* = 0.999836\dots \\
 a := 2^{-19} & \lambda_{2,a}^{[10]} = -6.87162\dots \times 10^{10} & \lambda_{2,a}^{[10]}/\lambda_{2,a}^* = 0.999952\dots \\
 a := 2^{-17} & \lambda_{2,a}^{[11]} = -4.29478\dots \times 10^9 & \lambda_{2,a}^{[11]}/\lambda_{2,a}^* = 0.999956\dots \\
 a := 2^{-18} & \lambda_{2,a}^{[11]} = -1.71753\dots \times 10^{10} & \lambda_{2,a}^{[11]}/\lambda_{2,a}^* = 0.999736\dots \\
 a := 2^{-19} & \lambda_{2,a}^{[11]} = -6.87149\dots \times 10^{10} & \lambda_{2,a}^{[11]}/\lambda_{2,a}^* = 0.999933\dots \\
 a := 2^{-17} & \lambda_{2,a}^{[12]} = -4.29477\dots \times 10^9 & \lambda_{2,a}^{[12]}/\lambda_{2,a}^* = 0.999955\dots \\
 a := 2^{-18} & \lambda_{2,a}^{[12]} = -1.6911\dots \times 10^{10} & \lambda_{2,a}^{[12]}/\lambda_{2,a}^* = 0.984353\dots \\
 a := 2^{-19} & \lambda_{2,a}^{[12]} = -6.87187\dots \times 10^{10} & \lambda_{2,a}^{[12]}/\lambda_{2,a}^* = 0.999989\dots
 \end{array}$$

Acknowledgements The authors want to thank the anonymous referees for their many insightful comments and suggestions.

References

- [1] T. M. Apostol, *Introduction to analytic number theory*, Undergraduate Texts in Mathematics, Springer, New York, 1976.
- [2] T. M. Apostol, *Formulas for higher derivatives of the Riemann zeta function*. Math. Comput. 44(1985), no. 169, 223–232.
- [3] B. C. Berndt, *On the Hurwitz zeta-function*. Rocky Mountain J. Math. 2(1972), no. 1, 151–157.
- [4] E. Bombieri and J. Lagarias, *Complements to Li’s criterion for the Riemann hypothesis*. J. Number Theory 77(1999), 274–287.
- [5] F. Brown, *Li’s criterion and zero-free regions of L -functions*. J. Number Theory 111(2005), 1–32.
- [6] M. W. Coffey, *Series representations for the Stieltjes constants*. Rocky Mountain J. Math. 44(2014), no. 2, 443–477.
- [7] P. Flajolet and L. Vepstas, *On differences of zeta values*. J. Comput. Appl. Math. 220(2008), nos. 1–2, 58–73.
- [8] J. C. Lagarias, *Li coefficients for automorphic L-functions*. Ann. Inst. Fourier (Grenoble) 57(2007), 1689–1740.
- [9] A. Laurinćikas and R. Garunkštis, *The Lerch zeta-function*, Kluwer Academic Publishers, Dordrecht, 2002.
- [10] X.-J. Li, *The positivity of a sequence of numbers and the Riemann hypothesis*. J. Number Theory 65(1997), 325–333.
- [11] K. Mazhouda and L. Smajlović, *Evaluation of the Li coefficients on function fields and applications*. Eur. J. Math. 5(2019), no. 2, 540–550.

- [12] K. Mazhouda and B. Sodaïgui, *The Li–Sekatskii coefficients for the Selberg class*. Int. J. Math. 33(2022), no. 12, Article no. 2250075, 23 pp.
- [13] T. Nakamura, *The functional equation and zeros on the critical line of the quadrilateral zeta function*. J. Number Theory 233(2022), 432–455.
- [14] T. Nakamura, *On Lerch’s formula and zeros of the quadrilateral zeta function*. Preprint, 2022, arXiv:2001.01981.
- [15] T. Nakamura, *On zeros of bilateral Hurwitz and periodic zeta and zeta star functions*. Rocky Mountain J. Math. 53(2023), no. 1, 157–176.
- [16] S. Omar and K. Mazhouda, *The Li criterion and the Riemann hypothesis for the Selberg class II*. J. Number Theory 130(2010), no. 4, 1109–1114.
- [17] L. Smajlović, *On Li’s criterion for the Riemann hypothesis for the Selberg class*. J. Number Theory 130(2010), no. 4, 828–851.
- [18] E. C. Titchmarsh, *Theory of the Riemann zeta-function*. 2nd ed., Clarendon Press, Oxford, 1986.
- [19] A. Voros, *Sharpenings of Li’s criterion for the Riemann hypothesis*. Math. Phys. Anal. Geom. 9(2006), 53–63.

Department of Mathematics, University of Prishtina, Mother Theresa, No. 5, 10000 Prishtina, Kosovo
e-mail: kajtaz.bllaca@uni-pr.edu

Higher Institute of Applied Sciences and Technology, University of Sousse, Sousse 4003, Tunisia INSA Hauts-De-France, University Polytechnique Hauts-De-France, FR CNRS 2037, CERAMATHS, F-59313 Valenciennes, France
e-mail: kamel.mazhouda@fsm.rnu.tn

Department of Liberal Arts, Faculty of Science and Technology, Tokyo University of Science, 2641 Yamazaki, Noda-shi, Chiba-ken 278-8510, Japan
e-mail: nakamuratakashi@rs.tus.ac.jp