

## ENUMERATION OF GROUPS IN SOME SPECIAL VARIETIES OF $A$ -GROUPS

ARUSHI  and GEETHA VENKATARAMAN  

(Received 3 March 2024; accepted 29 April 2024; first published online 27 August 2024)

### Abstract

We find an upper bound for the number of groups of order  $n$  up to isomorphism in the variety  $\mathfrak{S} = \mathfrak{A}_p\mathfrak{A}_q\mathfrak{A}_r$ , where  $p$ ,  $q$  and  $r$  are distinct primes. We also find a bound on the orders and on the number of conjugacy classes of subgroups that are maximal amongst the subgroups of the general linear group that are also in the variety  $\mathfrak{A}_q\mathfrak{A}_r$ .

2020 *Mathematics subject classification*: primary 20E10; secondary 20B15, 20B35, 20D10, 20E45, 20H30.

*Keywords and phrases*: group enumeration, variety of groups, general linear group, symmetric group, transitive subgroup, primitive subgroup, conjugacy class.

### 1. Introduction

A group is an  $A$ -group if its nilpotent subgroups are abelian. For any class of groups  $\mathfrak{B}$ , we denote the number of groups of order  $n$  up to isomorphism by  $f_{\mathfrak{B}}(n)$ . Computing  $f(n)$  becomes harder as  $n$  gets bigger. Thus, in the area of group enumerations, we attempt to approximate  $f(n)$ . When counting is restricted to the class of abelian groups,  $A$ -groups or groups in general, the asymptotic behaviour of  $f(n)$  varies significantly. Let  $f_{A,\text{sol}}(n)$  be the number of isomorphism classes of soluble  $A$ -groups of order  $n$ . Dickenson [2] showed that  $f_{A,\text{sol}}(n) \leq n^{c \log n}$  for some constant  $c$ . McIver and Neumann [7] showed that the number of nonisomorphic  $A$ -groups of order  $n$  is at most  $n^{\lambda+1}$ , where  $\lambda$  is the number of prime divisors of  $n$  including multiplicities. In the same paper, they stated the following conjecture based on a result of Higman [4] and Sims [12] on  $p$ -group enumerations.

**CONJECTURE 1.1.** Let  $f(n)$  be the number of (isomorphism classes of groups of) order  $n$ . Then  $f(n) \leq n^{(2/27+\epsilon)\lambda^2}$ , where  $\epsilon \rightarrow 0$  as  $\lambda \rightarrow \infty$ .

In 1993, Pyber [9] proved a powerful version of Conjecture 1.1: the number of groups of order  $n$  with specified Sylow subgroups is at most  $n^{75\mu+16}$ , where  $\mu$  is the



largest integer such that  $p^\mu$  divides  $n$  for some prime  $p$ . From the results of Higman and Sims, and Pyber,  $f(n) \leq n^{2\mu^2/27+O(\mu^{5/3})}$ . In [13], it was shown that  $f_{A,sol}(n) \leq n^{7\mu+6}$ .

The variety  $\mathfrak{A}_u\mathfrak{A}_v$  consists of all groups  $G$  with an abelian normal subgroup  $N$  of exponent dividing  $u$  such that  $G/N$  is abelian of exponent dividing  $v$ . (For more on varieties, see [8].) Let  $p, q$  and  $r$  be distinct primes. In this paper, we find a bound for  $f_\mathfrak{S}(n)$ , where  $\mathfrak{S} = \mathfrak{A}_p\mathfrak{A}_q\mathfrak{A}_r$ , and  $f_\mathfrak{S}(n)$  counts the groups in  $\mathfrak{S}$  of order  $n$  up to isomorphism. The idea behind studying the variety  $\mathfrak{S}$  is that enumerating within the varieties of  $A$ -groups might yield a better upper bound for the enumeration function for  $A$ -groups. The ‘best’ bounds for  $A$ -groups, or even soluble  $A$ -groups, still lack the correct leading term. It is believed that a correct leading term for the upper bound of  $A$ -groups would lead to the right error term for the enumeration of groups in general.

A few smaller varieties of  $A$ -groups have already been studied in [1, Ch. 18]. The class of  $A$ -groups for which the ‘best’ bounds exist was obtained by enumerating in such small varieties of  $A$ -groups, but this did not narrow the difference between the upper and lower bounds for  $f_{A,sol}(n)$  because these groups did not contribute a large enough collection of  $A$ -groups. Hence, a good lower bound could not be reached. To reduce the difference, we enumerate in the larger variety  $\mathfrak{S}$  of  $A$ -groups.

Throughout the paper,  $p, q, r$  and  $t$  are distinct primes. We assume that  $s$  is a power of  $t$ . We take logarithms to the base 2, unless stated otherwise, and follow the convention that  $0 \in \mathbb{N}$ . We use  $C_m$  to denote a cyclic group of order  $m$  for any positive integer  $m$ . Let  $O_{p'}(G)$  denote the largest normal  $p'$ -subgroup of  $G$ . The techniques we use are similar to those in [1, 9, 13].

The main result proved in this paper is the following theorem.

**THEOREM 1.2.** *Let  $n = p^\alpha q^\beta r^\gamma$ , where  $\alpha, \beta, \gamma \in \mathbb{N}$ . Then,*

$$f_\mathfrak{S}(n) \leq p^{6\alpha^2} 2^{\alpha-1+(23/6)\alpha \log \alpha + \alpha \log 6} (6^{1/2})^{(\alpha+\gamma)\beta + (\alpha+\beta)\gamma + \alpha(\alpha-1)/2} n^{\beta+\gamma}.$$

To prove Theorem 1.2, we prove a bound on the number of conjugacy classes of subgroups that are maximal amongst subgroups of  $GL(\alpha, s)$  and that are in the variety  $\mathfrak{A}_q\mathfrak{A}_r$  or  $\mathfrak{A}_r$ . We also prove results about the order of primitive subgroups of  $S_n$  that are in the variety  $\mathfrak{A}_q\mathfrak{A}_r$ , and show that they form a single conjugacy class. These results are stated below.

**THEOREM 1.3.** *Let  $q$  and  $r$  be distinct primes. Let  $G$  be a primitive subgroup of  $S_n$  that is in  $\mathfrak{A}_q\mathfrak{A}_r$  and let  $|G| = q^\beta r^\gamma$ , where  $\beta, \gamma \in \mathbb{N}$ . Let  $M$  be a minimal normal subgroup of  $G$ .*

- (i) *If  $\beta = 0$ , then  $|M|$  is a power of  $r$  and  $|G| = n = r$  with  $G \cong C_r$ .*
- (ii) *If  $\beta \geq 1$ , then  $|M| = q^\beta = n$  with  $\beta = \text{order } q \text{ mod } r$ . Further,  $G \cong M \rtimes C_r$  and  $|G| = nr < n^2$ .*
- (iii) *If  $\gamma = 0$ , then  $|M|$  is a power of  $q$  and  $|G| = n = q$  with  $G \cong C_q$ .*

**THEOREM 1.4.** *The primitive subgroups of  $S_n$  that are in  $\mathfrak{A}_q\mathfrak{A}_r$  and of order  $q^\beta r^\gamma$ , where  $\beta, \gamma \in \mathbb{N}$ , form a single conjugacy class.*

**THEOREM 1.5.** *There exist constants  $b$  and  $c$  such that the number of conjugacy classes of subgroups that are maximal amongst the subgroups of  $\text{GL}(\alpha, s)$  that are in  $\mathfrak{A}_q\mathfrak{A}_r$  is at most*

$$2^{(b+c)(\alpha^2/\sqrt{\log \alpha})+(5/6)\alpha \log \alpha + \alpha(1+\log 6)} s^{(3+c)\alpha^2},$$

where  $t, q$  and  $r$  are distinct primes,  $s$  is a power of  $t$ , and  $\alpha > 1$ .

Section 2 investigates primitive subgroups of  $S_n$  that are in  $\mathfrak{A}_r$  or  $\mathfrak{A}_q\mathfrak{A}_r$ . Sections 3 and 4 deal with subgroups of the general linear group. Theorem 1.2 is proved in Section 5.

### 2. Primitive subgroups of $S_n$ that are in $\mathfrak{A}_r$ or $\mathfrak{A}_q\mathfrak{A}_r$

In this section, we prove results that give us the structure of the primitive subgroups of  $S_n$  that are in  $\mathfrak{A}_r$  or  $\mathfrak{A}_q\mathfrak{A}_r$ . We also show that such subgroups form a single conjugacy class. Both Theorems 1.3 and 1.4 are proved in this section.

Theorem 1.3 provides the order of a primitive subgroup of  $S_n$  that is in the variety  $\mathfrak{A}_q\mathfrak{A}_r$ . By [13, Proposition 2.1], if  $G$  is a soluble  $A$ -subgroup of  $S_n$ , then  $|G| \leq (6^{1/2})^{n-1}$ . Indeed, this bound is determined primarily by considering primitive soluble  $A$ -subgroups of  $S_n$ . This bound would clearly hold for any subgroup of  $S_n$  that is in the variety  $\mathfrak{A}_q\mathfrak{A}_r$ . However, we show that when the subgroup is primitive and in the variety  $\mathfrak{A}_q\mathfrak{A}_r$ , we can do better.

**LEMMA 2.1.**  *$S_n$  has a primitive subgroup in  $\mathfrak{A}_r$  if and only if  $n = r$ . In this case, any primitive subgroup  $G$  that is in  $\mathfrak{A}_r$  will be cyclic of order  $r$ . All primitive subgroups of  $S_n$  that are in  $\mathfrak{A}_r$  form a single conjugacy class.*

**PROOF.** Let  $G$  be a primitive subgroup of  $S_n$  that is in  $\mathfrak{A}_r$ . Since  $G$  is soluble,  $M$  is an elementary abelian  $r$ -subgroup. By the O’Nan–Scott theorem [10],  $|M| = n = |G|$ , so  $G = M \cong C_r$  and  $n = r$ . Conversely, any transitive subgroup  $G$  of  $S_r$  is primitive [15, Theorem 8.3]. Since  $n$  is prime, any subgroup of order  $n$  in  $S_n$  will be generated by an  $n$ -cycle. Further, any two  $n$ -cycles are conjugate in  $S_n$ . Thus, the primitive subgroups of  $S_n$  that are also in  $\mathfrak{A}_r$  form a single conjugacy class.  $\square$

**PROOF OF THEOREM 1.3.** Let  $G$  be a subgroup of  $S_\Omega$ , where  $|\Omega| = n$ , and let  $G \in \mathfrak{A}_q\mathfrak{A}_r$ . Then  $G = Q \rtimes R$ , where  $Q$  is an elementary abelian Sylow  $q$ -subgroup,  $R$  is an elementary abelian Sylow  $r$ -subgroup and  $|G| = q^\beta r^\gamma$ , with  $\beta, \gamma \in \mathbb{N}$ . Let  $M$  be a minimal normal subgroup of  $G$ . Then  $M$  is an elementary abelian  $u$ -group. Clearly,  $|M| = u^k$  for some  $k > 1$  and for some prime  $u \in \{q, r\}$ .

Now  $F(G)$ , the Fitting subgroup of  $G$ , is an abelian normal subgroup of  $G$  and so, by the O’Nan–Scott theorem,  $n = |M| = |F(G)|$ . However,  $M \leq F(G)$ , therefore,  $M = F(G)$  and  $n = u^k$ . If  $\beta \geq 1$ , then  $Q \leq F(G)$  and we have  $n = q^\beta = u^k$  and  $M = F(G) = Q$ . Let  $H = G_\alpha$  be the stabiliser of an  $\alpha \in \Omega$ . By [1, Proposition 6.13],  $G$  is a semidirect product of  $M$  by  $H$  and  $H$  acts faithfully by conjugation on  $M$ . By Maschke’s theorem,

$M$  is completely reducible. However,  $M$  is a minimal normal subgroup of  $G$ , so  $M$  is a nontrivial irreducible  $\mathbb{F}_q H$ -module and  $H$  is an abelian group acting faithfully on  $M$ . By [14, Corollary 4.1],  $H \cong C_r$  and  $\beta = \dim M = \text{order } q \text{ mod } r$  and the result follows. If  $\gamma = 0$  or  $\beta = 0$ , then  $|G|$  is a power of  $u$ , where  $u \in \{q, r\}$ . Thus,  $G$  is a primitive subgroup that is also in  $\mathfrak{A}_u$  and the result follows by Lemma 2.1.  $\square$

It is clear from these results that if  $S_n$  has a primitive subgroup  $G$  of order  $q^\beta r^\gamma$  in  $\mathfrak{A}_q \mathfrak{A}_r$ , then  $n$  must be  $r$  or  $q$  and  $G$  is cyclic with  $|G| = n$ , or  $n = q^\beta$  and  $G$  is a semi-direct product of an elementary abelian  $q$ -group of order  $q^\beta$  by a cyclic group of order  $r$ . The limits imposed on  $n$  and the structure of such primitive subgroups gives the next result.

**PROOF OF THEOREM 1.4.** Let  $G$  be a primitive subgroup of  $S_\Omega$  that is in  $\mathfrak{A}_q \mathfrak{A}_r$ , where  $|\Omega| = n$ , and let  $|G| = q^\beta r^\gamma$ . Let  $M$  be a minimal normal subgroup of  $G$ . As seen in the proof of Theorem 1.3,  $M = F(G)$  and  $n = |M|$  is either a power of  $q$  or  $r$ . If  $\gamma = 0$  or  $\beta = 0$ , then  $|G|$  is a power of  $u$ , where  $u \in \{q, r\}$ . Thus,  $G$  is a primitive subgroup that is also in  $\mathfrak{A}_u$  and the result follows by Lemma 2.1.

We know the structure of  $G$  when  $\beta \geq 1$  from the proof of Theorem 1.3. Hence,  $H$  can be regarded as a soluble  $r$ -subgroup of  $GL(\beta, q)$  and it is not difficult to show that the conjugacy class of  $G$  in  $S_n$  is determined by the conjugacy class of  $H$  in  $GL(\beta, q)$ . Let  $S$  be a Singer subgroup of  $GL(\beta, q)$ , so that  $|S| = q^\beta - 1$ . Now,  $|H| = r$  and  $r$  divides  $|S|$ . Further,  $\gcd(|GL(\beta, q)|/|S|, r) = 1$  as  $\beta$  is the least positive integer such that  $r \mid q^\beta - 1$ . From [3, Theorem 2.11],  $H^x \leq S$  for some  $x \in GL(\beta, q)$ . Since all Singer subgroups are conjugate in  $GL(\beta, q)$ , the result follows.  $\square$

### 3. Subgroups of $GL(\alpha, s)$ that are in $\mathfrak{A}_r$

In this section, we prove results that give us a bound on the number of conjugacy classes of the subgroups that are maximal amongst subgroups of  $GL(\alpha, s)$  that are in  $\mathfrak{A}_r$ . The limits on the structure of such groups ensures that if they exist, they form a single conjugacy class.

**LEMMA 3.1.** *The number of conjugacy classes of irreducible subgroups of  $GL(\alpha, s)$  that are also in  $\mathfrak{A}_r$  is at most 1.*

**PROOF.** Let  $G$  be a nontrivial irreducible subgroup of  $GL(\alpha, s)$  that is also in  $\mathfrak{A}_r$ . Then  $G$  is an elementary abelian  $r$ -group of order  $r^\gamma$ , say, where  $\gamma \in \mathbb{N}$ . Since  $G$  is a faithful abelian irreducible subgroup of  $GL(\alpha, s)$  whose order is coprime to  $s$ , it follows that  $G$  is cyclic [14, Lemma 4.2]. Thus,  $|G| = r$  and  $\alpha = d$ , where  $d = \text{order } s \text{ mod } r$ . From [11, Theorem 2.3.3], the irreducible cyclic subgroups of order  $r$  in  $GL(\alpha, s)$  lie in a single conjugacy class.  $\square$

**PROPOSITION 3.2.** *The number of conjugacy classes of subgroups that are maximal amongst subgroups of  $GL(\alpha, s)$  that are also in  $\mathfrak{A}_r$  is at most 1.*

**PROOF.** Let  $G$  be maximal amongst subgroups of  $\text{GL}(\alpha, s)$  that are also in  $\mathfrak{A}_r$ . Since  $\text{char}(\mathbb{F}_p) = t \nmid |G|$ , by Maschke's theorem, we can find groups  $G_i$  such that  $G \leq G_1 \times G_2 \times \cdots \times G_k = \hat{G} \leq \text{GL}(\alpha, s)$ , where for each  $i$ , the group  $G_i$  is a (maximal) irreducible subgroup of  $\text{GL}(\alpha_i, s)$  that is also in  $\mathfrak{A}_r$ . Further,  $\alpha = \alpha_1 + \cdots + \alpha_k$ . Clearly,  $G_i \cong C_r$  and  $\alpha_i = d = \text{order } s \text{ mod } r$  for each  $i$ . Thus, we must have  $\alpha = dk$  and by the maximality of  $G$ , we have  $G = \hat{G}$ . Further, the conjugacy classes of  $G_i$  in  $\text{GL}(\alpha_i, s)$  determine the conjugacy class of  $G$  in  $\text{GL}(\alpha, s)$ .

So if  $d$  does not divide  $\alpha$ , then  $\text{GL}(\alpha, s)$  cannot have an elementary abelian  $r$ -subgroup. If  $d \mid \alpha$ , then any  $G$  that is maximal amongst subgroups of  $\text{GL}(\alpha, s)$  that are also in  $\mathfrak{A}_r$  must have order  $r^k$ , where  $k = \alpha/d$ . By Lemma 3.1, all such groups form a single conjugacy class. □

#### 4. Subgroups of $\text{GL}(\alpha, s)$ that are also in $\mathfrak{A}_q \mathfrak{A}_r$

We prove results that give a bound on the order of subgroups of  $\text{GL}(\alpha, s)$  that are in  $\mathfrak{A}_q \mathfrak{A}_r$  and also a bound for the number of conjugacy classes of subgroups that are maximal amongst subgroups of  $\text{GL}(\alpha, s)$  that are in  $\mathfrak{A}_q \mathfrak{A}_r$ . Theorem 1.5 is proved in this section.

**PROPOSITION 4.1.** *Let  $G$  be a subgroup of  $\text{GL}(\alpha, s)$  that is in  $\mathfrak{A}_q \mathfrak{A}_r$ .*

- (i) *Let  $m = |F(G)|$ . If  $G$  is primitive, then  $|G| \leq cm$ , where  $c = \text{order } s \text{ mod } m$  and  $c \mid \alpha$ . Further,  $m$  is either  $r$  or  $q$  or  $qr$ .*
- (ii)  *$|G| \leq (6^{1/2})^{\alpha-1} d^\alpha$ , where  $d = \min\{qr, s\}$ .*

**PROOF.** Let  $V = (\mathbb{F}_s)^\alpha$ . Let  $G$  be a primitive subgroup of  $\text{GL}(\alpha, s)$  that is in  $\mathfrak{A}_q \mathfrak{A}_r$  and let  $|G| = q^\beta r^\gamma$ , where  $\beta$  and  $\gamma$  are natural numbers. If  $\beta = 0$  or  $\gamma = 0$ , then the result follows from Lemma 3.1. Assume that  $\beta$  and  $\gamma$  are at least 1. Let  $F = F(G)$  be the Fitting subgroup of  $G$ . Since  $G \in \mathfrak{A}_q \mathfrak{A}_r$ , it follows that  $F$  is abelian and  $|F| = q^\beta r^{\gamma_1} = m$ , where  $\gamma_1 \leq \gamma$ . By Clifford's theorem, since  $G$  is primitive,  $V = X_1 \oplus X_2 \oplus \cdots \oplus X_a$  as an  $F$ -module, where the  $X_i$  are conjugates of  $X$ , an irreducible  $\mathbb{F}_s F$ -submodule of  $V$ . Note that  $F$  acts faithfully on  $X$ .

Let  $E$  be the subalgebra generated by  $F$  in  $\text{End}(V)$ . The  $X_i$  are conjugates of  $X$ , so  $E$  acts faithfully and irreducibly on  $X$  and  $E$  is commutative. By [1, Proposition 8.2 and Theorem 8.3],  $E$  is a field. Thus,  $E \cong \mathbb{F}_{s^c}$  as an  $\mathbb{F}_s F$ -module, where  $c = \dim(X)$  and  $\alpha = ac$ . Note that  $F$  is an abelian group of order  $m$  acting faithfully and irreducibly on  $X$ . Consequently,  $F$  is cyclic and  $c$  is the least positive integer such that  $m \mid s^c - 1$ . Clearly,  $m = q$  or  $m = qr$  and so  $\beta = 1$ . It is not difficult to show that  $G$  acts on  $E$  by conjugation. Hence, there exists a homomorphism from  $G$  to  $\text{Gal}_{\mathbb{F}_s}(E)$ . Let  $N$  be the kernel of this map. Then  $N = C_G(E) \leq C_G(F) \leq F$ . However,  $F \leq N$ . Hence,  $F = N$ . So  $G/F \leq \text{Gal}_{\mathbb{F}_s}(E) \cong C_c$  and  $|G| \leq cm$ .

Let  $G$  be an irreducible imprimitive subgroup of  $\text{GL}(\alpha, s)$  that is also in  $\mathfrak{A}_q \mathfrak{A}_r$ . Then  $G \leq G_1 \text{ wr } G_2 \leq \text{GL}(\alpha, s)$ , where  $G_1$  is a primitive subgroup of  $\text{GL}(\alpha_1, s)$  that is in  $\mathfrak{A}_q \mathfrak{A}_r$ , and the group  $G_2$  can be regarded as a transitive subgroup of  $S_k$  that is in

$\mathfrak{A}_q \mathfrak{A}_r$ . Further,  $\alpha = \alpha_1 k$ . By the previous part,  $|G_1| \leq c' m'$ , where  $c' = \text{order } s \text{ mod } m'$  and  $m' = |F(G_1)|$  is either  $r$  or  $q$  or  $qr$ . Also  $c' \mid \alpha_1$ . By [13, Proposition 2.1],  $|G_2| \leq (6^{1/2})^{k-1}$ . Using  $c' \leq 2^{c'-1} \leq (6^{1/2})^{c'-1}$ , we see that  $|G| \leq (6^{1/2})^{\alpha-1} (m')^k$ . Since  $m' \mid p^{c'} - 1$ , we have  $(m')^k \leq d^\alpha$ , where  $d = \min\{qr, s\}$ .

Since  $t$  does not divide  $q$  or  $r$ , by Maschke's theorem, any subgroup  $G$  of  $\text{GL}(\alpha, s)$  that is in  $\mathfrak{A}_q \mathfrak{A}_r$  will be completely reducible. Thus,  $G \leq G_1 \times \dots \times G_k \leq \text{GL}(\alpha, s)$ , where the  $G_i$  are irreducible subgroups of  $\text{GL}(\alpha_i, s)$  that are in  $\mathfrak{A}_q \mathfrak{A}_r$  and  $\alpha = \alpha_1 + \dots + \alpha_k$ . Hence,  $|G| \leq (6^{1/2})^{\alpha-1} d^\alpha$ , where  $d = \min\{qr, s\}$ .  $\square$

**PROPOSITION 4.2.** *There exist constants  $b$  and  $c$  such that the number of conjugacy classes of subgroups that are maximal amongst irreducible subgroups of  $\text{GL}(\alpha, s)$  that are in  $\mathfrak{A}_q \mathfrak{A}_r$  is at most  $2^{(b+c)(\alpha^2/\sqrt{\log \alpha} + (5/6)\log \alpha + \log 6)s^{(3+c)\alpha^2}}$  provided  $\alpha > 1$ .*

**PROOF.** Let  $G$  be a subgroup of  $\text{GL}(\alpha, s)$  that is maximal amongst irreducible subgroups of  $\text{GL}(\alpha, s)$  that are in  $\mathfrak{A}_q \mathfrak{A}_r$ . Let  $|G| = q^\beta r^\gamma$ , where  $\beta$  and  $\gamma$  are natural numbers. If  $\beta = 0$  or  $\gamma = 0$ , then the result follows from Lemma 3.1. Assume that  $\beta$  and  $\gamma$  are at least 1. Let  $V = (\mathbb{F}_s)^\alpha$  and  $F = F(G)$ , the Fitting subgroup of  $G$ . Then  $F = Q \times R_1$ , where  $Q$  is the unique Sylow  $q$ -subgroup of  $G$  and  $R_1 \leq R$ , where  $R$  is a Sylow  $r$ -subgroup of  $G$ . So  $F$  is abelian and  $|F| = q^{\beta} r^{\gamma_1} = m$ , where  $\gamma_1 \leq \gamma$ .

From Clifford's theorem, regarding  $V$  as an  $\mathbb{F}_s F$ -module,  $V = Y_1 \oplus Y_2 \oplus \dots \oplus Y_l$ , where  $Y_i = kX_i$  for all  $i$ , and  $X_1, \dots, X_l$  are irreducible  $\mathbb{F}_s F$ -submodules of  $V$ . Further, for each  $i, j$ , there exists  $g_{ij} \in G$  such that  $g_{ij} X_i = X_j$  and, for  $i = 1, \dots, l$ , the  $X_i$  form a maximal set of pairwise nonisomorphic conjugates. Also, the action of  $G$  on the  $Y_i$  is transitive. It is not difficult to check that  $C_F(Y_i) = C_F(X_i) = K_i$ , say. Thus,  $F/K_i$  acts faithfully on  $Y_i$  and when its action is restricted to  $X_i$ , it acts faithfully and irreducibly on  $X_i$ . Since  $X_i$  is a nontrivial irreducible faithful  $\mathbb{F}_s F/K_i$ -module, and  $t$  is coprime to  $q$  and  $r$ , it follows that  $F/K_i$  is cyclic and  $\dim_{\mathbb{F}_s}(X_i) = d_i$ , where  $d_i$  is the least positive integer such that  $m_i$  divides  $s^{d_i} - 1$ , and where  $m_i$  is the order of  $F/K_i$ . Since the  $X_i$  are conjugate,  $\dim_{\mathbb{F}_s}(X_i) = d_i = d$  for all  $i$ .

Let  $E_i$  be the subalgebra generated by  $F/K_i$  in  $\text{End}_{\mathbb{F}_s}(Y_i)$ . Note that  $E_i$  is commutative as  $F/K_i$  is abelian. Further,  $X_i$  is a faithful irreducible  $E_i$ -module. So  $E_i$  is simple and becomes a field such that  $E_i \cong \mathbb{F}_{s^{d_i}}$ . We also observe that  $\alpha = kld$ .

Let  $k, l, d$  be fixed such that  $\alpha = kld$ . Next we find the number of choices for  $F$  up to conjugacy in  $\text{GL}(V)$ . Clearly,

$$\begin{aligned} F &\leq F/K_1 \times F/K_2 \times \dots \times F/K_l \leq E_1^* \times E_2^* \times \dots \times E_l^* \\ &\leq \text{GL}(Y_1) \times \text{GL}(Y_2) \times \dots \times \text{GL}(Y_l) \leq \text{GL}(V), \end{aligned}$$

where  $E_i^*$  denotes the multiplicative group of the field  $E_i$ . Let  $E = E_1^* \times E_2^* \times \dots \times E_l^*$ . Then  $|E| = (s^d - 1)^l$ . Regarding  $V$  as an  $\mathbb{F}_s E$ -module,  $V = kX_1 \oplus kX_2 \oplus \dots \oplus kX_l$ , where  $E_i^*$  acts faithfully and irreducibly on  $X_i$  and  $\dim_{E_i}(X_i) = 1$  for all  $i$ . Further, for  $i \neq j$ ,  $E_i^*$  acts trivially on  $X_j$ . It is not difficult to show that there is only one conjugacy class of subgroups of type  $E$  in  $\text{GL}(V)$ .

So once  $k, l$  and  $d$  are chosen such that  $\alpha = kld$ , up to conjugacy, there is only one choice for  $E$ . Since  $E$  is a direct product of  $l$  isomorphic cyclic groups, any subgroup of  $E$  can be generated by  $l$  elements. In particular,  $F$  can be generated by  $l$  elements. So the number of choices for  $F$  as a subgroup of  $E$  is at most  $|E|^l = (s^d - 1)^{l^2}$ .

Since,  $G$  acts transitively on  $\{Y_1, \dots, Y_l\}$ , there exists a homomorphism  $\phi$  from  $G$  into  $S_l$ . Let  $N = \ker(\phi) = \{g \in G \mid gY_i = Y_i \text{ for all } i\}$ . Clearly,  $F \leq N$  and  $G/N$  is a transitive subgroup of  $S_l$  that is in  $\mathfrak{A}_r$ . If  $g \in N$ , then  $gE_i g^{-1} = E_i$ . Thus, there exists a homomorphism  $\psi_i : N \rightarrow \text{Gal}_{\mathbb{F}_s}(E_i)$ . This induces a homomorphism  $\psi$  from  $N$  to  $\text{Gal}_{\mathbb{F}_s}(E_1) \times \text{Gal}_{\mathbb{F}_s}(E_2) \times \dots \times \text{Gal}_{\mathbb{F}_s}(E_l)$  such that  $\ker(\psi) = \bigcap_{i=1}^l N_i = F$ , where  $N_i = \ker(\psi_i) = C_N(E_i)$ . So  $N/F$  is isomorphic to a subgroup of  $\text{Gal}_{\mathbb{F}_s}(E_1) \times \text{Gal}_{\mathbb{F}_s}(E_2) \times \dots \times \text{Gal}_{\mathbb{F}_s}(E_l)$ . Since  $\text{Gal}_{\mathbb{F}_s}(E_i) \cong C_d$  for every  $i$ , it follows that  $N/F$  can be generated by  $l$  elements.

Let  $T = \text{GL}(\alpha, s)$ . Let  $\hat{N} = \{x \in N_T(F) \mid xY_i = Y_i \text{ for all } i\}$ . Then  $F \leq N \leq \hat{N} \leq N_T(F)$ . We will find the number of choices for  $N$  as a subgroup of  $\hat{N}$ , given that  $F$  has been chosen. The group  $\hat{N}$  acts by conjugation on  $E_i$  and fixes the elements of  $\mathbb{F}_s$ . So we have a homomorphism  $\rho_i : \hat{N} \rightarrow \text{Gal}_{\mathbb{F}_s}(E_i)$  with kernel  $C_{\hat{N}}(E_i)$ . Define  $C = \bigcap_{i=1}^l C_{\hat{N}}(E_i)$ . Note that  $N \cap C = F$ . Also,  $\hat{N}/C$  is isomorphic to a subgroup of  $\text{Gal}_{\mathbb{F}_s}(E_1) \times \text{Gal}_{\mathbb{F}_s}(E_2) \times \dots \times \text{Gal}_{\mathbb{F}_s}(E_l)$ , where each  $\text{Gal}_{\mathbb{F}_s}(E_i)$  is isomorphic to  $C_d$ . So  $|\hat{N}/C| \leq d^l$ . Clearly,  $C$  centralises  $E_i$  for each  $i$ . Therefore, there exists a homomorphism from  $C$  into  $\text{GL}_{E_i}(Y_i)$  for each  $i$ . Hence,  $C$  is isomorphic to a subgroup of  $\text{GL}_{E_1}(Y_1) \times \text{GL}_{E_2}(Y_2) \times \dots \times \text{GL}_{E_l}(Y_l)$ . As  $\dim_{\mathbb{F}_i}(Y_i) = k$  and  $E_i \cong \mathbb{F}_{s^d}$  for all  $i$ , it follows that  $|C| \leq s^{dk^2l}$ . Hence,  $|\hat{N}| \leq d^l s^{dk^2l}$ .

Now  $NC/C \cong N/(N \cap C) = N/F$ . So  $NC/C$  can be generated by  $l$  elements since  $N/F$  can be generated by  $l$  elements. However,  $|\hat{N}/C| \leq d^l$ , therefore, there are at most  $d^{l^2}$  choices for  $NC/C$  as a subgroup of  $\hat{N}/C$ . Once we make a choice for  $NC/C$  as a subgroup of  $\hat{N}/C$ , we choose a set of  $l$  generators for  $NC/C$ . As  $N \cap C = F$ , we see that  $N$  is determined as a subgroup of  $\hat{N}$  by  $F$  and  $l$  other elements that map to the chosen generating set of  $NC/C$ . We have  $|C|$  choices for an element of  $\hat{N}$  that maps to any fixed element of  $\hat{N}/C$ . Thus, there are at most  $|C|^l$  choices for  $N$  as a subgroup of  $\hat{N}$  once  $NC/C$  has been chosen. So we have at most  $d^{l^2} (s^{dk^2l})^l = d^{l^2} s^{dk^2l^2}$  choices for  $N$  as a subgroup of  $\hat{N}$ , once  $F$  is fixed.

Next we find the number of choices for  $G$  given that  $F$  and  $N$  are fixed as subgroups of  $T$  and  $\hat{N} \leq T$ , respectively. Let  $\hat{Y} = \{y \in N_T(F) \mid y \text{ permutes the } Y_i\}$ . Then  $F \leq G \leq \hat{Y} \leq N_T(F) \leq \text{GL}(V)$ . Also there exists a homomorphism from  $\hat{Y}$  to  $S_l$  with kernel  $\{y \in \hat{Y} \mid yY_i = Y_i \text{ for all } i\} = \hat{N}$ . Thus,  $\hat{Y}/\hat{N}$  may be regarded as a subgroup of  $S_l$ . However,  $G \cap \hat{N} = N$ . Thus,  $G/N = G/(G \cap \hat{N}) \cong G\hat{N}/\hat{N}$ . So  $G/N \cong G\hat{N}/\hat{N} \leq \hat{Y}/\hat{N} \leq S_l$ . Note that  $G/N$  is a transitive subgroup of  $S_l$  that is in  $\mathfrak{A}_r$ . By [5, Theorem 1], there exists a constant  $b$  such that  $S_l$  has at most  $2^{b l^2 / \sqrt{\log l}}$  transitive subgroups for  $l > 1$ . Hence, the number of choices for  $G\hat{N}/\hat{N}$  as a subgroup of  $\hat{Y}/\hat{N}$  is at most  $2^{b l^2 / \sqrt{\log l}}$ .

By [6, Theorem 2], there exists a constant  $c$  such that any transitive permutation group of finite degree greater than 1 can be generated by  $\lfloor cl / \sqrt{\log l} \rfloor$  generators. Thus,  $G\hat{N}/\hat{N}$  can be generated by  $\lfloor cl / \sqrt{\log l} \rfloor$  generators for  $l > 1$ . Once a choice for  $G\hat{N}/\hat{N}$

is made as a subgroup of  $\hat{Y}/\hat{N}$  and  $\lfloor cl/\sqrt{\log l} \rfloor$  generators are chosen for  $G\hat{N}/\hat{N}$  in  $\hat{Y}/\hat{N}$ , then  $G$  is determined as a subgroup of  $\hat{Y}$  by  $\hat{N}$  and the elements of  $\hat{Y}$  that map to the  $\lfloor cl/\sqrt{\log l} \rfloor$  generators chosen for  $G\hat{N}/\hat{N}$ . So we have at most  $|\hat{N}|^{\lfloor cl/\sqrt{\log l} \rfloor}$  choices for  $G$  as a subgroup of  $\hat{Y}$  once a choice of  $G\hat{N}/\hat{N}$  in  $\hat{Y}/\hat{N}$  is fixed. Hence, there are

$$2^{bl^2/\sqrt{\log l}} (d^l s^{dk^2l})^{\lfloor cl/\sqrt{\log l} \rfloor} \leq 2^{bl^2/\sqrt{\log l}} d^{cl^2/\sqrt{\log l}} s^{cdk^2l^2/\sqrt{\log l}}$$

choices for  $G$  as a subgroup of  $\hat{Y}$  assuming that choices for  $F$  and  $N$  have been made. Putting together all these estimates, the number of conjugacy classes of subgroups that are maximal amongst irreducible subgroups of  $GL(\alpha, s)$  that are in  $\mathfrak{A}_q \mathfrak{A}_r$  is at most

$$\sum_{(k,l,d)} (s^d - 1)^l d^{l^2} s^{dk^2l^2} 2^{bl^2/\sqrt{\log l}} d^{cl^2/\sqrt{\log l}} s^{cdk^2l^2/\sqrt{\log l}},$$

where  $(k, l, d)$  ranges over ordered triples of natural numbers which satisfy  $\alpha = kld$  and  $l > 1$ . We simplify the above expression as follows. Writing  $\alpha = kld$ ,

$$(s^d - 1)^l d^{l^2} s^{dk^2l^2} s^{cdk^2l^2/\sqrt{\log l}} \leq s^{(3+c)\alpha^2}.$$

Since  $x/\sqrt{\log x}$  is increasing for  $x > e^{1/2}$ , we have  $l/\sqrt{\log l} \leq \alpha/\sqrt{\log \alpha}$  for  $l \geq 2$ . Thus,  $2^{bl^2/\sqrt{\log l}} d^{cl^2/\sqrt{\log l}} \leq 2^{(b+c)\alpha^2/\sqrt{\log \alpha}}$ .

There are at most  $2^{(5/6)\log \alpha + \log 6}$  choices for  $(k, l, d)$ . Thus, there exist constants  $b$  and  $c$  such that the number of conjugacy classes of subgroups that are maximal amongst irreducible subgroups of  $GL(\alpha, s)$  that are in  $\mathfrak{A}_q \mathfrak{A}_r$  is at most

$$2^{(b+c)(\alpha^2/\sqrt{\log \alpha}) + (5/6)\log \alpha + \log 6} s^{(3+c)\alpha^2}$$

provided  $\alpha > 1$ . □

Theorem 1.5 follows as a corollary to Proposition 4.2.

**PROOF OF THEOREM 1.5.** Let  $G$  be maximal amongst subgroups of  $GL(\alpha, s)$  that are also in  $\mathfrak{A}_q \mathfrak{A}_r$ . As the characteristic of  $\mathbb{F}_s = t$  and  $t \nmid |G|$ , by Maschke’s theorem,  $G \leq \hat{G}_1 \times \dots \times \hat{G}_k \leq GL(\alpha, s)$ , where the  $\hat{G}_i$  are maximal among irreducible subgroups of  $GL(\alpha_i, p)$  that are also in  $\mathfrak{A}_q \mathfrak{A}_r$ , and where  $\alpha = \alpha_1 + \dots + \alpha_k$ . By the maximality of  $G$ , we have  $G = \hat{G}_1 \times \dots \times \hat{G}_k$ .

The conjugacy classes of  $\hat{G}_i \in GL(\alpha_i, s)$  determine the conjugacy class of  $G \in GL(\alpha, s)$ . So by Proposition 4.2, the number of conjugacy classes of subgroups that are maximal amongst the subgroups of  $GL(\alpha, s)$  that are also in  $\mathfrak{A}_q \mathfrak{A}_r$  is at most

$$\sum_{(\alpha)} \prod_{i=1}^k 2^{(b+c)(\alpha_i^2/\sqrt{\log \alpha_i}) + (5/6)\log \alpha_i + \log 6} s^{(3+c)\alpha_i^2},$$

where the sum is over all unordered partitions  $\alpha_1, \dots, \alpha_k$  of  $\alpha$ . We assume that if  $\alpha_i = 1$  for some  $i$ , then the part of the expression corresponding to it in the product is 1. Since  $x/\sqrt{\log x}$  is increasing for  $x > e^{1/2}$  and  $\alpha = \alpha_1 + \dots + \alpha_k$ ,



$$\prod_{i=1}^k 2^{(b+c)(\alpha_i^2/\sqrt{\log \alpha_i})+(5/6) \log \alpha_i + \log 6} \leq 2^{(b+c)(\alpha^2/\sqrt{\log \alpha})+(5/6)\alpha \log \alpha + \alpha \log 6}.$$

It is not difficult to show that the number of unordered partitions of  $\alpha$  is at most  $2^{\alpha-1}$ . So the number of conjugacy classes of subgroups that are maximal amongst the subgroups of  $\text{GL}(\alpha, s)$  that are also in  $\mathfrak{A}_q \mathfrak{A}_r$  is at most

$$2^{(b+c)(\alpha^2/\sqrt{\log \alpha})+(5/6)\alpha \log \alpha + \alpha(1+\log 6)} 5^{(3+c)\alpha^2}$$

provided  $\alpha > 1$ . □

We end this section with the following remark that provides an alternative bound.

**REMARK 4.3.** We do not have an estimate for the constants  $b$  and  $c$  occurring in Theorem 1.5. If we use a weaker fact that any subgroup of  $S_n$  can be generated by  $\lfloor n/2 \rfloor$  elements for all  $n \geq 3$ , then we get a weaker result that the number of transitive subgroups of  $S_n$  that are in  $\mathfrak{A}_q \mathfrak{A}_r$  is at most  $6^{n(n-1)/4} 2^{(n+2) \log n}$ . Using this in the proof of Theorem 1.5 shows that the number of conjugacy classes of subgroups that are maximal amongst the subgroups of  $\text{GL}(\alpha, s)$  that are also in  $\mathfrak{A}_q \mathfrak{A}_r$  is at most

$$s^{5\alpha^2} 6^{\alpha(\alpha-1)/4} 2^{\alpha-1+(23/6)\alpha \log \alpha + \alpha \log 6},$$

where  $t, q$  and  $r$  are distinct primes,  $s$  is a power of  $t$ , and  $\alpha \in \mathbb{N}$ .

### 5. Enumeration of groups in $\mathfrak{A}_p \mathfrak{A}_q \mathfrak{A}_r$

In this section, we prove Theorem 1.2, namely,

$$f_{\in}(n) \leq p^{6\alpha^2} 2^{\alpha-1+(23/6)\alpha \log \alpha + \alpha \log 6} (6^{1/2})^{(\alpha+\gamma)\beta + (\alpha+\beta)\gamma + \alpha(\alpha-1)/2} n^{\beta+\gamma},$$

where  $n = p^\alpha q^\beta r^\gamma$  and  $\alpha, \beta, \gamma \in \mathbb{N}$ . We use techniques adapted from [9, 13, 14].

**PROOF OF THEOREM 1.2.** Let  $G$  be a group of order  $n = p^\alpha q^\beta r^\gamma$  in  $\mathfrak{A}_p \mathfrak{A}_q \mathfrak{A}_r$ . Then  $G = P \rtimes H$ , where  $P$  is the unique Sylow  $p$ -subgroup of  $G$  and  $H \in \mathfrak{A}_q \mathfrak{A}_r$ . So we can write  $H = Q \rtimes R$ , where  $|Q| = q^\beta$  and  $|R| = r^\gamma$ . Let  $G_1 = G/O_{p'}(G)$ ,  $G_2 = G/O_{q'}(G)$  and  $G_3 = G/O_{r'}(G)$ . Clearly, each  $G_i$  is a soluble  $A$ -group and  $G \leq G_1 \times G_2 \times G_3$  as a subdirect product. Further,  $O_{p'}(G_1) = 1 = O_{q'}(G_2) = O_{r'}(G_3)$ .

Since  $G_1 = G/O_{p'}(G)$ , we see that  $G_1 \in \mathfrak{A}_p \mathfrak{A}_q \mathfrak{A}_r$  and if  $P_1$  is the Sylow  $p$ -subgroup of  $G_1$ , then  $P_1 \cong P$ . Thus,  $|G_1| = p^\alpha q^{\beta_1} r^{\gamma_1}$  and we can write  $G_1 = P_1 \rtimes H_1$ , where  $H_1 \in \mathfrak{A}_q \mathfrak{A}_r$ . So  $H_1 = Q_1 \rtimes R_1$ , where  $Q_1 \in \mathfrak{A}_q$  and  $|Q_1| = q^{\beta_1}$ ,  $R_1 \in \mathfrak{A}_r$  and  $|R_1| = r^{\gamma_1}$ . Further,  $H_1$  acts faithfully on  $P_1$ . Hence, we can regard  $H_1 \leq \text{Aut}(P_1) \cong \text{GL}(\alpha, p)$ . Let  $M_1$  be a subgroup that is maximal amongst  $p'$ - $A$ -subgroups of  $\text{GL}(\alpha, p)$  that are also in  $\mathfrak{A}_q \mathfrak{A}_r$  and such that  $H_1 \leq M_1$ . Let  $\hat{G}_1 = P_1 M_1$ . The number of conjugacy classes of the  $M_1$  in  $\text{GL}(\alpha, p)$  is at most  $p^{5\alpha^2} 6^{\alpha(\alpha-1)/4} 2^{\alpha-1+(23/6)\alpha \log \alpha + \alpha \log 6}$  by Remark 4.3.

Since  $G_2 = G/O_{q'}(G)$ , we see that  $G_2 \in \mathfrak{A}_q \mathfrak{A}_r$  and if  $Q_2$  is the Sylow  $q$ -subgroup of  $G_2$ , then  $Q_2 \cong Q$ . Thus,  $|G_2| = q^{\beta} r^{\gamma_2}$  and we can write  $G_2 = Q_2 \rtimes H_2$ , where

$H_2 \in \mathfrak{A}_r$ . So  $|H_2| = r^{\gamma_2}$ . Also,  $H_2 \leq \text{Aut}(Q_2) \cong \text{GL}(\beta, q)$ . Let  $M_2$  be a subgroup that is maximal amongst  $q'$ - $A$ -subgroups of  $\text{GL}(\beta, q)$  that are also in  $\mathfrak{A}_r$  and such that  $H_2 \leq M_2$ . Let  $\hat{G}_2 = Q_2 M_2$ . The number of conjugacy classes of  $M_2$  in  $\text{GL}(\beta, q)$  is at most 1 by Proposition 3.2.

Since  $G_3 = G/O_{r'}(G)$ , we see that  $G_3 \in \mathfrak{A}_r \mathfrak{A}_q$  and if  $R_3$  is the Sylow  $r$ -subgroup of  $G_3$ , then  $R_3 \cong R$ . Thus,  $|G_3| = q^{\beta_3} r^{\gamma}$  and we can write  $G_3 = R_3 \rtimes H_3$ , where  $H_3 \in \mathfrak{A}_r$ . So  $|H_3| = q^{\beta_3}$ . Also,  $H_3 \leq \text{Aut}(R_3) \cong \text{GL}(\gamma, r)$ . Let  $M_3$  be a subgroup that is maximal amongst  $r'$ - $A$ -subgroups of  $\text{GL}(\gamma, r)$  that are also in  $\mathfrak{A}_q$  and such that  $H_3 \leq M_3$ . Let  $\hat{G}_3 = R_3 M_3$ . The number of conjugacy classes of the  $M_3$  in  $\text{GL}(\gamma, r)$  is at most 1 by Proposition 3.2.

Let  $\hat{G} = \hat{G}_1 \times \hat{G}_2 \times \hat{G}_3$ . Then  $G \leq \hat{G}$ . The choices for  $P_1, Q_2$  and  $R_3$  are unique, up to isomorphism. We enumerate the possibilities for  $\hat{G}$  up to isomorphism and then find the number of subgroups of  $\hat{G}$  of order  $n$  up to isomorphism. For the former, we count the number of  $\hat{G}_i$  up to isomorphism which depends on the conjugacy class of the  $M_i$ . Hence, the number of choices for  $\hat{G}$  up to isomorphism is  $\prod_{i=1}^3 \{\text{number of choices for } \hat{G}_i \text{ up to isomorphism}\}$ . Now we estimate the choices for  $G$  as a subgroup of  $\hat{G}$  using a method of ‘Sylow systems’ introduced by Pyber in [9].

Let  $\hat{G}$  be fixed. We count the number of choices for  $G$  as a subgroup of  $\hat{G}$ . Let  $S = \{S_1, S_2, S_3\}$  be a Sylow system for  $G$ , where  $S_1$  is the Sylow  $p$ -subgroup of  $G$ ,  $S_2$  is a Sylow  $q$ -subgroup of  $G$  and  $S_3$  is a Sylow  $r$ -subgroup of  $G$  such that  $S_i S_j = S_j S_i$  for all  $i, j = 1, 2, 3$ . Then  $G = S_1 S_2 S_3$ . By [1, Theorem 6.2, page 49], there exists  $\mathcal{B} = \{B_1, B_2, B_3\}$ , a Sylow system for  $\hat{G}$  such that  $S_i \leq B_i$ , where  $B_1$  is the Sylow  $p$ -subgroup of  $\hat{G}$ ,  $B_2$  is a Sylow  $q$ -subgroup of  $\hat{G}$  and  $B_3$  is a Sylow  $r$ -subgroup of  $\hat{G}$ . Note that  $|B_1| = p^\alpha$ . Further, any two Sylow systems for  $\hat{G}$  are conjugate. Hence, the number of choices for  $G$  as a subgroup of  $\hat{G}$  and up to conjugacy is at most

$$|\{S_1, S_2, S_3 \mid S_i \leq B_i, |S_1| = p^\alpha, |S_2| = q^\beta, |S_3| = r^\gamma\}| \leq |B_1|^\alpha |B_2|^\beta |B_3|^\gamma.$$

We observe that  $B_2 = T_{21} \times T_{22} \times T_{23}$ , where  $T_{2i}$  are Sylow  $q$ -subgroups of  $\hat{G}_i$  for  $i = 1, 2, 3$ . From [13, Proposition 3.1],  $|T_{21}| \leq |M_1| \leq (6^{1/2})^{\alpha-1} p^\alpha$  and  $|T_{23}| = |M_3| \leq (6^{1/2})^{\gamma-1} r^\gamma$ . Further,  $|T_{22}| = |Q_2| = q^\beta$ . Hence,  $|B_2| \leq (6^{1/2})^{\alpha+\gamma-2} p^\alpha q^\beta r^\gamma \leq (6^{1/2})^{\alpha+\gamma} n$  and so  $|B_2|^\beta \leq (6^{1/2})^{(\alpha+\gamma)\beta} n^\beta$ . Similarly, we can show that  $|B_3| \leq (6^{1/2})^{\alpha+\beta-2} p^\alpha q^\beta r^\gamma$ . So  $|B_3|^\gamma \leq (6^{1/2})^{(\alpha+\beta)\gamma} n^\gamma$ . Putting all the estimates together, the number of choices for  $G$  as a subgroup of  $\hat{G}$  up to conjugacy is at most  $|B_1|^\alpha |B_2|^\beta |B_3|^\gamma$ , which is at most

$$p^{\alpha^2} (6^{1/2})^{(\alpha+\gamma)\beta} n^\beta (6^{1/2})^{(\alpha+\beta)\gamma} n^\gamma \leq p^{\alpha^2} (6^{1/2})^{(\alpha+\gamma)\beta+(\alpha+\beta)\gamma} n^{\beta+\gamma}.$$

Therefore, the number of groups of order  $p^\alpha q^\beta r^\gamma$  in  $\mathfrak{A}_p \mathfrak{A}_q \mathfrak{A}_r$  up to isomorphism is at most

$$\begin{aligned} & p^{5\alpha^2} 6^{\alpha(\alpha-1)/4} 2^{\alpha-1+(23/6)\alpha \log \alpha + \alpha \log 6} p^{\alpha^2} (6^{1/2})^{(\alpha+\gamma)\beta+(\alpha+\beta)\gamma} n^{\beta+\gamma} \\ &= p^{6\alpha^2} 2^{\alpha-1+(23/6)\alpha \log \alpha + \alpha \log 6} (6^{1/2})^{(\alpha+\gamma)\beta+(\alpha+\beta)\gamma+\alpha(\alpha-1)/2} n^{\beta+\gamma}. \quad \square \end{aligned}$$

## References

- [1] S. R. Blackburn, P. M. Neumann and G. Venkataraman, *Enumeration of Finite Groups* (Cambridge University Press, Cambridge, 2007).
- [2] G. Dickenson, 'On the enumeration of certain classes of soluble groups', *Q. J. Math.* **20**(1) (1969), 383–394.
- [3] M. D. Hestenes, 'Singer groups', *Canad. J. Math.* **22**(3) (1970), 492–513.
- [4] G. Higman, 'Enumerating  $p$ -groups. I: Inequalities', *Proc. Lond. Math. Soc. (3)* **10**(3) (1960), 24–30.
- [5] A. Lucchini, 'Enumerating transitive finite permutation groups', *Bull. Lond. Math. Soc.* **30**(6) (1998), 569–577.
- [6] A. Lucchini, F. Menegazzo and M. Morigi, 'Asymptotic results for transitive permutation groups', *Bull. Lond. Math. Soc.* **32**(2) (2000), 191–195.
- [7] A. McIver and P. M. Neumann, 'Enumerating finite groups', *Q. J. Math.* **38**(2) (1987), 473–488.
- [8] H. Neumann, *Varieties of Groups* (Springer-Verlag, Berlin–Heidelberg, 1967).
- [9] L. Pyber, 'Enumerating finite groups of given order', *Ann. of Math. (2)* **137**(1) (1993), 203–220.
- [10] L. Scott, 'Representations in characteristic  $p$ ', in: *The Santa Cruz Conference on Finite Groups*, Proceedings of Symposia in Pure Mathematics, 37 (eds. B. Cooperstein and G. Mason) (American Mathematical Society, Providence, RI, 1981).
- [11] M. W. Short, *The Primitive Soluble Permutation Groups of Degree Less than 256* (Springer, Berlin–Heidelberg, 1992).
- [12] C. C. Sims, 'Enumerating  $p$ -groups', *Proc. Lond. Math. Soc. (3)* **15**(3) (1965), 151–166.
- [13] G. Venkataraman, 'Enumeration of finite soluble groups with Abelian Sylow subgroups', *Q. J. Math.* **48**(1) (1997), 107–125.
- [14] G. Venkataraman, *Enumeration of Finite Soluble Groups in Small Varieties of  $A$ -groups and Associated Topics*, Tech. Report, Centre for Mathematical Sciences, St. Stephen's College (University of Delhi, Delhi, 1999).
- [15] H. Wielandt, *Finite Permutation Groups* (Academic Press, London, 1964), translated from German by R. Bercov.

ARUSHI, School of Liberal Studies (Mathematics),  
Dr. B. R. Ambedkar University Delhi, Delhi 110006, India  
e-mail: [arushi.18@stu.aud.ac.in](mailto:arushi.18@stu.aud.ac.in), [arushi.garvita@gmail.com](mailto:arushi.garvita@gmail.com)

GEETHA VENKATARAMAN, School of Liberal Studies (Mathematics),  
Dr. B. R. Ambedkar University Delhi, Delhi 110006, India  
e-mail: [geetha@aud.ac.in](mailto:geetha@aud.ac.in), [geevenkat@gmail.com](mailto:geevenkat@gmail.com)