

JOURNAL
OF
HELMINTHOLOGY

Volume 56, 1982

Edited by
SHEILA WILLMOTT

**London School of Hygiene
and
Tropical Medicine
Keppel Street
London
WC1E 7HT**

JOURNAL OF HELMINTHOLOGY

Founded by R. T. Leiper in 1923

Former Editors:

R. T. Leiper 1923–46
J. J. C. Buckley 1946–72
R. Muller 1972–80

Published by:

London School of Hygiene and Tropical Medicine,
Keppel Street (Gower Street),
London WC1E 7HT, UK

ISSN 0022-149X

Printed by Taylor & Francis (Printers) Ltd, Rankine Road, Basingstoke, Hampshire RG24 0PR, U.K.

INSTRUCTION TO CONTRIBUTORS

The *Journal of Helminthology* publishes papers on all aspects of animal parasitic helminths, particularly those of medical or veterinary importance, but only in exceptional circumstances will systematic or taxonomic studies be acceptable.

Manuscripts, which must be in English or French (with an English summary), should be addressed to:

The Editor, Journal of Helminthology,
London School of Hygiene and Tropical Medicine,
Winches Farm Field Station,
395 Hatfield Road,
St Albans, Herts AL4 0XQ,
England.

Two copies of a typescript, on size A4 paper with double spacing, should be submitted. Papers should be preceded by a short abstract and will normally have the following sections: brief Introduction; Materials and Methods; Results; Discussion; Acknowledgements; References. However, the form of the paper may vary, depending on its subject matter; recent past issues should be consulted for a suitable form. Research Notes should also be preceded by a brief abstract. Illustrations should be drawn in Indian ink, preferably not more than double the final size. Care should be taken that all illustrations fit into the format of the Journal. The maximum size an illustration will be printed is 12.0 × 20.0 cm. Where many separate drawings are made, some indication of how they may be grouped to make a corporate plate without undue wastage of space, should be indicated. Some indication of scale (preferably a scale bar) should normally be given on the figure. Photocopies of illustrations should be enclosed for refereeing purposes. Lettering and numbering, which must be of a high standard, should be added by the author, with due regard for subsequent reduction.

Photographs should be glossy prints of the same size as they are to appear in the Journal (maximum size 12.0 × 20.0 cm). Composite prints must be mounted and can have the separate photographs abutting; they will then have a separating line inserted by the printers. All figures and letters on photographs must be inserted by the author.

Information should not be repeated in the text and in tables or figures. The legends to tables and to figures should be sufficiently detailed for the information to be understood without reference to the text.

References should be given in alphabetical order with the full title of the journal. The following are examples:

DUKE, B. O. L. (1971) The ecology of onchocerciasis in man and animals. In: *Ecology and Physiology of parasites* (editor, A. M. Fallis) pp. 213–222. Adam Hilger Ltd.: London.

JAMES, C. & WEBBE, G. (1973) A comparison of Egyptian and East African strains of *Schistosoma haematobium*. *Journal of Helminthology*, **47**, 49–59.

25 offprints are provided free of charge; additional copies may be ordered at the proof stage.

Contents

	Pages
Immunization of calves against the microfilariae of <i>Onchocerca lienalis</i> . SIMON TOWNSON and A. E. BIANCO	297-303
The effects of treatment with transfer factor, parasitic antigen and precipitated immunoglobulin on the parasitological and histopathological responses in <i>Haemonchus contortus</i> infection in lambs under six months of age. GRACE MACKENZIE, A. R. HUNTER and J. G. ROSS	305-313
<i>Spirometra mansonoides</i> : effects of plerocercoid infection on glycogen deposition in rats. C. K. PHARES and NGUYEN DUY AI	315-322
A method for determining the concentration of <i>Onchocerca volvulus</i> microfilariae in urine. (Research Note.) OLADELE O. KALE and OLUSOLA AYENI	323-326
Biochemistry of infective eggs of the cestode <i>Moniezia benedeni</i> . (Research Note.) A. VINAYAKAM	327-328
<i>Nematospiroides dubius</i> in the jird, <i>Meriones unguiculatus</i> : factors affecting the course of a primary infection. JOHN HANNAH and JERZEY M. BEHNKE	329-338
The nature of antigens common to both the metacestodes of <i>Taenia crassiceps</i> and its laboratory host. J. CHERNIN	339-344
Prevalence of <i>Angiostrongylus cantonensis</i> infection in commensal rats in Bombay. D. M. RENAPURKAR, M. K. BHOPALE, LALITA S. LIMAYE and K. S. SHARMA	345-349
The tail morphology of different developmental stages of <i>Ollulanus tricuspis</i> (Leuckart, 1865) (Nematoda: Trichostrongyloidea). (Research Note.) M.-A. HASSLINGER and F. X. WITTMANN	351-353
The anthelmintic effects of mebendazole on <i>Necator americanus</i> in rabbit model. (Research Note.) M. K. BHOPALE, SUSHILA MENON and VIJAYALAXMI R. KAMATH	355-356
Subject Index	357
Author Index	360
Contents of Volume	i-iii