

Conclusions: The bone obliteration technique combined with scar tissue and cartilage grafting saves time and effort in giving a dry and clean ear after recidivism. Most of the problems in a wet mastoid cavity are solved with this technique.

Learning Objectives: the video presentation gives a clear demonstration of the technique to be adopted by surgeons handling recidivism.

doi:10.1017/S0022215116003339

Update on bacteriology and the role of biofilms in chronic otitis media (K753)

ID: 753.1

Bacterial Biofilms & Chronic Otitis Media: Stuck in the middle

Presenting Author: **Luanne Hall-Stoodley**

Luanne Hall-Stoodley
The Ohio State University

Otitis media is a multifactorial disease, a result of complex host-microbial interactions. Understanding the pathogenesis of chronic otitis media (COM) is crucial for improving therapies. Direct detection of aggregated adherent otopathogenic bacteria on middle ear mucosal biopsies from children with COM demonstrated that biofilms were consistent with an infectious etiology in spite of culture-negative clinical data. This seminar will provide an overview of how biofilms contribute to chronic infections like COM, including problems in diagnosing the infectious agent in the polymicrobial context of the upper airway, the challenges of treatment and new therapeutic approaches on the horizon.

doi:10.1017/S0022215116003340

Biomaterials in middle ear reconstruction (R761)

ID: 761.1

Use of bone substitutes in mastoid obliteration

Presenting Author: **Daniele Bernardeschi**

Daniele Bernardeschi
Pitié-Salpêtrière Hospital

The purpose of this communication is to describe the indications, surgical technique and anatomical and function results of the mastoid and epitympanic obliteration using bone substitutes. This technique employed in our department since 2006, encompasses the use of synthetic biomaterials for the obliteration of mastoid and epitympanic spaces. Granules of biphasic ceramic have been used up to 2012 (n = 130) and, since 2013 (n = 74) we are using bioactive glass S53P4. Differences in composition and mechanism of action will be detailed, with particular attention to the antibacterial activity of the bioactive glass S53P4.

doi:10.1017/S0022215116003352

Biomaterials in middle ear reconstruction (R761)

ID: 761.2

SerenoCem™ - glass ionomeric granules in mastoid obliteration, a hidden problem!

Presenting Author: **Ian Bottrill**

Ian Bottrill
John Radcliffe Hospital, Oxford

Introduction: A common problem with canal-wall down mastoidectomy procedures is a discharging cavity. Many techniques for mastoid obliteration to reduce the cavity size have been described. Different biomaterials have been tried on the basis that they should be non-resorbable, non-reactive and integrate. This study aimed to assess the effectiveness of SerenoCem granules, a glass ionomeric cement, as a suitable biomaterial for mastoid obliteration and to review its longterm effects.

Methods: 16 patients with chronically discharging mastoid cavities were selected for mastoid obliteration. The subsequent procedures were performed between 2001 and 2003. The two main outcome measures were the number of attendances for aural care and the Glasgow Benefit Inventory (GBI). A secondary measure was the comparison of pre- and post-operative hearing thresholds. These patients were assessed in 2006 providing a minimum of 3 years follow up. As a result of recent chance finding following late revision surgery, a further review of implanted patients was undertaken in 2015/16.

Results: The need for aural care reduced in all but one patient. There was a significant difference in the number of aural visits pre and post operation. Benefit in quality of life was assessed using the GBI. In only one patient was there a negative score. The mean values indicate that there has been a positive benefit in quality of life. Complete pure tone average results were available for 13/16 patients. In 8/13 patients the hearing was improved, as intended by additional ossiculoplasty procedures. Of importance a reduction of hearing was noted in only 5 patients, the worst of which was 7.5 dB for the 4-tone average.

Conclusions: The initial results of this technique were promising, however, the recent chance review of one of these patients showed the granules may be inducing bone lysis in. All patients have been reviewed and the results will be presented.

doi:10.1017/S0022215116003364

Biomaterials in middle ear reconstruction (R761)

ID: 761.3

Titanium in mastoid reconstruction

Presenting Author: **Konrad Schwager**

Konrad Schwager
Klinikum Fulda, Philipps University of Marburg - Campus Fulda