



Potential role of targeted echocardiography as a screening test for select diagnoses in the paediatric population: bicuspid aortic valve and left ventricular hypertrophy

Original Article

Cite this article: Yang C, Edwards LA, Vernon MM, Conwell J, and Buddhe S (2025) Potential role of targeted echocardiography as a screening test for select diagnoses in the paediatric population: bicuspid aortic valve and left ventricular hypertrophy. *Cardiology in the Young* **35**: 921–925. doi: [10.1017/S1047951125001490](https://doi.org/10.1017/S1047951125001490)

Received: 17 August 2024
Revised: 24 January 2025
Accepted: 7 March 2025
First published online: 21 April 2025

Keywords:

bicuspid aortic valve; left ventricular hypertrophy; targeted single-view echocardiography; cardiac point-of-care ultrasound; goal-directed echocardiography

Corresponding author:

Christina Yang;
Email: christinayang721@gmail.com

Christina Yang¹ , Lindsay A. Edwards², Margaret M. Vernon², Jeffrey Conwell² and Sujatha Buddhe²

¹University of Washington, Department of Pediatrics, Seattle, WA, USA and ²Seattle Children's Hospital, Division of Pediatric Cardiology, Seattle, WA, USA

Abstract

Objective: We explore the role of targeted echocardiography as a screening tool for bicuspid aortic valve and left ventricular hypertrophy, specifically assessing the risk of missing significant cardiac findings that would otherwise be identified by comprehensive echocardiograms. **Method:** Children < 18 years at initial echocardiogram for indications of “family history of bicuspid aortic valve” and “left ventricular hypertrophy on electrocardiogram” were queried. Cardiology clinic notes and complete echocardiogram reports were reviewed for additional background histories and incidental findings. Follow-up clinic visits, if any, and management for those with incidental findings were reviewed. **Results:** Bicuspid aortic valve group included 138 patients, 71 (51%) males and mean age at comprehensive echo was 8.4 ± 4.8 years. Bicuspid aortic valve was found in 3.6%, incidental findings were found in 15 (11%), and follow-up was recommended in 4 (2.8%). Left ventricular hypertrophy group included 70 patients, 58 (83%) males and mean age at echo 10.9 ± 4.7 years. Left ventricular hypertrophy was found in 2.8%, incidental findings were found in 9 (13%), and follow-up was recommended in 2 (2.8%).

None of the follow-up group developed symptoms or required cardiac medications, exercise restrictions, or catheter or surgical-based interventions, except for one case of mild aortic root dilation who was restricted from heavy weightlifting. **Conclusion:** The risk of missing clinically important findings with targeted echocardiography that would have been identified with comprehensive echocardiography is extremely low for screening indications of isolated left ventricular hypertrophy on electrocardiogram or family history of bicuspid aortic valve, suggesting that targeted echocardiography could be an effective screening tool.

Introduction

Bicuspid aortic valve is one of the most common forms of CHD with an estimated prevalence of 1–2%.¹ Although commonly asymptomatic in childhood, it can be associated with serious complications including aortic valve stenosis or regurgitation and ascending aortic dilation, aneurysm, and dissection.² Early diagnosis is important and allows for monitoring and, in some cases, intervention before potentially catastrophic complications develop. Given the inheritable nature of bicuspid aortic valve, screening of first-degree family members is recommended with each new diagnosis.³ The gold standard for diagnosis is transthoracic echocardiography in children, which has an estimated sensitivity of 92% and specificity of 96%.⁴

Left ventricular hypertrophy is a common finding on paediatric electrocardiogram, especially in highly trained athletes.⁵ This finding is nonspecific with diverse aetiologies including structural heart disease, cardiomyopathy, and non-cardiac causes. Echocardiogram, the gold standard for diagnosis, is often performed with this electrocardiogram finding. In a study analysing 4637 encounters of patients 0–18 years of age with left ventricular hypertrophy comparing electrocardiogram and echocardiogram performed within 24 hours, electrocardiogram had high sensitivity ($\geq 90\%$) but poor specificity (43%), with low positive predictive value ($< 20\%$).⁵

Current standard of practice for initial echocardiographic assessment for all screening indications in paediatrics is comprehensive echocardiography. While comprehensive echocardiograms provide standardised and detailed information, targeted ultrasonography has been increasingly utilised in many healthcare settings with impressive efficiency and diagnostic results. Real-time cardiac point-of-care ultrasound interpretation performed by paediatric emergency medicine attendings had sensitivity and specificity of 100% and 99.5%, respectively, for both pericardial effusion and left ventricular systolic dysfunction in comparison to expert review.⁶ Cardiac point-of-care ultrasound has also been increasingly incorporated into

© The Author(s), 2025. Published by Cambridge University Press. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.

neonatal practice with application in assessment of umbilical venous catheter placement, pericardial effusion, and cardiac tamponade.⁷

Targeted echocardiogram may be a more cost-effective screening tool. A proof-of-concept pilot study performing cost-benefit analysis on 100 patients with increased cardiac risk showed that a targeted echocardiogram performed before elective non-cardiac surgery may be more cost-effective than conventional comprehensive echocardiogram.⁸ Targeted echocardiogram for low-yield screening indications may be more accessible and more cost-effective and may avoid identification of insignificant incidental findings that lead to costly follow-up visits, imaging studies, and potential stress for patients and their families.

Our study aims to explore the role of targeted echocardiography as a screening tool for bicuspid aortic valve and left ventricular hypertrophy in the paediatric age group. We specifically aim to assess the risk of missing significant cardiac diagnoses that would otherwise be identified by comprehensive echocardiograms.

Methods

Study design and patient selection. This retrospective observational study was approved by the Seattle Children's Hospital Institutional Review Board. The Picture Archiving and Communication Systems imaging system was queried for echocardiogram studies meeting study criteria. Inclusion criteria were: 1) < 18 years of age at the time of initial screening echocardiogram; 2) comprehensive transthoracic screening echocardiogram was performed; 3) the screening echocardiogram was performed between 2011 and 2020; and 4) indication for screening echocardiogram was bicuspid aortic valve or left ventricular hypertrophy as indicated by the echo screening indication codes below:

1. The indication for screening for bicuspid aortic valve included "family history of bicuspid aortic valve."
2. The indication for screening for left ventricular hypertrophy included "left ventricular hypertrophy on electrocardiogram" or "biventricular hypertrophy on electrocardiogram."

Patients with additional history or exam findings documented in the cardiology visit note meeting the "appropriate" or "may be appropriate" criteria for performing a comprehensive echocardiogram according to the 2014 paediatric appropriate use criteria guidelines were excluded.⁹ Patients with clinical histories or exam findings not specifically mentioned in the appropriate use criteria guidelines were reviewed by two independent reviewers and excluded from the study if full echocardiogram is deemed indicated by both reviewers.

Targeted echocardiogram views. As per institutional protocol, the parasternal short-axis view at the aortic valve portion was routinely used to diagnose bicuspid aortic valve. This view demonstrates loss of normal triangular opening shape of aortic valve morphology in systole in the presence of a bicuspid valve (Figure 1). Similarly, parasternal short-axis view at the level of the papillary muscles was routinely used to diagnose left ventricular hypertrophy as per the American Society of Echocardiography 2024 guideline recommendations to assess wall thickness in short-axis view by 2D or M-mode (Figure 2).¹⁴

Data collection: Echocardiogram reports were reviewed for findings. Findings other than bicuspid aortic valve or left ventricular hypertrophy were collected as they were likely to be missed if only the targeted echocardiogram view was performed.




Figure 1. Parasternal short axis at the aortic valve view demonstrating loss of normal triangular opening shape in systole in patient with bicuspid aortic valve.




Figure 2. Parasternal short-axis mid-cavity view allows for assessment of end-diastolic posterior and septal wall thicknesses and relative end-diastolic dimension.

We recorded all abnormal findings for both groups documented by the contemporary reader in the echocardiogram reports. If follow-up was recommended for the finding, follow-up clinic visits and echocardiogram reports were reviewed to assess for any disease progression or additional findings. Clinical parameters recorded included symptoms, family history, cardiac medications, physical activity restrictions, need for subacute bacterial endocarditis prophylaxis, catheter-/surgery-based interventions, and/or ongoing cardiology follow-up.

Results

Bicuspid aortic valve group

In the study period, 153 comprehensive screening echocardiograms for family history of bicuspid aortic valve were identified; 15 were excluded (Figure 3). Some common indications that met the "may be appropriate" criteria of the 2014 paediatric appropriate




Figure 3. Flow chart of patient selection in the bicuspid aortic valve group. Patients with family history of bicuspid aortic valve who met the 2014 paediatric appropriate use criteria guidelines for a full echocardiogram based on other indications were excluded.

use criteria for a full echocardiogram and were hence excluded from this study were: family history of other CHD such as coarctation of the aorta; hypoplastic aortic arch; and hypoplastic left heart syndrome. Of note, presumptively innocent murmur with no symptoms, signs, or findings of cardiovascular disease and a benign family history was not an indication for an echocardiogram according to the appropriate use criteria, and of the 138 patients selected, 24 had documented innocent murmurs on exam. No pathologic murmurs were documented. Four of the 138 patients reported intermittent non-exertional chest pain.

Of the final 138 included in the study, 71 (51%) were males and mean age at echo was 8.4 ± 4.8 years. Bicuspid aortic valve was diagnosed by comprehensive echocardiogram in 5 patients (3.6%).

Incidental findings were noted in 15 (11%) of this group, with four of the 15 patients already diagnosed with bicuspid aortic valve. Of the 11 patients without the diagnosis of bicuspid aortic valve, seven had findings that were determined to be clinically insignificant during their cardiology visit, and follow-up was not recommended (Appendix Table A). Follow-up clinic visit and echocardiogram were recommended for the remaining four; findings and management are shown in Table 1.

These incidental findings include mild aortic root or ascending aorta dilation, trivial insufficiency of aortic valve, high and leftward take-off of the right coronary artery, and one case of patent foramen ovale vs. small secundum atrial septal defect. One patient was lost to follow up but three were seen back at clinic and received either comprehensive or limited follow-up echocardiograms.

One patient without bicuspid aortic valve but with incidental finding of mild dilation of ascending aorta was restricted from heavy weightlifting. This patient had family history of paternal grandfather with aortic root dilation in addition to bicuspid aortic valve; there was a lack of clear family history of presence or absence of connective tissue disorder. The rest of the patients without bicuspid aortic valve who required follow up due to incidental findings resulted in no change in management at their follow up visits, including cardiac medications, physical activity restrictions, need for subacute bacterial endocarditis prophylaxis, and catheter-/surgery-based interventions.

Left ventricular hypertrophy group

In the study period, 125 comprehensive screening echocardiograms for left ventricular hypertrophy on electrocardiogram were identified; 54 were excluded (Figure 4). Some indications that met the definitive criteria of the 2014 paediatric appropriate use criteria for a full echocardiogram and were hence excluded from this study were: family history of hypertrophic cardiomyopathy; exertional chest pain or non-exertional chest pain with abnormal electrocardiogram; palpitations with abnormal electrocardiogram; systemic hypertension; marfanoid body habitus on exam; and family history of cardiac death before age 50. Of the 71 patients selected, 17 had documented innocent heart murmurs on exam. Of the final 71 included in the study, 58 (83%) were males and mean age at echo was 10.9 ± 4.7 years. Left ventricular hypertrophy was diagnosed in two patients (2.8%) in this screening group by comprehensive echocardiogram.

Incidental findings were noted in nine (13%) of this group, with two of the nine patients already diagnosed with left ventricular hypertrophy. Of the seven patients without the diagnosis of left ventricular hypertrophy, five had findings that were determined to be clinically insignificant during their cardiology visit, and follow-up was not recommended (Appendix Table B). Follow-up clinic visit and echocardiogram was recommended for the remaining two; findings and management are shown in Table 2.

These echo findings included one incidental finding of bicuspid aortic valve without family history, another with slightly increased trabeculations in the left ventricular apex, and possible left ventricular noncompaction. Both patients followed up in clinic and had a repeat echo, which resulted in no change in management, including cardiac medications, physical activity restrictions, need for subacute bacterial endocarditis prophylaxis, or catheter-/surgery-based interventions.

Discussion


Targeted echocardiography is a potentially effective screening tool for identifying bicuspid aortic valve and left ventricular hypertrophy in select indications. Most importantly, our study shows only a minor percent (2.8% for both groups) had incidental findings that prompted additional follow-up after the initial screening. Only one patient was found to have mild ascending aorta dilation and was recommended exercise restriction in the bicuspid aortic valve group. The rest of the follow-up group had insignificant incidental findings. Overall, our findings suggest that targeted single-view echocardiographic screening for patients for isolated indication of bicuspid aortic valve and left ventricular hypertrophy is unlikely to result in missed significant forms of heart disease.

We demonstrated a low prevalence of bicuspid aortic valve (3.6%) and left ventricular hypertrophy (2.8%) amongst those referred for isolated family history of bicuspid aortic valve and left ventricular hypertrophy on electrocardiogram, respectively. This estimate in the population we studied is slightly lower than reported by some other studies.⁵

Screening for bicuspid aortic valve is clinically important and highly cost-effective. In a study utilising an economic evaluation model performing a cost-effectiveness analysis of screening for first-degree relatives of patients with bicuspid aortic valve, it was found that screening with echocardiogram was less costly than no screening, leading to savings of 208 euros and gain of quality-

Table 1. Actionable incidental findings on comprehensive echocardiogram in patients who screened negative for bicuspid aortic valve

Incidental finding	Additional Family History	Follow-up Timeline	Follow-up Recommendation
Mild dilation of ascending aorta	Aortic root dilation in paternal grandfather; Aortic valve thickening in father;	2 years	Follow-up comprehensive echocardiogram showed stable ascending aorta diameter Avoid any heavy weightlifting. No meds/interventions. Additional follow up in Cardiac Genetics Clinic in 3 years with echo
Trileaflet aortic valve with trivial insufficiency	none	3 years	Lost to follow-up.
The right coronary artery has a high take-off and arises leftward along the right and left aortic cusp commissure. Cannot exclude right coronary artery arising from the left cusp	VSD in younger sister	2 years	A follow-up limited echo was performed with similar right coronary artery description. No meds/restrictions/interventions. Additional follow up recommended in 3-4 years with a CT scan to assess coronary arteries
PFO versus small secundum ASD with small left to right flow	none	2-3 years	Complete echo showed "small interatrial communication" with mild left-to-right shunt. No meds/restrictions/interventions. Additional follow-up recommended in 1 year with echo.

**Figure 4.** Flow chart of patient selection in the left ventricular hypertrophy group. Patients with left ventricular hypertrophy on electrocardiograms but who met the 2014 paediatric appropriate use criteria guidelines for a full echocardiogram based on other indications were excluded.

adjusted life years.¹⁵ The average cost of a comprehensive echocardiogram is much higher compared to a point-of-care cardiac ultrasound. For example, a complete transthoracic echocardiogram at some institutions is billed for \$1952.2 compared to a limited echocardiogram that is billed for \$722.5.¹⁶ Targeted echocardiography could be billed as a limited echocardiogram or point-of-care cardiac ultrasound. In addition to healthcare cost burden of initial screening, incidental findings leading to follow up clinic visits and echocardiograms add to the cost burden of comprehensive screening echocardiograms. Thus, targeted echocardiography may significantly reduce healthcare costs for the screening indications such as bicuspid aortic valve and left ventricular hypertrophy.

Targeted echocardiography may improve availability of care in resource-limited settings as this diagnosis can be mostly made in a single short-axis view image. There is growing evidence suggesting that point-of-care cardiac ultrasound can be learned quickly and performed effectively by non-cardiology medical professionals. Hellmann et al. demonstrated that medical residents learned to perform handheld cardiac ultrasonography at a reasonable rate with minimal formal training followed by one-on-one instruction by a certified technician.¹⁰ Engelman et al found that the quality of targeted cardiac ultrasound performed by a briefly trained health worker is adequate for diagnosis of rheumatic heart disease 96.6% of the time.¹¹ If targeted single-view could be successfully performed by nonexpert scanners with hand-held ultrasound and remote cardiology interpretation, this may greatly improve accessibility for screening echocardiography in low-resource settings. If coupled with artificial intelligence, an emerging tool for disease diagnosis in cardiac imaging, it has the potential to lead to instantaneous, automated diagnosis.¹²

In our study, incidental findings on comprehensive echocardiograms performed for screening for isolated indications of bicuspid aortic valve or left ventricular hypertrophy triggered recommendation for additional cardiology follow-up and testing without a change in management in 93 and 100% of patients with incidental findings in respective groups. While an overall low percentage of clinical significance, these patients may experience anxiety while awaiting results or during observation periods and incur the cost associated with follow-up visits and testing. In a 2019 study on "cascades of care" in adult patients after incidental findings, 99.4% of physicians reported experiencing cascades with resultant patient psychological harm (68.4%), physical harm (15.6%), financial burden (57.5%), frustration (52.5%), and anxiety (45.4%).¹³ Performing targeted single-view for specified indications may decrease both financial and non-financial costs associated with incidental findings on standard echocardiography.

One of the limitations of the study is that we did not examine the specific diagnostic accuracy, specificity, sensitivity, and

Table 2. Actionable incidental findings in patients who screened negative for left ventricular hypertrophy

Incidental finding	Additional Family History	Follow-up Timeline	Follow-up Recommendation
Bicuspid aortic valve	None	2 years	A comprehensive echo was performed and showed normal aortic valve function and no ascending aorta dilation. No meds/restrictions/interventions. Additional follow up in 2 years with echo.
Slightly increased trabeculations in the left ventricular apex; possible left ventricular noncompaction	None	1 year	A follow-up limited echo was performed with no changes compared to prior. No meds/restrictions/interventions. Additional follow up in 2 years with echo.

predictive value of targeted single-view echocardiography for screening for bicuspid aortic valve and left ventricular hypertrophy. In future studies, we recommend assessing the diagnostic accuracy by comparing diagnostic accuracy using single echocardiographic views to the gold standard of complete echocardiograms by expert review.

Conclusion

Our study suggests that the risk of missing clinically important incidental findings identifiable with comprehensive echocardiography is extremely low for screening indications of isolated family history such as bicuspid aortic valve or left ventricular hypertrophy on electrocardiogram. Hence, targeted echocardiography focused on short-axis view is potentially an effective screening tool for those with isolated screening indications. Targeted echocardiography would likely reduce the rate of minor clinically insignificant incidental findings, potentially resulting in reduced financial and non-financial healthcare costs. With established appropriate use criteria, it may improve access to screening in remote and resource-limited areas.

Future studies should focus on prospectively evaluating targeted single-view echocardiography sensitivity amongst various echocardiogram screening indications. Efficacy of protocols with hand-held device for screening for bicuspid aortic valve and left ventricular hypertrophy as well as other indications should be explored. This study should serve as a stepping stone for further development of Artificial Intelligence driven automated and hand-held device technology for cardiac screening.

Acknowledgements. None.

Financial support. This research received no specific grant from any funding agency, commercial, or not-for-profit sectors.

Competing interests. The author(s) declare none.

Ethical standard. The authors assert that all procedures contributing to this work comply with the ethical standards of the relevant national guidelines on human experimentation and with the Helsinki Declaration of 1975, as revised in 2008, and has been approved by the institutional committees Institutional Review Boards at the Seattle Children's Hospital.

References

- Ward C. Clinical significance of the bicuspid aortic valve. *Heart* 2000; 83: 81–85.
- Michelena HI, Desjardins VA, Avierinos JF, et al. Natural history of asymptomatic patients with normally functioning or minimally dysfunctional bicuspid aortic valve in the community. *Circulation* 2008; 117 (21): 2776–2784.
- Jain R, Ammar KA, Kalvin L, et al. Diagnostic accuracy of bicuspid aortic valve by echocardiography. *Echocardiography* 2018; 35 (12): 1932–1938.
- Chan KL, Stinson WA, Veinot JP. Reliability of transthoracic echocardiography in the assessment of aortic valve morphology: pathological correlation in 178 patients. *Can J Cardiol* 1999; 15 (1): 48–52.
- Tague L, Wiggs J, Li Q, et al. Comparison of left ventricular hypertrophy by Electrocardiography and echocardiography in children using analytics tool. *Pediatr Cardiol* 2018; 39 (7): 1378–1388.
- Miller AF, Arichai P, Gravel CA, et al. Use of cardiac point-of-care ultrasound in the pediatric emergency department. *Pediatr Emerg Care* 2022; 38 (1): e300–e305.
- Singh Y, Bhombal S, Katheria A, Tissot C, Fraga MV. The evolution of cardiac point of care ultrasound for the neonatologist. *Eur J Pediatr* 2021; 180 (12): 3565–3575.
- Neale D, Williams D, Canty D. Benefit analysis of focused pre-operative transthoracic echocardiography in the pre-operative clinic in patients at increased cardiac risk presenting for non-cardiac surgery - a pilot study. *Int J Anesthetic Anesthesiology* 2015; 2: 023.
- Campbell RM, Douglas PS, Eidem BW, Lai WW, Lopez L, Sachdeva R. ACC/AAP/AHA/ASE/HRS/SCAI/SCCT/SCMR/SOPE 2014 appropriate use criteria for initial transthoracic echocardiography in outpatient pediatric cardiology: a report of the American College of Cardiology Appropriate Use Criteria Task Force, American Academy of Pediatrics, American Heart Association, American Society of Echocardiography, Heart Rhythm Society, Society for Cardiovascular Angiography and Interventions, Society of Cardiovascular Computed Tomography, Society for Cardiovascular Magnetic Resonance, and Society of Pediatric Echocardiography. *J Am Coll Cardiol* 2014; 64 (19): 2039–2060.
- Hellmann DB, Whiting-O'Keefe Q, Shapiro EP, Martin LD, Martire C, Ziegelstein RC. The rate at which residents learn to use hand-held echocardiography at the bedside. *Am J Med* 2005; 118 (9): 1010–1018.
- Engelman D, Kado JH, Reményi B, et al. Screening for rheumatic heart disease: quality and agreement of focused cardiac ultrasound by briefly trained health workers. *BMC Cardiovasc Disord* 2016; 16 (30): 30.
- Barry T, Farina JM, Chao C-J, et al. The role of artificial intelligence in echocardiography. *J Imaging* 2023; 9 (2): 50.
- Ganguli I, Simpkin AL, Lupo C, et al. Cascades of care after incidental findings in a US national survey of physicians. *JAMA Netw Open* 2019; 2 (10): e1913325.
- Lopez L, Saurers DL, Piers CA, Barker PCA, et al. Guidelines for performing a comprehensive pediatric transthoracic echocardiogram: recommendations from the American society of echocardiography. *J Am Soc Echocardiogr* 2024; 37 (2): 119–170. doi: [10.1016/j.echo.2023.11.015](https://doi.org/10.1016/j.echo.2023.11.015).
- Tessler I, Leshno M, Shmueli A, Shpitz S, Durst R, Gilon D. Cost-effectiveness analysis of screening for first-degree relatives of patients with bicuspid aortic valve. *Eur Heart J Qual Care Clin Outcomes* 2021; 7 (5): 447–457. doi: [10.1093/ehjqco/qcab047](https://doi.org/10.1093/ehjqco/qcab047).
- Insurance and cost of care. Seattle Children's Hospital. Accessed December 28, 2024. <http://www.seattlechildrens.org/clinics/paying-for-care/insurance>.