

Comparison of standardised dietary folate intake across ten countries participating in the European Prospective Investigation into Cancer and Nutrition

Jin Young Park^{1*}, Genevieve Nicolas¹, Heinz Freisling¹, Carine Biessy¹, Augustin Scalbert¹, Isabelle Romieu¹, Véronique Chajès¹, Shu-Chun Chuang², Ulrika Ericson³, Peter Wallström⁴, Martine M. Ros^{5,6}, Petra H. M. Peeters⁷, Amalia Mattiello⁸, Domenico Palli⁹, José María Huerta^{10,11}, Pilar Amiano^{11,12}, Jytte Halkjær¹³, Christina C. Dahm^{14,15}, Antonia Trichopoulou^{16,17}, Philippos Orfanos¹⁶, Birgit Teucher¹⁸, Silke Feller¹⁹, Guri Skeie²⁰, Dagrun Engeset²⁰, Marie-Christine Boutron-Ruault²¹, Françoise Clavel-Chapelon²¹, Francesca Crowe²², Kay-Tee Khaw²³, Paolo Vineis² and Nadia Slimani¹

¹Section of Nutrition and Metabolism, International Agency for Research on Cancer, 150 cours Albert-Thomas, 69372 Lyon Cedex 08, France

²School of Public Health, Imperial College London, London, UK

³Diabetes and Cardiovascular Disease, Genetic Epidemiology, Department of Clinical Sciences, Lund University, Malmö, Sweden

⁴Nutrition Epidemiology Research Group, Department of Clinical Sciences, Lund University, Malmö, Sweden

⁵National Institute for Public Health and the Environment, Bilthoven, The Netherlands

⁶Department of Epidemiology, Biostatistics and HTA, Radboud University Nijmegen Medical Centre, Nijmegen, The Netherlands

⁷Julius Centre, University Medical Centre Utrecht, Utrecht, The Netherlands

⁸Department of Clinical and Experimental Medicine, Federico II University, Naples, Italy

⁹Molecular and Nutritional Epidemiology Unit, Cancer Research and Prevention Institute, Florence, Italy

¹⁰Department of Epidemiology, Murcia Regional Health Authority, Murcia, Spain

¹¹CIBER Epidemiología y Salud Pública, Spain

¹²Public Health Division of Gipuzkoa, Institute Investigation IIS BioDonostia, Basque Country Region, Spain

¹³Danish Cancer Society, Institute of Cancer Epidemiology, Copenhagen, Denmark

¹⁴Department of Cardiology, Aalborg Hospital, Aarhus University Hospital, Aarhus, Denmark

¹⁵Department of Epidemiology, School of Public Health, Aarhus University, Aarhus, Denmark

¹⁶WHO Collaborating Centre for Food and Nutrition Policies, Department of Hygiene, Epidemiology and Medical Statistics, University of Athens Medical School, Athens, Greece

¹⁷Hellenic Health Foundation, Athens, Greece

¹⁸German Cancer Research Centre, Heidelberg, Germany

¹⁹Department of Epidemiology, German Institute of Human Nutrition Potsdam-Rehbruecke, Nuthetal, Germany

²⁰Department of Community Medicine, University of Tromsø, Tromsø, Norway

²¹Inserm, ERI 20, Institut Gustave Roussy, Villejuif, France

²²Cancer Epidemiology Unit, Nuffield Department of Clinical Medicine, University of Oxford, Oxford, UK

²³Department of Public Health and Primary Care, University of Cambridge, Cambridge, UK

(Submitted 24 May 2011 – Final revision received 22 August 2011 – Accepted 20 September 2011 – First published online 1 November 2011)

Abbreviations: 24-HDR, 24h dietary recall; ENDB, European Prospective Investigation into Cancer Nutrient DataBase; EPIC, European Prospective Investigation into Cancer and Nutrition.

* **Corresponding author:** J. Y. Park, fax +33 4 72 73 83 61, email parkjy@fellows.iarc.fr

Abstract

Folate plays an important role in the synthesis and methylation of DNA as a cofactor in one-carbon metabolism. Inadequate folate intake has been linked to adverse health events. However, comparable information on dietary folate intake across European countries has never been reported. The objective of the present study was to describe the dietary folate intake and its food sources in ten countries in the European Prospective Investigation into Cancer and Nutrition (EPIC) study. A cross-sectional analysis was conducted in 36 034 participants (aged 35–74 years) who completed a single 24 h dietary recall using a computerised interview software program, EPIC-Soft[®] (International Agency for Research on Cancer, Lyon). Dietary folate intake was estimated using the standardised EPIC Nutrient DataBase, adjusted for age, energy intake, weight and height and weighted by season and day of recall. Adjusted mean dietary folate intake in most centres ranged from 250 to 350 $\mu\text{g}/\text{d}$ in men and 200 to 300 $\mu\text{g}/\text{d}$ in women. Folate intake tended to be lower among current smokers and heavier alcohol drinkers and to increase with educational level, especially in women. Supplement users (any types) were likely to report higher dietary folate intake in most centres. Vegetables, cereals and fruits, nuts and seeds were the main contributors to folate intake. Nonetheless, the type and pattern of consumption of these main food items varied across the centres. These first comparisons of standardised dietary folate intakes across different European populations show moderate regional differences (except the UK health conscious group), and variation by sex, educational level, smoking and alcohol-drinking status, and supplement use.

Key words: Dietary folate intake: 24 h Dietary recall: European Prospective Investigation into Cancer and Nutrition: EPIC-Soft[®]: Europe

Folate is a water-soluble B vitamin, which plays an important role in the synthesis and methylation of DNA together with other B vitamins including vitamin B₂, vitamin B₆ and vitamin B₁₂ as crucial cofactors in one-carbon metabolism⁽¹⁾. Dietary inadequacies of these nutrients can contribute to negative health outcomes. For example, inadequate folate intake has been linked to the risk of anaemia⁽²⁾, neuropsychiatric disorders⁽³⁾ and neural tube defects⁽⁴⁾. It has also been shown that insufficient intakes of folate, vitamin B₆ or vitamin B₁₂ are associated with elevated plasma homocysteine concentrations, a potential risk factor for CVD^(5,6). Furthermore, folate deficiency leads to the disruption of DNA synthesis, repair and methylation⁽⁷⁾, which may increase the risk of developing some cancers, notably colorectal cancer⁽⁸⁾.

Foods rich in folate include vegetables, especially green leafy vegetables, such as spinach and beet leaf, and dried peas and lentils as well as fruits, nuts and seeds⁽⁹⁾. Animal sources include offal such as beef and chicken liver⁽⁹⁾. The synthetic form of the vitamin, folic acid, is widely used for the purpose of food fortification and food supplements. In some countries, folic acid fortification of many cereal-based food products became mandatory, e.g. in the USA⁽¹⁰⁾, Canada⁽¹¹⁾ and Chile⁽¹²⁾, after consideration of compelling evidence for a protective effect of periconceptual folic acid supplementation against neural tube defects⁽¹³⁾. However, so far, no European countries have introduced mandatory folic acid fortification, although voluntary fortification is accepted in some countries including France and the UK⁽¹⁴⁾.

Despite the suggested beneficial effects of folate on many different diseases, there has been growing concern regarding possible adverse effects of excessive levels of folate intake^(13,15–17). Increased folic acid intake has been shown to have the potential for masking the diagnosis of a vitamin B₁₂ deficiency, particularly in the elderly^(18,19). In addition, animal and human studies have reported that a high dose of folic acid may promote the progression of already existing premalignant and malignant lesions, particularly in colorectal cancer settings^(20–22). It is therefore important to monitor folate status of those groups of people who are at an increased risk of cancer and who are more likely to be exposed to

higher doses of folic acid intakes⁽²³⁾ besides those who are at a risk of low folate intake.

Deharveng *et al.*⁽²⁴⁾ reported earlier a lack of clarity and consistency in the terminology and definitions used for folate information in the food composition tables available in nine countries participating in the European Prospective Investigation into Cancer and Nutrition (EPIC) study. More recently, an evaluation of folate data in eighteen European and international databases concluded that a lack of comparability still exists between countries⁽¹⁴⁾. Consequently, studies that estimated dietary folate intakes across European countries to date relied on each country's own food composition data which tend to be heterogeneous^(25–27). Thus, there is a great need to develop a more accurate and standardised database to enhance comparability and consistency of folate data across Europe.

In the absence of a reference European nutrient database, the EPIC Nutrient DataBase (ENDB) project was initiated to harmonise separate nutrient databases using common procedures and guidelines, with support from the local national compilers in ten countries in the EPIC⁽²⁸⁾. The ENDB was first completed for twenty-six priority components to provide a standardised reference instrument for calibrating the EPIC dietary measurements at the nutrient level⁽²⁹⁾. This work has been extended to cover other nutrients including folate, and has just been completed⁽³⁰⁾. In the present descriptive study, we present and compare the levels and food sources of dietary folate intake obtained by means of a single 24 h dietary recall (24-HDR) collected from a representative sample of the EPIC cohort using standardised folate concentration data, recently compiled as an extension of the ENDB.

Materials and methods

Study population, design and dietary assessment

The rationale and methods of the EPIC study have previously been described in detail^(31–33). The EPIC cohort consists of twenty-three subcohorts in ten European countries (Denmark, France, Greece, Germany, Italy, The Netherlands, Norway,

Spain, Sweden and UK), providing a wide range of cancer occurrence rates, lifestyle and dietary habits. EPIC participants were mostly recruited from the general population residing within defined geographical areas between 1992 and 2000, with some exceptions: women members of a health insurance for school employees (France); women attending breast cancer screening (Utrecht, The Netherlands); blood donors (some centres in Italy and Spain) and a cohort consisting predominantly of vegetarians ('health conscious' cohort in Oxford, UK). In France, Norway, the Utrecht centre of The Netherlands and in the Naples centre of Italy, all participants were women. For the purposes of the present study, the initial twenty-three EPIC administrative centres have been redefined into twenty-seven geographical regions relevant to the analysis of dietary consumption patterns⁽³⁴⁾. Approval for the study was obtained from the ethical review boards of the International Agency for Research on Cancer (Lyon, France) and from all local recruiting institutes. All EPIC participants provided informed consent.

Within the design of the EPIC study, a subsample of each study centre was randomly (age, sex stratified) chosen for the application of a standardised 24-HDR assessment gathered by means of a computerised software (EPIC-Soft[®]; International Agency for Research on Cancer, Lyon, France)^(35,36). This subcohort is referred to as the EPIC Calibration substudy and was undertaken between 1995 and 2000. Each participant provided a single 24-HDR in a face-to-face interview⁽³⁵⁾, except in Norway where it was obtained by a telephone interview⁽³⁷⁾. In total, complete 24-HDR information was available from 36 994 participants (13 486 men and 23 508 women), representing approximately 8% of the entire EPIC cohort. This sample has been shown to be a reasonably representative sample of the entire EPIC cohort⁽³⁴⁾. A total of 36 034 participants with 24-HDR data were included in the present analysis, after exclusion of 960 participants under 35 or over 74 years of age because of low participation in these age categories. Using the EPIC-Soft[®], information on the intake of all foods and beverages was collected, described, quantified and coded according to common rules. The classification of the EPIC-Soft[®] food groups and food subgroups used in the calibration study is derived from a system described in detail elsewhere⁽³⁴⁾.

Dietary folate intake was estimated using the updated ENDB⁽³⁰⁾. Although the ENDB values were obtained from country-specific food composition tables, they were standardised as much as possible across the EPIC countries by matching of the EPIC foods to the national databases according to the recommendation from the recent review⁽¹⁴⁾. In particular, a microbiological assay was chosen as the reference analytical method for folate values in the ENDB. Folate values of unavailable foods were derived by values either from recipe calculation or borrowed from similar foods⁽³⁰⁾. During the ENDB compilation for folate, particular attention has been given to the issue of fortification of breakfast cereals, particularly in the UK and France as their cereal consumption was substantially higher compared with other EPIC countries. In Scandinavian countries and in The Netherlands, fortification was not allowed at the time of data collection. In other EPIC countries, breakfast cereal consumption was very low and

the information on folic acid-fortified foods was not always available⁽¹⁴⁾. It was therefore decided not to adopt the dietary folate equivalent conversion, which considers different bioavailability of naturally occurring folate and synthetic folic acid.

Data on age as well as body weight and height in most centres were self-reported by the participants during the 24-HDR interview. Data on other lifestyle factors, including total physical activity, educational level, smoking history and alcohol intake, considered in the present analysis were collected at baseline through standardised questionnaires and clinical examinations, and have been described elsewhere^(31,34,38–40). The physical activity questions being asked in the Umeå (Sweden) and Norwegian centres were different from the EPIC core questions⁽⁴⁰⁾ and the information was omitted from the analyses of the present study. The mean time interval between these baseline questionnaire measures and the 24-HDR interview varied by country, from 1 d to 3 years after⁽³⁴⁾.

The folate data described here come from food intakes only, as our main interest is in dietary folate levels in European populations (not from dietary supplements). However, we present the level of dietary folate intake of those who reported being dietary supplement users compared with those who did not, as they may differ from each other with regard to dietary characteristics⁽⁴¹⁾. Dietary supplement information was obtained at the end of the 24-HDR with a question, 'Did you take any dietary supplements?' with yes or no answers, followed by further questions on the name of the supplements, the frequency of use on the recalled day and the number of units taken per consumption occasion⁽⁴²⁾. For the present study, a dietary supplement user was defined as any subject who reported taking at least one dietary supplement on the recalled day. A folic acid-containing supplement user was defined as any subject who reported taking at least one dietary supplement containing folic acid on the recalled day. Nevertheless, it should be noted that quantitative nutrient data derived from food supplements were not available for the present study.

Statistical methods

Folate intake ($\mu\text{g}/\text{d}$) was calculated as least square means and standard errors by sex, age (10-year categories from 35–74 years) and EPIC centre (ordered according to a geographical south–north gradient). The mean intake was adjusted for age, total energy intake, weight and height and weighted by season of the 24-HDR collection (spring, summer, autumn and winter) and day of the week of recall (Monday to Thursday; Friday to Sunday) to control for different sampling procedures of the 24-HDR interviews across seasons and days of the week.

Stratified analyses were performed to describe differences in folate intake levels according to level of physical activity (inactive, moderately inactive, moderately active and active; data completeness > 86%), smoking status (never smoker, former smoker and current smoker; completeness > 98%), daily alcohol intake (abstainers, < 12 g/d, 12 to < 24 g/d and \geq 24 g/d,



Table 1. Mean intake of folate ($\mu\text{g/d}$)* by centre ordered from south to north, sex and age group†
(Number of participants, mean values with their standard errors)

Country and centre	Men											Women												
	n	All		35–44		45–54		55–64		65–75		P_{trend}	n	All		35–44		45–54		55–64		65–75		P_{trend}
		Mean	SE	Mean	SE	Mean	SE	Mean	SE	Mean	SE			Mean	SE	Mean	SE	Mean	SE	Mean	SE	Mean	SE	
Greece	1311	330.2	3.7	291.5	11.0	309.2	7.5	342.2	6.8	350.8	6.0	0.02	1373	257.6	3.1	238.1	8.1	256.8	5.2	264.6	5.4	261.2	6.2	0.16
Spain																								
Granada	214	309.1	9.0	–‡	–‡	340.3	18.8	300.8	12.1	303.2	20.2	0.61	300	259.2	6.4	275.0	16.4	274.4	10.7	237.5	10.0	265.0	20.4	0.51
Murcia	243	346.3	8.4	405.4	26.1	341.6	15.1	327.4	11.7	393.2	29.2	0.83	304	264.5	6.3	264.8	12.7	269.3	10.6	254.4	10.5	–‡	–‡	0.66
Navarra	444	316.8	6.3	277.8	27.0	299.1	10.4	332.6	8.9	319.2	19.1	0.15	271	253.7	6.7	208.4	17.5	268.0	10.9	255.6	10.2	–‡	–‡	0.41
San Sebastian	490	313.5	6.0	304.2	14.1	310.0	8.2	317.5	11.3	248.4	29.8	0.35	244	258.3	7.0	241.7	15.0	268.1	11.4	257.1	11.8	–‡	–‡	0.98
Asturias	386	304.9	6.7	269.7	25.1	284.7	11.2	316.0	10.0	334.8	18.1	0.01	324	240.1	6.1	236.3	14.8	225.6	9.9	244.4	9.8	290.7	22.4	0.18
Italy																								
Ragusa	168	300.3	10.1	–‡	–‡	309.0	15.0	286.3	15.8	–‡	–‡	0.51	138	219.1	9.4	175.5	15.5	214.8	17.4	258.2	16.7	–‡	–‡	0.005
Naples													403	229.7	5.5	205.6	17.8	219.6	8.7	247.5	8.5	213.3	17.8	0.64
Florence	271	301.1	7.9	275.4	25.0	313.9	13.6	297.8	11.3	–‡	–‡	0.79	784	255.2	3.9	273.5	13.2	251.5	6.8	256.1	5.4	240.5	15.3	0.11
Turin	676	304.8	5.1	301.7	16.3	286.5	8.4	317.1	7.2	287.3	19.2	0.89	392	255.9	5.5	251.1	17.5	261.9	9.2	253.4	7.7	–‡	–‡	0.32
Varese	327	263.1	7.2	–‡	–‡	306.4	16.1	252.7	8.7	262.9	24.3	0.84	794	222.6	3.9	207.5	12.5	212.1	6.5	239.7	5.9	200.0	11.8	0.96
France																								
South coast													620	286.9	4.4	–‡	–‡	282.4	7.3	300.2	6.9	273.4	9.2	0.79
South													1425	280.7	2.9	–‡	–‡	279.1	4.5	289.6	4.6	268.3	6.5	0.66
North-east													2059	274.9	2.5	–‡	–‡	267.4	3.8	281.2	3.8	280.2	5.6	0.37
North-west													631	276.6	4.4	–‡	–‡	287.2	6.9	275.5	6.6	257.6	10.5	0.08
Germany																								
Heidelberg	1034	270.5	4.1	241.6	10.9	265.3	6.5	276.2	6.0	–‡	–‡	0.01	1087	234.2	3.4	237.2	5.7	221.6	6.1	238.4	5.6	–‡	–‡	0.27
Potsdam	1233	258.9	3.7	237.6	10.7	243.2	7.5	269.9	4.9	242.2	14.5	0.64	1061	229.2	3.4	219.7	6.7	230.3	6.6	232.1	5.0	221.6	21.3	0.85
The Netherlands																								
Bilthoven	1024	292.9	4.2	273.5	8.0	295.5	6.4	287.1	7.1	–‡	–‡	0.40	1086	242.3	3.4	231.1	5.9	241.4	5.2	249.9	6.4	–‡	–‡	0.01
Utrecht													1870	259.3	2.6	–‡	–‡	255.8	4.4	260.7	3.9	266.3	5.2	0.03
United Kingdom																								
General population	402	344.1	6.5	331.5	21.4	364.2	11.6	336.7	11.9	337.4	11.8	0.91	570	285.3	4.6	281.1	13.8	281.9	7.5	295.2	8.3	280.1	9.8	0.81
Health conscious	114	479.1	12.3	–‡	–‡	448.1	20.0	503.6	18.9	–‡	–‡	0.40	197	362.5	7.8	331.1	24.7	367.7	12.8	378.7	12.3	320.9	21.6	0.91
Denmark																								
Copenhagen	1356	296.6	3.6	–‡	–‡	290.5	5.8	301.7	4.6	283.7	17.8	0.76	1484	249.7	2.9	–‡	–‡	252.9	4.7	248.5	3.7	252.9	13.5	0.99
Aarhus	567	304.0	5.5	–‡	–‡	304.2	7.8	303.0	7.8	–‡	–‡	0.32	510	258.6	4.9	–‡	–‡	266.1	6.9	252.5	7.0	–‡	–‡	0.11
Sweden																								
Malmö	1421	235.6	3.7	–‡	–‡	238.9	10.4	245.2	5.5	240.1	4.9	0.88	1711	201.1	2.7	–‡	–‡	198.6	5.5	208.3	4.4	200.9	4.2	0.85
Umeå	1344	252.9	3.6	242.9	12.1	255.0	6.7	255.6	4.9	250.6	10.6	0.48	1574	217.3	2.8	211.9	6.7	218.2	4.9	221.5	4.2	206.5	8.9	0.76
Norway																								
South and East													1004	227.4	3.5	218.0	8.4	226.2	4.3	232.5	8.6	–‡	–‡	0.05
North and West													793	214.5	4.0	200.0	8.9	214.9	4.8	219.5	10.2	–‡	–‡	0.19

Dietary folate intake in European countries

* Adjusted for age (when not stratified for age), total energy intake, weight and height and weighted by season and day of recall.
 † P_{trend} was derived from allocating scores 1, 2, 3 and 4 for all participants in the age group of 35–44, 45–54, 55–64 and 65–75 years, respectively.
 ‡ If fewer than twenty persons are present in a certain age group, mean intake is not presented.

as presented in an earlier report⁽⁴³⁾; no missing data), educational level (none/primary, technical/secondary, university or higher; completeness > 98%) and season (spring, summer, autumn and winter; no missing data). When analyses were stratified by season, the mean intake of folate was not weighted by season. Further analyses were stratified by dietary supplement use (yes, no). In the stratified analysis, sex- and centre-specific mean intakes are presented across variables of interest in main tables. Tests for trend were performed by allocating scores for participants in each category of lifestyle factors. *P* values were derived from *t* tests for continuous variables. The main food groups contributing to folate intake are presented as the mean percentage of intake (percentage of total intake, derived from the unadjusted 24-HDR data); the contribution of a subgroup was given as a percentage of the food group. The categorisation into food groups and food subgroups is common across the centres and is adapted from the EPIC-Soft[®] food classification system as described elsewhere^(36,44). All analyses were conducted using SAS statistical software (version 9.1, SAS Institute, Cary, NC, USA).

Results

Intake of folate

Table 1 shows the adjusted mean folate intake ($\mu\text{g}/\text{d}$) presented by each centre and age categories at recruitment in men and women. Daily folate intake in all centres, except in the UK health conscious group, ranged from 250 to

350 $\mu\text{g}/\text{d}$ in men and 200 to 300 $\mu\text{g}/\text{d}$ in women. Intake was higher in centres in Spain, France and in the UK general population and was relatively low in the Swedish centres, especially in Malmö in both men and women, but with no obvious geographical gradient (Fig. 1). In the UK health conscious group, where participants are mainly vegetarians or vegans, folate intake was markedly higher, averaging approximately 480 $\mu\text{g}/\text{d}$ in men and approximately 360 $\mu\text{g}/\text{d}$ in women. There was neither a consistent nor a strong trend observed in folate intake across age categories, although it tended to increase with age in men from Greece, Asturias (Spain) and Heidelberg (Germany) and women from Ragusa (Italy), The Netherlands and South and East Norway. Additional adjustment for smoking status and alcohol consumption did not materially alter the results in men and women (data not shown).

Intake of folate stratified by lifestyle factors, educational level and season

Tables 2 and 3 present adjusted folate intake in each centre according to lifestyle factors in men and women. There were no systematic differences observed in the levels of folate intake when participants were stratified according to different physical activity levels (Table 2). Compared with those with lower educational level, participants with university or higher degree were likely to report higher folate intake, with the tendency being clearer in women (Table 2). Folate intake varied according to smoking status and level of alcohol




Fig. 1. Mean intake of folate ($\mu\text{g}/\text{d}$) in men (■) and women (▒), stratified by centre ordered from south to north, adjusted for age, total energy intake, weight and height and weighted by season and day of recall.

Table 2. Mean intake of folate ($\mu\text{g/d}$)* by centre, according to levels of physical activity and education†
(Mean values with their standard errors)

Country and centre	Physical activity																		
	Men									Women									
	Inactive		Moderately inactive		Moderately active		Active		P_{trend}	Inactive		Moderately inactive		Moderately active		Active		P_{trend}	
	Mean	SE	Mean	SE	Mean	SE	Mean	SE		Mean	SE	Mean	SE	Mean	SE	Mean	SE		
Greece	326.1	9.6	318.6	6.2	343.8	6.0	326.4	13.7	0.69	243.2	12.1	254.2	7.2	258.0	4.0	264.4	9.8	0.02	
Spain																			
Granada	298.4	20.4	319.8	16.7	298.8	15.1	330.7	26.5	0.39	277.4	44.9	257.0	24.1	256.3	7.1	311.3	35.4	0.49	
Murcia	346.0	17.0	331.8	15.6	367.3	15.9	334.3	23.3	0.99	233.7	27.4	281.9	16.6	257.8	7.7	320.0	27.0	0.18	
Navarra	311.8	14.7	295.0	12.0	320.5	10.5	354.0	16.3	0.21	223.2	29.6	230.2	20.2	259.7	7.9	239.1	32.1	0.37	
San Sebastian	301.6	12.9	323.7	12.6	311.6	9.9	320.8	15.4	0.41	299.0	26.7	264.7	16.8	250.1	9.3	262.9	22.0	0.24	
Asturias	301.2	16.4	293.7	12.1	311.8	12.1	315.2	15.9	0.21	259.2	27.9	247.6	17.8	235.3	7.4	253.9	21.8	0.65	
Italy																			
Ragusa	303.9	19.4	263.4	18.8	318.9	18.5	350.5	34.3	0.30	285.0	33.6	226.5	22.5	213.6	12.3	176.2	31.7	0.03	
Naples										247.5	15.6	227.6	9.2	225.3	8.9	225.8	22.2	0.19	
Florence	336.2	15.8	281.4	14.1	296.1	14.8	286.4	27.5	0.30	261.8	12.8	262.8	8.4	252.6	5.4	239.1	14.6	0.08	
Turin	311.2	10.3	304.7	8.9	290.4	9.4	332.4	16.4	0.64	262.1	17.0	254.2	13.1	260.8	7.7	227.8	16.5	0.22	
Varese	286.9	19.3	248.8	12.1	254.4	12.0	318.3	24.9	0.60	209.2	13.3	230.8	9.5	222.7	5.1	214.8	13.5	0.88	
France																			
South coast										301.4	12.0	287.1	5.8	275.5	10.6	275.8	29.1	0.07	
South										278.2	7.8	280.7	3.8	285.8	7.3	247.8	21.4	0.35	
North-east										280.8	6.0	270.6	3.2	280.5	6.2	311.9	20.9	0.26	
North-west										288.9	12.0	274.9	5.6	273.3	11.3	259.8	36.2	0.04	
Germany																			
Heidelberg	254.1	9.4	276.3	7.6	271.4	6.6	281.9	13.2	0.16	238.3	8.9	241.2	6.3	228.0	5.3	231.7	10.8	0.30	
Potsdam	250.6	10.2	263.6	8.1	259.5	5.2	256.0	12.0	0.71	238.0	7.7	223.4	6.5	227.4	5.2	240.2	17.0	0.83	
The Netherlands																			
Bilthoven	280.2	12.1	291.8	9.0	293.8	6.3	299.9	9.6	0.04	242.4	12.4	252.8	7.6	238.4	4.7	241.9	8.8	0.67	
Utrecht										263.9	11.2	263.0	5.5	258.2	3.7	256.1	6.1	0.03	
United Kingdom																			
General population	326.5	17.0	365.0	12.6	329.3	10.7	359.4	16.8	0.59	286.9	12.0	293.1	8.7	279.1	7.3	289.0	14.4	0.83	
Health conscious	431.7	32.4	413.9	23.8	510.0	19.6	574.2	33.0	0.09	364.2	16.8	345.4	16.0	373.4	12.0	339.0	39.0	0.62	
Denmark																			
Copenhagen	293.1	7.1	303.2	7.1	286.0	6.7	309.9	9.3	0.60	251.3	5.6	246.1	5.0	249.8	5.5	259.9	11.7	0.35	
Aarhus	303.8	11.4	300.4	10.5	310.0	9.7	294.4	15.8	0.63	253.8	10.4	248.6	8.3	269.9	8.9	273.0	19.9	0.15	
Sweden																			
Malmö	233.5	8.4	227.2	5.7	243.1	6.3	253.9	18.3	0.15	203.9	6.6	199.3	4.4	201.4	4.6	203.6	13.1	0.92	
Umeå																			
Norway																			
South and East																			
North and West																			

Dietary folate intake in European countries

Country and centre	Education													
	Men							Women						
	Primary		Secondary/ technical		University		P_{trend}	Primary		Secondary/ technical		University		P_{trend}
	Mean	SE	Mean	SE	Mean	SE		Mean	SE	Mean	SE	Mean	SE	
Greece	339.7	5.4	324.5	7.9	317.5	6.6	0.13	268.4	4.2	241.8	6.2	241.3	6.2	0.32
Spain														
Granada	302.2	12.0	331.4	22.5	301.2	17.1	0.98	251.2	6.8	297.5	22.9	364.0	29.1	0.07
Murcia	338.8	9.9	331.1	26.3	380.2	19.4	0.43	269.8	7.3	218.5	20.0	266.1	15.5	0.96
Navarra	310.6	7.4	339.6	13.5	297.6	23.2	0.80	258.1	7.3	200.8	22.1	244.9	23.9	0.86
San Sebastian	304.5	7.6	323.9	10.9	342.1	19.4	0.01	261.3	8.2	242.8	15.5	273.2	25.4	0.75
Asturias	312.9	8.4	283.2	14.0	290.8	17.9	0.49	239.4	6.8	243.5	17.9	228.2	21.4	0.50
Italy														
Ragusa	293.6	14.9	300.1	16.1	312.4	25.5	0.11	237.7	13.3	185.7	14.7	253.4	26.9	0.86
Naples								213.7	8.4	233.2	8.4	257.4	13.4	0.04
Florence	278.9	12.8	315.2	11.8	319.4	19.2	0.27	253.6	5.6	253.7	6.5	261.1	9.7	0.33
Turin	299.5	8.1	305.7	6.9	321.1	16.5	0.15	250.8	7.3	257.8	9.7	275.2	16.6	0.15
Varese	252.0	10.4	274.4	10.4	260.8	34.1	0.75	220.1	4.8	220.7	7.5	245.3	14.2	0.32
France														
South coast								238.2	12.1	305.6	6.2	276.2	7.5	0.62
South								274.9	9.0	282.6	4.1	280.0	4.9	0.55
North-east								271.1	6.9	271.6	3.5	280.5	3.9	0.30
North-west								263.1	11.9	276.5	5.8	289.3	8.1	0.01
Germany														
Heidelberg	268.6	6.9	268.7	6.8	276.0	7.4	0.32	208.2	6.4	240.5	4.7	254.7	6.9	0.14
Potsdam	244.3	8.2	254.4	6.6	268.1	5.3	0.05	222.9	6.6	231.1	4.8	232.9	6.7	0.22
The Netherlands														
Bilthoven	291.2	11.2	288.6	5.4	304.0	7.9	0.43	247.5	8.9	237.9	4.2	257.2	7.0	0.66
Utrecht								238.3	5.1	262.3	3.3	280.7	6.6	0.05
United Kingdom														
General population	339.4	17.9	355.8	9.5	334.0	13.3	0.85	287.2	10.8	286.5	6.8	274.7	10.9	0.30
Health conscious	–	–	512.0	23.9	455.3	17.0		–	–	381.7	13.1	351.2	11.2	
Denmark														
Copenhagen	283.8	6.6	298.5	5.6	307.1	6.4	0.09	231.3	5.5	253.9	3.6	268.4	7.9	0.08
Aarhus	297.0	9.7	305.1	8.3	312.0	10.9	0.03	241.0	8.9	265.8	6.1	270.7	16.2	0.24
Sweden														
Malmö	229.4	5.3	235.0	6.2	248.9	7.3	0.15	188.5	4.2	205.4	4.3	213.6	5.6	0.13
Umeå	238.0	5.9	256.0	5.5	272.7	7.7	0.01	199.6	5.2	215.4	4.1	238.7	5.2	0.07
Norway														
South and East								211.5	8.5	228.7	4.3	247.8	8.6	0.02
North and West								208.5	8.3	218.1	4.8	215.9	11.0	0.48

J. Y. Park *et al.*

* Adjusted for age, total energy intake, weight and height and weighted by season and day of recall.

† P_{trend} was derived by allocating scores for participants in each category of lifestyle factors.

Table 3. Mean intake of folate ($\mu\text{g/d}$)* by centre, according to smoking and alcohol status†
(Mean values with their standard errors)

Country and centre	Smoking													
	Men							Women						
	Never		Former		Current		P_{trend}	Never		Former		Current		P_{trend}
	Mean	SE	Mean	SE	Mean	SE		Mean	SE	Mean	SE	Mean	SE	
Greece	336.3	6.9	338.1	6.2	316.6	6.5	0.38	261.2	3.6	255.8	12.0	246.0	7.1	0.11
Spain														
Granada	329.5	15.3	301.3	13.8	295.2	18.1	0.23	259.6	6.9	259.6	28.0	257.8	20.3	0.34
Murcia	371.0	15.3	349.4	15.4	325.5	13.2	0.02	269.6	7.2	252.6	23.3	249.9	15.8	0.25
Navarra	330.7	10.5	307.3	14.3	310.2	9.1	0.41	261.9	7.9	254.9	19.7	220.8	15.8	0.23
San Sebastian	319.1	10.4	330.9	11.7	297.0	9.2	0.56	257.6	8.3	264.5	18.6	254.3	18.9	0.80
Asturias	321.0	11.3	299.7	11.5	292.9	11.8	0.18	242.7	7.0	238.0	23.1	229.5	14.7	0.11
Italy														
Ragusa	298.2	18.5	290.3	14.6	322.3	21.2	0.49	231.2	14.4	210.3	20.7	209.5	15.2	0.31
Naples								237.3	7.8	233.5	12.2	215.9	9.9	0.23
Florence	283.3	15.3	309.3	11.5	302.9	16.1	0.48	253.1	5.6	269.9	8.0	246.2	7.4	0.82
Turin	302.7	9.6	305.4	7.6	303.1	9.8	0.91	258.9	7.1	255.2	11.7	245.8	13.5	0.16
Varese	250.0	12.1	271.9	11.4	266.3	14.9	0.49	229.0	4.7	213.7	10.5	202.2	9.8	0.05
France														
South coast								294.0	5.1	277.6	10.9	266.5	17.7	0.07
South								285.4	3.5	269.0	6.4	270.1	11.3	0.37
North-east								272.1	2.9	282.2	5.7	289.9	9.4	0.05
North-west								280.4	5.3	272.9	9.4	261.1	15.9	0.08
Germany														
Heidelberg	267.5	7.2	285.1	6.0	243.9	8.6	0.61	248.1	4.8	222.0	5.9	216.8	7.4	0.23
Potsdam	267.7	6.9	261.0	5.3	242.8	7.9	0.17	230.4	4.3	238.1	6.8	207.1	8.8	0.49
The Netherlands														
Bilthoven	306.7	8.4	295.2	6.6	280.4	6.9	0.04	256.5	5.7	243.2	5.9	225.8	5.8	0.05
Utrecht								268.7	3.8	260.1	4.4	237.8	5.8	0.16
United Kingdom														
General population	348.1	10.8	350.3	9.7	326.3	15.7	0.39	285.0	5.9	291.2	8.6	265.3	14.7	0.48
Health conscious	463.6	19.3	479.6	19.1	509.1	28.2	0.11	370.6	9.6	350.1	13.8	280.1	51.5	0.20
Denmark														
Copenhagen	303.3	6.5	296.1	6.8	291.6	6.3	0.08	256.5	4.1	256.0	5.5	228.8	5.8	0.32
Aarhus	302.5	10.7	319.6	8.9	288.0	9.2	0.70	268.5	7.1	255.8	9.2	243.3	9.5	0.004
Sweden														
Malmö	239.3	6.3	242.6	5.4	222.6	7.0	0.43	203.1	3.9	207.7	5.0	192.0	5.5	0.52
Umeå	261.5	5.0	249.1	6.5	235.5	8.8	0.02	221.0	3.4	216.2	6.7	204.9	6.7	0.15
Norway														
South and East								227.6	5.8	230.4	6.0	219.5	6.8	0.49
North and West								220.5	6.7	218.1	6.7	200.5	7.5	0.26

Dietary folate intake in European countries

Country and centre	Alcohol															
	Men									Women						
	Abstainers		1–12 g		12.1–24 g		> 24 g		P_{trend}	Abstainers		1–12 g		> 12 g		P_{trend}
	Mean	SE	Mean	SE	Mean	SE	Mean	SE		Mean	SE	Mean	SE	Mean	SE	
Greece	348.3	6.0	338.7	8.6	326.7	9.4	308.7	6.5	0.01	269.4	4.0	245.0	5.4	247.0	8.9	0.38
Spain																
Granada	332.6	20.5	333.2	16.1	313.3	22.1	273.2	15.0	0.09	264.3	8.5	264.5	10.2	205.6	24.3	0.33
Murcia	391.1	19.6	367.2	17.0	290.1	21.9	333.2	12.7	0.26	284.9	10.8	263.5	9.3	234.4	13.7	0.06
Navarra	340.2	14.5	354.9	13.6	317.1	15.8	289.0	9.1	0.14	279.3	8.7	232.3	12.0	183.0	19.8	0.01
San Sebastian	358.2	15.0	314.1	14.3	312.3	15.1	297.7	8.2	0.09	286.1	11.3	257.5	10.7	205.2	15.8	0.11
Asturias	333.4	13.5	318.2	15.1	320.2	17.4	275.9	10.2	0.12	244.8	8.5	241.7	10.4	223.4	15.6	0.25
Italy																
Ragusa	289.8	18.4	311.2	19.6	289.5	27.0	304.1	18.2	0.75	219.0	13.0	237.5	16.2	184.0	23.2	0.55
Naples										234.6	8.8	229.1	9.6	226.4	10.1	0.12
Florence	309.6	18.4	301.5	16.1	325.1	17.0	281.3	13.0	0.57	264.7	6.6	259.4	6.1	236.1	7.9	0.22
Turin	342.7	13.2	305.9	12.5	310.2	11.9	291.7	6.9	0.11	276.1	10.4	253.0	8.8	243.2	9.6	0.15
Varese	308.9	20.3	271.8	16.4	258.0	19.7	249.5	9.7	0.05	242.3	6.8	216.5	5.8	208.3	8.2	0.19
France																
South coast										305.5	7.3	279.4	7.7	270.6	7.8	0.18
South										295.5	4.7	287.8	5.3	253.9	5.2	0.22
North-east										290.0	4.2	271.1	4.4	261.7	4.0	0.12
North-west										304.2	7.7	275.1	7.9	252.9	7.0	0.05
Germany																
Heidelberg	293.8	10.3	273.3	8.6	288.9	9.9	254.6	6.0	0.26	254.2	7.2	235.9	5.4	221.3	5.2	0.04
Potsdam	263.3	8.2	261.9	7.9	251.0	7.7	259.0	6.3	0.44	241.4	6.3	223.7	5.0	226.6	6.4	0.43
The Netherlands																
Bilthoven	300.9	7.0	304.0	13.1	281.9	10.5	285.8	6.5	0.20	256.6	4.7	232.8	8.0	223.2	5.9	0.15
Utrecht										269.5	3.7	250.7	6.2	248.5	4.3	0.27
United Kingdom																
General population	343.9	10.7	340.0	15.6	366.9	16.9	337.7	11.5	0.92	302.7	6.7	292.7	10.0	258.0	7.9	0.20
Health conscious	515.0	18.0	379.1	29.4	485.2	38.9	482.3	23.2	0.98	381.2	11.5	376.7	16.1	322.3	14.0	0.29
Denmark																
Copenhagen	315.7	7.5	311.5	14.0	298.8	8.8	286.1	4.8	0.02	250.7	4.6	272.2	8.4	243.2	4.0	0.84
Aarhus	322.1	10.0	298.4	25.8	308.4	13.3	290.5	7.8	0.20	265.2	7.5	252.4	13.8	251.8	7.1	0.31
Sweden																
Malmö	238.6	5.6	240.7	6.8	242.8	9.2	222.1	7.8	0.35	200.3	3.7	203.9	5.1	200.5	5.8	0.96
Umeå	249.7	4.8	265.3	7.5	250.0	11.3	247.9	10.0	0.67	218.1	3.4	222.4	5.8	204.4	7.6	0.48
Norway																
South and East										231.8	4.3	229.2	9.6	211.0	7.4	0.26
North and West										215.2	4.6	218.0	11.7	208.0	9.2	0.51

* Adjusted for age, total energy intake, weight and height and weighted by season and day of recall.
 † P_{trend} was derived by allocating scores for participants in each category of lifestyle factors.

intake (Table 3). Though not entirely consistent in all centres, the mean intake of folate was lower among current smokers and among heavier alcohol drinkers. No systematic variations were observed for folate intake according to the season of 24-HDR collection, although folate intake in southern European countries seemed lowest in summer (see Table S1 of the supplementary material, available online at <http://www.journals.cambridge.org/bjn>).

Dietary supplement use and folate intake

On average, approximately 22% of men and 34% of women reported using any type of dietary supplements during the recalled day, of which 33 and 25%, respectively, reported using folic acid-containing supplements (Table 4). Total dietary folate intake was generally higher in users of any type of supplement in most centres. An exception to this was men in the UK health conscious group, where more than 50% of the participants were supplement users. The proportion of folic acid-containing supplement users was low, especially in participants from the southern European countries (Table 4). Dietary folate intake among those who consumed folic acid-containing supplements varied across the countries.

Common dietary sources of folate intake

Table 5 shows the main food groups contributing to the intake of folate in men and women in each country. Among all the food groups, vegetables, cereals and fruits, nuts and seeds were the main contributors of dietary folate intake in all centres (Fig. 2 and see Table S2 of the supplementary material, available online at <http://www.journals.cambridge.org/bjn>). On the one hand, vegetables in Greece and Italy accounted for approximately 30% of folate intake in men and more than 30% in women. There was a clear north–south difference in the source of vegetable contribution to folate intake, with leafy and fruiting vegetables being the major vegetable source of folate in the southern countries and cabbages and root vegetables being the major vegetable source in the northern countries (Fig. 3).

On the other hand, cereals were the most important dietary source of folate intake in both men and women in the UK general population, where breakfast cereals accounted for 43% of cereal and cereal products as source of folate (Fig. 4). This was quite distinctive from other centres where bread was the main source of folate intake among cereal products, contributing more than 80%. An exception to this was Italy, where pasta, rice and dough such as pizza together with bread were the major cereal-based source of folate intake.

Both non-alcoholic (e.g. fruit and vegetable juices) and alcoholic beverages (e.g. wine and beer) were also some of the main sources of folate intake in the northern countries (Table 5). In the UK health conscious group, the condiments and sauces and miscellaneous food groups (consisting primarily of special vegan/vegetarian food items including soya-based products and non-dairy cheeses) were found to account for 15% of folate in men and 13% in women while

these food groups contribute less than 2% of intake in all other centres (Fig. 2).

Discussion

In the present study, we compared standardised dietary folate intake across ten European countries according to selected lifestyle factors and major contributing food sources. Dietary folate intake did not differ greatly between centres apart from the UK centres, especially the UK health conscious group where a much higher intake was observed compared with other centres. The average dietary folate intake observed in the present study without the UK health conscious group was 307 $\mu\text{g}/\text{d}$ for men and 252 $\mu\text{g}/\text{d}$ for women, after adjusting for participants' age, total energy intake, weight and height and the effect of season and day of recall.

There have been few published data that investigated the level of dietary folate intake at the national level across European countries. A study by de de Bree *et al.*⁽²⁵⁾ reported mean dietary folate intake in 18–64-year-old adults of 291 $\mu\text{g}/\text{d}$ (range 197–326) for men and 247 $\mu\text{g}/\text{d}$ (range 168–320) for women in nine European countries by reviewing their national food consumption surveys. In 2005, a final report on the European project on folate estimated that average dietary folate intakes in participants aged 18–90 years were 283 $\mu\text{g}/\text{d}$ (range 218–352) for men and 238 $\mu\text{g}/\text{d}$ (range 207–284) for women in eight European countries on the basis of each country's food consumption and food composition data⁽²⁷⁾. It is not clear in these studies whether folate values were adjusted for energy intake and other covariates. More recently, the European Nutrition and Health Report has summarised that dietary folate equivalent (i.e. 1 μg food folate = 0.5 μg folic acid = 0.6 μg folic acid taken with meals, values unadjusted) ranged from 203 to 494 $\mu\text{g}/\text{d}$ in men and 131 to 392 $\mu\text{g}/\text{d}$ in women aged 19–64 years in twenty-one participating countries⁽²⁶⁾.

While these studies provide an overview on the folate status in European countries attempting to use nationally representative data, none of these used a standardised nutrient database across countries, limiting the comparability of the studies included in each report due to the use of different dietary assessment methods, different years and periods of data collection and different age classifications. It has also been pointed out that quantification methods, terminologies and mode of expression used in folate data varied across food composition tables in different countries in the absence of a reference European nutrient database⁽¹⁴⁾. We describe for the first time the dietary folate intake in ten European countries using a folate database that has been standardised across those countries.

In the present study, we observed a relatively high dietary folate intake in both of the UK centres. In the UK, folic acid has been added to most breakfast cereals since 1987, with the amount of added folic acid being further increased in 1994, followed by a considerable increase in dietary folate intake⁽⁴⁵⁾ as well as in blood folate levels⁽⁴⁶⁾. Considering that cereal and cereal products, notably breakfast cereals, were the main contributor of folate intake, high folate

Table 4. Mean intake of folate ($\mu\text{g/d}$)* by country, according to use of supplements (any types† and folic acid-containing supplement‡)§
(Number of participations, mean values with their standard errors)

	Men							Women							
	Yes			No				P	Yes			No			
	Folate intake			Folate intake					Folate intake			Folate intake			
	n	Mean	SE	n	Mean	SE	n		Mean	SE	n	Mean	SE	P	
Supplement user (any types)															
Greece	26	410.8	27.3	1285	329.0	3.8	0.003	92	259.7	11.1	1281	257.7	3.2	0.86	
Spain	105	330.6	12.9	1672	314.5	3.3	0.23	174	266.0	8.3	1269	253.4	3.1	0.15	
Italy	96	309.1	13.1	1346	292.4	3.6	0.22	316	254.1	6.3	2195	237.0	2.4	0.01	
France								1524	283.6	2.9	3211	276.2	2.0	0.03	
Germany	469	284.3	6.0	1798	258.2	3.1	<0.001	580	233.5	4.8	1568	231.0	2.8	0.65	
The Netherlands	164	304.4	10.2	860	288.8	4.6	0.16	949	265.7	3.6	2007	246.8	2.5	<0.001	
United Kingdom															
General population	146	359.1	11.0	256	336.6	8.1	0.10	271	302.8	6.7	299	269.7	6.3	<0.001	
Health conscious	59	440.6	18.5	55	509.1	16.4	0.006	102	366.9	11.1	95	358.2	11.1	0.58	
Denmark	980	301.9	4.3	943	296.2	4.2	0.33	1312	259.1	3.1	682	238.4	4.2	<0.001	
Sweden	843	254.2	4.6	1922	241.5	3.1	0.02	1392	211.3	3.0	1893	207.1	2.5	0.27	
Norway								1088	225.3	3.5	709	215.9	4.1	0.07	
Folic acid-containing supplement user															
Greece	1	668.1		1310	330.3	3.7	–	2	329.5	62.6	1371	257.8	3.1	–	
Spain	3	195.3	74.5	1774	316.1	3.2	–	8	239.7	35.6	1435	255.3	3.0	–	
Italy	2	267.3	104.8	1440	293.8	3.5	–	5	269.7	54.0	2506	239.1	2.2	–	
France								121	286.8	9.7	4614	278.1	1.7	0.38	
Germany	7	267.7	51.6	2260	263.8	2.8	–	22	237.4	25.4	2126	231.8	2.4	0.83	
The Netherlands	40	299.3	20.4	984	291.4	4.3	0.71	174	260.3	8.3	2782	252.5	2.1	0.36	
United Kingdom															
General population	21	320.2	27.7	381	345.7	6.7	0.37	43	296.6	16.5	527	284.3	4.8	0.48	
Health conscious	12	516.6	40.4	102	475.0	12.9	0.33	14	323.0	26.7	183	366.1	8.2	0.12	
Denmark	608	303.6	5.5	1315	296.9	3.6	0.30	773	258.8	4.0	1221	247.5	3.1	0.02	
Sweden	255	245.8	8.4	2510	244.7	2.7	0.91	494	212.6	5.0	2791	208.0	2.1	0.39	
Norway								285	225.6	6.5	1512	220.9	2.9	0.51	

* Adjusted for age, total energy intake, weight and height and weighted by season and day of recall.

† Any type of supplement user was defined as any subject who reported taking at least one dietary supplement on the recalled day.

‡ Folic acid-containing supplement user was defined as any subject who reported taking at least one dietary supplement containing folic acid on the recalled day.

§ P_{value} was derived from *t* tests and is not presented for comparisons with fewer than ten persons.

Table 5. Percentage contribution of main food groups to the intake of folate by country in men and women

Country	Dietary food groups (percentage contribution of each food group to total intake of the nutrient)												Total
	Potatoes/ other tubers	Vegetables	Legumes	Fruits, nuts and seeds	Dairy products	Cereal and cereal products	Meat and meat products	Fish/ shellfish	Egg and egg products	Sugar, confectionery/ cakes, biscuits	Non alcoholic beverages	Alcoholic beverages	
Men													
All men	8.1	20.8	3.1	10.0	10.6	20.4	7.0	1.6	2.3	3.9	4.7	4.2	96.6
Greece	4.5	29.7	6.8	9.9	8.5	27.7	2.9	1.6	1.5	1.2	2.3	1.5	98.0
Spain	6.3	24.6	11.4	14.0	6.4	12.6	6.7	3.3	4.3	2.2	2.2	1.9	96.0
Italy	4.0	30.0	1.8	12.4	6.1	24.9	7.4	1.2	1.5	3.9	1.7	1.4	96.2
Germany	8.0	19.3	0.4	10.7	10.7	20.1	6.1	0.7	1.8	4.0	7.1	7.7	96.5
The Netherlands	8.4	19.2	0.4	9.9	13.1	16.4	10.7	0.7	2.0	2.3	7.0	6.3	96.5
United Kingdom	7.8	17.8	2.7	7.8	8.2	26.3	2.7	0.8	1.2	3.4	9.9	3.0	91.6
General population	8.5	17.4	2.0	6.0	9.8	27.2	3.4	1.0	1.4	3.7	10.7	3.4	94.6
Health conscious	5.9	18.8	4.6	12.7	3.9	24.0	0.8	0.1	0.6	2.4	7.9	1.8	83.6
Denmark	8.2	14.8	0.2	6.5	15.9	21.1	11.6	1.7	2.3	4.5	5.4	6.1	98.4
Sweden	14.4	14.5	0.6	8.0	13.4	19.4	5.8	1.6	2.5	7.5	4.9	3.9	96.5
Women													
All women	6.1	25.2	1.4	12.7	13.6	16.3	5.4	1.4	2.1	4.2	6.9	1.0	96.3
Greece	4.2	33.6	5.0	11.2	10.3	22.9	2.1	1.4	1.6	2.0	3.9	0.5	98.5
Spain	5.8	27.0	6.9	18.3	9.7	9.9	4.9	2.7	3.4	3.2	3.3	0.8	95.9
Italy	3.2	32.0	2.0	15.6	8.1	20.1	5.1	1.0	1.8	4.8	2.4	0.5	96.6
France	4.6	28.5	1.0	12.8	16.4	12.6	6.2	1.5	1.8	3.3	5.8	0.6	95.1
Germany	6.7	25.6	0.2	13.4	12.8	16.9	3.8	0.6	1.8	4.1	8.9	1.7	96.9
The Netherlands	6.9	21.6	0.4	12.4	16.5	13.1	7.4	0.6	2.1	2.9	12.0	0.9	96.7
United Kingdom	6.4	22.4	2.6	9.7	9.3	22.7	2.2	0.9	1.2	3.5	11.0	0.6	92.6
General population	6.9	21.1	2.0	8.7	10.9	24.6	3.0	1.0	1.5	3.6	11.7	0.6	95.6
Health conscious	5.4	25.3	3.9	12.2	5.7	18.5	0.4	0.5	0.7	3.3	9.4	0.5	85.8
Denmark	6.2	19.9	0.2	10.7	15.9	20.8	6.3	1.5	2.3	4.8	7.5	2.1	98.3
Sweden	10.5	19.7	0.4	11.4	13.7	16.2	4.9	1.6	2.6	7.7	5.9	1.6	96.2
Norway	6.8	20.3	0.2	10.9	12.5	19.4	6.0	2.2	2.6	5.6	9.8	1.2	97.5

Dietary folate intake in European countries

intake in the UK centres may be partly explained by voluntary fortification of breakfast cereals. Indeed, in the other EPIC countries, breakfast cereals were not a big contributor of folate intake, as they are neither much consumed nor widely fortified. In addition to the relatively high cereal consumption, the vegetarian diet practised in the UK health

conscious group resulted in an exceptionally high dietary folate intake compared with the other centres included in the present study. Vegetarian (legume/soya-based) dishes are a good source of folate and the consumption of these food items (categorised as miscellaneous foods in the present study) in this health conscious group was four (e.g. com-




Fig. 2. Percentage contribution of all food groups (■, miscellaneous; □, soups, bouillons; ▨, sugar and confectionery; ▩, cakes and biscuits; ▪, condiments and sauces; ■, egg and egg products; □, fish and shellfish; ▨, alcoholic beverages; ▩, non-alcoholic beverages; ▪, potatoes and other tubers; ▫, meat and meat products; ▬, legumes; ▭, dairy products; ▮, fruits, nuts and seeds; ▯, cereal and cereal products; ▰, vegetables) to the intake of folate by centre ordered from south to north in (a) men and (b) women.




Fig. 3. Percentage contribution of vegetable sources (■ mixed salad, mixed vegetables; ▨ stalk vegetables, sprouts; □ grain and pod vegetables; ▤ mushrooms; ▥ onion, garlic; ▦ root vegetables; ▧ cabbages; ▨ fruiting vegetables; ■ leafy vegetables) to the intake of folate by country ordered from south to north in (a) men and (b) women.

pared with The Netherlands or France) to forty times (compared with Greece) as high as the one in other EPIC countries (data not shown).

We also showed some different patterns in the dietary folate intake between the southern and northern European countries. While the intake of folate did not differ statistically

significantly according to the season that the 24-HDR was collected, folate intake in Greece, Spain and Italy appeared to be lower in summer compared with the northern countries. This may be partly due to the fact that the green leafy vegetables, one of the major dietary source of folate in those countries, are broadly known as an early-spring or




Fig. 4. Percentage contribution of cereal sources (□, salty biscuits, aperitif biscuits, crackers; ■, flour, flakes, starches, semolina; ▨, dough and pastry (puff, short-crust, pizza); ▩, pasta, rice, other grains; ▤, breakfast cereals; ■, bread, crispbread, rusks) to the intake of folate by country ordered from south to north in (a) men and (b) women.

a late-autumn crop⁽⁴⁷⁾. Furthermore, both alcoholic and non-alcoholic beverages contributed to a greater proportion of folate intake in the northern countries. It should be noted that there was an approximately 10-fold difference in non-alcoholic beverage consumption between the southern and northern countries (data not shown). A higher proportion of folate intake was accounted for by alcoholic beverages

in countries such as Germany, The Netherlands, the UK and Denmark, where beer, a source of folate, is the main source of alcohol consumption⁽⁴³⁾.

Men, in general, in the present study had a substantially higher folate intake than did women. This may be attributed to the fact that men tend to have larger body size and to consume more food compared with women. These differences in

folate intake between men and women seemed slightly larger (approximately 45%) in the southern European centres compared with those in the northern centres. This may reflect to some extent a different dietary pattern which is specific to each sex and to each centre. For example, consumption of vegetables, especially leafy vegetables, one of the biggest source of dietary folate, was much higher in men compared with women in the southern countries such as Greece, Spain and Italy, while little difference by sex was observed in northern countries⁽⁴⁸⁾. Similarly, consumption of cereal, meat and certain alcoholic beverages, which were also the sources of folate intake, was considerably higher in men from the southern countries compared with women (data not shown). In the case of beer and wine consumption, there was up to a 5-fold difference in the level of consumption in men and women in those countries⁽⁴³⁾.

The present results also showed that folate intake tended to be lower among current smokers and heavier alcohol drinkers. Previous studies have found that smokers have lower plasma and erythrocyte folate concentrations compared with non-smokers^(49–52) even after adjusting for dietary folate intake^(50,52). Similarly, heavy alcohol drinkers have been observed to have lower circulating concentration of folate^(53,54) as well as lower dietary folate intake⁽⁵⁵⁾. Chronic ethanol intake has been shown to interfere with folate metabolism^(54,56). This indicates that smokers and heavy alcohol drinkers whose folate bioavailability is low and who may have additionally lower dietary folate intake are at an increased risk of suboptimal folate status and therefore need to be carefully monitored.

To our knowledge, this is the first and largest study that evaluated dietary folate intake using standardised methods across European countries. Dietary folate intake in the present study was derived from the ENDB, a nutrient database that has been standardised across all the countries participating in the EPIC study⁽²⁸⁾. In addition, the 24-HDR was validated against independent biomarkers in the EPIC^(57,58) and the interview procedures were standardised, with interviewers receiving substantial training in the use of the software⁽³⁶⁾. This comparison also benefits from the unique setting offered by the EPIC, a Europe-wide study with heterogeneity in dietary habits and in other lifestyle factors across countries⁽³³⁾.

However, some limitations of the present study need to be discussed. First, in the present study, only one set of 24-HDR was collected from a large population sample with the ultimate purpose to have a good and sufficiently valid estimate of mean population intakes, and this approach was not designed to provide true long-term individual intakes due to the lack of information on intra-individual variability⁽⁵⁹⁾ or intake distribution (several repeated 24-HDR would have been required for this aim). Second, the 24-HDR was undertaken between 1995 and 2000 and may not reflect the most up-to-date information on folate intake in Europe. For example, voluntary fortification of foods with folic acid has been more widely practised in the European Union since 2006 under the regulation on the addition of vitamins and minerals to foods⁽⁶⁰⁾. Nonetheless, the present study still provides valid and important information on the level of dietary

folate intake in European countries before voluntary fortification was widespread. Further, this standardised information across European countries improves our understanding on the differences in dietary folate exposure across European countries and its relationship to cancer and other chronic diseases through studies carried out within the EPIC.

Another limitation is that not all of the EPIC centres applied population-based sampling, and thus a direct comparison of the present results to the general population of each region should be made with caution⁽³¹⁾. Nevertheless, the subsample used for the present study has been shown to be representative of the entire EPIC cohort⁽³⁴⁾.

In the present study, we did not consider the differential bioavailability of food folate and folic acid added to fortified foods using the dietary folate equivalent conversion. However, apart from the UK and France, folic acid fortification was not a common practice in Europe at the time of the data collection, resulting in considerably low consumption of the folic acid-fortified foods in those countries. Similarly, as mentioned earlier, the contribution of folic acid added to supplements has not been taken into account in estimating total folate intakes because although we had data for types of supplements consumed, quantitative data on dietary supplements were not available⁽⁴²⁾. Nonetheless, folic acid-containing supplements were not one of the most frequently consumed types of supplements in this study⁽⁴²⁾. Moreover, the primary aim of the present study was to investigate dietary folate intake from food sources in European populations with diverse dietary practices. Given the controversial role of high folic acid intakes, older adults, especially those who may have pre-existing lesions or be at risk for a marginal vitamin B₁₂, should be carefully examined in total amounts of folate uptake and should be encouraged to meet the folate requirement by food sources.

In summary, we described dietary folate intake across the EPIC cohorts in ten European countries using a recently developed standardised nutrient database. The present results showing sex- and region-specific differences in folate intake may serve as a good basis for epidemiological research of cancer and other chronic diseases in relation to folate status. Considering that any dietary intake assessment methods are associated with measurement errors to some extent⁽⁶¹⁾, future priorities should be given to the use of biomarker data of folate status, in addition to the standardised dietary data, in the beneficial context of international settings.

Acknowledgements

We thank all of the study participants for their cooperation and all of the persons who participated in the fieldwork studies in each EPIC centre. We particularly thank Jérôme Vignat for his expertise in the compilation of the folate nutrient database. This study was undertaken during a tenure of a Postdoctoral Fellowship from the International Agency for Research on Cancer, partially supported by the European Commission FP7 Marie Curie Actions – People – Co-funding of regional, national and international programmes (J. Y. P.), and was supported by the World Cancer Research Fund (grant no. 2008/

51), and by the 'Europe Against Cancer' Programme of the European Commission. The cohort studies included in the EPIC study received funding from the following: Public Health and Consumer Protection Directorate 1993–2004; Research Directorate-General 2005; Ligue contre le Cancer (France); Société 3M (France); Mutuelle Générale de l'Education Nationale; Institut National de la Santé et de la Recherche Médicale (INSERM); Institut Gustave Roussy; German Cancer Aid; German Cancer Research Centre; German Federal Ministry of Education and Research; Danish Cancer Society; Health Research Fund of the Spanish Ministry of Health; Spanish Regional Governments of Andalucía, Asturias, Basque Country, Murcia and Navarra and the Catalan Institute of Oncology; ISCIII RETIC (RD06/0020), Spain; Cancer Research UK; Medical Research Council, UK; Stavros Niarchos Foundation and the Hellenic Health Foundation (Greece); Italian Association for Research on Cancer; Italian National Research Council, Regione Sicilia (Sicilian government); Associazione Iblea per la Ricerca Epidemiologica – ONLUS (Hyblean association for epidemiological research, NPO); Dutch Ministry of Public Health, Welfare and Sports; Dutch Prevention Funds; LK Research Funds; Dutch ZON (Zorg Onderzoek Nederland); World Cancer Research Fund (WCRF); Swedish Cancer Society; Swedish Research Council; Regional Government of Skane and the County Council of Vasterbotten, Sweden; Norwegian Cancer Society; the Norwegian Research Council and the Norwegian Foundation for Health and Rehabilitation. The authors' responsibilities were as follows: J. Y. P., G. N., H. F., I. R., A. S., C. B., V. C., S.-C. C., P. V. and N. S. constituted the writing group, conducted the statistical analysis and prepared the manuscript; G. N. compiled the folate nutrient database; N. S. supervised the preparation of the manuscript; all other authors contributed to and/or supervised the collection and analysis of dietary data and/or provided comments and suggestions on the intermediate and final manuscripts. All authors read and approved the final manuscript. None of the authors had a personal or financial conflict of interest.

References

1. Kim YI (2004) Folate and DNA methylation: a mechanistic link between folate deficiency and colorectal cancer? *Cancer Epidemiol Biomarkers Prev* **13**, 511–519.
2. Carmel R (2008) Nutritional anemias and the elderly. *Semin Hematol* **45**, 225–234.
3. Stanger O, Fowler B, Piertz K, *et al.* (2009) Homocysteine, folate and vitamin B₁₂ in neuropsychiatric diseases: review and treatment recommendations. *Expert Rev Neurotherapeutics* **9**, 1393–1412.
4. Wolff T, Witkop CT, Miller T, *et al.* (2009) Folic acid supplementation for the prevention of neural tube defects: an update of the evidence for the U.S. Preventive Services Task Force. *Ann Intern Med* **150**, 632–639.
5. Homocysteine Studies Collaboration (2002) Homocysteine and risk of ischemic heart disease and stroke. *JAMA* **288**, 2015–2022.
6. Ntaios G, Savopoulos C, Grekas D, *et al.* (2009) The controversial role of B-vitamins in cardiovascular risk: an update. *Arch Cardiovasc Dis* **102**, 847–854.

7. Blount BC, Mack MM, Wehr CM, *et al.* (1997) Folate deficiency causes uracil misincorporation into human DNA and chromosome breakage: implications for cancer and neuronal damage. *Proc Natl Acad Sci U S A* **94**, 3290–3295.
8. Kim YI (2006) Folate: a magic bullet or a double edged sword for colorectal cancer prevention? *Gut* **55**, 1387–1389.
9. Gnagnarella P, Salvini S & Parpinel M (2008) Food Composition Database for Epidemiological Studies in Italy, version 1. <http://www.ieo.it/bda> (accessed December 2010).
10. Food and Drug Administration (1996) Food standards: amendment of standards of identity for enriched grain products to require addition of folic acid. Final rule. 21 CFR Parts 136, 137, and 139. *Fed Regist* **61**, 8781–8807.
11. De Wals P, Tairou F, Van Allen MI, *et al.* (2007) Reduction in neural-tube defects after folic acid fortification in Canada. *N Engl J Med* **357**, 135–142.
12. Hirsch S, de la Maza P, Barrera G, *et al.* (2002) The Chilean flour folic acid fortification program reduces serum homocysteine levels and masks vitamin B-12 deficiency in elderly people. *J Nutr* **132**, 289–291.
13. Kim YI (2004) Will mandatory folic acid fortification prevent or promote cancer? *Am J Clin Nutr* **80**, 1123–1128.
14. Bouckaert KP, Slimani N, Nicolas G, *et al.* (2010) Critical evaluation of folate data in European and international databases: recommendations for standardization in international nutritional studies. *Mol Nutr Food Res* **55**, 166–180.
15. Osterhues A, Holzgreve W & Michels KB (2009) Shall we put the world on folate? *Lancet* **374**, 959–961.
16. Shane B (2003) Folate fortification: enough already? *Am J Clin Nutr* **77**, 8–9.
17. Hubner RA, Houlston RD & Muir KR (2007) Should folic acid fortification be mandatory? No. *BMJ* **334**, 1253.
18. Morris MS, Jacques PF, Rosenberg IH, *et al.* (2007) Folate and vitamin B-12 status in relation to anemia, macrocytosis, and cognitive impairment in older Americans in the age of folic acid fortification. *Am J Clin Nutr* **85**, 193–200.
19. Reynolds E (2006) Vitamin B₁₂, folic acid, and the nervous system. *Lancet Neurol* **5**, 949–960.
20. Cole BF, Baron JA, Sandler RS, *et al.* (2007) Folic acid for the prevention of colorectal adenomas. *JAMA* **297**, 2351–2359.
21. Kim YI (2004) Folate, colorectal carcinogenesis, and DNA methylation: lessons from animal studies. *Environ Mol Mutagen* **44**, 10–25.
22. Ulrich CM & Potter JD (2007) Folate and cancer-timing is everything. *JAMA* **297**, 2408–2409.
23. Ulrich CM & Potter JD (2006) Folate supplementation: too much of a good thing? *Cancer Epidemiol Biomarkers Prev* **15**, 189–193.
24. Deharveng G, Charrondiere UR, Slimani N, *et al.* (1999) Comparison of nutrients in the food composition tables available in the nine European countries participating in EPIC. European Prospective Investigation into Cancer and Nutrition. *Eur J Clin Nutr* **53**, 60–79.
25. de Bree A, van Dusseldorp M, Brouwer IA, *et al.* (1997) Folate intake in Europe: recommended, actual and desired intake. *Eur J Clin Nutr* **51**, 643–660.
26. Elmadfa I (volume editor) (2009) *European Nutrition and Health Report*. Basel: Karger.
27. Finglas PM (2005) Folate: From Food to Functionality and Optimal Health. Folate Func Health. <http://www.ifr.ac.uk/folate/> (accessed December 2010).
28. Slimani N, Deharveng G, Unwin I, *et al.* (2007) The EPIC nutrient database project (ENDB): a first attempt to standardize nutrient databases across the 10 European countries participating in the EPIC study. *Eur J Clin Nutr* **61**, 1037–1056.

29. Slimani N & Margetts B (editors) (2009) Nutrient intake and patterns in the EPIC cohorts from 10 European countries. *Eur J Clin Nutr* **63**, Suppl. 4, S1–S2.
30. Nicolas G, Withöft, Vignat J, *et al.* (2011) Standardization of folate database for international nutritional research: report of experience from the EPIC study. (In the Press).
31. Riboli E, Hunt KJ, Slimani N, *et al.* (2002) European Prospective Investigation into Cancer and Nutrition (EPIC): study populations and data collection. *Public Health Nutr* **5**, 1113–1124.
32. Riboli E & Kaaks R (1997) The EPIC Project: rationale and study design. European Prospective Investigation into Cancer and Nutrition. *Int J Epidemiol* **26**, S6–S14.
33. Bingham S & Riboli E (2004) Diet and cancer – the European Prospective Investigation into Cancer and Nutrition. *Nat Rev Cancer* **4**, 206–215.
34. Slimani N, Kaaks R, Ferrari P, *et al.* (2002) European Prospective Investigation into Cancer and Nutrition (EPIC) calibration study: rationale, design and population characteristics. *Public Health Nutr* **5**, 1125–1145.
35. Slimani N, Deharveng G, Charrondiere RU, *et al.* (1999) Structure of the standardized computerized 24-h diet recall interview used as reference method in the 22 centres participating in the EPIC project. European Prospective Investigation into Cancer and Nutrition. *Comput Methods Programs Biomed* **58**, 251–266.
36. Slimani N, Ferrari P, Ocke M, *et al.* (2000) Standardization of the 24-hour diet recall calibration method used in the European Prospective Investigation into Cancer and Nutrition (EPIC): general concepts and preliminary results. *Eur J Clin Nutr* **54**, 900–917.
37. Brustad M, Skeie G, Braaten T, *et al.* (2003) Comparison of telephone vs face-to-face interviews in the assessment of dietary intake by the 24 h recall EPIC SOFT program – the Norwegian calibration study. *Eur J Clin Nutr* **57**, 107–113.
38. Friedenreich C, Cust A, Lahmann PH, *et al.* (2007) Anthropometric factors and risk of endometrial cancer: the European Prospective Investigation into Cancer and Nutrition. *Cancer Causes Control* **18**, 399–413.
39. Haftenberger M, Lahmann PH, Panico S, *et al.* (2002) Overweight, obesity and fat distribution in 50- to 64-year-old participants in the European Prospective Investigation into Cancer and Nutrition (EPIC). *Public Health Nutr* **5**, 1147–1162.
40. Haftenberger M, Schuit AJ, Tormo MJ, *et al.* (2002) Physical activity of subjects aged 50–64 years involved in the European Prospective Investigation into Cancer and Nutrition (EPIC). *Public Health Nutr* **5**, 1163–1177.
41. Kirk SF, Cade JE, Barrett JH, *et al.* (1999) Diet and lifestyle characteristics associated with dietary supplement use in women. *Public Health Nutr* **2**, 69–73.
42. Skeie G, Braaten T, Hjartaker A, *et al.* (2009) Use of dietary supplements in the European Prospective Investigation into Cancer and Nutrition calibration study. *Eur J Clin Nutr* **63**, S226–S238.
43. Sieri S, Agudo A, Kesse E, *et al.* (2002) Patterns of alcohol consumption in 10 European countries participating in the European Prospective Investigation into Cancer and Nutrition (EPIC) project. *Public Health Nutr* **5**, 1287–1296.
44. Slimani N, Fahey M, Welch A, *et al.* (2002) Diversity of dietary patterns observed in the European Prospective Investigation into Cancer and Nutrition (EPIC) project. *Public Health Nutr* **5**, 1311–1328.
45. Henderson L, Gregory J, Irving K, *et al.* (2003) *The National Diet & Nutrition Survey: Adults Aged 19 to 64 years. Vitamin and Mineral Intake and Urinary Analytes*. London: TSO.
46. Clarke R, Sherliker P, Hin H, *et al.* (2008) Folate and vitamin B₁₂ status in relation to cognitive impairment and anaemia in the setting of voluntary fortification in the UK. *Br J Nutr* **100**, 1054–1059.
47. Caron D & Walker D (2011) *University of Delaware Cooperative Extension – Green Leafy Vegetables*. Newark, DE: College of Agriculture & Natural Resources, University of Delaware.
48. Agudo A, Slimani N, Ocke MC, *et al.* (2002) Consumption of vegetables, fruit and other plant foods in the European Prospective Investigation into Cancer and Nutrition (EPIC) cohorts from 10 European countries. *Public Health Nutr* **5**, 1179–1196.
49. Piyathilake CJ, Macaluso M, Hine RJ, *et al.* (1994) Local and systemic effects of cigarette smoking on folate and vitamin B-12. *Am J Clin Nutr* **60**, 559–566.
50. Mannino DM, Mulinare J, Ford ES, *et al.* (2003) Tobacco smoke exposure and decreased serum and red blood cell folate levels: data from the Third National Health and Nutrition Examination Survey. *Nicotine Tob Res* **5**, 357–362.
51. Ulvik A, Ebbing M, Hustad S, *et al.* (2010) Long- and short-term effects of tobacco smoking on circulating concentrations of B vitamins. *Clin Chem* **56**, 755–763.
52. Walmsley CM, Bates CJ, Prentice A, *et al.* (1999) Relationship between cigarette smoking and nutrient intakes and blood status indices of older people living in the UK: further analysis of data from the National Diet and Nutrition Survey of people aged 65 years and over, 1994/95. *Public Health Nutr* **2**, 199–208.
53. Gloria L, Cravo M, Camilo ME, *et al.* (1997) Nutritional deficiencies in chronic alcoholics: relation to dietary intake and alcohol consumption. *Am J Gastroenterol* **92**, 485–489.
54. Cravo ML, Gloria LM, Selhub J, *et al.* (1996) Hyperhomocysteinemia in chronic alcoholism: correlation with folate, vitamin B-12, and vitamin B-6 status. *Am J Clin Nutr* **63**, 220–224.
55. Manari A, Preedy V & Peters T (2003) Nutritional intake of hazardous drinkers and dependent alcoholics in the UK. *Addict Biol* **8**, 201–210.
56. Mason JB & Choi SW (2005) Effects of alcohol on folate metabolism: implications for carcinogenesis. *Alcohol* **35**, 235–241.
57. Slimani N, Bingham S, Runswick S, *et al.* (2003) Group level validation of protein intakes estimated by 24-hour diet recall and dietary questionnaires against 24-hour urinary nitrogen in the EPIC Calibration Study. *Cancer Epidemiol Biomarkers Prev* **12**, 784–795.
58. Al-Delaimy WK, Ferrari P, Slimani N, *et al.* (2005) Plasma carotenoids as biomarkers of intake of fruits and vegetables: individual-level correlations in EPIC. *Eur J Clin Nutr* **59**, 1387–1396.
59. Willett WC (1998) *Nutritional Epidemiology*, 2nd ed. Oxford: Oxford University Press.
60. European Food Safety Authority (2009) ESCO report prepared by the EFSA Scientific Cooperation Working Group on Analysis of Risks and Benefits of Fortification of Food with Folic Acid. <http://www.efsa.europa.eu/fr/supporting/pub/3e.htm> (accessed August 2011).
61. Bingham SA (2002) Biomarkers in nutritional epidemiology. *Public Health Nutr* **5**, 821–827.