

11. *a.* Six parallelograms, whose diagonals intersect at  $M$  are  $HOVO'$ ,  $KOVO'$ ,  $LOWO'$ ;  $HKUV$ ,  $KLVW$ ,  $LHWU$ .  
*b.* Six parallelograms whose diagonals intersect at  $J$  are  $HIUI'$ ,  $KIVI'$ ,  $LIWI'$ ;  $HKUV$ ,  $KLVW$ ,  $LHWU$ .  
*c.* Six parallelograms whose diagonals intersect at  $J_1$  are  $HI_1UI_1'$ ,  $KI_1VI_1'$ ,  $LI_1WI_1'$ ;  $HKUV$ ,  $KLVW$ ,  $LHWU$ .
12. *a.*  $HWKULV$  is a hexagon whose opposite sides are parallel, and respectively  $= \frac{1}{2}O'A$ ,  $\frac{1}{2}O'B$ ,  $\frac{1}{2}O'C$ .  
*b.*  $HWKULV$  is a hexagon whose opposite sides are parallel, and respectively  $= \frac{1}{2}I'A$ ,  $\frac{1}{2}I'B$ ,  $\frac{1}{2}I'C$ .  
*c.*  $HWKULV$  is a hexagon whose opposite sides are parallel, and respectively  $= \frac{1}{2}I_1'A$ ,  $\frac{1}{2}I_1'B$ ,  $\frac{1}{2}I_1'C$ .
13. *a.*  $AO'$ ,  $BO'$ ,  $CO'$  pass through the points where the circumscribed circle of  $\triangle HKL$  cuts the sides of  $\triangle ABC$ .  
*b.*  $AI'$ ,  $BI'$ ,  $CI'$  pass through the points where the inscribed circle of  $\triangle HKL$  touches the sides of  $\triangle HKL$ .  
*c.*  $AI_1'$ ,  $BI_1'$ ,  $CI_1'$  pass through the points where the first escribed circle of  $\triangle HKL$  touches the sides of  $\triangle HKL$ .

---

On Determinants with  $p$ -termed elements.

By THOMAS MUIR, M.A., F.R.S.E.

This paper will be found in the *Messenger of Mathematics* for January 1884, Vol. xiii, New Series.

---

Construction for Euclid II. 9, 10.

By R. W. M'ARTHUR.

Take line  $AB$  divided in  $C$  and  $D$  as in Euclid. On  $AD$  describe the rectangle  $Aefd$  having  $AE$ ,  $DF$  each equal to  $AC$  or  $CB$ . According as  $D$  is in  $AB$ , or in  $AB$  produced, from  $DF$  or  $DF$  produced cut off  $FG$  equal to  $DB$ ; and join  $CG$ ,  $GE$ ,  $EC$ .

---

Mr JAMES TAYLOR gave a proof of the known theorem:—"If two sides of a skew quadrilateral  $ABDC$  inscribed in a circle be produced to meet in  $E$ , and  $FEG$  be drawn perpendicular to the diameter passing through  $E$ , the two other sides produced make equal intercepts on  $FEG$ ." Mr Taylor's object was to call attention to the desirability of obtaining a simpler mode of demonstration.