

Exploring under-five child malnutrition in Bangladesh: Analysis using the Extended Composite Index of Anthropometric Failure (ECIEF)

Farzana Akhter Bornee^{1#}, Mohammad Rocky Khan Chowdhury^{2#}, Badrun Naher Siddique², Baki Billah², Farjana Akter³, Md Nazmul Karim²

¹Department of Pediatrics, Bangabandhu Sheikh Mujib Medical University, Dhaka, Bangladesh

²Department of Epidemiology and Preventive Medicine, School of Public Health and Preventive Medicine, Faculty of Medicine, Nursing and Health Sciences, Monash University, Melbourne, Australia.

³Department of Sociology and Social Work, The people's University of Bangladesh, Dhaka, Bangladesh

Farzana Akhter Bornee and Mohammad Rocky Khan Chowdhury equally contributed in the paper.

Corresponding author: Md Nazmul Karim, Senior Lecturer, Department of Epidemiology and Preventive Medicine, Monash University. Email: Nazmul.Karim@monash.edu

Short title: Under-five child malnutrition in Bangladesh



This is an Accepted Manuscript for Public Health Nutrition. This peer-reviewed article has been accepted for publication but not yet copyedited or typeset, and so may be subject to change during the production process. The article is considered published and may be cited using its DOI 10.1017/S1368980025000138

Public Health Nutrition is published by Cambridge University Press on behalf of The Nutrition Society. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.

Acknowledgements: We thank MEASURE DHS for granting us permission to use the data. We also acknowledge the support of Bangabandhu Sheikh Mujib Medical University, Dhaka, Bangladesh and Monash University, Melbourne, Australia, where this study was conducted.

Financial Support: This research did not receive any fund.

Conflicts of Interest: None

Authorship: M.R.K.C., M.N.K., and F.A.B. designed the research plan. M.R.K.C. analysed the data. F.A.B., B.N.S. and F.A. prepared the first draft. M.R.K.C reviewed and edited the first draft. M.N.K. and B.B. critically reviewed the second draft. All authors reviewed and commented on final version of the manuscript.

Ethical standards disclosure: The 2017-18 BDHS was carried out by the National Institute of Population Research and Training (NIPORT) under the Ministry of Health and Family Welfare. Mitra and Associates, a Bangladeshi firm, were responsible for the survey's implementation, while technical support was provided by ICF International from the USA as part of their international Demographic and Health Surveys Program. Interviews were only conducted if the respondents verbally consented after being presented with an informed consent statement by the interviewers. It's worth noting that the current research, utilizing this data as a secondary source, does not require ethical approval.

Abstract

Objective: The current study is an attempt to explore under-five child malnutrition in a low-income population setting using the Extended Composite Index of Anthropometric Failure (ECIAF).

Design: Data from the Bangladesh Demographic and Health Survey 2017-18 were analyzed. Malnutrition using ECIAF was estimated using stunting, wasting underweight and overweight. Multilevel logistic regression models identified factors associated with malnutrition. Geospatial analysis was conducted using R programming.

Setting: Bangladesh.

Participants: Children under five years of age.

Results: In Bangladesh, as indicated by the ECIAF, approximately 40.8% (95% Confidence interval (CI): 39.7, 41.9) of children under-five experience malnutrition where about 3.3% (95% CI: 2.9, 3.7) were overweight. Children of parents with no formal education (56.3%, 95% CI: 50.8, 61.8), underweight mothers (53.4%, 95% CI: 50.4, 56.3), belonging to the lowest socio-economic strata (50.6%, 95% CI: 48.3, 53.0), residing in rural areas (43.3%, 95% CI: 41.9, 44.6), and aged below three years (47.7%, 95% CI: 45.2, 50.2) demonstrated a greater age and sex adjusted prevalence of malnutrition. The Sylhet division (Eastern region) exhibited a higher prevalence of malnutrition (>55.0%). Mothers with no formal education (Adjusted odds ratio (AOR): 1.51, 95% CI: 1.08, 2.10), underweight mother (AOR: 1.54, 95% CI: 1.03, 1.83), poorest socio-economic status (AOR: 2.14, 95% CI: 1.64, 2.81), children age 24-35 months of age (AOR: 2.37, 95% CI: 1.97, 2.85), and fourth and above birth order children (AOR: 1.41, 95% CI: 1.16, 1.72) were identified key factors associated with childhood malnutrition while adjusting community and household level variations.

Conclusion: In Bangladesh, two out of five children were malnourished and one in 35 children was overweight. Continuous monitoring of the ECIAF over time would facilitate tracking changes in the prevalence of different forms of malnutrition, helping to plan interventions and assess the effectiveness of interventions aimed at addressing both undernutrition and overweight.

Keywords: Bangladesh, children, malnutrition, undernutrition, overweight, composite index

Introduction

Child malnutrition, classified as indicators of anthropometric failure (AF) such as stunting, wasting, underweight, and overweight, often refers to the inadequate or excessive intake of essential nutrients necessary for the healthy growth and development of children.(1) Child malnutrition continues to be a significant public health issue for children under five years in many low- and middle-income countries.(2) According to current reports, approximately 148.1 million children under five suffer from stunting, 45 million experience wasting, and 82 million are underweight worldwide.(2) Furthermore, around 37 million children in this age group are affected by overweight.(2) Undernutrition is responsible for approximately half of all under-five child deaths worldwide.(1) Moreover, deaths associated with overweight surpass those related to underweight in the majority of the world's population.(3) A possible reason could be that global estimates of under-five child malnutrition are likely to be underestimated. Additionally, many existing tools for assessing malnutrition primarily focus on undernutrition (i.e., stunting, wasting, and underweight), neglecting the opposite end of the spectrum—overnutrition.(4-9) The failure to include the entire spectrum has the potential to underestimate the true impact of nutritional disorders, particularly the synergistic effects of the coexistence of multiple forms and the double burden of malnutrition (i.e., stunting, wasting, underweight, and overweight). These effects include increased susceptibility to infections, chronic diseases, impaired physical growth and cognitive development, higher healthcare costs, and reduced economic productivity.(10)

The standard estimates or single indicators often overlap, despite reflecting distinct biological processes. As a result, they are unable to provide an accurate and complete assessment of undernourishment among subjects. For example, some stunted children might also be wasted, while others may simultaneously experience both stunting and being underweight. In some cases, children may suffer from all three conditions concurrently. Consequently, these traditional metrics do not adequately capture the overall burden of malnutrition.(11) In nutritional assessment, several composite indices are used. These indices combine multiple indicators of AF, along with their interactions, into a single summary measure, providing a more comprehensive and accurate picture of the nutritional status of a population than any single indicator alone.(12-15) Recent literature reveals a multitude of efforts to use composite indices.(16-19) However, the majority of these indices were not able to address the "elephant in the room." Specifically, they have overlooked overweight, which has increasingly become

a concern in low-resource settings. According to the World Health Organization (WHO), many nations are grappling with the dual burden of malnutrition. This phenomenon highlights the simultaneous presence of inadequate nutrition (e.g., stunting) and excessive weight (e.g., overweight and obesity).(20, 21) The prevalence of undernutrition, once prominent in numerous low- and middle-income countries, is now giving way to elevated levels of overweight and obesity. This shift can be attributed to transitions in nutrition, epidemiology, and demographics.(10)

The composite index proposed by Bejarano et al. (2019) adopts a holistic approach by viewing the spectrum of nutritional disorders, from undernutrition to overnutrition, under the umbrella of malnutrition.(22) They reported approximately a threefold increase in the prevalence of malnutrition compared to undernutrition when malnutrition was assessed using the proposed index. The use of such tools in the assessment of malnutrition is likely to reveal the true extent of the malnutrition problem in low-income countries like Bangladesh, where policy planning has historically been heavily focused on undernutrition.(23) Despite a significant apparent reduction in under-five child undernutrition, the prevalence of child malnutrition, addressing both undernutrition and overnutrition in Bangladesh, is likely underestimated, as these are measured using available indices that primarily focus on undernutrition.(4, 12, 17) This study attempts to explore under-five child malnutrition in Bangladesh using one such tool, the Extended Composite Index of Anthropometric Failure (ECIAF).

Methods

Data Source: Bangladesh Demographic and Health Survey (BDHS)

We analyzed data from the Bangladesh Demographic and Health Survey (BDHS) 2017–2018. The data were collected between October 24, 2017, and March 15, 2018. The survey collected demographic data on Bangladeshi adults, information on maternal health, reproductive health, and maternal and child nutritional status from all eight administrative regions (divisions): Barisal (southern region), Chittagong (southeastern region), Dhaka (central region), Khulna (western region), Mymensingh (upper-central region), Rajshahi (midwestern region), Rangpur (northwestern region), and Sylhet (eastern region). The survey adopted a multistage cluster sampling technique. In the first stage, 675 primary sampling

units (PSUs) were selected using probability proportional to size, with 250 PSUs from urban areas and 425 PSUs from rural areas. The PSUs were based on enumeration areas (clusters) listed in the 2011 population census conducted by the Bangladesh Bureau of Statistics.(24) In the second stage, an average of 30 households was selected from each PSU using an equal probability systematic sampling technique. The multistage sampling and corresponding sampling weights were designed to reduce potential sampling bias. Additionally, all ever-married women aged 15–49 years (with or without children under five years) from the preselected households were interviewed without replacement or changes in the implementation stage to prevent selection bias. A total of 20,127 women aged 15–49 years were interviewed from 19,457 households, achieving a response rate of 99%. In BDHS 2017–18, a total of 8,759 children under five were listed, and 7,902 children were found eligible for analysis (**Supplementary Figure 1**). Detailed methodology is described in the BDHS 2017–18 report.(24)

Malnutrition assessment

Nandy et al. (2005) introduced the concept of Composite Index of Anthropometric Failure (CIAF) aimed at assessing undernutrition. This index encompassed seven distinct AF groups: A) absence of failure, B) only wasting, C) co-occurrence of wasting and underweight, D) simultaneous presence of wasting, stunting, and underweight, E) stunting and underweight, F) only stunting, and G) only underweight.(16) Later, Bejarano et al. (2019) expanded upon this concept by introducing ECIAF to gauge child malnutrition. They proposed a broader range of nine AF categories: A) absence of failure, B) only wasting, C) co-occurrence of wasting and underweight, D) simultaneous presence of wasting, stunting, and underweight, E) stunting and underweight, F) only stunting, G) only underweight, H) only overweight, and I) co-occurring overweight and stunting.(22, 23) The ECIAF is computed by subtracting the count of group A children from the total sample, and it served as a tool to measure the outcome, under-five child malnutrition in this study (**Supplementary Table 1**).

Four commonly used AF measures of child malnutrition are, stunting, wasting, underweight and overweight. A child was considered to be stunted (short stature for age), wasted (dangerously thin) and underweight (low weight for age) if the height-for-age, weight-for-height and weight-for-age indices were 2 standard deviations or more below the respective median of the WHO reference population. Conversely, a child was considered to be

overweight if the weight-for-height index was 2 standard deviations or more above the respective median of the WHO reference population.(25, 26)

Operational definitions of demographic and other characteristics

Parental characteristics, children characteristics, household, home environment and contextual factors were considered in the study are pooled from relevant literatures.(13, 17, 19, 27, 28) The parental characteristics includes maternal age in years (15–19, 20–24, 25–29, 30–34, 35–39, ≥ 40); parents' education (both parents uneducated, only father was uneducated when mother was educated, only mother was uneducated when father was educated, both parents educated); mother's working status (not working or currently working); mother's body size (normal, underweight and overweight); mother's attitudes toward wife-beating (justified, not justified); mother's decision-making autonomy (participated, not participated); father's occupational status (currently not working, manual labourer, service holder, businessman). Household factor includes mass media exposure (no, ye) and source of drinking water (improved, unimproved). Home environmental factors are use of cooking fuel (non-solid, solid); and type of toilet facilities (improved, unimproved). Contextual factors comprise wealth index (poorest, poorer, middle, richest and richer); place of residence (urban, rural); and region of residence (Barisal, Chittagong, Dhaka, Khulna, Mymensingh, Rajshahi, Rangpur, Sylhet). Factors related to children are children's age (0-11 months, 12-23 months, 24-35 months, 36-47 months and 48-59 months), sex of child (male, female); birth order (first, second, third and fourth and above); diarrhea (no, yes); and acute respiratory infection (ARI) (no, yes). Operational definitions of selected factors were presented in **Supplementary Table 3**

Statistical analysis

Descriptive statistics were used to present the background characteristics of the children. Bivariate analysis (Chi-square test) was used to explore the significant associations (significance level set at $p < 0.05$) between the independent variables and malnutrition as per the ECIAF among under-five children. Initially, multivariable multilevel binary logistic regression model with a random intercept term at community- and household-level were performed to identify significant determinants of childhood malnutrition. In DHS, individuals are nested within households and households are nested within communities, which indicate

that individuals, households, and communities are not independent of each other.(4, 29) We employed multilevel random intercept model to account for variation in different levels (community, household and individual level). Further, multivariable binary logistic regression analysis (fixed effect model) was performed to identify the determinants of malnutrition. Variables that were found to be significant at the level $p < 0.25$ in bivariate analysis were included in the multivariable analysis.(30-33) The level of significance for multivariable regression analysis was set at $p < 0.05$, and an odds ratio (OR) with 95% confidence interval (CI) was estimated to determine the associated factors. Interclass correlation coefficient (ICC) was generated out and identified 25% for random effect model which indicated low ICC. Further, the estimated value of Akaike's information criterion (AIC) and Bayesian information criterion (BIC) for multilevel model was 10206.2 and 10408.3 and slightly lower compare to fixed effect regression model (**Supplementary Table 4**). Therefore, random effect model could be considered as better model compared to fixed effect model. Stata version 17 (StataCorp LP, College Station, Texas) was used for all analyses. To control the effect of the complex survey design, all bivariate and multivariable analyses of this study were performed using Stata's "svyset" and melogit commands respectively. Geospatial analysis was carried out using R programming language (4.3.3) to visualize the regional prevalence of malnutrition as per the ECIAF and CIAF, and regional prevalence of childhood overweight.

Results

A significant proportion (62.8%) of the mothers fell within the younger age bracket (20 to 29 years). Around 7.1% of the children's mothers had received no formal education (with 4.0% having both parents uneducated and 3.1% being solely uneducated mothers). Around 15% of the mothers were classified as underweight. Approximately 47.8% of the children were of female and around 40% of the children were from socio-economically disadvantaged backgrounds, and a substantial majority (65.9%) of all children were residents of rural areas. For more comprehensive information, please see Table 1 which outlines these background characteristics in detail.

Prevalence and geospatial distribution of child malnutrition

In Bangladesh, as per the ECIAF, the prevalence of under-five child malnutrition was 40.8% (95% Confidence Interval (CI): 39.7, 41.9), while the CIAF indicates that around 38.3% (95% CI: 37.2, 39.5) of children experience undernourishment, as illustrated in Figure 1. Moreover, approximately 3.3% (95% CI: 2.9, 3.7) of children face overweight conditions alongside other forms of undernutrition, as highlighted in Figure 1.

Notably, the prevalence of malnutrition is higher in the Sylhet division (eastern region), reaching over 55%, compared to the lower prevalence observed in the Khulna division (western region) (below 38%). Similarly, the prevalence of undernutrition according to the CIAF is also elevated in Sylhet division (eastern region) (around 50%). On the other hand, the Dhaka division (central region) exhibits a higher prevalence of under-five child overweight (over 5%), followed by Barisal (southern) and Sylhet (eastern) divisions, each at around 4%, as depicted in Figure 2.

The age and sex adjusted prevalence of child malnutrition

Among children, the age- and sex-adjusted prevalence of malnutrition was significantly higher at 56.3% (95% CI: 50.8, 61.8) in households where mothers had no formal education. Similarly, a heightened prevalence was observed among children born to underweight mothers (53.4%, 95% CI: 50.4, 56.3), those in families with a fourth or higher birth order (52.3%, 95% CI: 48.1, 54.4), children from the most socio-economically poorest (50.6%, 95% CI: 48.3, 53.0), and those aged 24–35 months (47.7%, 95% CI: 45.2, 50.2), as detailed in Table 2.

Factors associated with child malnutrition

Table 3 reveals that simple logistic regression analysis yielded significant associations between under-five child malnutrition and various factors, including parental education, mother currently working, mother's body size, use of cooking fuel, type of toilet facilities, mass media exposure, wealth index, place of residence, children age, and birth order.

For adjusted model, random effect model was preferred over fixed effect model to find out the association between outcome and exposures (Table 3). The national level data consists of

multiple entities (e.g., states, regions, households) where each entity is assumed to have its unique and uncorrelated characteristics. Random effect model focuses on understanding how the effects of explanatory variables differ across various entities and aims to estimate entity-specific variances whereas fixed effect model assumes unobserved entity-specific effects which are correlated with the explanatory variables.

In Table 3, children had higher likelihood of being malnourished in mothers with no formal education (Adjusted Odds Ratio-random (AORr): 1.51, 95% CI: 1.08, 2.10)) compare to children with both parents educated. Further, children of underweight mothers (AORr: 1.54, 95%: 1.03, 1.83); fathers being manual labourer (AORr: 1.17, 95% CI: 1.01, 1.36); poorest socio-economic status (AORr: 2.14, 95% CI: 1.64, 2.81); children age 24-35 months of age (AORr: 2.37, 95% CI: 1.97, 2.85) and fourth and above birth order children (AORr: 1.41, 95% CI: 1.16, 1.72) had significant effects on under-five child malnutrition when considering community and household level variations. In comparison, the findings of fixed effect model were presented in **Supplementary Table 4**.

In the sensitivity analysis, a multilevel logistic regression model (random effects model) was utilized, incorporating four additional variables into the primary model. These variables were minimum acceptable diet, low birth weight, antenatal care, and prenatal care, focused on children aged 6-23 months (**Supplementary Table 5**). The analysis revealed that children born with low birth weight had a higher likelihood of malnutrition (AORr: 2.16, 95% CI: 1.29, 3.59) compared to children of normal birth weight.

Discussion

While the CIAF was utilized to provide a comprehensive view of the undernutrition status among children under the age of five in Bangladesh, revealing a prevalence of 38.3%, the ECIAF was employed to broaden the analysis by including overweight prevalence (3.3%). This resulted in a more inclusive measurement of malnutrition, with a prevalence of 40.8%. This approach offers a more thorough evaluation of the nutritional well-being of children, spanning the entire spectrum from undernutrition to overnutrition. Although undernutrition rates in Bangladesh have been declining, the gradual rise in childhood overweight poses a significant concern. The prevalence of childhood overweight in the country increased from 1.6% in 2004 to 2.3% in 2014 and further escalated to 3% in 2019.(34) A comparative

analysis across South Asian nations indicates that Bangladesh's childhood overweight rate (3.3%) is lower than that of Maldives (5.4%) and Pakistan (4.9%), yet surpasses that of India (2.8%) and Nepal (1.4%).(35)

Among all the administrative divisions in Bangladesh, the Sylhet region (located in the eastern hilly part) exhibited the highest prevalence of malnutrition as indicated by the ECIAF, mirroring the pattern observed for undernutrition based on the CIAF. These findings align with the results of prior studies conducted in Bangladesh.(4, 17) The Sylhet region includes the Haor basin, characterized by its distinctive wetland landscape resembling an oxbow lake. This terrain necessitates specific agricultural approaches.(36) Residents of various Haor areas frequently experience food insecurity due to sudden pre-monsoon floods and the reliance on cultivating only one crop. Additionally, the region faces pronounced education- and wealth-based inequalities.(37) While child undernutrition was prevalent among economically disadvantaged and less-educated parents, issues of childhood overweight were observed among children from more affluent backgrounds in the same area. To address these challenges, strategies to reduce poverty in the most vulnerable communities could include measures such as enhancing employment opportunities, ensuring a stable food supply, and improving female literacy rates. At the same time, strengthening awareness programs across all social strata could help mitigate childhood malnutrition.(38-40) The current study revealed that the prevalence of overweight in children under five was highest in Dhaka (located in the central plain region) at 5.6%, followed by Barisal (southern region) at 4.2%, and Sylhet (eastern region) at 4.2%. Dhaka, being a central region, faces rapid urbanization, which has led to limited play spaces for children.(41) This scarcity contributes to inadequate physical activity, increased screen time, unhealthy dietary patterns, and greater access to processed and calorie-dense foods. These factors collectively drive the rising incidence of childhood overweight.

This study also found that malnutrition was highly prevalent among children of uneducated parents, underweight mothers, and children of fourth or higher birth order. Furthermore, mothers with no formal education, underweight mothers, poorest socio-economic status, children aged 24–35 months, and children of fourth or higher birth order were identified as key factors associated with childhood malnutrition, while accounting for community- and household-level variations. Nonetheless, comparable factors exhibited significance in relation to malnutrition, albeit with varying odds ratios when analyzed using a fixed-effects model.

These crucial determinants have been previously established as significant contributors to childhood undernutrition in numerous studies conducted in Bangladesh and other developing regions.(13, 15, 17, 27, 28) It is worth noting that these past investigations employed a fixed-effects model, unlike ours, which may have underestimated the variations present at different levels or failed to account for potential heterogeneity among groups, such as urban and rural children.

Previous studies in Bangladesh focused on under-five child undernutrition have highlighted various causative factors. For instance, the condition of uneducated and impoverished women often leads to maternal underweight, which consequently contributes to child undernutrition.(4, 12) Additionally, inappropriate and insufficient initiation of complementary feeding, as well as the prioritization of older children within a household when multiple children are present, can adversely affect the nutritional status of younger children.(42) These studies have proposed potential solutions to address undernutrition, including strategies such as enhancing educational attainment, improving access to health and nutrition services, ensuring proper maternal healthcare utilization to improve maternal health, implementing effective interventions targeted at vulnerable groups (particularly children from the most socioeconomically disadvantaged backgrounds), fostering parental engagement through awareness initiatives, implementing multisectoral healthcare programs to reduce the prevalence of undernutrition, and reinforcing family planning efforts to delay childbirth and reduce higher-order births.(4, 5, 12, 13, 17, 42) However, the government of Bangladesh and other relevant stakeholders have already undertaken many undernutrition alleviation strategies, such as the National Adaptation Plan of Bangladesh (2023–2050), Meeting the Undernutrition Challenge (MUCH), and the National Food and Nutrition Security Policy Plan of Action (2021–2030), among others.(43-45) Despite these concerted efforts, child undernutrition remains high in Bangladesh. A lack of multisectoral planning and coordination (involving government, non-governmental, social, cultural, and religious institutions), gaps in food and nutrition information, and the absence of quality epidemiological data are some of the complex reasons preventing significant improvements in the nutritional status of children in Bangladesh.(46, 47)

Future direction and policy implication

To address childhood malnutrition effectively, it is crucial to establish and monitor cross-sectoral coordination (involving government, nongovernmental, social, cultural, and religious institutions). Ensuring proper maternal nutrition before, during, and after pregnancy is essential for a child's health. Focus should be directed toward the critical first thousand days of a child's life, promoting nutritious, varied, and safe dietary practices from the outset to prevent both undernutrition and obesity.(48) This includes fostering a healthy environment with access to essential health services, clean water, proper sanitation, and opportunities for safe physical activity. Improving living conditions and enhancing maternal nutritional knowledge can significantly reduce rates of childhood malnutrition.(49) Elevating the status of women should also be a priority. To tackle both undernourishment and overnutrition, authorities must enhance the quantity and quality of available food while ensuring it remains affordable. Community-level education programs and public awareness campaigns are vital for encouraging healthy lifestyles and dietary habits. Further research, particularly among preschool children in rural areas and across different ethnic groups, should be conducted using the ECIAF model to better understand the scope of malnutrition. The ECIAF can provide valuable epidemiological information for monitoring the progress of public health policies.(22, 23) Considering the findings from current investigations, including the distribution of childhood malnutrition, education and social status, and family planning efforts to delay childbirth and reduce higher-order births, can inform the development of improved health and nutritional policies. Additionally, reshaping existing strategies and interventions using the double-duty action proposed by Hawkes et al. (2020) may help address the double burden of malnutrition.(50) Last but not least, policies, programs, and resources cannot be successfully adopted, implemented, or sustained without strong commitment.

Limitation of using the ECIAF

The ECIAF which typically aggregates multiple indicators into a single value, may not always be the best tool for diagnosing the nutritional status of individuals. By combining different malnutrition indicators such as stunting, wasting, underweight, and overweight into a single measure, the composite index may overly simplify the complexities of nutritional issues, potentially obscuring specific problems like micronutrient deficiencies. People have

unique health conditions, genetic predispositions, and lifestyle factors that influence their nutritional needs. A composite index standardized for a general population might not accurately reflect these individual differences. Composite indices are often designed to capture broad trends rather than diagnose specific health conditions. They may not be sensitive enough to detect mild or emerging nutritional problems in an individual. When indicators are combined, high values in one indicator can offset low values in another, potentially creating a misleading picture of an individual's health. This can lead to underestimation of risk or need in specific nutritional areas. Nutritional factors also differ widely across cultural and regional contexts, and a universal index might not accurately capture local dietary practices, economic conditions, or health priorities, thereby limiting its effectiveness.

Composite indices are often used in research and policy-making to provide a general overview of health or nutritional status across populations or groups. They simplify complex data, making it easier to communicate and make broad comparisons. However, heavy reliance on a single composite index for policymaking could skew directions and resource allocations, overlooking crucial aspects necessary to tackle the root causes of malnutrition. To overcome these limitations, it is essential to adopt more nuanced approaches that integrate local contexts and specific nutritional issues into health assessments and interventions.

Strength and limitation of the study

The large sample size and robust statistical techniques are key strengths of the study. Furthermore, this is a comprehensive assessment of under-five child malnutrition, covering both undernutrition and overnutrition, using nationally representative samples. One limitation of the study is that the cross-sectional nature of the data does not allow for the determination of causal relationships. The data were collected retrospectively and were self-reported, thus there is a possibility of underreporting and recall bias. In addition, other important variables, such as complementary feeding practices, antenatal care, postnatal care, and birth weight, were not adjusted for due to a large number of missing observations. Nutrient intake and micronutrient status could not be adjusted in the model due to a lack of data

Conclusion

The findings revealed that two out of five children in Bangladesh were malnourished, with a higher prevalence in the eastern region (Sylhet division) and a higher prevalence of overweight in the central region (Dhaka division). Children born to mothers with no formal education, underweight mothers, those from the poorest socio-economic backgrounds, children aged 24–35 months, and those born as the fourth child or above are at greater risk of developing malnutrition. Despite its limited occurrence in Bangladesh, the inclusion of overweight was considered a significant component in the composite index, designed as the ECIAF, to assess the comprehensive state of malnutrition.

Continuous monitoring of the ECIAF over time has the potential to facilitate tracking changes in the prevalence of different forms of malnutrition, helping to assess the effectiveness of interventions aimed at addressing both undernutrition and overweight. Further research on composite indices is important to enhance their reliability and applicability.

References

1. WHO (2020) Malnutrition. *World Health Organization (WHO)*.
2. Group UWWB (2023) Levels and trends in child malnutrition. UNICEF / WHO / World Bank Group Joint Child Malnutrition Estimates 2023 edition.
3. WHO (2021) Obesity and overweight. *World Health Organization (WHO)*.
4. Chowdhury MRK, Rahman MS, Khan MMH *et al.* (2016) Risk factors for child malnutrition in Bangladesh: a multilevel analysis of a nationwide population-based survey. *The Journal of pediatrics* 172, 194-201. e191.
5. Das S, Baffour B Richardson A (2022) Prevalence of child undernutrition measures and their spatio-demographic inequalities in Bangladesh: an application of multilevel Bayesian modelling. *BMC Public Health* 22, 1-21.
6. Sahu SK, Kumar SG, Bhat BV *et al.* (2015) Malnutrition among under-five children in India and strategies for control. *Journal of natural science, biology, and medicine* 6, 18.
7. Asim M & Nawaz Y (2018) Child malnutrition in Pakistan: evidence from literature. *Children* 5, 60.

8. Delisle H, Shrimpton R, Blaney S *et al.* (2017) Capacity-building for a strong public health nutrition workforce in low-resource countries. *Bulletin of the World Health Organization* 95, 385.
9. Abdul-Rahman T, Ghosh S, Lukman L *et al.* (2023) Inaccessibility and low maintenance of medical data archive in low-middle income countries: Mystery behind public health statistics and measures. *Journal of Infection and Public Health* 16, 1556-1561.
10. Wells JC, Sawaya AL, Wibaek R *et al.* (2020) The double burden of malnutrition: aetiological pathways and consequences for health. *The Lancet* 395, 75-88.
11. Ziba M, Kalimbira AA, Kalumikiza Z (2018) Estimated burden of aggregate anthropometric failure among Malawian children. *South African Journal of Clinical Nutrition* 31, 20-23.
12. Chowdhury MRK, Rahman MS, Billah B *et al.* (2023) Prevalence and factors associated with severe undernutrition among under-5 children in Bangladesh, Pakistan, and Nepal: a comparative study using multilevel analysis. *Scientific Reports* 13, 10183.
13. Chowdhury MRK, Khan HT, Mondal M (2021) Differences in the socio-demographic determinants of undernutrition in children aged < 5 years in urban and rural areas of Bangladesh measured by the composite index of anthropometric failure. *Public Health* 198, 37-43.
14. Al-Sadeeq AH, Bukair AZ, Al-Saqladi A-WM (2018) Assessment of undernutrition using Composite Index of Anthropometric Failure among children aged < 5 years in rural Yemen. *Eastern Mediterranean Health Journal* 24, 1119-1126.
15. Permatasari TAE & Chadirin Y (2022) Assessment of undernutrition using the composite index of anthropometric failure (CIAF) and its determinants: A cross-sectional study in the rural area of the Bogor District in Indonesia. *BMC nutrition* 8, 133.
16. Nandy S & Svedberg P (2012) The Composite Index of Anthropometric Failure (CIAF): An alternative indicator for malnutrition in young children. In *Handbook of anthropometry: Physical measures of human form in health and disease*, pp. 127-137: Springer.

17. Islam MS & Biswas T (2020) Prevalence and correlates of the composite index of anthropometric failure among children under 5 years old in Bangladesh. *Maternal & child nutrition* 16, e12930.
18. NASSAR MF, EMAM EK, EL-KHOLY H *et al.* (2022) Composite Index of Anthropometric Failure Burden Among Hospitalized Pediatric Patients. *The Medical Journal of Cairo University* 90, 409-415.
19. Boregowda G, Soni G, Jain K *et al.* (2015) Assessment of under nutrition using composite index of anthropometric failure (CIAF) amongst toddlers residing in urban slums of Raipur City, Chhattisgarh, India. *Journal of clinical and diagnostic research: JCDR* 9, LC04.
20. Khaliq A, Wraith D, Miller Y *et al.* (2021) Prevalence, trends, and socioeconomic determinants of coexisting forms of malnutrition amongst children under five years of age in Pakistan. *Nutrients* 13, 4566.
21. Bejarano IF, Oyhenart EE, Torres MF *et al.* (2019) Extended composite index of anthropometric failure in Argentinean preschool and school children. *Public health nutrition* 22, 3327-3335.
22. Bejarano IF, Oyhenart EE, Torres MF *et al.* (2019) Extended composite index of anthropometric failure in Argentinean preschool and school children. *Public health nutrition* 22, 3327-3335.
23. Bose K (2018) The Concept of Composite Index of Anthropometric Failure (CIAF): Revisited and Revised.
24. NIPORT I (2020) Bangladesh demographic and health survey 2017-18. *Dhaka and Maryland*.
25. De Onis M (2006) WHO Child Growth Standards Length/height-for-age, weight-for-age, weight-for-length, weight-for-height and body mass index-for-age. *Methods and development*.
26. WHO (2023) Child malnutrition: Overweight prevalence among children under 5 years of age. *World Health Organization (WHO)*.
27. Balogun OS, Asif AM, Akbar M *et al.* (2021) Prevalence and potential determinants of aggregate anthropometric failure among pakistani children: findings from a community health survey. *Children* 8, 1010.
28. Porwal A, Acharya R, Ashraf S *et al.* (2021) Socio-economic inequality in anthropometric failure among children aged under 5 years in India: evidence from the

- Comprehensive National Nutrition Survey 2016–18. *International journal for equity in health* 20, 1-10.
29. Chowdhury MRK, Khan MMH, Khan HT *et al.* (2020) Prevalence and risk factors of childhood anemia in Nepal: A multilevel analysis. *Plos one* 15, e0239409.
 30. Bursac Z, Gauss CH, Williams DK *et al.* (2008) Purposeful selection of variables in logistic regression. *Source code for biology and medicine* 3, 1-8.
 31. Thiese MS, Ronna B Ott U (2016) P value interpretations and considerations. *Journal of thoracic disease* 8, E928.
 32. Kim J (2015) How to choose the level of significance: A pedagogical note.
 33. Chowdhury MRK, Khan HT, Rashid M *et al.* (2022) Prevalence and correlates of severe under-5 child anthropometric failure measured by the Composite Index of Severe Anthropometric Failure in Bangladesh. *Frontiers in Pediatrics* 10.
 34. Shawon MSR, Hossain FB, Thomson B *et al.* (2020) Trends in the prevalence of overweight among Bangladeshi children aged 24–59 months (2004–2014) by sex and socioeconomic status. *International Journal of Obesity* 44, 664-674.
 35. Bishwajit G & Yaya S (2020) Overweight and obesity among under-five children in South Asia. *Child and Adolescent Obesity* 3, 105-121.
 36. Kazal MMH, Rahman S Hossain MZ (2017) Poverty profiles and coping strategies of the haor (ox-bow lake) households in Bangladesh.
 37. Rahman MS, Chowdhury MRK, Islam MR *et al.* (2023) Determinants and Projections of Minimum Acceptable Diet among Children Aged 6–23 Months: A National and Subnational Inequality Assessment in Bangladesh. *International Journal of Environmental Research and Public Health* 20, 2010.
 38. Farzana FD, Choudhury N, Haque MA *et al.* (2022) Type of terrain and infant and young child feeding practices: cross-sectional study findings on children below 2 years of age from northern Bangladesh. *BMJ open* 12, e056593.
 39. Ahmed M (2010) Multiple providers and access to primary education: The case of Bangladesh. *Prospects* 40, 393-415.
 40. Oxfam G (2006) Girls; Education in South Asia: Education and Gender Equality Series. *Programme Insights Paper developed by Beyond Access Project.*
 41. Das SK, Chisti MJ, Huq S *et al.* (2013) Changing trend of overweight and obesity and their associated factors in an urban population of Bangladesh.

42. Chowdhury MRK, Rahman MS Khan MMH (2016) Levels and determinants of complementary feeding based on meal frequency among children of 6 to 23 months in Bangladesh. *BMC public health* 16, 1-11.
43. MoEFCC (2022) National Adaptation Plan of Bangladesh (2023-2050). *Ministry of Environment, Forest and Climate Change (MoEFCC), Government of the People's Republic of Bangladesh*
44. FAO (2022) Evaluation of the “Meeting the Undernutrition Challenge (MUCH): Strengthening the enabling environment for food security and nutrition” programme. Programme Evaluation Series, 05/2022. Rome. *Food and Agriculture Organization (FAO) of the United Nation.*
45. FPMU (2021) National Food and Nutrition Security Policy. Plan of Action (2021-2030). *Food Planning and Monitoring Unit (FPMU), Ministry of Food, Government of the People's Republic of Bangladesh.*
46. Dikoda (2021) How to bridge the food nutrition information gap in low and middle-income countries.
47. SUN (2014) Undernutrition in Bangladesh. A Common Narrative. *Scaling Up Nutrition.*
48. Likhari A & Patil MS (2022) Importance of maternal nutrition in the first 1,000 days of life and its effects on child development: a narrative review. *Cureus* 14.
49. Keats EC, Das JK, Salam RA *et al.* (2021) Effective interventions to address maternal and child malnutrition: an update of the evidence. *The Lancet Child & Adolescent Health* 5, 367-384.
50. Hawkes C, Ruel MT, Salm L *et al.* (2020) Double-duty actions: seizing programme and policy opportunities to address malnutrition in all its forms. *The Lancet* 395, 142-155.

Table 1 Background characteristic of respondents

Factors	Number (N)	Unweighted percentage (%)	Weighted percentage (%)
Maternal age (in years)			
15-19	985	12.5	13.0
20-24	2,753	34.8	34.7
25-29	2,210	28.0	28.1
30-34	1,326	16.7	16.7
34-39	497	6.3	6.0
≥ 40	131	1.7	1.6
Parents' education			
Both parents uneducated	318	4.0	4.1
Only father uneducated	987	12.5	12.4
Only mother uneducated	248	3.1	3.1
Both parents educated	6,349	80.4	80.4
Mother's currently working			
No working	4,701	59.5	59.6
Currently working	3,201	40.5	40.4
Mother's body size (BMI)			
Normal	4,688	59.3	60.3
Underweight	1,167	14.8	14.0
Overweight	2,047	25.9	25.8
Mother's attitude towards wife beating			
Not Justified	6,462	81.8	81.2
Justified	1,440	18.2	18.8
Mothers' decision-making autonomy			
Not practiced	1,222	15.5	15.2
Practiced	6,680	84.5	84.8
Fathers' occupational status			
Currently not working	212	2.7	2.6
Manual labourer	5,529	70.0	71.3

Service holder	481	6.1	5.4
Businessman	1,680	21.3	20.7
Source of drinking water			
Improved	6,867	86.9	86.9
Not improved	1,035	13.1	13.1
Use of cooking fuel			
Non-solid	2,286	28.9	29.7
Solid	5,616	71.1	70.3
Type of toilet facilities			
Improved	4,487	56.8	56.8
Not improved	3,415	43.2	43.2
Mass media exposure			
No	2,863	36.2	35.0
Yes	5,039	63.8	65.0
Wealth index			
Poorest	1,767	22.4	21.8
Poorer	1,597	20.2	20.5
Middle	1,430	18.1	19.2
Richer	1,565	19.8	20.0
Richest	1,543	19.5	18.5
Place of residence			
Urban	2,694	34.1	26.4
Rural	5,208	65.9	73.6
Children age			
0-11 months	1,736	22.0	21.8
12-23 months	1,617	20.5	20.7
24-35 months	1,530	19.3	19.6
36-47 months	1,459	18.5	18.7
48-59 months	1,560	19.7	19.2
Sex of child			
Male	4,125	52.2	52.3
Female	3,777	47.8	47.7
Birth order			

First	3,019	38.2	38.2
Second	2,573	32.6	32.5
Third	1,327	16.8	16.9
Fourth and above	983	12.4	12.4
Diarrhea prior two weeks			
No	7,507	95.0	95.2
Yes	395	5.0	4.8
ARI prior two weeks			
No	6,878	87.0	86.7
Yes	1,024	13.0	13.3
Total	7,902	100.0	100.0

ARI, acute respiratory infection

Table 2 Prevalence of under-five child malnutrition as per the ECIAF

Factors	Number (N)	Crude prevalence (95% CI)	Age and sex adjusted prevalence (95% CI)	p-values
Overall	3262	40.6 (39.3, 42.4)	40.8 (39.7, 41.9)	
Maternal age (in years)				
15-19	420	41.2 (37.6, 44.9)	44.0 (40.8, 47.2)	0.927
20-24	1097	40.1 (37.7, 42.5)	39.6 (37.8, 41.5)	
25-29	912	41.0 (38.6, 43.5)	41.0 (38.9, 43.0)	
30-34	559	41.3 (38.1, 44.6)	41.6 (38.9, 44.2)	
34-39	213	41.0 (36.1, 46.1)	42.6 (38.2, 46.9)	
≥ 40	61	45.4 (35.6, 55.5)	45.4 (36.9, 54.0)	
Parents' education				
Both parents uneducated	180	55.8 (49.3, 62.1)	56.3 (50.8, 61.8)	<0.001
Only father uneducated	490	49.9 (46.4, 53.5)	49.6 (46.4, 52.7)	
Only mother uneducated	133	51.6 (44.2, 59.0)	53.9 (47.7, 60.1)	
Both parents educated	2459	38.3 (36.7, 39.8)	38.6 (37.4, 39.8)	
Mother's working status				
No	1891	39.6 (37.7, 41.6)	40.3 (39.0, 41.8)	0.034
Yes	1371	42.6 (40.5, 44.8)	42.3 (40.6, 44.0)	
Mother's body size				
Normal	1959	41.2 (39.2, 43.1)	43.3 (41.6, 45.0)	<0.001
Underweight	621	51.6 (48.4, 54.7)	53.4 (50.4, 56.3)	
Overweight	682	34.2 (31.8, 36.8)	34.5 (32.9, 36.2)	
Mother's attitude towards wife beating				
Not Justified	2640	40.3 (38.7, 41.9)	40.7 (39.5, 41.9)	0.096
Justified	622	43.0 (40.0, 46.1)	43.1 (40.6, 45.7)	
Mothers' decision-making				

autonomy					
Not practiced	518	41.2 44.3)	(38.1, 42.3	(39.5, 45.0)	0.824
Practiced	2744	40.8 42.4)	(39.1, 40.9	(39.8, 42.1)	
Fathers' occupational status					
Currently not working	96	45.6 53.2)	(38.3, 45.0	(38.2, 51.7)	<0.001
Manual labourer	2417	43.3 45.0)	(41.6, 43.6	(42.3, 44.9)	
Service holder	133	26.4 30.8)	(22.4, 27.6	(23.5, 31.6)	
Businessman	616	35.6 38.4)	(32.8, 36.4	(34.1, 38.7)	
Source of drinking water					
Improved	2848	41.1 42.7)	(39.5, 41.0	(40.0, 42.4)	0.225
Not improved	414	39.0 42.3)	(35.7, 40.8	(37.7, 43.8)	
Use of cooking fuel					
Non-solid	807	36.0 38.7)	(33.4, 35.5	(33.5, 37.4)	<0.001
Solid	2455	42.9 44.6)	(41.1, 43.4	(42.2, 44.8)	
Type of toilet facilities					
Improved	1718	38.1 40.1)	(36.1, 38.0	(36.6, 39.4)	<0.001
Not improved	1544	44.4 46.5)	(42.4, 45.3	(43.6, 47.0)	
Mass media exposure					
No	1352	46.4 48.8)	(44.2, 47.4	(45.5, 49.2)	<0.001
Yes	1910	37.8 39.7)	(36.0, 37.6	(36.3, 39.0)	
Wealth index					
Poorest	897	50.0 52.8)	(47.2, 50.6	(48.3, 53.0)	<0.001
Poorer	753	45.6 48.6)	(42.5, 47.2	(44.7, 49.7)	
Middle	565	38.3 41.3)	(35.4, 39.4	(36.9, 42.0)	
Richer	606	39.0 42.0)	(36.2, 38.7	(36.2, 41.1)	
Richest	441	29.4 32.6)	(26.4, 28.2	(25.9, 30.40)	
Place of residence					
Urban	1003	37.2 40.1)	(34.4, 37.1	(35.2, 38.9)	0.005
Rural	2259	42.1	(40.3, 43.3	(41.9, 44.6)	

			43.9)		
Children age					
0-11 months	563	32.0 34.7)	(29.5, 32.5 (30.3, 34.7)		<0.001
12-23 months	713	42.6 (39.8,45.6)	43.9 (41.5, 46.3)		
24-35 months	728	47.8 50.8)	(44.7, 47.7 (45.2, 50.2)		
36-47 months	645	43.6 46.7)	(40.6, 44.1 (41.5, 46.6)		
48-59 months	613	39.1 42.0)	(36.3, 39.3 (36.9, 41.7)		
Sex of child					
Male	1731	41.8 43.7)	(39.9, 41.8 (40.3, 43.3)		0.119
Female	1531	39.8 41.8)	(37.8, 40.4 (38.8, 42.0)		
Birth order					
First	1177	39.0 41.2)	(37.0, 38.7 (36.9, 40.0)		<0.001
Second	1013	39.3 41.7)	(36.9, 39.4 (37.5, 41.3)		
Third	570	41.4 44.5)	(38.4, 42.8 (40.0, 45.4)		
Fourth and above	502	49.6 53.6)	(45.7, 51.3 (48.1, 54.4)		
Diarrhea prior two weeks					
No	3110	40.9 42.5)	(39.4, 41.3 (40.2, 42.4)		0.536
Yes	395	39.1 44.9)	(33.6, 38.2 (33.3, 43.0)		
ARI prior two weeks					
No	2,833	40.8 42.4)	(39.2, 41.0 (39.8, 42.2)		0.924
Yes	429	41.0 44.3)	(37.7, 42.3 (39.2, 45.3)		

ARI, acute respiratory infection; CI, confidence interval

Table 3 Determinants of under-five child malnutrition as per the ECIAF

Factors	COR	(95% CI)	p-values	AORr (95% CI)	p-values
Parents' education					
Both parents uneducated	1.88	(1.53, 2.29)	<0.001	1.43 (1.05, 1.95)	0.021
Only father uneducated	1.57	(1.37, 1.82)	<0.001	1.28 (1.07, 1.54)	0.008
Only mother uneducated	1.83	(1.41, 2.36)	<0.001	1.51 (1.08, 2.10)	0.016
Both parents educated (RC)	1.00			1.00	
Mother's working status					
No	1.00			1.00	
Yes	1.11	(1.02, 1.22)	0.021	0.91 (0.81, 1.03)	0.156
Mother's body size					
Normal (RC)	1.00			1.00	
Underweight	1.58	(1.39, 1.80)	<0.001	1.54 (1.30, 1.83)	<0.001
Overweight	0.70	(0.62, 0.77)	<0.001	0.79 (0.69, 0.90)	<0.001
Mother's attitude towards wife beating					
Not Justified (RC)	1.00			1.00	
Justified	1.10	(0.98, 1.23)	0.103	0.99 (0.86, 1.15)	0.942
Fathers' occupational status					
Currently not working	1.43	(1.07, 1.91)	0.015	1.14 (0.79, 1.67)	0.476
Manual labourer	1.34	(1.20, 1.50)	<0.001	1.17 (1.01, 1.36)	0.031
Service holder	0.66	(0.52, 0.86)	<0.001	0.88 (0.66, 1.17)	0.386

	0.82)			
Businessman (RC)	1.00		1.00	
Use of cooking fuel				
Non-solid (RC)	1.00		1.00	
Solid	1.42	(1.29, <0.001	0.97 (0.83, 1.14)	0.750
	1.57)			
Type of toilet facilities				
Improved (RC)	1.00		1.00	
Not improved	1.33	(1.21, <0.001	1.05 (0.92, 1.21)	0.425
	1.45)			
Mass media exposure				
No	1.46	(1.33, <0.001	1.04 (0.90, 1.20)	0.584
	1.61)			
Yes (RC)	1.00		1.00	
Wealth index				
Poorest	2.58	(2.23, <0.001	2.14 (1.64, 2.81)	<0.001
	2.98)			
Poorer	2.23	(1.92, <0.001	1.98 (1.54, 2.54)	<0.001
	2.58)			
Middle	1.63	(1.40, <0.001	1.52 (1.21, 1.91)	<0.001
	1.90)			
Richer	1.58	(1.36, <0.001	1.56 (1.26, 1.92)	<0.001
	1.83)			
Richest (RC)	1.00		1.00	
Place of residence				
Urban (RC)	1.00		1.00	
Rural	1.29	(1.17, <0.001	0.98 (0.84, 1.15)	0.864
	1.42)			
Children age				
0-11 months (RC)	1.00		1.00	
12-23 months	0.61	(0.53, <0.001	1.89 (1.58, 2.25)	<0.001
	0.70)			
24-35 months	1.16	(1.01, 0.032	2.37 (1.97, 2.85)	<0.001

	1.34)				
36-47 months	1.01	(0.87, 0.918	2.05 (1.70, 2.46)	<0.001	
	1.16)				
48-59 months	0.82	(0.71, 0.008	1.55 (1.29, 1.85)	<0.001	
	0.95)				
Sex of child					
Male (RC)	1.00		1.00		
Female	0.94	(0.86, 0.192	0.93 (0.83, 1.04)	0.192	
	1.03)				
Birth order					
First (RC)	1.00		1.00		
Second	1.02	(0.91, 0.769	1.06 (0.92, 1.21)	0.416	
	1.13)				
Third	1.19	(1.03, 0.014	1.15 (0.97, 1.36)	0.100	
	1.34)				
Fourth and above	1.63	(1.41, <0.001	1.41 (1.16, 1.72)	0.001	
	1.89)				

AORr, adjusted odds ratio-random; CI, confidence interval; COR, crude odds ratio; RC, reference category




Figure 1 Prevalence of various indicators of anthropometric failures (malnutrition's indicators)

Abbreviation: ECIAF: Extended Composite Index of Anthropometric Failure; CIAF: Composite Index of Anthropometric Failure




Figure 2 Geospatial distribution of prevalence of malnutrition, undernutrition and overweight
 Abbreviation: BDHS: Bangladesh Demographic and Health survey; CIAF: Composite Index of Anthropometric Failure ECI AF: Extended Composite Index of Anthropometric Failure;