

# ON LATTICE COMPLEMENTS

by ROBERT BUMCROT

(Received 6 January, 1964)

Let  $(L, \leq)$  be a distributive lattice with first element 0 and last element 1. If  $a, b$  in  $L$  have complements, then these must be unique, and the De Morgan laws provide complements for  $a \vee b$  and  $a \wedge b$ . We show that the converse statement holds under weaker conditions.

**THEOREM 1.** *If  $(L, \leq)$  is a modular lattice with 0 and 1 and if  $a, b$  in  $L$  are such that  $a \vee b$  and  $a \wedge b$  have (not necessarily unique) complements, then  $a$  and  $b$  have complements.*

*Proof.* We indicate by  $=*$  the steps at which the modular condition is used. Choosing any complements  $(a \vee b)'$  and  $(a \wedge b)'$  of  $a \vee b$  and  $a \wedge b$  respectively, let

$$x = (a \vee b)' \vee [(a \wedge b)' \wedge b].$$

Then

$$\begin{aligned} a \vee x &= a \vee (a \wedge b) \vee (a \vee b)' \vee [(a \wedge b)' \wedge b] \\ &= * a \vee (a \vee b)' \vee \{[(a \wedge b) \vee (a \wedge b)'] \wedge b\} = a \vee (a \vee b)' \vee b = 1, \end{aligned}$$

and

$$\begin{aligned} a \wedge x &= a \wedge (a \vee b) \wedge \{(a \vee b)' \vee [(a \wedge b)' \wedge b]\} \\ &= * a \wedge \{0 \vee [(a \wedge b)' \wedge b]\} = a \wedge (a \wedge b)' \wedge b = 0, \end{aligned}$$

so  $x$  is a complement of  $a$ . Similarly  $y = (a \vee b)' \vee [(a \wedge b)' \wedge a]$  is a complement of  $b$ .

This provides an extension of the De Morgan laws to modular lattices.

**THEOREM 2.** *If  $(L, \leq)$  is a modular lattice with 0 and 1, if  $a, b$  in  $L$  have unique complements  $a', b'$  respectively, and if  $a \vee b$  and  $a \wedge b$  have complements, then  $a' \vee b'$  is a complement of  $a \wedge b$  and  $a' \wedge b'$  is a complement of  $a \vee b$ .*

*Proof.* Form  $x$  and  $y$  as in the above proof. By dualization,  $\bar{x} = (a \wedge b)' \wedge [(a \vee b)' \vee b]$  and  $\bar{y} = (a \wedge b)' \wedge [(a \vee b)' \vee a]$  are also complements of  $a$  and  $b$  respectively. Then since  $a'$  and  $b'$  are unique,  $a' = x = \bar{x}$  and  $b' = y = \bar{y}$ . Now

$$\begin{aligned} a' \vee b' \vee (a \wedge b) &= x \vee y \vee (a \wedge b) \\ &= (a \vee b)' \vee [(a \wedge b)' \wedge b] \vee [(a \wedge b)' \wedge a] \vee (a \wedge b) \\ &= *(a \vee b)' \vee \{b \wedge [(a \wedge b)' \vee (a \wedge b)]\} \vee \{a \wedge [(a \wedge b)' \vee (a \wedge b)]\} \\ &= (a \vee b)' \vee a \vee b = 1, \end{aligned}$$

and

$$\begin{aligned} (a' \vee b') \wedge (a \wedge b) &= (\bar{x} \vee \bar{y}) \wedge (a \wedge b) \\ &= \{[(a \wedge b)' \wedge [(a \vee b)' \vee b]] \vee [(a \wedge b)' \wedge [(a \vee b)' \vee a]]\} \wedge (a \wedge b) \\ &= *(a \wedge b)' \wedge \{(a \vee b)' \vee b \vee [(a \wedge b)' \wedge [(a \vee b)' \vee a]]\} \wedge (a \wedge b) = 0, \end{aligned}$$

so  $a' \vee b'$  is a complement of  $a \wedge b$ . Dually  $a' \wedge b'$  is a complement of  $a \vee b$ .

Figure 1 shows a modular lattice of order seven which contains uniquely complemented elements  $a$  and  $b$  such that  $a \vee b$  has no complement. Contrary to Exercise 4, p. 153 of [1],

there are no lattices of order six or less, modular or otherwise, in which the complemented elements fail to form a sublattice.

If  $(L, \leq)$  is a *complemented* modular lattice, then, by a theorem of J. von Neumann [1, p. 124], the uniquely complemented elements constitute the center of the lattice, which in any lattice with 0 and 1 is a sublattice [1, p. 27]. But [1, p. 120] in a complemented modular




FIG. 1.




FIG. 2.

lattice every central element is neutral, and hence [1, p. 28] distributes with every pair of elements of the lattice. Thus in a complemented modular lattice those elements which have unique complements form a Boolean algebra, and hence satisfy the De Morgan laws. Some questions in this direction remain unanswered. For example, if  $(L, \leq)$  is modular, if  $a$  and  $b$  have unique complements in  $L$ , and if  $a \vee b$  and  $a \wedge b$  have complements, are these complements necessarily unique?

Finally, Dilworth [2] has shown, without giving an example, the existence of non-distributive lattices in which each element has a unique complement. Such lattices are non-modular by what we have said above. It seems doubtful that the De Morgan laws would hold in all such lattices. Figure 2 shows a complemented, though not uniquely complemented, non-modular lattice containing elements  $a$  and  $b$  such that  $a, b, a \vee b$  and  $a \wedge b$  have unique complements, but both De Morgan laws fail.

REFERENCES

1. G. Birkhoff, *Lattice theory* (American Mathematical Society Colloquium Publications, Vol. 25; 2nd edition, 1948).
2. R. P. Dilworth, Lattices with unique complements, *Trans. Amer. Math. Soc.* 57 (1945), 123-154.

OHIO STATE UNIVERSITY  
COLUMBUS, OHIO