

TABLE II. ANNUAL ACCUMULATION, MELTING, NET LOSS OR GAIN AT VARIOUS ELEVATIONS

Stake No.	Elevation (metres)	Centimetres of water					
		Melt started	Melt ended	Total accumulation	Melting and evaporation	Net loss	Net gain
1	440	14 May	4 Sept.	38	268	222	—
2	750	19 May	1 Sept.	40	154	106	—
A3	1010	23 June	1 Sept.	41	118	70	—
4	1060	"	"	36	97	52	—
5	1300	"	"	40	66	18	—
6	1380	"	"	40	48	0	0
7	1600	"	"	37	?	—	>12
A2	1920	"	"	63	?	—	63
8	1630	"	"	39	?	—	39
9	1940	"	"	60	?	—	60
A1	2050	"	"	43	10-15	—	43

The 1952-53 accumulation, equal to ablation, at the equilibrium line is 40 cm. water.

ACKNOWLEDGEMENTS

The authors wish to acknowledge the valuable assistance of B. Bonnlander, S. Orvig and H. Röthlisberger in maintaining observations. One of us (W. H. W.) wishes to acknowledge the help from the Royal Society for a Government Grant-in-Aid and from the Department of Scientific and Industrial Research for providing facilities for carrying out the study.

MS. received 2 January 1954

REFERENCE

1. Baird, P. D., Ward, W. H., and Orvig, S. The glaciological studies of the Baffin Island expedition, 1950. *Journal of Glaciology*, Vol. 2, No. 11, 1952, p.2.

THE INTERNATIONAL ASSOCIATION ON QUATERNARY RESEARCH, 1953

THE Fourth Meeting of the International Association for Quaternary Research (INQUA) was held in Rome and Pisa from 30 August to 10 September 1953. It was attended by some 300 students of the various aspects of the Quaternary.

The papers fell naturally into two main groups, those dealing with the archaeological aspect of the period and those dealing with the stratigraphy of the Quaternary in its broadest sense.

Little time was given to the study of modern glaciers, but several important papers were read on the divisions of the glacial period and the problems of correlation between widely separated areas. This was particularly evident in the discussions on the correlation of the various episodes of the last (4th) glacial.

During the congress numerous whole and half-day excursions were arranged. These either visited the famous Upper Palaeolithic sites of Italy or were planned to show the members the stratigraphy of the sites upon which the boundary between the Pliocene and Pleistocene was drawn at the top of the Plaisancian-Astian and base of the Calabrian-Villafranchien.

Department of Geology,
Cambridge University

W. B. R. KING