

ARTICLE

Effect of humidity and temperature on hatching success, time to hatch, and lifespan of the first-instar larvae of the hemlock looper (Lepidoptera: Geometridae)

Lucie Royer 

Natural Resources Canada, Canadian Forest Service, Laurentian Forestry Centre, City of Québec, Quebec, Canada
Email: lucie.royer@nrcan-rncan.gc.ca

(Received 25 April 2025; accepted 12 August 2025)

Abstract

Climate change affects not only temperatures but also rainfall patterns, which can either accentuate or mitigate the effects of warming on water stress in terrestrial insects. Water stress is more likely to affect eggs and larvae due to their small size. Hemlock looper (Lepidoptera: Geometridae) overwinters as eggs, and first-instar larvae must move to settle on host trees in the spring. Their survival depends both on their energy and water reserves that remain after overwintering and on the abiotic conditions present each spring. The effects of humidity (40, 60, and 80%) on the hatching success, time to hatch, and lifespan of unfed first-instar hemlock looper larvae were assessed at two temperatures, at 10 °C and at 22 °C. Lower humidity levels reduced hatching success and increased time to hatch, suggesting that humidity modulates development. On the other hand, higher temperatures reduced hatching success and time to hatch. The survival probability of unfed first-instar larvae was not influenced by ambient humidity but was longer for larvae from eggs reared at high humidity and 10 °C, suggesting that the physiological state of larvae at the time of hatching influences their survival. The ecological significance of these results and how they can influence our management tools are discussed.

Résumé

Le changement climatique affecte non seulement les températures, mais aussi les précipitations qui peuvent accentuer ou atténuer l'effet du réchauffement sur le stress hydrique des insectes terrestres. Les oeufs et les larves, en raison de leur petite taille, sont particulièrement vulnérables à ce stress. L'arpenteuse de la pruche (Lepidoptera: Geometridae) hiverne sous forme d'oeufs, et les larves de premier stade doivent se déplacer pour s'établir sur les arbres hôtes au printemps. Leur survie dépend des réserves d'énergie et d'eau après l'hivernage et des conditions abiotiques printanières. Les effets de l'humidité (40, 60 et 80 %) sur le succès et le temps d'éclosion, ainsi que la durée de vie des larves de premier stade non nourries de l'arpenteuse, ont été évalués à deux températures, à 10 °C et à 22 °C. Les faibles humidités ont réduit le succès d'éclosion et augmenté le temps d'éclosion. En revanche, les températures élevées ont diminué le succès et le temps d'éclosion. La survie des larves de premier stade n'a pas été influencée par l'humidité ambiante, mais elle a été prolongée chez les larves issues d'oeufs élevés à 10 °C et une humidité élevée. Les implications écologiques de ces résultats et leur impact sur les outils de gestion sont discutés.

Introduction

Humidity is one of the key abiotic factors that influence the abundance and distribution of terrestrial insects (Cahill *et al.* 2013; Rozen-Rechels *et al.* 2019), for which maintaining water

Subject editor: Maya Evenden

© Natural Resources Canada, 2025. Published by Cambridge University Press on behalf of Entomological Society of Canada. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.

balance is a constant challenge (Barton-Browne 1964). As climate warms, rainfall patterns also change (Dore 2005; Hicke *et al.* 2022), which could accentuate or dampen the effects of warming on water stress in terrestrial insects (Cahill *et al.* 2013; Rozen-Rechels *et al.* 2019). However, little recent research on insect responses to a changing environment has examined the impact of water stress (Bale *et al.* 2002; Chown *et al.* 2011).

Many species of forest Lepidoptera overwinter as eggs (Danks 1987). The eggs have only the fixed reserve of energy and water provided by their mother in the fall to survive overwintering, embryonic development, hatching, and settlement on the host plant. As a result, the eggs and first-instar larvae of these forest Lepidoptera are particularly vulnerable to water stress due to their limited water reserves and high surface-area-to-volume ratios (Potter and Woods 2012; Klockmann and Fischer 2017; Kühnel *et al.* 2017). In the spring, egg hatching usually coincides with the early stages of foliage expansion, a high-quality food source (Feeny 1970; Leather 1986; van Asch and Visser 2007). This foliage is often the sole source of nutrients and water available to first-instar larvae, which need to replenish their reserves after overwintering (Zalucki *et al.* 2002). Climate change can affect insect and plant phenology differently, which may result in a mismatch between egg hatching and the most suitable host phenology (Parmesan 2007; van Asch and Visser 2007). A similar mismatch may occur if moisture alters the insect's growth rate, as suggested by Tauber *et al.*'s (1998) "development modulator" hypothesis. In addition, young larvae may die of starvation or desiccation while searching for feeding sites (Zalucki *et al.* 2002). This mortality of young larvae before they settle on the host plant may be accentuated by low ambient humidity or by the initial hydration state of the larvae, as suggested by Woods's (2010) "neonate desiccation" hypothesis. However, the impact of water stress on the embryonic development and its potential cross-stage effect in the forest Lepidoptera remain largely unexplored.

The hemlock looper, *Lambdina fiscellaria* (Guenée) (Lepidoptera: Geometridae), is a native univoltine moth whose larvae periodically defoliate the coniferous forests of Canada (Martineau 1984). In September, females lay their eggs singly on sites, such as soil, mosses, lichens, and tree branches and stems, where humidity conditions are highly variable (Carroll 1956). The eggs undergo an obligatory diapause that lasts about three months, after which they remain in quiescence until spring warming triggers the resumption of embryonic development (Delisle *et al.* 2009). Egg hatching occurs from mid-May to late June in eastern Canada and generally coincides with an elongation of 25–35% of the maximum annual growth of balsam fir, *Abies balsamea* (Linnaeus) Miller (Pinaceae) (Butt *et al.* 2010). This synchronism between hatching time and the phenology of its preferred host tree results in optimal insect survival and fitness (Butt *et al.* 2010), provided that the first-instar looper larvae can reach the new shoots. The first-instar larvae must travel from the laying sites to the current year foliage to feed, a movement facilitated by the larvae's responses to gravity and light (Royer *et al.* 2021). Because the consumption of new foliage is crucial to young looper larvae survival, high mortality can occur when eggs are laid far from preferred feeding sites and abiotic conditions are unfavourable (Carroll 1956). Therefore, the ambient humidity and the initial hydration state of the larvae could influence their ability to survive and settle on host trees.

The hemlock looper mainly occurs in a band of about 10° latitude straddling the Canada–United States of America border from coast to coast, from Newfoundland and Labrador to British Columbia in Canada and from Maine to Washington State in the United States of America (Martineau 1984; McGuffin 1987). The species can also be found in remote areas, such as Labrador, Alaska, Georgia, Oregon, and California (Torgersen and Baker 1971; Martineau 1984; McGuffin 1987; Turnquist 1991; Crummey 2007). In Canada, major outbreaks occur in Newfoundland, the Maritime provinces, Quebec, Ontario, and British Columbia, mainly along coasts and in humid inland forests (Carroll 1956; Otvos *et al.* 1979; Turnquist 1991). Although humidity seems to play a key role in hemlock looper outbreaks, to date, no studies have investigated the effect of humidity on the performance of the looper stages that may be most vulnerable to water stress, such as eggs and first-instar larvae.

Table 1. Mean relative humidity and temperature recorded in the various treatments during the experiment

Salt sugar solution	Relative humidity $\pm$ standard deviation (%)	Temperature $\pm$ standard deviation ($^{\circ}$ C)
K ₂ CO ₃ .1.5H ₂ O	43.4 $\pm$ 3.4	10.2 $\pm$ 0.9
Glucose	61.2 $\pm$ 9.5	10.3 $\pm$ 0.8
KCl	85.6 $\pm$ 1.4	10.2 $\pm$ 0.9
K ₂ CO ₃ .1.5H ₂ O	44.3 $\pm$ 1.7	22.2 $\pm$ 0.2
Glucose	66.9 $\pm$ 1.7	22.3 $\pm$ 0.2
KCl	79.8 $\pm$ 3.8	22.0 $\pm$ 0.7

The present study aims to quantify the effect of air humidity (40, 60, and 80%) on hemlock looper hatching success, time to hatch, and lifespan of unfed first-instar larvae in spring. Experiments were carried out at two temperatures (10 $^{\circ}$ C and 22 $^{\circ}$ C) to determine the extent to which this factor can attenuate or exacerbate the impact of humidity. First, the effect of humidity and temperature on hatching success was quantified to test the null hypothesis that these two abiotic factors have no influence. Second, the time to hatch was quantified to test the development modulator hypothesis, assuming that humidity has no effect. Finally, the effect of humidity at the egg and larval stages on the lifespan of unfed first-instar looper larvae was assessed to test the neonate desiccation hypothesis. No cross-stage effect of humidity was expected.

Materials and methods

In winter 2001, diapausing eggs of the Newfoundland hemlock looper were obtained from the Canadian Forest Service's Insect Production and Quarantine Laboratories (Sault Ste. Marie, Ontario, Canada). Upon receipt, the eggs were stored in total darkness at 2.5 ± 0.5 $^{\circ}$ C and $95 \pm 5\%$ relative humidity until they were used in spring experiments.

To obtain the targeted humidity of 40, 60, and 80% (Table 1), supersaturated solutions were prepared by dissolving salts or sugars in distilled water heated to boiling point (Winston and Bates 1960). The solutions were cooled and decanted into plastic boxes (11.4-L storage box; Rubbermaid Inc., Wooster, Ohio, United States of America). One box for each humidity was then placed in growth chambers at 10.0 ± 0.5 $^{\circ}$ C or 22.0 ± 0.5 $^{\circ}$ C. Historically (1991–2020), mean daily relative humidity during the month preceding egg hatching on the west coast of the island of Newfoundland ranges from 56.3% to 82.4%, and mean daily temperatures vary from 6.9 $^{\circ}$ C to 12.1 $^{\circ}$ C, with maximums reaching 18.6 $^{\circ}$ C (Environment Canada 2025a). However, daily mean temperatures in the Lac Saint-Jean region of Quebec, where outbreaks are less frequent, vary from 10.2 $^{\circ}$ C to 16.0 $^{\circ}$ C, with maximums of 22.3 $^{\circ}$ C, and mean relative humidity ranges from 47.5% to 78.5% (Environment Canada 2025b). These data show that the temperatures selected for the present study are encountered in the current range of the species during the month preceding egg hatching. The low humidity tested in the study was slightly lower than the natural conditions observed, but it was chosen to assess the insect's ability to adjust its physiology.

To determine the combined effect of humidity and temperature on hatching success and time to hatch, hemlock looper eggs were placed individually in Beem capsules (size 3; SPI Supplies, West Chester, Pennsylvania, United States of America). Capsule lids were partially closed to prevent egg loss, leaving an open space for internal and external humidity equilibration. Groups of 50 randomly selected capsules were put in 127-mL plastic cups (model TP400; Solo Cup Co, Lake Forest, Illinois, United States of America), which were pierced with 0.5-cm holes to balance the

Table 2. Number of hemlock looper first-instar larvae whose lifespan was monitored to determine the effect of humidity (40, 60, and 80%) at the egg and larval stages at two temperatures (10 °C and 22 °C)

T (°C)	Humidity during egg stage (%)	Humidity during larval stage (%)		
		40	60	80
10	40	57	36	36
	60	115	140	130
	80	153	151	183
22	40	86	84	83
	60	102	97	101
	80	107	102	111

internal humidity of the cup with the external ambient conditions. The cups were then transferred to wire grids suspended two centimetres off the supersaturated solution in the treatment boxes that were placed in the growth chamber at the desired temperature. The actual relative humidity and temperature conditions of each treatment box were recorded using data loggers (Model HOBO® H08-004-02; Hoskin Scientific Ltd, Saint-Laurent, Quebec, Canada) and are presented in Table 1. Eggs were checked daily to estimate hatching success and time to hatch. The time spent away from treatment was minimised as much as possible and was equivalent for each treatment. To calculate the time to hatch, day 0 corresponded to the day eggs were taken out of cold storage and transferred to growth chambers. At the end of the experiment, unhatched eggs were classified according to three categories: no apparent development, collapsed, and fully developed first instar inside the egg, hereafter called “pharate” larvae (Stacey *et al.* 1975). A percentage of each category of unhatched eggs was calculated based on total mortality for each temperature and humidity. Ten cups of 50 eggs were tested for each humidity and temperature treatment.

To assess the effect of humidity during the egg and larval stages on the survival of unfed first-instar larvae, two cups of 50 eggs were reared in each combination of humidity and temperature because high mortality was expected in some treatments of the previous experiment. Larvae from these additional cups were added to those from the previous experiment. Upon hatching, neonates were randomly reassigned to one of three humidity treatments – 40, 60, and 80% humidity – but were maintained in their initial temperature treatment (10 °C or 22 °C). This resulted in a total of nine treatments per temperature. Table 2 shows the number of larvae per treatment, and the survival of which was checked daily. The lifespan of 1001 and 873 first-instar larvae was assessed at 10 °C and at 22 °C, respectively.

Statistical analyses

The appropriate distribution or transformation of data on hatching success, percentage of dead eggs that reached the pharate larval stage, and time to hatch was determined through an Individual Distribution Identification procedure, based on an Anderson–Darling test. Analyses of variance were then used to estimate the effect of humidity and temperature treatments, as well as of their interaction on hatching success and the squared percentage of dead eggs that reached the pharate larval stage. Following the procedure to identify the distribution or transformation, the natural logarithm of the time to hatch was averaged for each cup (Delisle *et al.* 2009). Each log-scale mean was weighted by the number of eggs that hatched in the cup, and the variance of these means was analysed using a model that had as fixed effects temperature, humidity, and their interactions (Delisle *et al.* 2009). All these analyses were followed by a comparison of all pairs of means using the Bonferroni method, with an overall confidence level of 95%. The means of the




Figure 1. Combined effect of humidity (40, 60, and 80%) and temperature (10 °C and 22 °C) on the mean hatching success ($\pm$ standard error of the mean) of hemlock looper eggs under a 16:8-hour light:dark photoperiod. Means followed by different letters are significantly different at $P=0.05$ (analysis of variance followed by a Bonferroni test).




Figure 2. Combined effect of humidity (40, 60, and 80%) and temperature (10 °C and 22 °C) on the mean percentage ($\pm$ standard error of the mean) of dead hemlock looper eggs having reached the pharate larval stage under a 16:8-hour light:dark photoperiod. Means followed by different letters are significantly different at $P=0.05$ (analysis of variance followed by a Bonferroni test).




Figure 3. Combined effect of humidity (40, 60, and 80%) and temperature (10 °C and 22 °C) on the mean time to hatch ($\pm$ standard error of the mean) of hemlock looper eggs under a 16:8-hour light:dark photoperiod. Means followed by different letters are significantly different at $P=0.05$ (analysis of variance followed by a Bonferroni test).

original untransformed data and their standard errors of the means are presented in Figures 1 to 3, and all interpretations were based on the original untransformed data. In addition, linear relationships between hatching success and time to hatch per cup were established for each temperature using the 10 replicates per humidity treatment.

To assess the effect of humidity on the probability of larval survival at each temperature, a Distribution ID Plot procedure was first conducted to identify the probability distribution that best matched the survival data of first-instar larvae. The least-squares method was used to estimate the population parameters, fit a regression line that minimises the sum of squared deviations (least-squares error), and calculate a Pearson correlation coefficient. The Weibull distribution was selected because it obtained the highest correlation coefficients ($R^2_{10\text{ °C}} > 0.96$, $R^2_{22\text{ °C}} > 0.89$). A regression with life data (reliability/survival) analysis for each temperature was then conducted to determine whether humidity during egg or larval stages affected the probability of survival of unfed first-instar larvae, assuming a Weibull distribution. The full model included humidity during the egg and larval stages and their interaction as factors. The probability plot for the Cox–Snell residuals produced by this analysis confirmed the chosen Weibull distribution assumption. All statistical tests were performed using Minitab Statistical Software, version 18 (Minitab; eBase Solutions Inc., Vaughan, Ontario, Canada).

Table 3. Summary of variance analyses assessing the influence of humidity and temperature on hatching success, squared percentage of dead eggs reaching the pharate larval stage, and natural logarithm of time to hatch per cup of hemlock looper eggs

Variables	Source	df	F-value	P-value
Hatching success (%)	Humidity	2	149.21	< 0.0001
	Temperature	1	20.94	< 0.0001
	Humidity*temperature	2	67.72	< 0.0001
Pharate larva (%)	Humidity	2	5.20	0.009
	Temperature	1	58.45	< 0.0001
	Humidity*temperature	2	6.06	0.004
Time to hatch (day)	Humidity	2	221.29	< 0.0001
	Temperature	1	74654.28	< 0.0001
	Humidity*temperature	2	97.05	< 0.0001

Results

The hatching success of hemlock looper eggs was lower at 22 °C than at 10 °C under all humidity treatments except 40% humidity, when the opposite was observed (Table 3; Fig. 1). Decreased humidity reduced the hatching success, but this effect was more pronounced at 10 °C than at 22 °C. The percentage of dead eggs that reached the pharate larval stage did not differ with humidity at 10 °C and corresponded to more than 89% of dead eggs (Table 3; Fig. 2). The remaining dead eggs either showed no development ($8.2 \pm 1.6\%$) or collapsed ($1.6 \pm 0.6\%$). At 22 °C, the percentage of dead eggs reaching the pharate larval stage rose from 69% to 85% when humidity decreased (Table 3; Fig. 2). The other eggs died with no apparent development.

Time to hatch was five times longer at 10 °C than at 22 °C (50.5 days *versus* 8.9 days at 40% humidity and 38.6 days *versus* 8.4 days at 80% humidity) and was inversely correlated with humidity (Table 3; Fig. 3). The humidity effect was more pronounced at 10 °C, when eggs reared at 40% humidity took 31% longer to hatch than those reared at 80% humidity (50.5 and 38.6 days, respectively). Time to hatch explained 92% of the variation in hatching success at 10 °C ($F_{1, 28} = 307.50$; $P < 0.0001$; $y = -4.97x + 275.92$; $R^2 = 0.92$) and 45% of the variation in hatching success at 22 °C ($F_{1, 28} = 22.70$; $P < 0.0001$; $y = -19.66x + 218.30$; $R^2 = 0.45$).


Humidity during the larval stage did not adversely affect the survival probability of unfed first-instar larvae at 10 °C or at 22 °C (Table 4). However, humidity during the egg stage influenced the survival probability of larvae at low temperature but not at high temperature (Table 4). An increase in humidity at the egg stage prolonged the survival probability of larvae kept at 10 °C (Fig. 4A), as predicted by Woods’s (2010) neonate desiccation hypothesis. Indeed, the average 50% survival probability of unfed first-instar larvae was 3.8 ± 0.1 , 4.4 ± 0.1 , and 5.2 ± 0.1 days when eggs were reared at 40, 60, and 80% humidity, respectively. In contrast, the 50% survival probability of larvae reared at 22 °C was only 2.6 ± 0.1 days on average, regardless of humidity treatments (Fig. 4B). However, daily observations may not have been sufficiently frequent to detect small differences in this temperature. Therefore, a small humidity impact cannot be ruled out.

Discussion

In the context of climate change, warmer, drier springs and summers are predicted in some areas of eastern Canada (Williamson *et al.* 2009; Hicke *et al.* 2022), and marked rainfall deficits have already been observed (World Meteorological Organisation 2024, 2025). This study shows

Table 4. Summary of regressions assessing the effect of humidity during egg (RH_{egg}) and larval (RH_{larva}) stages on the survival probability of hemlock looper first-instar larvae at 10 °C and at 22 °C

Temperature	Source	Z-value	P-value
10 °C	Intercept	5.75	< 0.0001
	RH_{egg}	2.20	0.028
	RH_{larva}	-0.56	0.575
	$RH_{\text{egg}} \times RH_{\text{larva}}$	0.39	0.693
22 °C	Intercept	8.59	< 0.0001
	RH_{egg}	-0.14	0.891
	RH_{larva}	-0.54	0.590
	$RH_{\text{egg}} \times RH_{\text{larva}}$	0.97	0.332

**Figure 4.** Effect of humidity (40, 60, and 80%) at the egg stage on survival probabilities of hemlock looper first-instar larvae **A**, at 10 °C and **B**, at 22 °C. Survival data of first-instar larvae subjected to the different humidity levels during the larval stage were pooled. Error bars represent the 95% confidence interval.

that (1) spring mortality of hemlock looper eggs increases with increasing temperature and decreasing humidity, (2) warm temperatures speed up embryonic development, but low humidities slow it down, and (3) humidity at the egg stage has a greater influence on survival of subsequent larvae than does humidity during the larval stage.

The sessile eggs of hemlock looper cannot alleviate thermal or water stresses by moving. As a result, their survival depends on the limited reserves of energy and water provided by their mother in the fall and the microhabitat where she chose to lay her eggs. In the present study, looper hatching success was reduced at high temperatures and low humidity levels, suggesting that warm and dry springs are likely to affect the population densities of this species. The high mortality rate of looper eggs in warm and dry environments could be explained by exchange mechanisms between the eggs and their environment, similar to those demonstrated in *Manduca sexta* (Linnaeus) (Lepidoptera: Sphingidae). In this species, eggs adjust their shell conductance to delay water loss under dry conditions, but these changes are accompanied by a decrease in oxygen and carbon dioxide exchanges with the environment (Ludwig and Anderson 1942; Woods *et al.* 2005;

Zrubek and Woods 2006). High temperatures amplify the vapour pressure gradient between the eggs and the environment (Wharton and Richards 1978) and increase the metabolic rate of eggs (Woods and Singer 2001), which increases the eggs' oxygen demand and water loss (Woods and Hill 2004). However, water loss and reduced gas exchange cause a decrease in the metabolic rate, which prolongs embryonic development and leads to death if unfavourable conditions persist (Chaplin 2001; Woods and Hill 2004; Woods *et al.* 2005; Zrubek and Woods 2006; Rozen-Rechels *et al.* 2019). In the present study, the loopers' time to hatch did indeed increase at lower humidities under both temperature treatments, which is in line with Tauber *et al.*'s (1998) development modulator hypothesis. Hatching success was also inversely correlated with the time to hatch in the present study, suggesting that the effects of humidity depend on exposure time, similar to what Godfrey and Holtzer (1991) observed in *Ostrinia nubilalis* (Hübner) (Lepidoptera: Crambidae). However, in the present study, the effect of humidity was less pronounced at 22 °C than at 10 °C because the time to hatch was shorter at this temperature. At the two temperatures studied here, mortality of hemlock looper eggs occurred late in embryonic development, with more than 69% of dead eggs reaching the pharate larval stage. Possible reasons for this late mortality include desiccation, hypoxia, or both (Ludwig and Anderson 1942; Woods *et al.* 2005; Zrubek and Woods 2006), weakening of the larvae by resource depletion, desiccation-caused chorion hardening, and the resultant prevention of hatching (Buxton 1932; Clark 1935), or a combination of these factors.

In successful hatchlings, the probability of survival was unaffected by humidity at the larval stage, regardless of temperature regime. The cuticle of Lepidoptera larvae is generally impermeable to water (Wigglesworth 1945), and water loss can be reduced by regulating gas exchange at the spiracles (Buxton 1932; Buck 1962). However, larvae also face a trade-off between metabolic gas exchange and water conservation. The probability of survival for 50% of unfed larvae was greater at 10 °C than at 22 °C, as the latter temperature accelerated their metabolism. The effect of humidity during the egg stage on larval survival was only detected at 10 °C. At this temperature, the probability of survival decreased by about 15% for each 20% decrease in humidity during the egg stage. This result suggests that the physiological state of the looper at hatching affects the probability of subsequent larval survival, which is consistent with Woods's (2010) neonate desiccation hypothesis. Such a cross-stage effect could be widespread in forest insects that overwinter as eggs and exploit the trees in early spring. Spring conditions that shorten the survival of unfed first instars reduce the time during which hemlock looper larvae can search for suitable feeding sites, thereby also decreasing the likelihood they will settle on host trees (Schneider 1980; Leather 1986). Overall, drier and warmer springs, as are predicted under climate change, are expected to reduce the survival of looper eggs and larvae. Further studies are recommended to quantify the simultaneous effect of temperature and humidity gradients on water loss and gas exchange in looper eggs and larvae, with particular attention paid to the impact on time to hatch because any change could disrupt the trophic interactions of the hemlock looper in areas of its range affected by unfavourable conditions.

The timing of hemlock looper egg hatch is crucial because larval survival and growth depend on the host plant's phenology. Hatching of looper eggs is usually synchronised with a 25–35% elongation of balsam fir shoots, which allows larvae to achieve the highest fitness (Butt *et al.* 2010). Spring drought can compromise this optimal timing, with low humidity prolonging the embryonic development of the hemlock looper, as seen in the present study. Indeed, the hatching time at 10 °C was extended by 12 days at 40% humidity compared to hatching time at 80% humidity, which may be enough to shift hatching beyond the most suitable foliage window for looper larval development. Such a mismatch leads to reduced survival, size, and fecundity (Carroll 1999; Butt *et al.* 2010). Warmer temperatures may also cause mismatches. Bud development in mature conifers in boreal forests is mainly driven by soil temperature and photoperiod cues (Delpierre *et al.* 2016), whereas embryonic development of hemlock looper eggs depends on air temperature (Delisle *et al.* 2009). However, in spring, the air tends to warm faster

than the soil does. Warmer temperatures may thus shift hatching earlier than the most suitable looper–foliage window. Butt *et al.* (2010) found that only 30% of the first-instar larvae placed on balsam fir before bud break survive, and survivors pay a high fitness cost. Any decoupling between the phenological phases of the insect and those of its host plant may contribute to a decline of hemlock looper populations.

Spring conditions can also affect hemlock looper interactions with its natural enemies. The prolongation of embryonic development as humidity falls can lengthen the window of time during which eggs are vulnerable to parasitoid attack in spring. Adding to this risk, natural enemies seem less sensitive to changes in temperature and humidity than their hosts or prey species are (Thackeray *et al.* 2016). Among parasitoids of hemlock looper eggs, *Telenomus coloradensis* Crawford (Hymenoptera: Scelionidae) is the most abundant (Pelletier and Piché 2003; Carleton *et al.* 2009), and its attack depends on host density (Carleton *et al.* 2010). *Telenomus* adults are active at temperatures as low as 4 °C and can live for more than 55 days at 24 °C (Legault *et al.* 2012). They are also able to mitigate the effect of low air humidity by drinking dewdrops or choosing moist microhabitats. Any increased overlap between host and parasitoid can significantly reduce hemlock looper populations, as *Telenomus* can cause them to collapse (Hébert *et al.* 2001). On the other hand, the period of egg vulnerability to *Telenomus* attack is expected to decrease when temperatures warm because the time to hatch was about five times shorter at 22 °C than at 10 °C. Such a change in the window of egg vulnerability may also impact hemlock looper populations.

The present study shows that dry springs can lead to high mortality of looper eggs, prolong looper embryonic development, and reduce the survival of pre-feeding looper larvae, thereby decreasing their chances of settling on the host tree. Drier springtime conditions could also cause the timing of looper–host development to decouple and increase the eggs’ vulnerability to its dominant egg parasitoid. Warmer temperatures may exacerbate the effects of dry springs on looper mortality but could mitigate the impacts of low humidity on larvae–host mismatches. In the context of climate change, these results suggest that a multidimensional approach that includes temperature and humidity would be more appropriate to predict changes in hemlock looper range and abundance. In addition, phenological models based solely on temperature can wrongly predict looper hatching time, which is delayed significantly under low-humidity conditions. Integration of a humidity component would help to improve the models used to plan control interventions for this pest, as Godfrey and Holtzer (1991), Ortega-López *et al.* (2014), and Liu *et al.* (2015) have demonstrated with other species. The new, integrated models that take account of water conditions would help to improve hemlock looper management and our understanding of the pest’s seasonal and demographic responses to climate change.

Acknowledgements. The author thanks Amanda Follett and Julie Robbins for technical assistance, as well as Anne-Marie Dion, Alain Labrecque, and three anonymous reviewers for their constructive comments on earlier versions of this manuscript. Natural Resources Canada, Canadian Forest Service funded this research.

Competing interests. The author declares that she has no competing interests.

References

- Bale, J.S., Masters, G.J., Hodkinson, I.D., Awmack, C., Bezemer, T.M., Brown, V.K., *et al.* 2002. Herbivory in global climate change research: direct effects of rising temperature on insect herbivores. *Global Change Biology*, **8**: 1–16. <https://doi.org/10.1046/j.1365-2486.2002.00451.x>.
- Barton-Browne, L.B. 1964. Water regulation in insects. *Annual Review of Entomology*, **9**: 63–82. <https://doi.org/10.1146/annurev.en.09.010164.000431>.
- Buck, J. 1962. Some physical aspects of insect respiration. *Annual Review of Entomology*, **7**: 27–56. <https://doi.org/10.1146/annurev.en.07.010162.000331>.

- Butt, C., Quiring, D., Hébert, C., Delisle, J., Berthiaume, R., Bauce, E., and Royer, L. 2010. Influence of balsam fir (*Abies balsamea*) budburst phenology on hemlock looper (*Lambdina fiscellaria*). *Entomologia Experimentalis et Applicata*, **134**: 220–226. <https://doi.org/10.1111/j.1570-7458.2009.00963.x>.
- Buxton, P.A. 1932. Terrestrial insects and the humidity of the environment. *Biological Reviews*, **7**: 275–320. <https://doi.org/10.1111/j.1469-185X.1962.tb01044.x>.
- Cahill, A.E., Aiello-Lammens, M.E., Fisher-Reid, M.C., Hua, X., Karanewsky, C.J., Ryu, H.Y., *et al.* 2013. How does climate change cause extinction? *Proceedings of the Royal Society B: Biological Sciences*, **280**: 20121890. <https://doi.org/10.1098/rspb.2012.1890>.
- Carleton, D., Quiring, D., Heard, S., Hébert, C., Delisle, J., Berthiaume, R., *et al.* 2010. Density-dependent and density-independent responses of three species of *Telenomus* parasitoids of hemlock looper eggs. *Entomologia Experimentalis et Applicata*, **137**: 296–303. <https://doi.org/10.1111/j.1570-7458.2010.01065.x>.
- Carleton, D., Royer, L., Hébert, C., Delisle, J., Berthiaume, R., Bauce, E., and Quiring, D. 2009. Seasonal parasitism of hemlock looper (Lepidoptera: Geometridae) eggs in eastern Canada. *The Canadian Entomologist*, **141**: 614–618. <https://doi.org/10.4039/n09-040>.
- Carroll, A.L. 1999. Physiological adaptation to temporal variation in conifer foliage by a caterpillar. *The Canadian Entomologist*, **131**: 659–669. <https://doi.org/10.4039/Ent131659-5>.
- Carroll, W.J. 1956. History of the hemlock looper, *Lambdina fiscellaria fiscellaria* (Guen.) (Lepidoptera: Geometridae), in Newfoundland, and notes on its biology. *The Canadian Entomologist*, **88**: 587–599. <https://doi.org/10.4039/Ent88587-10>.
- Chaplin, M.F. 2001. Water: its importance to life. *Biochemistry and Molecular Biology Education*, **29**: 54–59. [https://doi.org/10.1016/S1470-8175\(01\)00017-0](https://doi.org/10.1016/S1470-8175(01)00017-0).
- Chown, S.L., Sørensen, J.G., and Terblanche, J.S. 2011. Water loss in insects: an environmental change perspective. *Journal of Insect Physiology*, **57**: 1070–1084. <https://doi.org/10.1016/j.jinsphys.2011.05.004>.
- Clark, N. 1935. The effect of temperature and humidity upon the eggs of the bug, *Rhodnius prolixus* (Heteroptera, Reduviidae). *Journal of Animal Ecology*, **4**, 82–87. <https://doi.org/10.2307/1215>.
- Crummey, H. 2007. Status of important forest insect activity in 2006 and outlook for 2007 – Newfoundland and Labrador. In *Proceedings of the Forest Pest Management Forum 2006*, December 5–6–7, 2006, Ottawa, Ontario, Canada. Natural Resources Canada, Canadian Forest Service, Québec, Quebec, Canada. Pp. 167–178.
- Danks, H.V. 1987. Insect Dormancy: An Ecological Perspective. Biological Survey of Canada (Terrestrial Arthropods), Ottawa, Ontario, Canada.
- Delisle, J., Royer, L., Bernier-Cardou, M., Bauce, E., and Labrecque, A. 2009. The combined effect of photoperiod and temperature on egg dormancy in an island and a mainland population of the hemlock looper, *Lambdina fiscellaria*. *Entomologia Experimentalis et Applicata*, **133**: 232–243. <https://doi.org/10.1111/j.1570-7458.2009.00931.x>.
- Delpierre, N., Vitasse, Y., Chuine, I., Guillemot, J., Bazot, S., Rutishauser, T., and Rathgeber, C.B.K. 2016. Temperate and boreal forest tree phenology: from organ-scale processes to terrestrial ecosystem models. *Annals of Forest Science*, **73**: 5–25. <https://doi.org/10.1007/s13595-015-0477-6>.
- Dore, M.H.I. 2005. Climate change and changes in global precipitation patterns: what do we know? *Environment International*, **31**: 1167–1181. <https://doi.org/10.1016/j.envint.2005.03.004>.
- Environment Canada. 2025a. Canadian climate normals 1991–2020 data for Deer Lake, Newfoundland and Labrador. Available from https://climate.weather.gc.ca/climate_normals/results_1991_2020_e.html?searchType=stnName_1991&txtStationName_1991=Deer+Lake&searchMethod=contains&txtCentralLatMin=0&txtCentralLatSec=0&txtCentralLongMin=0&txtCentralLongSec=0&stnID=61000000&dispBack=1 [accessed 26 August 2025].

- Environment Canada. 2025b. Canadian climate normals 1991–2020 data for Bagotville, Quebec. Available from https://climate.weather.gc.ca/climate_normals/results_1991_2020_e.html?searchType=stnName_1991&txtStationName_1991=Bagotville&searchMethod=contains&txtCentralLatMin=0&txtCentralLatSec=0&txtCentralLongMin=0&txtCentralLongSec=0&stnID=83000000&dispBack=1 [accessed 26 August 2025].
- Feeny, P. 1970. Seasonal changes in oak leaf tannins and nutrients as a cause of spring feeding by winter moth caterpillars. *Ecology*, **51**: 565–581. <https://doi.org/10.2307/1934037>.
- Godfrey, L.D. and Holtzer, T.O. 1991. Influence of temperature and humidity on European corn borer (Lepidoptera: Pyralidae) egg hatchability. *Environmental Entomology*, **20**: 8–14. <https://doi.org/10.1093/ee/20.1.8>.
- Hébert, C., Berthiaume, R., Dupont, A., and Auger, M. 2001. Population collapses in a forecasted outbreak of *Lambdina fiscellaria* (Lepidoptera: Geometridae) caused by spring egg parasitism by *Telenomus* spp. (Hymenoptera: Scelionidae). *Environmental Entomology*, **30**: 37–43. <https://doi.org/10.1603/0046-225X-30.1.37>.
- Hicke, J.A., Lucatello, S., Mortsch, L.D., Dawson, J., Domínguez Aguilar, M., Enquist, C.A.F., *et al.* 2022. North America. Chapter 14. *In* Climate Change 2022: Impacts, Adaptation and Vulnerability. Contribution of Working Group II to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change. *Edited by* H.-O. Pörtner, D.C. Roberts, M. Tignor, E.S. Poloczanska, K. Mintenbeck, A. Alegría, *et al.* Cambridge University Press, Cambridge, United Kingdom and New York, New York, United States of America. Pp. 1929–2042. Available from <https://www.ipcc.ch/report/ar6/wg2/chapter/chapter-14/> [accessed 26 August 2025].
- Klockmann, M. and Fischer, K. 2017. Effects of temperature and drought on early life stages in three species of butterflies: mortality of early life stages as a key determinant of vulnerability to climate change? *Ecology and Evolution*, **7**: 10871–10879. <https://doi.org/10.1002/ece3.3588>.
- Kühse, S., Brückner, A., Schmelzle, S., Heethoff, M., and Blüthgen, N. 2017. Surface area–volume ratios in insects. *Insect Science*, **24**: 829–841. <https://doi.org/10.1111/1744-7917.12362>.
- Leather, S.R. 1986. The effect of neonatal starvation on the growth, development and survival of larvae of the pine beauty moth, *Panolis flammea* (D & S). *Oecologia*, **71**: 90–93. <https://doi.org/10.1007/BF00377325>.
- Legault, S., Hébert, C., Blais, J., Berthiaume, R., Bause, E., and Brodeur, J. 2012. Seasonal ecology and thermal constraints of *Telenomus* spp. (Hymenoptera: Scelionidae), egg parasitoids of the hemlock looper (Lepidoptera: Geometridae). *Environmental Entomology*, **41**: 1290–1301. <https://doi.org/10.1603/en12129>.
- Liu, B., Li, H.Q., Ali, A., Li, H.B., Liu, J., Yang, Y.Z., and Lu, Y.H. 2015. Effects of temperature and humidity on immature development of *Lygus pratensis* (L.) (Hemiptera: Miridae). *Journal of Asia-Pacific Entomology*, **18**: 139–143. <https://doi.org/10.1016/j.aspen.2014.12.011>.
- Ludwig, D. and Anderson, J.M. 1942. Effects of different humidities, at various temperatures, on the early development of four saturniid moths (*Platysamia cecropia* Linnaeus, *Telea polyphemus* Cramer, *Samia walkeri* Felder and Felder, and *Callosamia promethea* Drury), and on the weights and water contents of their larvae. *Ecology*, **23**: 259–274. <https://doi.org/10.2307/1930667>.
- Martineau, R. 1984. *Insects Harmful to Forest Trees*. Multiscience Publications, Ottawa, Ontario, Canada.
- McGuffin, W.C. 1987. Guide to the Geometridae of Canada (Lepidoptera), II. Subfamily Ennominae. 4. *Memoirs of the Entomological Society of Canada*, **119**: 5–182. <https://doi.org/10.4039/entm119138fv>.
- Ortega-López, V., Amo-Salas, M., Ortiz-Barredo, A., and Díez-Navajas, A.M. 2014. Male flight phenology of the European grapevine moth, *Lobesia botrana* (Lepidoptera: Tortricidae), in different wine-growing regions in Spain. *Bulletin of Entomological Research*, **104**: 566–575. <https://doi.org/10.1017/S0007485314000339>.

- Otvos, I.S., Clarke, L.J., and Durling, D.S. 1979. A history of recorded eastern hemlock looper outbreaks in Newfoundland. Information Report N-X-179. Environment Canada, Forestry Service, Newfoundland Forest Research Centre, St. John's, Newfoundland and Labrador, Canada. 46 pp. Available from <https://ostrnrcan-dostrncan.canada.ca/handle/1845/239780> [accessed 26 August 2025].
- Parmesan, C. 2007. Influences of species, latitudes and methodologies on estimates of phenological response to global warming. *Global Change Biology*, **13**: 1860–1872. <https://doi.org/10.1111/j.1365-2486.2007.01404.x>.
- Pelletier, G. and Piché, C. 2003. Species of *Telenomus* (Hymenoptera: Scelionidae) associated with the hemlock looper (Lepidoptera: Geometridae) in Canada. *The Canadian Entomologist*, **135**: 23–39. <https://doi.org/10.4039/n02-035>.
- Potter, K.A. and Woods, H.A. 2012. No evidence for the evolution of thermal or desiccation tolerance of eggs among populations of *Manduca sexta*. *Functional Ecology*, **26**: 112–122. <https://doi.org/10.1111/j.1365-2435.2011.01912.x>.
- Royer, L., Delisle, J., and Labrecque, A. 2021. Ontogenetic changes in the behavioral response to light and gravity in a geometrid larva. *Journal of Insect Behavior*, **34**: 61–69. <https://doi.org/10.1007/s10905-021-09768-y>.
- Rozen-Rechels, D., Dupoué, A., Lourdais, O., Chamaillé-Jammes, S., Meylan, S., Clobert, J., and Le Galliard, J.F. 2019. When water interacts with temperature: ecological and evolutionary implications of thermo-hydroregulation in terrestrial ectotherms. *Ecology and Evolution*, **9**: 10029–10043. <https://doi.org/10.1002/ece3.5440>.
- Schneider, J.C. 1980. The role of parthenogenesis and female aptery in microgeographic, ecological adaptation in the fall cankerworm, *Alsophila pometaria* Harris (Lepidoptera: Geometridae). *Ecology*, **61**: 1082–1090. <https://doi.org/10.2307/1936827>.
- Stacey, L., Roe, R., and Williams, K. 1975. Mortality of eggs and pharate larvae of the eastern tent caterpillar, *Malacosoma americana* (F.) (Lepidoptera: Lasiocampidae). *Journal of the Kansas Entomological Society*, **48**: 521–523. Available from <http://www.jstor.org/stable/25078485> [accessed 26 August 2025].
- Tauber, M.J., Tauber, C.A., Nyrop, J.P., and Villani, M.G. 1998. Moisture, a vital but neglected factor in the seasonal ecology of insects: hypotheses and tests of mechanisms. *Environmental Entomology*, **27**: 523–530. <https://doi.org/10.1093/ee/27.3.523>.
- Thackeray, S.J., Henrys, P.A., Hemming, D., Bell, J.R., Botham, M.S., Burthe, S., *et al.* 2016. Phenological sensitivity to climate across taxa and trophic levels. *Nature*, **535**: 241–245. <https://doi.org/10.1038/nature18608>.
- Torgersen, T.R. and Baker, B.H. 1971. The occurrence of the hemlock looper (*Lambdina fiscellaria* (Guenée)) (Lepidoptera: Geometridae) in southeast Alaska, with notes on its biology. Research Notes No. PNW-61. United States Department of Agriculture, Forest Service, Pacific Northwestern Forest and Range Experiment Station, Portland, Oregon, United States of America. 6 pp. Available from https://archive.org/details/nfsl_1_13231 [accessed 26 August 2025].
- Turnquist, R. 1991. Western Hemlock Looper in British Columbia. Forest Insect and Disease Survey Report 91-08. Forestry Canada, Pacific Forestry Centre, Victoria, British Columbia, Canada. 42 pp.
- van Asch, M. and Visser, M.E. 2007. Phenology of forest caterpillars and their host trees: the importance of synchrony. *Annual Review of Entomology*, **52**: 37–55. <https://doi.org/10.1146/annurev.ento.52.110405.091418>.
- Wharton, G.W. and Richards, A.G. 1978. Water vapor exchange kinetics in insects and acarines. *Annual Review of Entomology*, **23**: 309–328. <https://doi.org/10.1146/annurev.en.23.010178.001521>.

- Wigglesworth, V.B. 1945. Transpiration through the cuticle of insects. *Journal of Experimental Biology*, **21**: 97–114. Available from <https://jeb.biologists.org/content/jexbio/21/3-4/97.full.pdf> [accessed 26 August 2025].
- Williamson, T.B., Colombo, S.J., Duinker, P.N., Gray, P.A., Hennessey, R.J., Houle, D., *et al.* 2009. Climate Change and Canada's Forests: From Impacts to Adaptation. Sustainable Forest Management Network and Natural Resources Canada, Canadian Forest Service, Northern Forestry Centre, Edmonton, Alberta, Canada. 104 pp.
- Winston, P.W. and Bates, D.H. 1960. Saturated solutions for the control of humidity in biological research. *Ecology*, **41**: 232–237. <https://doi.org/10.2307/1931961>.
- Woods, H.A. 2010. Water loss and gas exchange by eggs of *Manduca sexta*: trading off costs and benefits. *Journal of Insect Physiology*, **56**: 480–487. <https://doi.org/10.1016/j.jinsphys.2009.05.020>.
- Woods, H.A., Bonnacaze, R.T., and Zrubek, B. 2005. Oxygen and water flux across eggshells of *Manduca sexta*. *Journal of Experimental Biology*, **208**: 1297–1308. <https://doi.org/10.1242/jeb.01525>.
- Woods, H.A. and Hill, R.I. 2004. Temperature-dependent oxygen limitation in insect eggs. *Journal of Experimental Biology*, **207**: 2267–2276. <https://doi.org/10.1242/jeb.00991>.
- Woods, H.A. and Singer, M.S. 2001. Contrasting responses to desiccation and starvation by eggs and neonates of two Lepidoptera. *Physiological and Biochemical Zoology*, **74**: 594–606. <https://doi.org/10.1086/322169>.
- World Meteorological Organisation. 2024. State of the Global Climate 2023. Report WMO-No. 1347. World Meteorological Organisation, Geneva, Switzerland. 53 pp. Available from <https://library.wmo.int/idurl/4/68835> [accessed 26 August 2025].
- World Meteorological Organisation. 2025. State of the Global Climate 2024. Report WMO-No. 1368. World Meteorological Organisation, Geneva, Switzerland. 42 pp. Available from <https://library.wmo.int/idurl/4/69455> [26 August 2025].
- Zalucki, M.P., Clarke, A.R., and Malcolm, S.B. 2002. Ecology and behavior of first-instar larval Lepidoptera. *Annual Review of Entomology*, **47**: 361–393. <https://doi.org/10.1146/annurev.ento.47.091201.145220>.
- Zrubek, B. and Woods, H.A. 2006. Insect eggs exert rapid control over an oxygen–water tradeoff. *Proceedings of the Royal Society B: Biological Sciences*, **273**: 831–834. <https://doi.org/10.1098/rspb.2005.3374>.