

THE LINEARITY OF THE COSMIC EXPANSION FIELD AND THE VALUE OF THE HUBBLE CONSTANT

G. A. TAMMANN

*Astronomisches Institut der Universität Basel,
Venusstr. 7, CH-4102 Binningen, Switzerland*

Abstract. A linearity test shows H_0 to decrease by 7% out to 18 000 km s⁻¹. The value at 10 000 km s⁻¹ is a good approximation to the mean value of H_0 over very large scales. The construction of the extragalactic distance scale is discussed. Field galaxies, cluster distances relative to Virgo, and blue supernovae of type Ia yield H_0 (cosmic) with increasing weight; they give consistently $H_0 = 57 \pm 7$ (external error). This value is supported by purely physical distance determinations (SZ effect, gravitational lenses, MWB fluctuations). Arguments for $H_0 > 70$ are discussed and shown to be flawed.

1. Introduction

The calibration of the cosmic expansion rate H_0 consists of two steps. The first step is an investigation of the cosmic expansion field. How *linear* is the expansion? How large are *systematic* deviations from linearity in function of distance? What is the *scatter* of individual objects due to peculiar motions about the mean expansion? Only after these questions are solved can the second step be tackled, i. e. the calibration of the expansion rate in absolute terms.

The procedural difference between the two steps is that only redshifts and *relative* distances are needed for an investigation of the characteristics of the expansion field, while the calibration of the present large-scale expansion rate H_0 requires in addition the *true* distance of at least one object which demonstratably partakes of the mean expansion.

Much confusion about the expansion rate has arisen from equating the velocity-distance ratio of a subjectively chosen object with H_0 . The determination of H_0 from the Virgo, Fornax, or Coma clusters, for instance,

is meaningful only if it is demonstrated that they reflect at their moderate distances the mean cosmic expansion. The long-standing problem of correcting the observed mean velocity of the Virgo cluster into the frame of the cosmic expansion field has become a classic (cf. Section 3.2). The Fornax cluster with a velocity of $v \approx 1200 \text{ km s}^{-1}$ cannot be used for the determination of H_0 , even if a useful distance was known for it, because its unknown peculiar velocity may well be as high as 20% of its observed velocity. And the Coma cluster at $v \approx 7000 \text{ km s}^{-1}$, which is sometimes used for the determination of H_0 , may still have a peculiar velocity component of 10%, as the peculiar velocity of 630 km s^{-1} with respect to the MWB of one other supercluster, i. e. the Local Supercluster, would suggest.

The present paper outlines this two-step procedure. In Section 2 the available data are used to map the expansion rate in function of distance well beyond $30\,000 \text{ km s}^{-1}$, i. e. out to distances where the truly cosmic character of H_0 cannot be questioned. Section 3 gives a summary of the various methods of determining distances of field galaxies, of the Virgo cluster, and — most decisively for H_0 — of distant blue SNe Ia. Methods leading seemingly to $H_0 > 70$ are critically discussed in Section 4. A brief outlook is given in Section 5.

2. The Cosmic Expansion Field

Hubble (1929) in his discovery paper plotted recession velocities versus *linear* distances to infer the expansion of the Universe. All subsequent papers on the subject have used instead a plot of $\log cz$ versus $\log(\text{distance})$, the advantage being that only distance *ratios* are needed. The resulting diagrams have become known as Hubble diagrams. As a measure of $\log(\text{distance})$ the apparent magnitude of standard candles, the apparent diameters of standard rods, or relative distance moduli $\Delta(m - M)$ can be used. Using the Hubble diagrams of different objects the overall linearity of the cosmic expansion field has been proven without doubt.

In the case of linear expansion the regression line of the Hubble diagram has slope 0.2 if apparent magnitudes of standard candles or relative distance moduli are used. Their Hubble diagrams allow therefore three additional tests:

- (1) If a specific data set yields a slope different from 0.2 it is an unailing indication that the distance scale is incorrect. (Uncorrected selection effects, i. e. Malmquist bias, always yield too steep slopes and a spurious increase of H_0 with distance).
- (2) The scatter about the regression line is due to a combination of the effects of peculiar motions and errors of the relative distances. If the latter are under control one can determine the mean relative size $\Delta v/v_c$ of the

peculiar motions, where v_c is the cosmic velocity required by the mean Hubble line.

(3) If the relative peculiar motions $\Delta v/v_c$ are plotted against the distance r (or in sufficient approximation against the recession velocity v) one can test for *local* deviations from perfect Hubble flow. If $H_i = (v_c + \Delta v)/r$ is the perturbed value of the Hubble ratio of the i -th object at distance r , and $H_0 = \langle H_i \rangle = v_c/r$ the true Hubble constant, then $\Delta H/H_0 = (H_i - H_0)/H_0 = \Delta v/v_c$. This test of relative variation of H_0 is performed in the following.

Three independent data sets are used for the test, viz. the Hubble diagram of first-ranked cluster galaxies (Sandage *et al.*, 1976), the Hubble diagram of blue SNe Ia (Saha *et al.*, 1997; slightly updated by Parodi and Tammann, 1998) and the Hubble diagram with 31 relative cluster distances (Federspiel *et al.*, 1998). In each Hubble diagram the residuals $\Delta v/v_c$ are read and combined within 5000 km s^{-1} bins. Sliding means in 2500 km s^{-1} steps are plotted in Fig. 1 against redshift.

Inspection of Fig. 1 strongly suggests that the Hubble constant decreases from 1000 km s^{-1} to about 18000 km s^{-1} by $\sim 7\%$. This trend is independently supported by the first-ranked cluster galaxies of Lauer and Postman (1994), which have not been used here. Beyond 18000 km s^{-1} the scatter becomes large, leaving the possibility of local $\pm 10\%$ variations of H_0 , but the distant overall *mean* of H_0 lies close to the value found at 10000 km s^{-1} .

The consequences for the calibration of H_0 are clear. Full-sky samples with $v > 1000 \text{ km s}^{-1}$ yield the cosmic value of H_0 to within $\pm 5\%$. The best mean cosmic value is found near 10000 km s^{-1} or from sufficiently large samples beyond 20000 km s^{-1} .

3. The Calibration of H_0

The interlaced construction of the distance scale out to 30000 km s^{-1} and beyond is schematically shown in Fig. 2. Brief comments on the individual steps, labelled ① to ⑨, proceed as follows.

① The distance of LMC is the first fundamental step outward. The best values come from the purely geometrical distance of the ring of SN 1987A (Panagia *et al.*, 1996), from Cepheids in various passbands, and RR Lyr stars. The zeropoint of the Cepheid P-L relation is based on Cepheids in Galactic clusters (Sandage and Tammann, 1971; Feast, 1995), on stellar radii (Di Benedetto, 1997), on the Baade-Becker-Wesselink method (Laney and Stobie, 1992) and on Hipparcos trigonometric parallaxes (Madore and Freedman, 1998; Sandage and Tammann, 1998). The zeropoint of the RR Lyr star P-L relation in function of metallicity comes from physical considerations (Sandage, 1993a), the Baade-Becker-Wesselink method (Sandage




Figure 1. The variation of H_0 with redshift derived from relative distances. a) Out to $35\,000\text{ km s}^{-1}$ using relative cluster distances, SNe Ia, and first-ranked cluster galaxies. b) Out to $72\,500\text{ km s}^{-1}$; beyond $35\,000\text{ km s}^{-1}$ the data depend only on first-ranked cluster galaxies.

and Cacciari, 1990), and from globular clusters fitted to the Hipparcos-calibrated main sequence of subdwarfs (Reid, 1998; Gratton *et al.*, 1997; Pont *et al.*, 1998). Individual LMC moduli are compiled in Federspiel *et al.* (1998) and give $(m - M) = 18.54 \pm 0.03$. This value is also in good agreement with the position of the red-giant tip (Lee *et al.*, 1993; Tammann, 1996). Conservatively $(m - M) = 18.50$ has been adopted in the following. This value is secure to within $\pm 5\%$ and is uncontroversial.

② The distance of LMC combined with the good photometry of many of its Cepheids yields the calibrated P-L relation at different wavelengths (Sandage and Tammann, 1971). For the *HST* observations in *I* and *V* the P-L relations of Madore and Freedman (1991) are generally used; this has the advantage that any divergence in the derived value of H_0 cannot be blamed on the use of different P-L relation.




Figure 2. Schematic presentation of the interlaced extragalactic distance scale

Typical errors of individual Cepheid distances derived from *HST* data are $\pm 10\%$ due to the width of the instability strip and restricted sample size and due to absorption. For the luminosity calibration of SNe Ia (cf. ⑧) the errors are smaller because only apparent distance moduli are used. — Attempts to improve Cepheid distances with the help of a period-luminosity-color (PLC) relation (e.g. Kochanek, 1997) are doomed because stellar evolution models combined with a pulsation code show that the basic assumptions going into a PLC relation are not met (Saio and Gaitschy, 1998). The same models show that the much-discussed metallicity has minimal effect on the $P-L_{bol}$ relation; remaining metallicity effects enter only

through the bolometric correction.

③ The Cepheid distances from *HST*, most relevant for H_0 , concern the six nearby galaxies which have produced seven well studied blue (i. e. “Branch - normal”) SNe Ia (cf. ⑧). Their mean luminosity is (Saha *et al.*, 1997; Saha, 1998) $M_B(\text{max}) = -19.54$, $M_V(\text{max}) = -19.50$ with an impressively small scatter of 0^m16 which – in addition to the arguments presented in ⑧ – demonstrates that blue SNe Ia are highly useful standard candles.

The luminosity of SNe Ia can also be determined from physical explosion models. The state-of-the-art light curve fitting method by Höflich and Khokhlov (1996) give $M_B(\text{max}) = -19.45$ for the 16 SNe Ia to which sufficiently blue models can be fitted. In a review of all physical determinations of $M(\text{max})$ Branch (1998) concludes $M_B(\text{max}) \approx M_V(\text{max}) \approx -19.4 - 19.5$. The close agreement of the astronomical and physical calibrations of blue SNe Ia is most encouraging.

④ The calibration of the 21 cm line width-luminosity (Tully-Fisher, TF) relation in the B-band can now be based on 18 Cepheid distances most of which are due to *HST*. They determine the zeropoint of the relation with a statistical error of only 0^m05 , but they are still not enough for the determination of the slope. The latter must be taken from a *complete* sample of 49 inclined spirals of the Virgo cluster (Federspiel *et al.*, 1998).

In principle redder wavelengths than the B band offer the advantage of smaller inclination-dependent absorption corrections. However, this advantage is entirely offset by the steeper slope of the TF relation at longer wavelengths (Schröder, 1996). Moreover the restricted number of available magnitudes at longer wavelengths and their possible inhomogeneity, particularly in the *I* and *H* bands, preclude the all-important construction of *complete* samples. Therefore absolute TF distances of the Virgo cluster and of field galaxies are problematic if derived from magnitudes other than *B*.

⑤ The distance of the Virgo cluster can be derived, as shown in Fig. 2, in six different ways:

a) Cepheid distances give so far only a rather poorly determined cluster distance. The reason is the important depth effect of the cluster. Four of the five galaxies with distances from Cepheids have been selected because they are well or even extremely well (NGC 4571) resolved and *expected* to lie on the near side of the cluster. Their mean distance is 16.1 ± 0.4 Mpc (Freedman *et al.*, 1998). One galaxy, NGC 4639, has been selected irrespective of resolution and its distance is 25.1 ± 2.5 Mpc (Saha *et al.*, 1997). The position of the four resolved galaxies on the near side and of the poorly resolved NGC 4639 on the far side of the cluster is independently confirmed by their relative TF distances (Federspiel *et al.*, 1998). One can therefore infer only that the *center* of the cluster lies roughly at $(m - M) = 31.5 \pm 0.5$ (20 Mpc). A preliminary way out is to use the mean distance of the Leo

TABLE 1. The Virgo cluster modulus from various methods

Method	$(m - M)_{\text{Virgo}}$	Hubble type
Cepheids	31.52 ± 0.21	S
SNe Ia	31.39 ± 0.17	E, S
Tully-Fisher	31.58 ± 0.24	S
Globular Clusters	31.67 ± 0.15	E
$D_n - \sigma$	31.85 ± 0.19	S0, S
Novae	31.46 ± 0.40	E
Mean:	31.60 ± 0.08	($\Rightarrow 20.9 \pm 0.8$ Mpc)

group of $(m - M) = 30.27 \pm 0.12$, based now on three galaxies with Cepheid distances, and to step up this value by the modulus difference of $\Delta(m - M) = 1.25 \pm 0.13$ (Tammann and Federspiel, 1997) between the Leo Group and the Virgo cluster, giving $(m - M)_{\text{Virgo}} = 31.52 \pm 0.21$.

b) Eight SNe Ia with known maximum magnitudes are available. Three of these have occurred in E/S0 galaxies; they are known to be underluminous by 0^m18 . If the latter are adjusted to SNe Ia in spirals to conform with the calibrating SNe Ia, the mean apparent magnitude of the eight SNe Ia becomes $\langle m_B(\text{max}) \rangle = 11.91 \pm 0.16$ and $\langle m_V(\text{max}) \rangle = 11.84 \pm 0.17$. With the calibration in ③ the mean cluster modulus becomes then 31.39 ± 0.17 .

c) The TF relation in B magnitudes of a *complete* sample of 49 cluster galaxies, combined with the TF calibration in ④, gives $(m - M)_{\text{Virgo}} = 31.58 \pm 0.24$ (Federspiel *et al.*, 1998). The relatively large error allows for a number of systematic effects like sample selection, propagation of observational errors, and inclination and color differences between calibrators and cluster galaxies.

d-f) Other useful determinations of the Virgo cluster distance come from the luminosity function of globular clusters, the $D_n - \sigma$ relation of the bulges of S0 and spiral galaxies, and from novae. Because of the restricted space here the reader is referred to Sandage and Tammann (1997) and Tammann and Federspiel (1997). The results of the six methods a)–f) are compiled in Table 1.

The surface brightness fluctuation (SBF) method is sometimes advertised as a distance indicator. However, its reliability has not been sufficiently demonstrated yet for distances beyond 10 Mpc. The SBF distance of NGC 7331 (Tonry, 1997) is 0^m55 smaller than its Cepheid distance (Hughes, 1997), and the proposed Virgo cluster modulus (Tonry, 1997) is smaller than that in Table 1 by the same amount. The applicability of the method will still depend on a large sample of Virgo cluster members to decide if

the same relation applies for ellipticals and bulges of spiral galaxies. If not, the zeropoint calibration of the method will depend on an adopted Virgo cluster distance.

It has also been suggested that the luminosity function of the shells of planetary nebulae (PNe) in the light of the λ 5004 Å line had a magic cutoff luminosity which could be used as a universal distance indicator. However, it has been shown that the available data are at least equally consistent with a roughly exponential bright tail of the luminosity function such that the brightness of the brightest PNe depends on the sample size, i. e. on the luminosity of the galaxy under consideration (Bottinelli *et al.*, 1991; Tammann, 1993). This conclusion has been buttressed by model calculations by Soffner *et al.* (1996). The recent proposal of a Virgo modulus of $(m - M) = 30.79 \pm 0.16$ (Ciardullo *et al.*, 1998), based on a simple cutoff assumption, is significantly smaller than the Cepheid distance of even the nearest, highly resolved Virgo spirals (cf. ⑤a) and is therefore self-defeating.

⑥ The route to H_0 through *field galaxies* is the most difficult one. The crux is selection bias. Its origin is the fact that astronomers work with magnitude-limited catalogs of field galaxies, in which case the mean luminosity of the catalogued objects increases with distance. The disastrous consequences of selection (Malmquist) bias are illustrated, for instance, by a realistic model calculation by Hendry and Simmons (1990). They show that denying the selection bias can lead to $H_0 = 80$, while proper allowance for bias — depending on the luminosity scatter of the galaxies — yields values of $H_0 = 56$ or even $H_0 = 44$. Two fundamental facts emerge from this. Neglect of the selection bias always leads to too high values of H_0 , and the severity of the error is a strong function of the luminosity scatter. In the case of cluster distances one can overcome the problem by working with *complete* samples within a given volume (cf. ⑦). Blue SNe Ia offer the very important advantage that their luminosity scatter is so small that H_0 will not be significantly overestimated (cf. ⑧).

Analytical corrections for Malmquist bias require knowledge of the true scatter, which can only be derived from very deep samples, and must neglect the clumpy distribution of galaxies. Practical ways to correct for bias are, e. g., by Bottinelli *et al.* (1986a; 1986b), Lynden-Bell *et al.* (1988), Sandage (1988), Federspiel *et al.* (1994), Sandage *et al.* (1995), Giovanelli (1997a), and Theureau *et al.* (1997). A tutorial on the subject is given by Sandage (1995).

Recent determinations of H_0 from bias-corrected field galaxies are compiled in Table 2. Field galaxies offer the advantage of full-sky coverage outside the zone of avoidance. But they are not only the most difficult route to H_0 , but also the least satisfactory, having their main thrust as close as $1000 - 3000 \text{ km s}^{-1}$, i. e. at a distance where H_0 (local) may still be

TABLE 2. H_0 from bias corrected field galaxies

Method	H_0	Source
Tully-Fisher	< 60	Sandage 1994
M 101 look-alike diameters	43 ± 11	Sandage 1993b
M 31 look-alike diameters	45 ± 12	Sandage 1993c
Spirals with luminosity classes	56 ± 5	Sandage 1996a
M 101, M 31 look alike luminosities	55 ± 5	Sandage 1996b
Tully-Fisher	55 ± 5	Theureau <i>et al.</i> 1997
Galaxy diameters	50 – 55	Goodwin <i>et al.</i> 1997
Tully-Fisher	60 ± 5	Federspiel 1998
mean	53 ± 3	

a few percent higher than H_0 (cosmic) (cf. Fig. 1).

⑦ H_0 (cosmic) from clusters out to 11 000 km s⁻¹. Cluster distances *relative* to the Virgo cluster have been compiled from the literature (Jerjen and Tammann, 1993). The sample is increased by *relative I-band TF* distances of clusters (Giovanelli, 1997b). These relative distances are excellent as seen from their small scatter about the Hubble line of slope 0.2 in Fig. 3. The scatter is in fact so small that it imposes stringent limits on the radial component of the peculiar motion of cluster centers (Jerjen and Tammann, 1993).

The best fit to the data in Fig. 3 is

$$\log cz = 0.2 [(m - M)_{\text{Cluster}} - (m - M)_{\text{Virgo}}] + (3.070 \pm 0.024) \quad (1)$$

(Federspiel *et al.*, 1998). From this follows directly

$$\log H_0(\text{cosmic}) = -0.2 (m - M)_{\text{Virgo}} + (8.070 \pm 0.024). \quad (2)$$

Inserting $(m - M)_{\text{Virgo}} = 31.60 \pm 0.08$ yields

$$H_0(\text{cosmic}) = 56 \pm 3 \quad (\text{internal error}). \quad (3)$$

Note that no use of any velocity of the Virgo cluster has been made. The value of H_0 holds out to $\sim 10\,000$ km s⁻¹ where its value is very close to the large-scale value (Fig. 1).

⑧ H_0 (cosmic) from blue SNe Ia. The Hubble diagram (in V) of 35 blue SNe Ia, mainly due to the heroic efforts of the Cerro Tololo group (Hamuy *et al.*, 1996), is shown in Fig. 4. The small scatter $\sigma(m_V)$ of 0^m24 confirms the conclusion in ③ that blue SNe Ia are extremely useful standard candles.




Figure 3. Hubble diagram of 31 clusters with known relative distances. Asterisks are data from Jerjen & Tammann (1993). Open circles are from Giovanelli (1997a). Filled circles are the average of data from both sources.




Figure 4. The Hubble diagram of 35 blue SNe Ia with photometry after 1985. The SNe Ia in elliptical galaxies are shifted by 0^m18 .

This is also supported by the absence of Malmquist bias, i. e. the Hubble line does not steepen with distance. A fit to the Hubble diagrams in B and V gives, after simple transformation,

$$\log H_0 = 0.2 M_B + (5.651 \pm 0.011) \quad (4)$$

$$\log H_0 = 0.2 M_V + (5.669 \pm 0.019). \quad (5)$$

These equations hold for SNe Ia in spirals. Those in E/S0 galaxies are fainter on average by $0^m18 \pm 0^m05$ in B and V and have been shifted by this amount. Equations (4) and (5) can therefore directly be compared with the calibrators in spiral galaxies from ③. Thus one obtain a mean value from B and V data of

$$H_0(\text{cosmic}) = 57 \pm 3. \quad (6)$$

Red SNe Ia with $(B_{\max} - V_{\max}) > 0.20$ have been excluded from the experiment because they are reddened or have peculiar spectra (Branch *et al.*, 1993). Their unjustified inclusion leads to second-parameter corrections of the calibration in ③, but H_0 remains in all cases < 65 (for details see Saha *et al.* 1997; cf. also 4.6). Within the claimed accuracy all second-parameter effects of blue SNe Ia have been taken care of by allowing for the underluminosity of 0^m18 of the SNe Ia in E/S0 galaxies.

Of all present methods to derive H_0 , the route through SNe Ia deserves the highest weight.

⑨ H_0 (cosmic) from *Physical Methods*. One distinguishes between astronomical and physical distance determinations. The former depend always on some adopted distance of a celestial body, be it only the Astronomical Unit in the case of trigonometric parallaxes. Physical methods derive the distance solely from the observed physical or geometrical properties of a specific object.

In the foregoing reference has been made to physical luminosity and distance determinations of the SN 1987A remnant, Cepheids, RR Lyr stars, and SNe Ia. But in addition there are a number of physical distance determinations which lead to the value of H_0 over very large scales. They are still model-dependent, but as the number of objects increases and the models improve, their weight is steadily increasing.

For brevity the most recent physical determinations of H_0 are compiled in Table 3.

Following Rephaeli and Yankovitch (1997) all previous values of H_0 from the SZ effect should be lowered by ~ 10 units due to relativistic effects.

The overall impression from the values in Table 3 is that H_0 will settle around $H_0 \approx 56$.

TABLE 3. H_0 from Physical Methods

Method	H_0	Source ^a
Sunyaev-Zeldovich effect		
for cluster A 2218	45 ± 20	(1)
for 6 other clusters	60 ± 15	(2)
cluster A 2163	$78 (+54, -28)$	(3)
2 clusters	42 ± 10	(4)
3 clusters	54 ± 14	(5)
incl. relativ. effects	44 ± 7	(6)
Gravitational lenses		
QSO 0957 + 561	62 ± 7	(7)
B 0218 + 357	$52 - 82$	(8)
PG 1115 + 080	60 ± 17	(9)
"	52 ± 14	(10)
"	62 ± 20	(11)
MWB fluctuation spectrum	58 ± 11	(12)
"	47 ± 6	(13)

^aSources: (1) McHardy *et al.* 1990; Birkinshaw and Hughes 1994; Lasenby and Hancock 1995 (2) Rephaeli 1995; Herbig *et al.* 1995 (3) Holzapfel *et al.* 1997 (4) Lasenby and Jones 1997 (5) Myers *et al.* 1997 (6) Rephaeli and Yankovitch 1997 (7) Falco *et al.* 1997 (8) Nair 1996 (9) Keeton and Kochanek 1997 (10) Kundić *et al.* 1997 (11) Schechter *et al.* 1997 (12) Lineweaver 1998 (13) Webster *et al.* 1998

4. $H_0 = 73$?

Freedman *et al.* (1997) have proposed seven arguments why H_0 should be 73 (cf. also Freedman *et al.* 1998). These arguments are briefly discussed in the following.

4.1 $H_0 = 80 \pm 17$ from the Virgo cluster. This high value can only be suggested by combining a low cluster distance of 17.8 Mpc, relying heavily on Cepheids of near-side cluster members, and an outdated mean cluster velocity of 1404 km s^{-1} . The best cluster velocity, corrected for all local effects, follows from equation (1) by setting $(m - M)_{\text{Cluster}} - (m - M)_{\text{Virgo}} = 0$, i. e. $1175 \pm 85 \text{ km s}^{-1}$. This with 17.8 Mpc gives $H_0 = 66$ and with a more realistic cluster distance from Table 1 $H_0 = 56 \pm 4$.

4.2 $H_0 = 77 \pm 16$ from Coma via Virgo. The value depends, of course, on the adopted distance of the Virgo cluster. With a Virgo distance of 20.9 ± 0.8 (Table 1) one obtains $H_0 = 66 \pm 10$ with a large error due to the peculiar motion of the Coma cluster and to some extent also due to the

error of the adopted relative distance Coma-Virgo.

4.3 $H_0 = 72 \pm 18$ from the Fornax cluster. The result comes from equating the single Cepheid distance of the spiral galaxy NGC 1365 of 18.4 ± 1.0 Mpc (Silbermann *et al.*, 1998) with the *mean* cluster distance. There is no basis for this ad hoc assumption. A compilation of 30 distance determinations of the Fornax cluster over the last 20 years actually indicates that the E/S0 members are (1.17 ± 0.05) times more distant than the spiral galaxies (Tammann and Federspiel, 1997). Taking this at face value one finds 21.5 ± 1.5 Mpc (cf. also 4.6) for the E/S0 galaxies and hence for the cluster center. This reduces the proposed value of $H_0 = 77 \pm 16$ to $H_0 = 66 \pm 15$ with a large uncertainty due to the unknown peculiar motion of the cluster (cf. Section 1).

4.4 $H_0 = 72 \pm 8$ from local data. The solution is dominated by the Virgo cluster (Sec. 4.1) and Fornax cluster (Sec. 4.3). The remaining three groups, NGC 2403, NGC 1023, and Leo, lie within 12 Mpc and are irrelevant for the *cosmic* value of H_0 .

4.5 $H_0 = 72 \pm 8$ from the Jerjen-Tammann clusters. The cluster sample has since been increased by the relative cluster distances of Giovanelli (1997; cf. Fig. 3 and equations (1) and (2)). With an (untenable) Virgo cluster modulus of $(m - M) = 31.25 \pm 0.20$ from Sec. 4.1, equation (2) yields $H_0 = 66 \pm 7$ (not 72!), but even this is too high because the inserted Virgo modulus is too small.

4.6 $H_0 = 67 \pm 8$ from SNe Ia. The authors follow in principle route ③ + ⑧, but they reject two calibrating SNe Ia on the ground of their photographic photometry (much of observational astronomy had to be discarded on that ground), *and* they speculate that SN 1980 and SN 1992A are at the same distance as NGC 1365. The two SNe Ia have occurred in the Fornax cluster E/S0 galaxies NGC 1316 and NGC 1380, respectively. As argued under 4.3, there is independent evidence that the E/S0 galaxies in this cluster are more distant than the spiral NGC 1365. But for the sake of the argument the speculation of a common distance is taken up here. Since blue SNe Ia in early-type galaxies are fainter by $0^m.18$ in *B* and *V* than their counterparts in spirals (cf. ⑧), the mean absolute magnitude of SNe 1980 and 1992 becomes $M_B = -18.95$ and $M_V = -19.00$ if they lied at the distance of NGC 1365 and if they had occurred in spirals. This averaged in with the four remaining calibrators in ③ gives $\langle M_B(\text{max}) \rangle = -19.31 \pm 0.12$ and $\langle M_V(\text{max}) \rangle = -19.33 \pm 0.11$, and inserted into equations (4) and (5), which hold for blue SNe Ia in *spirals*, yields a mean value of $H_0 = 62 \pm 6$. But even this rather low value is still internally inconsistent because the luminosity distribution of the six calibrators used becomes highly non-Gaussian, defying the basic conclusion of standard candles. It violates also the models of Höflich and Khokhlov (1996) for blue SNe Ia. It is much more plausible that SNe 1980N

and 1992A have the same standard luminosity as found in ③, adjusted for early-type parent galaxies. In that case they require for the Fornax E/S0 galaxies $(m - M) = 31.85$ (23.4 Mpc) in agreement with the independent value under 4.3.

4.7 $H_0 = 73 \pm 7$ from the TF method. The authors base their claim on *I*- and *H*-band TF cluster distances of Mould *et al.* (1997) and Giovanelli *et al.* (1997). These authors consider highly incomplete and hence necessarily biased cluster samples, which may be useful for *relative* cluster distances (cf. also Kraan-Korteweg *et al.* 1988). In fact the *relative* distances of Giovanelli *et al.*, who include a correction for *differential* bias, are found to be excellent (cf. Fig. 3). However, it is inadmissible to tie these *biased* subpopulations directly to the *distance-limited* sample of calibrators with known Cepheids. This necessarily leads to an underestimate of the cluster distances (Teerikorpi, 1987). It has been demonstrated, for instance, that the 25 sufficiently inclined Virgo galaxies with known *H*-magnitudes yield a cluster modulus $\sim 0^m6$ smaller than the true value derived from a complete Virgo sample (Kraan-Korteweg *et al.* 1988; their Fig. 4). This discrepancy, which corresponds to a distance factor of ~ 1.3 , perpetuates then through all cluster distances and immediately gives $H_0 \approx 55$. The intermediate step through the Virgo cluster, i.e. from local calibrators to Virgo and from Virgo to more distant clusters, is necessary because the Virgo cluster is so far the only cluster for which a large and *complete* sample of spirals is available as well as extensive photometry and 21 cm data.

4.8 $H_0 = 73 \pm 7$ from physical models of SNe II. The authors cite the work of Schmidt *et al.* (1994; the also cited paper by R. P. Kirshner has not appeared). The result depends strongly on how the bolometric luminosity is distributed over the spectrum. The so-called dilation factor is a major stumbling block. On different assumptions Baron *et al.* (1995; 1996) have obtained $H_0 \leq 50$. Obviously the method cannot be used at present for a quantitative discussion of H_0 (cf. also Nadyozhin 1998).

4.9 $H_0 = 73 \pm 6$ from the $D_n - \sigma$ method. Mould *et al.* (1997) have calibrated the $D_n - \sigma$ data of Faber *et al.* (1989) using the Leo Group and the controversial Virgo and Fornax clusters as a zeropoint. Only the Leo group with three Cepheid distances is secure, but it provides with only two $D_n - \sigma$ distances a shaky basis. If one adopts the Virgo cluster distance from Table 1 and $(m - M) = 31.85$ as Fornax cluster modulus (4.3), an alternative calibration is obtained, leading to $H_0 = 63 \pm 6$. — It may be noted that the $D_n - \sigma$ method applied to the bulges of S0 and S galaxies yields a *high* distance of the Virgo cluster (Table 1).

A recapitulation of Section 4.1-4.9 gives the following picture. 4.4 and 4.8 should be excluded as being too local and too controversial, respectively. 4.1, 4.2 and 4.5 depend entirely on the adopted small Virgo cluster

distance, i.e. on the high weight given to the highly resolved galaxies on the near side of the cluster (4.1 depends also on the adopted high Virgo cluster velocity). 4.3 stands and falls with the assumption that the Cepheid distance of a single spiral (NGC 1365) provides a useful *mean* distance of the E/S0 galaxies of the Fornax cluster. The same assumption affects 4.6 by about 15%. The small Virgo *and* Fornax distances are essential for the high value of H_0 in 4.9. The remaining Section 4.7 is a textbook illustration of Malmquist bias.

5. Conclusions

A Test for the variation of H_0 with distance suggests a decrease by $\sim 7\%$ from $1000 < v \lesssim 18\,000 \text{ km s}^{-1}$. At $v = 10\,000 \text{ km s}^{-1}$ H_0 goes through a value close to the mean over very large scales.

A system of three interconnected distance scales (field galaxies, cluster distances relative to the Virgo cluster, and most significantly blue SNe Ia) give H_0 (cosmic) = 57 ± 7 (external error). Physical distance determinations from the SZ effect, gravitationally lensed quasars, and MWB fluctuations scatter about the same value.

A discussion of proposed high values of H_0 shows that disagreement focuses on two topics: 1) the true distance of the Virgo cluster, and 2) the appreciation of the Malmquist bias. One may add as item 3) the distance of the E/S0 galaxies in the Fornax cluster; the latter has lower priority because the peculiar motion of this cluster is unknown, and it is poorly tied into the relative distance scale of other clusters.

Acknowledgement: Financial support of the Swiss National Science Foundation is gratefully acknowledged. The author thanks his colleagues in the *HST* team for the luminosity calibration of SNe Ia, i.e. Dres. A. Sandage, A. Saha, L. Labhardt F.D. Macchetto, and N. Panagia, as well as the many collaborators behind the scenes at the STScI; much of the present understanding of H_0 depends on their work. He also thanks Mr. Bernd Reindl for his excellent help in all computational and technical matters.

References

- Baron, E., Hauschildt, P. H., Branch, D., Austin, S., Garnavich, P., Ann, H. B., Wagner, R. M., Filippenko, A. V., Matheson, T. and Liebert, J.: 1995, *ApJ* **441**, 170.
Baron, E., Hauschildt, P. H., Nugent, P. and Branch, D.: 1996, *MNRAS* **283**, 297.
Birkinshaw, M. and Hughes, J. P.: 1994, *ApJ* **420**, 33.
Bottinelli, L., Gouguenheim, L., Paturel, G. H. and Teerikorpi, P.: 1986a, *A&A* **156**, 157.
Bottinelli, L., Gouguenheim, L., Paturel, G. H. and Teerikorpi, P.: 1986b, *A&A* **166**, 39.
Bottinelli, L., Gouguenheim, L., Paturel, G. H. and Teerikorpi, P.: 1991, *A&A* **252**, 550.
Branch, D.: 1998, *ARAA*, preprint.
Branch, D., Fisher, A. and Nugent, P.: 1993, *AJ* **106**, 2383.

- Ciardullo, R., Jacoby, G. H., Feldmeier, J. J. and Bartlett, R. E.: 1998, *ApJ* **492**, 62.
- Di Benedetto, G. P.: 1997, *ApJ* **486**, 60.
- Faber, S., Wegner, G., Burstein, D., Davies, R., Dressler, A., Lynden-Bell, D. and Terlevich, R.: 1989, *ApJS* **69**, 763.
- Falco, E. E., Lehár, J. and Shapiro, I. I.: 1997, *AJ* **113**, 540.
- Feast, M.: 1995, *Astrophys. Applications of Stellar Pulsations, ASP Conf. Series*, **83**, p. 209.
- Federspiel, M.: 1998, in preparation.
- Federspiel, M., Sandage, A. and Tammann, G. A.: 1994, *ApJ* **430**, 29.
- Federspiel, M., Tammann, G. A. and Sandage, A.: 1998, *ApJ* **495**, 115.
- Freedman, W. L., Madore, B. F. and Kennicutt, C.: 1997, in: *The Extragalactic Distance Scale*, eds. M. Livio, M. Donahue and N. Panagia, Cambridge: Cambridge University Press, p. 171.
- Freedman, W. L., Mould, J. R., Kennicutt, R. C. and Madore, B. F.: 1998, this volume.
- Giovanelli, R.: 1997a, in: *The Extragalactic Distance Scale*, eds. M. Livio, M. Donahue and N. Panagia, Cambridge: Cambridge University Press, p. 113.
- Giovanelli, R.: 1997b, private communication.
- Giovanelli, R., Haynes, M., da Costa, L. N., Freudling, W., Salzer, J. J. and Wegner, G.: 1997, *ApJ* **477**, L1.
- Goodwin, S. P., Gribbin, J. and Hendry, M. A.: 1997, *AJ* **114**, 2212.
- Gratton, R. G., Fusi Perci, F., Carretta, E., Clementini, G., Corsi, C. F. and Lattanzi, M.: 1997, *ApJ* **491**, 749.
- Hamuy, M., Phillips, M., Maza, J., Suntzeff, N., Schommer, R. and Aviles, R.: 1996, *AJ* **112**, 2398.
- Hendry, M. A. and Simmons, J. F. L.: 1990, *A&A* **237**, 275.
- Herbig, T., Lawrence, C. R. and Readhead, A. C. S.: 1995, *ApJ* **449**, L5.
- Höflich, P. and Khokhlov, A.: 1996, *ApJ* **457**, 500.
- Holzappel, W. L. *et al.*: 1997, *ApJ* **480**, 449.
- Hubble, E.: 1929, *Proc. Nat. Acad. Sci.* **15**, 168.
- Hughes, S. M.: 1997, private communication.
- Jerjen, H. and Tammann, G. A.: 1993, *A&A* **276**, 1.
- Keeton, C. R. and Kochanek, C. S.: 1997, *ApJ* **487**, 42.
- Kochanek, C. S.: 1997, *ApJ* **491**, 13.
- Kraan-Korteweg, R. C., Cameron, I. M. and Tammann, G. A.: 1988, *ApJ* **331**, 620.
- Kundić, T., Cohen, J. G., Blanford, R. D. and Lubin, L. M.: 1997, *AJ* **114**, 507.
- Laney, C. D. and Stobie, R. S.: 1992, in: *Variable Stars and Galaxies*, ed. B. Warner, *ASP Conf. Series*, **30**, p. 119.
- Lasenby, A. N. and Hancock, S.: 1995, in: *Current Topics in Astrofundamental Physics: The Early Universe*, eds. N. Sánchez and A. Zichichi, *NATO Advanced Science Institutes Series*, **467**, Dordrecht: Kluwer Academic Publishers, p. 327.
- Lasenby, A. N. and Jones, M. E.: 1997, in: *The Extragalactic Distance Scale*, eds. M. Livio, M. Donahue and N. Panagia, Cambridge: Cambridge University Press, p. 76.
- Lauer, T. R. and Postman, M.: 1994, *ApJ* **425**, 418.
- Lee, M. G., Freedman, W. L. and Madore, B. F.: 1993, *ApJ* **417**, 553.
- Lineweaver, C. H.: 1998, this volume.
- Lynden-Bell *et al.*: 1988, *ApJ* **326**, 19.
- Madore, B. and Freedman, W. L.: 1991, *PASP* **103**, 933.
- Madore, B. and Freedman, W. L.: 1998, *ApJ* **492**, 110.
- McHardy, L. M., Stewart, G. C., Edge, A. C., Cooke, B. A., Yamashita, K. and Hatsukade, I.: 1990, *MNRAS* **242**, 215.
- Mould, J., Sakai, S., Hughes, S. and Han, M.: 1997, in: *The Extragalactic Distance Scale*, eds. M. Livio, M. Donahue and N. Panagia, Cambridge: Cambridge University Press, p. 158.
- Myers, S. T., Baker, J. E., Readhead, A. C. S., Leitch, E. M. and Herbig, T.: 1997, *ApJ* **485**, 1.

- Nadyozhin, D. K.: 1998, preprint.
- Nair, S.: 1996, in: *Astrophysical Applications of Gravitational Lensing*, eds. C. S. Kochanek and J. N. Hewitt, Dordrecht: Kluwer, p. 197.
- Panagia, N. *et al.*: 1996, in: *The Extragalactic Distance Scale, Poster Papers*, eds. M. Livio, M. Donahue and N. Panagia, Baltimore: STScI, p. 54.
- Parodi, B. and Tammann, G. A.: 1998, to be published.
- Pont, F., Mayor, M., Turon, C. and Vandenberg, D. A.: 1998, *A&A* **329**, 87.
- Reid, I. N.: 1998, *AJ* **115**, 204.
- Rephaeli, Y.: 1995, *ARA&A* **33**, 541.
- Rephaeli, Y. and Yankovitch, D.: 1997, *ApJ* **481**, L55.
- Saha, A.: 1998, to be published.
- Saha, A., Sandage, A., Labhardt, L., Tammann, G. A., Macchetto, F. D. and Panagia, P.: 1997, *ApJ* **486**, 1.
- Saio, H. and Gautschi, A.: 1998, *ApJ*, in press.
- Sandage, A.: 1988, *PASP* **100**, 935.
- Sandage, A.: 1993a, *ApJ* **106**, 687 and 703.
- Sandage, A.: 1993b, *ApJ* **402**, 3.
- Sandage, A.: 1993c, *ApJ* **404**, 419.
- Sandage, A.: 1994, *ApJ* **430**, 13.
- Sandage, A.: 1995, in: *The Deep Universe*, eds. B. Binggeli and R. Buser, Berlin: Springer, p. 1.
- Sandage, A.: 1996a, *AJ* **111**, 1.
- Sandage, A.: 1996b, *AJ* **111**, 18.
- Sandage, A. and Cacciari, C.: 1990, *ApJ* **350**, 645.
- Sandage, A., Kristian, J. and Westphal, J. A.: 1976, *ApJ* **205**, 688.
- Sandage, A. and Tammann, G. A.: 1971, *ApJ* **167**, 293.
- Sandage, A. and Tammann, G. A.: 1997, in: *Critical Dialogues on Cosmology Conference*, ed. N. Turok, Princeton: Princeton University Press, p. 130.
- Sandage, A. and Tammann, G. A.: 1998, *MNRAS* **293**, L23.
- Sandage, A., Tammann, G. A. and Federspiel, M.: 1995, *ApJ* **452**, 1.
- Schechter, P. L. *et al.*: 1997, *ApJ* **475**, L85.
- Schmidt, B. P. *et al.*: 1994, *ApJ* **432**, 42.
- Schröder, A.: 1996, PhD thesis, University of Basel.
- Silbermann, N. *et al.*: 1998, *ApJ*, in press.
- Soffner, T. *et al.*: 1996, *A&A* **306**, 9.
- Tammann, G. A.: 1993, in: *Planetary Nebulae*, eds. R. Weinberger and A. Acker, *IAU Symp.*, **155**, Dordrecht: Kluwer, p. 515.
- Tammann, G. A.: 1996, in: *Reviews in Modern Astronomy 9*, ed. R. E. Schielicke, Hamburg: Astronomische Gesellschaft, p. 139.
- Tammann, G. A. and Federspiel, M.: 1997, in: *The Extragalactic Distance Scale*, eds. M. Livio, M. Donahue and N. Panagia, Cambridge: Cambridge University Press, p. 137.
- Teerikorpi, P.: 1987, *A&A* **173**, 39.
- Theureau, G. *et al.*: 1997, *A&A* **322**, 730.
- Tonry, J. L.: 1997, in: *The Extragalactic Distance Scale*, eds. M. Livio, M. Donahue and N. Panagia, Cambridge: Cambridge University Press, p. 297.
- Webster, M., Hobson, M. P., Lasenby, A. N., Lahav, O. and Rocha, G.: 1998, *ApJL*, submitted.