

Long-term physical health conditions and youth anxiety and depression: Is there a causal link?

Original Article

Cite this article: Shakeshaft, A., Mundy, J. R., Pedersen, E. M., Dennison, C. A., Riglin, L., Bragantini, D., Corfield, E. C., Thapar, A. K., Andreassen, O. A., Stergiakouli, E., Davey Smith, G., Hannigan, L., Musliner, K. L., Havdahl, A., & Thapar, A. (2025). Long-term physical health conditions and youth anxiety and depression: Is there a causal link? *Psychological Medicine*, **55**, e7, 1–11 <https://doi.org/10.1017/S0033291724003271>

Received: 29 November 2023

Revised: 12 November 2024

Accepted: 26 November 2024

Keywords:

Anxiety; Depression; The Norwegian Mother, Father, and Child Cohort Study; MoBa; Long-term physical health condition; Pediatrics

Corresponding author:

Amy Shakeshaft;
Email: shakeshafta@cardiff.ac.uk

Amy Shakeshaft^{1,2} , Jessica R. Mundy³, Emil M. Pedersen⁴ ,
Charlotte A. Dennison^{1,2} , Lucy Riglin^{1,2} , Daniela Bragantini^{5,6} ,
Elizabeth C. Corfield^{5,6,8} , Ajay K. Thapar^{1,2}, Ole A. Andreassen⁷ ,
Evie Stergiakouli⁸ , George Davey Smith⁸ , Laurie Hannigan^{5,6,8} ,
Katherine L. Musliner³ , Alexandra Havdahl^{5,6,9}  and Anita Thapar^{1,2} 

¹Wolfson Centre for Young People's Mental Health, Division of Psychological Medicine and Clinical Neuroscience, Cardiff University, UK; ²Centre for Neuropsychiatric Genetics and Genomics, Division of Psychological Medicine and Clinical Neurosciences, Cardiff University, Cardiff, UK; ³Department for Clinical Medicine, Aarhus University, Denmark; ⁴National Centre for Register-based Research, Department of Public Health, Aarhus University, Denmark; ⁵Nic Waals Institute, Lovisenberg Diaconal Hospital, Oslo, Norway; ⁶PsychGen Center for Genetic Epidemiology and Mental Health, Norwegian Institute of Public Health, Norway; ⁷Norwegian Centre for Mental Disorders Research (NORMENT), Institute of Clinical Medicine, University of Oslo and Oslo University Hospital, Norway; ⁸Population Health Sciences and MRC Integrative Epidemiology Unit, Bristol Medical School, University of Bristol, UK and ⁹PROMENTA Research Centre, Department of Psychology, University of Oslo

Abstract

Background. The prevalence of youth anxiety and depression has increased globally, with limited causal explanations. Long-term physical health conditions (LTCs) affect 20–40% of youth, with rates also rising. LTCs are associated with higher rates of youth depression and anxiety; however, it is uncertain whether observed associations are causal or explained by unmeasured confounding or reverse causation.

Methods. Using data from the Norwegian Mother, Father, and Child Cohort Study (MoBa) and Norwegian National Patient Registry, we investigated phenotypic associations between childhood LTCs, and depression and anxiety diagnoses in youth (<19 years), defined using ICD-10 diagnoses and self-rated measures. We then conducted two-sample Mendelian Randomization (MR) analyses using SNPs associated with childhood LTCs from existing genome-wide association studies (GWAS) as instrumental variables. Outcomes were: (i) diagnoses of major depressive disorder (MDD) and anxiety disorders or elevated symptoms in MoBa, and (ii) youth-onset MDD using summary statistics from a GWAS in iPSYCH2015 cohort.

Results. Having any childhood LTC phenotype was associated with elevated youth MDD (OR = 1.48 [95% CIs 1.19, 1.85], $p = 4.2 \times 10^{-4}$) and anxiety disorder risk (OR = 1.44 [1.20, 1.73], $p = 7.9 \times 10^{-5}$). Observational and MR analyses in MoBa were consistent with a causal relationship between migraine and depression (IVW OR = 1.38 [1.19, 1.60], $p_{FDR} = 1.8 \times 10^{-4}$). MR analyses using iPSYCH2015 did not support a causal link between LTC genetic liabilities and youth-onset depression or in the reverse direction.

Conclusions. Childhood LTCs are associated with depression and anxiety in youth, however, little evidence of causation between LTCs genetic liability and youth depression/anxiety was identified from MR analyses, except for migraine.

Introduction

It is estimated that 20–40% of children worldwide have a long-term physical health condition (LTC) (Finning, Neochoriti Varvarigou, Ford, Panagi, & Ukoumunne, 2022; Panagi, White, et al., 2022b; Panagi, Newlove-Delgado, et al., 2022a), and the prevalence of certain childhood LTCs, such as asthma, obesity, and diabetes, has increased in recent years (Abarca-Gómez et al., 2017; Dharmage, Perret, & Custovic, 2019; Royal College of Paediatrics and Child Health, 2020; Van Cleave, Gortmaker, & Perrin, 2010). It is well-established that children with LTCs are more likely to experience depression and anxiety than those without LTCs, particularly those with neurological illnesses such as epilepsy, migraine, and chronic fatigue syndrome (Finning et al., 2022; Glazebrook, Hollis, Heussler, Goodman, & Coates, 2003; Piquart & Shen, 2011a, 2011b; Suryavanshi & Yang, 2016).

This association between LTCs and anxiety/depression has been attributed to many different mediators including social stigma (Bakula et al., 2019), poor self-esteem (Piquart, 2013b), difficulties with peer relationships and bullying (Piquart, 2017; Pittet, Berchtold, Akre, Michaud, & Suris, 2010), difficulties in the family environment (Piquart, 2013a) (Qiu et al., 2021), and stress caused by the LTC itself (Compas, Jaser, Dunn, & Rodriguez, 2012). These are in

© The Author(s), 2025. Published by Cambridge University Press. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.

addition to the potential physiological impacts of LTCs on the brain, particularly for neurological disorders such as epilepsy (Agrawal & Govender, 2011). Other exacerbating factors may include increased school absenteeism (Finning et al., 2022) and poorer academic performance/educational attainment (Hughes et al., 2021; Lum et al., 2017). However, observational studies also indicate that children with LTCs have similar levels of life satisfaction as their peers (Blackwell et al., 2019).

Residual confounding from unmeasured confounders remains a possibility in all observational studies, as well as pleiotropy (genetic overlap between LTCs and mental health conditions (Zhu et al., 2019)). Additionally, for some LTCs, such as migraine, the direction of association is unclear, that is, whether migraine causes an increased risk of anxiety/depression, or whether anxiety/depression increases the risk of migraine (Dyb, Stensland, & Zwart, 2015; Falla et al., 2022). Therefore, it is currently unclear whether LTCs have causal effects on youth anxiety and depression. Identifying causal mechanisms that underlie the development of youth anxiety and depression is important for developing effective prevention and early intervention strategies (Thapar, Eyre, Patel, & Brent, 2022).

Mendelian randomization (MR) is one method for inferring causation that has some analogies with a randomized controlled trial (Smith & Ebrahim, 2003). It uses randomly allocated (via meiosis) genetic variants, identified from genome-wide association studies (GWAS), as instrumental variables (IVs) for an exposure to test for a causal relationship between IVs (for the exposure) and outcome (Sanderson et al., 2022). When observational studies are supported by other designs, including MR, this triangulation improves the inference of causation between an exposure and outcome, given the assumptions of MR are met.

We aim to assess whether the relationship between LTCs and depression/ anxiety in youth is causal, by: (i) directly testing for longitudinal associations in a large cohort and (ii) using MR. We hypothesized that childhood LTCs, particularly neurological conditions such as migraine and epilepsy, would show a causal relationship with youth depression and anxiety.

Methods

Sample

We used data from the Norwegian Mother, Father, and Child Cohort Study (MoBa) (Magnus et al., 2016; Paltiel et al., 2014), a population-based pregnancy cohort study conducted by the Norwegian Institute of Public Health. Participants were recruited from Norway between 1999 and 2008. Women consented to participation in 41% of the pregnancies. The cohort includes ~114,500 children, 95,200 mothers, and 75,200 fathers. The current study is based on version 12 of the quality-assured data files released for research in January 2019. The establishment of MoBa and initial data collection was based on a license from the Norwegian Data Protection Agency and approval from The Regional Committees for Medical and Health Research Ethics. The MoBa cohort is regulated by the Norwegian Health Registry Act. The current study was approved by The Regional Committees for Medical and Health Research Ethics (2016/1702). All procedures contributing to this work comply with the ethical standards of the relevant national and institutional committees on human experimentation and with the Helsinki Declaration of 1975, as revised in 2008. Questionnaires were sent to mothers at 0.5, 1.5, 3, 5, 7, and 8 years and to mothers and children at 14 years. Blood samples were obtained from both parents during pregnancy and

from mothers and children (umbilical cord) at birth (Paltiel et al., 2014). For MoBa genotyping information, including quality control, see Corfield et al. (2022).

LTCs definition

To identify children with LTCs, we used definitions established in previous consensus (Finning et al., 2022; Mokkink et al., 2008; Panagi, et al., 2022a; Panagi et al., 2022b), whereby a disease or condition is considered to be a chronic/long-term condition in childhood if: (1) it occurs in children aged 0–18 years; (2) the diagnosis is based on medical scientific knowledge and can be established using reproducible and valid methods or instruments according to professional standards; (3) it is not (yet) curable and (4) it has been present for longer than 3 months or it will, very probably, last longer than 3 months, or it has occurred three times or more during the past year and will probably reoccur. Using this definition and the data available we included the following conditions: (i) arthritis, (ii) asthma, (iii) cerebral palsy, (iv) chronic fatigue syndrome, (v) coeliac disease, (vi) diabetes (type 1), (vii) epilepsy, (viii) migraine, and (ix) reduced hearing/hearing loss. In MoBa, all disorders were reported by the mother at age 14, except for cerebral palsy which was reported by mothers at age 5. We also included obesity, which was defined as BMI \geq 95th percentile, calculated using the child's reported height and weight at age 14. See Table 1 for *N*.

Mental health outcomes

Anxiety and depression were primarily defined in MoBa using the linked Norwegian Patient Registry, providing health records from MoBa participants from 2008 up to the date of 23 June 2021. We used ICD-10 codes of F32 (unipolar depressive episode) and F33 (recurrent depressive disorder), to define a major depressive disorder (MDD) diagnosis, and F40 (phobic anxiety disorders) and F41 (other anxiety disorders, including panic disorder and generalized anxiety disorder) to define an anxiety disorder. Since our focus is on youth depression and anxiety, outcomes were defined as a diagnosis before 19 years of age. Because MoBa has rolling recruitment (participants recruited between 1999 and 2008), not

Table 1. Prevalence of LTCs

LTC	Total		Females	Males
	% with LTC	N with LTC	% with LTC	% with LTC
Asthma	12.0%	2097/17483	10.6%	13.3%
Migraine	8.7%	1520/17507	9.9%	7.5%
Obesity	5.1%	936/18525	5.0%	5.1%
Reduced hearing	2.3%	408/17501	2.2%	2.5%
Coeliac disease	1.9%	329/17493	2.3%	1.4%
Epilepsy	0.9%	153/17522	0.9%	0.8%
Diabetes	0.7%	126/17472	0.7%	0.8%
Chronic fatigue syndrome	0.3%	61/17506	0.5%	0.2%
Arthritis	0.3%	57/17523	0.4%	0.2%
Cerebral Palsy	0.2%	77/41233	0.2%	0.2%
Any LTC specified above	28.4%	5039/17729	28.5%	28.1%

all participants reached 19 years of age by the time of administrative censoring. Therefore, descriptive frequencies of depression and anxiety disorders in youth were limited to children born before 2003 (minimum of 18.5 years of age at time of censoring, since exact birth dates were not available).

Mother- and self-reported depression and anxiety symptoms in children at age 14 were used as secondary outcomes. Self-reported depression symptoms were measured using the 13-item Short Mood and Feelings Questionnaire (SMFQ) (Angold, Costello, Messer, & Pickles, 1995) and the 5-item Screen for Child Anxiety Related Disorders (SCARED) (Birmaher et al., 1997; Birmaher et al., 1999), respectively. Total symptom scores were calculated for each scale (range: SMFQ = 0–26, SCARED = 0–10), and recommended cut-points were used to estimate probable depression (SMFQ ≥12) (Eyre et al., 2021), and anxiety disorder (SCARED ≥3) (Birmaher et al., 1999). Mother-report depression was measured using the shortened 6-item SMFQ (range 0–12).

Statistical analysis

All analyses were conducted in R, version 4.2.1 (R Core Team, 2022). For an overview of the study design see Figure 1.

We determined the prevalence of each childhood LTC in the cohort and the combined prevalence of any childhood LTC at age 14. Chi-squared tests were used to establish whether LTC prevalence differed by sex. Associations between each separate LTCs were estimated using logistic regressions. We estimated the frequency of MDD or anxiety disorder diagnosis in youth, including only individuals who had reached the age of 18 at the time of administrative censoring.

Phenotypic associations

We tested for associations between childhood LTCs and mental health outcomes using logistic regressions. Birth year was included

as a covariate in all analyses. Additionally, where there was a sex difference in the prevalence of childhood LTCs, sex was included as a covariate.

Two-sample Mendelian randomization

MR analyses are reported using STROBE-MR guidelines (Skrivankova et al., 2021).

Exposures We identified genetic instruments from published childhood-onset specific GWAS for the following childhood LTCs: juvenile arthritis (Hinks et al., 2013), childhood-onset asthma (Ferreira et al., 2019), type 1 diabetes (Forgetta et al., 2020), childhood obesity (Bradfield et al., 2019), and genetic generalized epilepsy (International League Against Epilepsy Consortium on Complex Epilepsies et al., 2022). For migraine, where age-at-onset was not defined, the most recent and largest GWAS was used (Hautakangas et al., 2022). Descriptions of the samples used in each GWAS are presented in Supplementary Table S1. SNPs for use in MR analyses were identified as those associated with the phenotype at a genome-wide significant level ($p < 5 \times 10^{-8}$). Genetic instruments for eczema (Paternoster et al., 2015) were also included as an exposure of interest despite not being included in the phenotype analysis (since the relevant phenotypes were not available in MoBa). No suitable genetic instruments were available for cerebral palsy, chronic fatigue syndrome, coeliac disease, or hearing loss because of a lack of appropriate GWASs.

Outcomes We used two approaches for defining outcomes in MR analyses. First, we used the same outcomes as in the phenotype analyses (MDD/anxiety diagnosis before 19 years and self-reported depression and anxiety symptoms at 14 years of age) in MoBa as outcomes for MR analyses. Here, we estimated the association between genetic instruments for each exposure with these phenotype outcomes in the MoBa genotyped cohort, including the first five ancestral principal components and genotyping batch number




Figure 1. Investigating the association between LTC in childhood and youth anxiety/depression – study design. According to MR design, ? is the causal relationship of interest, where ? = ?/δ. MR assumptions: IV1, Relevance = instruments are robustly associated with exposure; IV2, Independence = instruments are independent of any confounding variables; IV3, Exclusion restriction = instruments are independent of the outcome given exposure. LTC: long-term physical health conditions; SNPs: single nucleotide polymorphisms.

as covariates. Using this approach only one direction of association could be estimated (LTC exposures \rightarrow MH outcomes).

Second, we performed two-sample bidirectional MR between childhood LTCs and youth-onset depression, using summary statistics from a GWAS of youth-onset MDD of 7,896 cases (diagnosed in a Danish psychiatric hospital, as an inpatient, outpatient, or in emergency settings, before the age of 19 years) and 23,590 controls (after controlling for relatedness and ancestry filtering) conducted in the Danish iPSYCH2015 cohort (Bybjerg-Grauholm *et al.*, 2020; Pedersen *et al.*, 2018). iPSYCH2015 is a large Danish registry-based genotyped case cohort established for the study of psychiatric disorders, see [Supplementary material](#) for more information. The threshold for identifying genetic instruments for the alternate direction MR (MDD \rightarrow LTCs) was $p < 5 \times 10^{-6}$ and F -statistic > 10 (although in practice no SNPs had F -statistic < 20 , indicating results are unlikely to suffer from weak instrument bias), leaving 15 SNPs as IVs.

For both approaches, we harmonized the outcome estimates with the exposure variants, so the effect estimates were expressed per effect allele increase. Where effect allele frequencies were not available in exposure/outcome data and harmonization was not possible because of being palindromic, SNPs were excluded from analyses. We used inverse-variance weighted (IVW) regression as the primary MR method though estimates were also generated using weighted median, weighted mode, MR-Egger (Bowden, Davey Smith, & Burgess, 2015) and MR-PRESSO (the MR pleiotropy residual sum and outlier, Verbanck, Chen, Neale, and Do (2018)) to assess horizontal pleiotropy and MR assumptions (Slob & Burgess, 2020). For details of MR methods see [Supplementary material](#). A consistent effect across all methods provides evidence for a causal effect (Lawlor, Tilling, & Davey Smith, 2016). Cochran's Q statistic was used to test for heterogeneity in instrument effects, whereby if $Q >$ degrees of freedom, this provides evidence for heterogeneity and invalid instruments. Steiger tests of directionality (Hemani, Tilling, & Davey Smith, 2017) were also run to test directionality in the causal effect by examining the variance explained by IVs on exposures and outcomes.

MR analysis was carried out using the TwoSampleMR (Hemani *et al.*, 2018) and MR-PRESSO (Verbanck, Chen, Neale, & Do, 2018) R packages.

We used the false discovery rate (FDR) to correct for multiple comparisons within phenotypic and within MR analysis, across exposures tested against each outcome ($n = 10$ for phenotypic analyses and $n = 7$ for MR). A q -value of 0.05 was used to define the FDR threshold.

Sensitivity analysis

To assess temporal ordering, we tested associations between the same childhood LTCs and *subsequent* depression/anxiety diagnoses in MoBa, by restricting cases to those with a depression/anxiety diagnosis between the ages of 14 and 18 years (i.e. subsequent to mother-reporting of LTC), using the same method as in the main analysis. We also performed additional analyses testing for associations between childhood LTCs and total (continuous) self-reported depression and anxiety symptom scores and mother-reported depression symptom scores at age 14.

MR analyses using the largest GWAS for depression (Als *et al.*, 2023) (excluding UK biobank) and anxiety (Purves *et al.*, 2020) were also performed and reported in the [Supplementary Material](#). These were not used in the primary analysis since they focus on

depression and anxiety in adults, and evidence suggests that the genetic architecture of these disorders may differ across age-at-onset (Harder *et al.*, 2022; Nguyen *et al.*, 2023; Thapar & Riglin, 2020). Purves *et al.* (2020) anxiety disorder GWAS was conducted using data from the UK biobank study, from which cohort members were also included in some of the exposure GWAS for childhood LTCs (see [Supplementary Table S1](#); migraine and childhood-onset asthma GWAS). Therefore, the type 1 error rate may be increased for these overlapping-sample MR sensitivity analyses.

Results

Descriptive information

Overall, 28% of children in the sample had at least one LTC reported at age 14. The numbers with and without each LTC are presented in [Table 1](#). There was a male preponderance for asthma ($p = 5.2 \times 10^{-8}$), and a female preponderance for migraine ($p = 3.0 \times 10^{-8}$), coeliac disease ($p = 1.1 \times 10^{-5}$), and chronic fatigue syndrome ($p = 8.9 \times 10^{-4}$). Associations between the presence of each LTC with another are presented in [Supplementary Table S2](#).

In the total cohort, 4.5% (716/15,958) had MDD, 4.9% (783/15,974) had an anxiety disorder, and 1.7% (262/15,827) had comorbid anxiety and depression diagnoses, with age-at-onset by 19 years as recorded in the linked healthcare registry data. Using questionnaire data at age 14, 26.8% (5,785/21,549) reported elevated anxiety symptoms, and 21.5% (4554/21,229) reported elevated depression symptoms. All anxiety and depression outcomes were more common in females ([Supplementary Table S3](#)).

Phenotypic associations between childhood LTCs and youth anxiety/depression

Having any LTC was associated with anxiety (OR = 1.44 [1.20, 1.73], $p = 7.9 \times 10^{-5}$) and depression diagnoses in youth (OR = 1.48 [1.19, 1.85], $p = 4.2 \times 10^{-4}$), as well as with elevated anxiety (OR = 1.25 [1.15, 1.36], $p = 1.4 \times 10^{-7}$) and depression symptoms (OR = 1.39 [1.27, 1.51], $p = 7.5 \times 10^{-13}$) ([Figure 2](#)). Specifically, the following childhood LTCs were associated with an increased risk of anxiety diagnosis in youth: arthritis, cerebral palsy, epilepsy, migraine, and obesity. Similarly, epilepsy, migraine, obesity, and additionally asthma, were also associated with elevated self-reported anxiety symptoms at age 14. Increased risk of MDD diagnosis was found for chronic fatigue syndrome and obesity. Asthma, migraine, and obesity were associated with elevated self-reported depressive symptoms.

MR analysis

In MR analyses using MoBa data for outcomes, there was some evidence of causality between genetic liability to childhood-onset asthma and risk of MDD (IVW OR = 1.11 [1.02, 1.21], $p = 0.02$, $p_{FDR} = 0.13$) and anxiety disorder diagnoses (IVW OR = 1.11 [1.03, 1.21], $p = 0.008$, $p_{FDR} = 0.05$). There was little evidence of causal links between genetic liability to other LTCs and MDD/anxiety disorders ([Figure 3](#), [Supplementary Table S5](#)). We found stronger causal evidence of genetic liability to migraine and our secondary outcome measures: elevated self-reported depression symptoms (IVW OR = 1.38 [1.19, 1.60], $p = 2.6 \times 10^{-5}$, $p_{FDR} = 1.8 \times 10^{-4}$, see [Supplementary Figure S1](#) for MR scatter plot) and elevated




Figure 2. Phenotypic associations between LTCs at age 14 and anxiety diagnosis (top, purple) and symptoms (top, pink) measured by the SCARED questionnaire, and major depressive disorder diagnosis (bottom, green) and depression symptoms (bottom, blue) measured by the SMFQ questionnaire. LTCs with ^ indicate sex was included as a covariate in the analysis. Cerebral palsy was reported by mothers when children were aged 5 years. * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$ (FDR adjusted p-values across 10 LTCs). Test statistics are presented in Supplementary Table S4.

self-reported anxiety symptoms (IVW OR = 1.20 [1.04, 1.38], $p = 0.01$, $p_{FDR} = 0.08$) at age 14.

We found limited evidence of horizontal pleiotropy using MR Egger (Supplementary Table S6), and the estimates of MR Egger, weighted median, and mode MR estimates agreed with IVW for these results (Supplementary Table S5). Results from MR-PRESSO global tests for estimates where we found evidence of potential causal links (childhood-onset asthma and MDD/anxiety, and migraine and anxiety/depression symptoms) did not

indicate outliers (Supplementary Table S7). There was limited evidence for heterogeneity in IV effects for the same estimates, see Supplementary Table S8. F statistics for all exposures are presented in Supplementary Table S9. Steiger directionality tests indicated the correct causal direction for MR analyses where we found evidence of potential causal links (Supplementary Table S10).

Bidirectional MR analysis using GWAS of youth-onset depression (in the iPSYCH2015 cohort) did not indicate strong evidence




Figure 3. MR analyses testing for evidence of causality between childhood LTCs and anxiety diagnosis (top, purple) and symptoms (top, pink) measured by the SCARED questionnaire, and major depressive disorder diagnosis (bottom, green) and depression symptoms (bottom, blue) measured using SMFQ in the MoBa dataset. IVW OR = Inverse variance weighted MR odds ratio estimate. *** FDR corrected $p < 0.001$. See [Supplementary Table S5](#) for all test statistics.

for causality between genetic liability to any childhood LTC and youth-onset depression (Table 2), nor in the alternate direction (Supplementary Table S11). MR Egger intercepts indicated limited evidence of horizontal pleiotropy for exposures/outcomes (Supplementary Table S12). Estimates from sensitivity analysis agreed with the main IVW analysis. Details of F statistics for all instruments are presented in Supplementary Table S9 and results from heterogeneity tests in Supplementary Table S13.

Sensitivity analysis

Phenotypic associations. Associations between childhood LTCs and subsequent anxiety and MDD (specifically, diagnosed after the LTC, between ages 14 and 18 years) showed a similar pattern of results as the main phenotype analysis, with evidence for associations between any childhood LTC and both anxiety (OR = 1.37 [1.09, 1.71], $p = 0.006$) and depression (OR = 1.55 [1.21, 1.97],

$p = 4.7 \times 10^{-4}$) diagnoses. For associations of individual LTCs see Supplementary Table S14.

There were similar associations of childhood LTC with total (continuous) depression and anxiety symptom scores (Supplementary Table S15), with stronger evidence of an association between cerebral palsy and elevated self-reported anxiety symptom scores at age 14 ($b = 1.37 \pm 0.43$, $p = 0.001$), as well as between arthritis and elevated self-reported depressive symptom scores at age 14 ($b = 2.15 \pm 0.95$, $p = 0.02$).

There were associations between asthma, chronic fatigue syndrome, diabetes, epilepsy migraine, obesity, and reduced hearing and mother-reported depressive symptom scores at age 14 (see Supplementary Table S16), although mother- and self-reported SMFQ scores at age 14 were only modestly correlated (Spearman's $r = 0.36$, $p < 2.2 \times 10^{-6}$).

MR using adult anxiety and depression GWAS. Two-sample MR testing for causality between childhood LTCs and depression/anxiety using the largest available GWASs from adult

Table 2. Results from Mendelian Randomisation (MR) analysis of childhood LTCs and youth-onset depression using iPSYCH2015 outcome data

Exposure	Outcome	MR method	N SNPs	OR	95% CI		P-value	P _{FDR}
					Lower	Upper		
Childhood obesity	Youth-onset depression	Inverse variance weighted	8	0.99	0.9	1.09	0.87	0.99
		MR Egger	8	0.89	0.57	1.40	0.64	
		Weighted median	8	0.98	0.87	1.12	0.80	
		Weighted mode	8	0.92	0.75	1.11	0.41	
Childhood-onset asthma	Youth-onset depression	Inverse variance weighted	86	0.98	0.94	1.02	0.33	0.74
		MR Egger	86	0.95	0.86	1.05	0.31	
		Weighted median	86	0.99	0.93	1.06	0.86	
		Weighted mode	86	1.00	0.92	1.10	0.95	
Eczema	Youth-onset depression	Inverse variance weighted	10	0.99	0.88	1.12	0.88	0.99
		MR Egger	10	1.20	0.73	1.98	0.48	
		Weighted median	10	0.96	0.84	1.10	0.60	
		Weighted mode	10	0.96	0.78	1.17	0.67	
Genetic generalized epilepsy	Youth-onset depression	Inverse variance weighted	19	1.12	0.91	1.37	0.28	0.74
		MR Egger	19	0.79	0.25	2.54	0.70	
		Weighted median	19	1.07	0.85	1.35	0.54	
		Weighted mode	19	1.07	0.74	1.54	0.73	
Juvenile idiopathic arthritis	Youth-onset depression	Inverse variance weighted	4	1.01	0.98	1.05	0.43	0.74
		MR Egger	4	1.09	0.99	1.20	0.22	
		Weighted median	4	1.02	0.98	1.06	0.30	
		Weighted mode	4	1.03	0.99	1.07	0.27	
Migraine	Youth-onset depression	Inverse variance weighted	96	1	0.91	1.1	0.99	0.99
		MR Egger	96	0.79	0.61	1.01	0.07	
		Weighted median	96	0.96	0.83	1.11	0.56	
		Weighted mode	96	0.96	0.78	1.19	0.72	
Type 1 diabetes	Youth-onset depression	Inverse variance weighted	40	0.99	0.97	1.01	0.28	0.74
		MR Egger	40	0.97	0.94	1.00	0.07	
		Weighted median	40	0.98	0.96	1.01	0.19	
		Weighted mode	40	0.99	0.96	1.01	0.26	

samples (Als et al., 2023; Purves et al., 2020) indicated some evidence that genetic liability to migraine is causally linked with depression, and genetic liability to type 1 diabetes and decreased risk of an anxiety disorder (Supplementary Table S17); however, these estimates did not survive FDR correction. There was also limited evidence of pleiotropy for these estimates (Supplementary Table S18).

Discussion

With the prevalence of youth anxiety and depression increasing globally (Thapar et al., 2022), it is vital to identify potentially causal risk factors. Rates of LTCs have also risen across a similar time period (Van Cleave et al., 2010) indicating a potential causal factor, although social changes are also important (Collishaw, 2015). In this study, we investigated the association between LTCs and anxiety/depression in young people using phenotypic and genomic

data from a large cohort with linked medical registry data. Consistent with previous literature, we observed phenotypic associations between several childhood LTCs and depression and anxiety disorders and symptoms, with neurological conditions such as epilepsy and migraine, as well as obesity, demonstrating the highest risk. We then used MR as a test of causality. However, our MR findings did not indicate causal relationships between most LTCs and youth anxiety and depression, with the exception of migraine.

Our observed phenotypic associations between childhood LTCs and depression/anxiety are comparable to estimates from previous literature, demonstrating associations between a range of LTCs and clinically diagnosed anxiety/depression as well as elevated anxiety and depression symptoms in youth. Overall, we saw a 48% increase in MDD and a 44% increase in anxiety disorder diagnosis in young people with any LTC compared with those without. In line with previous evidence, we observed associations between arthritis (Fair, Rodriguez, Knight, & Rubinstein, 2019), asthma (Dudeny, Sharpe,

Jaffe, Jones, & Hunt, 2017), chronic fatigue syndrome (Loades *et al.*, 2021), epilepsy (Ekinici, Titus, Rodopman, Berkem, & Trevathan, 2009), migraine (Falla *et al.*, 2022), and obesity (Lindberg, Hagman, Danielsson, Marcus, & Persson, 2020) with elevated anxiety. For depression, we observed associations with asthma (Chen *et al.*, 2014), chronic fatigue syndrome (Bould, Collin, Lewis, Rimes, & Crawley, 2013), migraine (Falla *et al.*, 2022), and obesity (Lindberg *et al.*, 2020). Unlike previous studies we did not see strong evidence of worse mental health in young people with type 1 diabetes (Buchberger *et al.*, 2016; Liu *et al.*, 2022), reduced hearing/hearing loss (Stevenson, Kreppner, Pimperton, Worsfold, & Kennedy, 2015), or coeliac disease (Coburn, Puppa, & Blanchard, 2019), although previous evidence for coeliac disease is varied.

However, MR analyses did not support a causal interpretation of the observational associations between the majority of childhood LTCs and youth anxiety/depression, with the only consistent association seen in both observational and MR estimates being for migraine and depression symptoms as well as weaker evidence for anxiety symptoms. There was also some evidence that genetic liability to child-onset asthma could have causal effects on depression and anxiety disorders using MR analysis in MoBa data, however, these associations were not present in the observational analysis.

There are several potential explanations for our findings. First, the observed phenotypic associations between childhood LTCs and anxiety/depression are not causal. Previous literature indicates a number of potential confounders, including shared risk factors for LTCs and anxiety/depression (e.g. pre-/peri-natal factors (Fitzallen, Sagar, Taylor, & Bora, 2021, Heikkila *et al.*, 2021), family adversities (Davies *et al.*, 2016; Kaasboll, Skokauskas, Lydersen, & Sund, 2021; Kinnunen *et al.*, 2021; Sieh, Meijer, Oort, Visser-Meily, & Van der Leij, 2010; Vejrup, Hillesund, Agnihotri, Helle, & Øverby, 2023), lifestyle factors (Liu *et al.*, 2022; Sundell & Angelhoff, 2021), and poverty (Lai *et al.*, 2019, Najman *et al.*, 2010, Royal College of Paediatrics and Child Health, 2020, Thapar *et al.*, 2022). To our knowledge, a thorough investigation of the effect of these confounding factors on observed associations has not yet been undertaken. However, it has been shown that residual confounding remains a problem in all observational designs (Fewell, Davey Smith, & Sterne, 2007; Thapar & Rutter, 2019). For this reason, it is important to use alternate designs to infer causality. Additionally, there is the possibility of other shared genetic risk factors (independent of genetic instruments used in the current study which did not show evidence of horizontal pleiotropy) contributing both to LTCs and depression/anxiety, since previous evidence suggests genetic overlap for some LTCs (Zhu *et al.*, 2019).

Longitudinal and MR estimates did not suggest evidence for reverse causation (that genetic liability to anxiety/depression could lead to childhood LTCs), nor does the normal temporal pattern of condition emergence support this, with many childhood LTCs onset in young childhood (such as cerebral palsy and asthma), whereas the typical age of onset for depression and many types of anxiety is during adolescence (McGrath *et al.*, 2023, Thapar *et al.*, 2022). For LTCs with a more variable age-of-onset, such as migraine, type 1 diabetes, and chronic fatigue syndrome, we were able to use bidirectional MR analyses to test for the presence of reverse causation, where we observed little causal evidence in the direction of youth-onset depression to LTC.

Our MR findings suggested that LTCs, excluding migraine, do not appear to be a causal risk factor for youth anxiety or depression. This is important as it suggests that emotional disorders are not unavoidable in children and adolescents diagnosed with an LTC,

since mechanisms other than a causal relationship may explain observed associations (e.g. poverty, early lifestyle factors, and family environment). It also suggests that LTCs are unlikely to explain the recent increase in rates of anxiety and depression in young people, although the causal relationship should be assessed using other designs as there are limitations to MR.

An exception to our findings was the potentially causal relationship between migraine and increased risk of depression/anxiety at age 14, with MR estimates in agreement with observational estimates. These results were supplemented with our MR analysis using the largest adult depression GWAS as an outcome (Als *et al.*, 2023), which also indicated weaker evidence of a causal relationship between genetic liability to migraine and lifelong depression risk. However, we did not find evidence consistent with a causal effect of genetic liability to migraine on a clinical MDD diagnosis in analyses using both MoBa and iPSYCH2015 data, potentially indicating different effects of migraine on depression symptoms versus a clinical diagnosis in young people. There is a large body of research on the links between migraine and depression, suggesting a variety of biopsychosocial mechanisms that might contribute to the association. These have included shared genetic risk, brain neurobiology, and environmental factors such as stress (Baksa, Gonda, & Juhasz, 2017; Wachowska *et al.*, 2023).

These results should be considered in light of several methodological points. The first is the potential of low statistical power/measurement error for MR analyses because of the limited availability of robust genetic instruments for some exposures, meaning that the lack of an observed causal effect between exposures and outcomes does not rule out the presence of a causal relationship. We recommend that other designs are used to triangulate evidence from this study including traditional genetic studies such as twin designs. We were stringent about our selection of genetic instruments in that we prioritized GWAS with a child/adolescent focus, which could be considered a limitation since larger (non-age-specific) GWAS may provide more genetic instruments. However, we opted for this approach because the genetic architecture of LTCs with childhood-onset are not the same as typically adult-onset LTCs, for example childhood versus adult-onset asthma (Ferreira *et al.*, 2019) and type 1 versus type 2 diabetes (Aylward, Chiou, Okino, Kadakia, & Gaulton, 2018), and similarly is the case for youth versus adult-onset depression (Nguyen *et al.*, 2023).

A further limitation is that we used mother-reporting of offspring LTCs because of the limited availability of LTC diagnostic data. This meant that we were not able to capture all childhood LTCs, only those included in the questionnaires. Further, we cannot be sure that mother-reporting was completely accurate. However, evidence from developmental disorders and other child health outcomes indicate that parent reporting can generally be considered reliable (DiLalla, Trask, Casher, & Long, 2020; Miller, Perkins, Dai, & Fein, 2017; Orzan *et al.*, 2021; Shaikh, Nettiksimmons, Bell, Tancredi, & Romano, 2012), and prevalence rates of LTCs in this study are congruent with published rates. Further, we were unable to stratify offspring LTC by illness severity which may be an important contributor to mental health outcomes (Brady, Deighton, & Stansfeld, 2021). Despite these limitations, we observed phenotypic associations.

Although these results do not provide support for a causal link between most childhood LTCs and anxiety/depression in youth, it has been consistently shown, across international cohorts, that young people with LTCs have higher rates of mental disorders (Finning *et al.*, 2022; Pinquart & Shen, 2011b; Suryavanshi & Yang,

2016). Therefore, this should be considered in the provision of services and support for these children, especially in light of the increasing rates of both physical and mental health conditions in young people globally (Abarca-Gómez et al., 2017; Dharmage et al., 2019; Royal College of Paediatrics and Child Health, 2020; Thapar et al., 2022; Van Cleave et al., 2010).

Conclusion

Evidence from this study substantiates previous evidence that children and young people with LTCs experience higher rates of depression and anxiety than their physically healthy counterparts. However, using MR analysis, we saw limited evidence to support causal relationships between common childhood LTCs and anxiety and depression, highlighting the need for further investigations to understand the observed associations between childhood LTCs and youth mental health.

Supplementary material. To view supplementary material for this article, please visit <http://doi.org/10.1017/S0033291724003271>.

Data availability statement. Data from the Norwegian Mother, Father, and Child Cohort Study and the Medical Birth Registry of Norway used in this study are managed by the national health register holders in Norway (Norwegian Institute of Public Health) and can be made available to researchers, provided approval from the Regional Committees for Medical and Health Research Ethics (REC), compliance with the EU General Data Protection Regulation (GDPR) and approval from the data owners. The consent given by the participants does not open for storage of data on an individual level in repositories or journals. Researchers who want access to data sets for replication should apply through helsedata.no. Access to data sets requires approval from The Regional Committee for Medical and Health Research Ethics in Norway and an agreement with MoBa.

Acknowledgments. This work was supported by the Wolfson Centre for Young People's Mental Health, established with support from the Wolfson Foundation. The Norwegian Mother, Father, and Child Cohort Study is supported by the Norwegian Ministry of Health and Care Services and the Ministry of Education and Research. We are grateful to all the participating families in Norway who take part in this ongoing cohort study. We thank the Norwegian Institute of Public Health (NIPH) for generating high-quality genomic data. This research is part of the HARVEST collaboration, supported by the Research Council of Norway (RCN) (#229624). We also thank the NORMENT Centre for providing genotype data, funded by RCN (#223273), South East Norway Health Authority, and KG Jebsen Stiftelsen. Further, we thank the Centre for Diabetes Research, the University of Bergen for providing genotype data and performing quality control and imputation of the data funded by the ERC AdG project SELECTIONPREDISPOSED, Stiftelsen Kristian Gerhard Jebsen, Trond Mohn Foundation, NRC, the Novo Nordisk Foundation, the University of Bergen, and the Western Norway health Authorities (Helse Vest). Part of this work was performed on the TSD (Tjeneste for Sensitive Data) facilities, owned by the University of Oslo, operated and developed by the TSD service group at the University of Oslo, IT-Department (USIT) (tsd-drift@usit.uio.no). The analyses were also performed on resources provided by Sigma2 – the National Infrastructure for High-Performance Computing and Data Storage in Norway. Data from the Norwegian Patient Registry has been used in this publication. The interpretation and reporting of these data are the sole responsibility of the authors, and no endorsement by the Norwegian Patient Registry is intended nor should be inferred. AH was supported by the Research Council of Norway (#274611) and the South-Eastern Norway Regional Health Authority (#2020022, #2019097, and #2021045), and the European Union's Horizon Europe Research and Innovation Programme (FAMILY, grant agreement No 101057529). DB and LJH were supported by the South-Eastern Norway Regional Health Authority (#2922083). EC was supported by the South-Eastern Norway Regional Health Authority (#2021045) and the Research Council of Norway (#274611).

iPSYCH is supported by the Lundbeck Foundation, as well as Aarhus University, the Capital Region of Denmark, Statens Serum Institut, Aarhus University Hospital, the Stanley Centre at Broad Institute, Simons Foundation, The National Institute of Mental Health, and the Novo Nordisk Foundation. Analysis involving iPSYCH data was performed on GenomeDK servers. The authors also acknowledge the International Headache Genetics Consortium for sharing GWAS summary statistics. GDS and ES work within the MRC Integrative Epidemiology Unit at the University of Bristol, which is supported by the Medical Research Council (MC_UU_00011/1). Views and opinions expressed are however those of the author(s) only and do not necessarily reflect those of the funding agents. The granting authorities cannot be held responsible for them.

Conflicts of interest. The authors report no conflicts of interest.

In accordance with the consent structure of iPSYCH and Danish law, individual-level genotype and phenotype data are not able to be shared publicly.

Information on GWAS summary statistics for all LTCs studied is detailed in the original studies (see Supplementary Table S1).

References

- Abarca-Gómez, L., Abdeen, Z. A., Hamid, Z. A., Abu-Rmeileh, N. M., Acosta-Cazares, B., Acuin, C., ... Ezzati, M. (2017). Worldwide trends in body-mass index, underweight, overweight, and obesity from 1975 to 2016: a pooled analysis of 2416 population-based measurement studies in 128.9 million children, adolescents, and adults. *The Lancet*, **390**, 2627–2642.
- Agrawal, N., & Govender, S. (2011). Epilepsy and neuropsychiatric comorbidities. *Advances in Psychiatric Treatment*, **17**, 44–53.
- Als, T. D., Kurki, M. I., Grove, J., Voloudakis, G., Therrien, K., Tasanko, E., ... Borglum, A. D. (2023). Depression pathophysiology, risk prediction of recurrence and comorbid psychiatric disorders using genome-wide analyses. *Nature Medicine* **29**, 1832–1844.
- Angold, A., Costello, E. J., Messer, S. C., & Pickles, A. (1995). Development of a short questionnaire for use in epidemiological studies of depression in children and adolescents. *International Journal of Methods in Psychiatric Research*, **5**(4), 237–249.
- Aylward, A., Chiou, J., Okino, M.-L., Kadakia, N., & Gaulton, K. J. (2018). Shared genetic risk contributes to type 1 and type 2 diabetes etiology. *Human Molecular Genetics*.
- Baksa, D., Gonda, X., & Juhasz, G. (2017). Why are migraineurs more depressed? A review of the factors contributing to the comorbidity of migraine and depression. *Neuropsychopharmacologia Hungarica*, **19**, 37–44.
- Bakula, D. M., Sharkey, C. M., Perez, M. N., Espeleta, H. C., Hawkins, M., Chaney, J. M., & Mullins, L. L. (2019). The role of stigma in the relationship between illness intrusiveness and adjustment in adolescents and young adults: A path model. *Journal of Pediatric Psychology*, **44**, 611–619.
- Birmaher, B., Brent, D. A., Chiappetta, L., Bridge, J., Monga, S., & Baugher, M. (1999). Psychometric properties of the Screen for Child Anxiety Related Emotional Disorders (SCARED): A replication study. *The Journal of the American Academy of Child & Adolescent Psychiatry*, **38**, 1230–1236.
- Birmaher, B., Khetarpal, S., Brent, D., Cully, M., Balach, L., Kaufman, J., & Neer, S. M. (1997). The Screen for Child Anxiety Related Emotional Disorders (SCARED): scale construction and psychometric characteristics. *The Journal of the American Academy of Child & Adolescent Psychiatry*, **36**, 545–553.
- Blackwell, C. K., Elliott, A. J., Ganiban, J., Herbstman, J., Hunt, K., Forrest, C. B., & Camargo, C. A., Jr. (2019). General Health and Life Satisfaction in Children With Chronic Illness. *Pediatrics*, **143**.
- Bould, H., Collin, S. M., Lewis, G., Rimes, K., & Crawley, E. (2013). Depression in paediatric chronic fatigue syndrome. *Archives of Disease in Childhood* **98**, 425–428.
- Bowden, J., Davey Smith, G., & Burgess, S. (2015). Mendelian randomization with invalid instruments: Effect estimation and bias detection through Egger regression. *International Journal of Epidemiology*, **44**, 512–528.
- Bradfield, J. P., Vogelesang, S., Felix, J. F., Chesni, A., Helgeland, O., Horikoshi, M., ... Grant, S. F. A. (2019). A trans-ancestral meta-analysis of genome-wide association studies reveals loci associated with childhood obesity. *Human Molecular Genetics*, **28**, 3327–3338.

- Brady, A. M., Deighton, J., & Stansfeld, S. (2021). Chronic illness in childhood and early adolescence: A longitudinal exploration of co-occurring mental illness. *Development and Psychopathology*, **33**, 885–898.
- Buchberger, B., Huppertz, H., Krabbe, L., Lux, B., Mattivi, J. T., & Sifarikas, A. (2016). Symptoms of depression and anxiety in youth with type 1 diabetes: A systematic review and meta-analysis. *Psychoneuroendocrinology*, **70**, 70–84.
- Bybjerg-Grauholm, J., Pedersen, C. B., Bækvad-Hansen, M., Pedersen, M. G., Adamsen, D., Hansen, C. S., ... Mortensen, P. B. (2020). The iPSYCH2015 Case-Cohort sample: updated directions for unravelling genetic and environmental architectures of severe mental disorders. *medRxiv*, 2020.11.30.20237768.
- Chen, M. H., Su, T. P., Chen, Y. S., Hsu, J. W., Huang, K. L., Chang, W. H., ... Bai, Y. M. (2014). Higher risk of developing major depression and bipolar disorder in later life among adolescents with asthma: A nationwide prospective study. *Journal of Psychiatric Research*, **49**, 25–30.
- Coburn, S. S., Puppa, E. L., & Blanchard, S. (2019). Psychological comorbidities in childhood celiac disease: A systematic review. *Journal of Pediatric Gastroenterology and Nutrition*, **69**, e25–e33.
- Collishaw, S. (2015). Annual research review: Secular trends in child and adolescent mental health. *Journal of Child Psychology and Psychiatry*, **56**, 370–393.
- Compas, B. E., Jaser, S. S., Dunn, M. J., & Rodriguez, E. M. (2012). Coping with chronic illness in childhood and adolescence. *Annual Review of Clinical Psychology*, **8**, 455–480.
- Corfield, E. C., Frei, O., Shadrin, A. A., Rahman, Z., Lin, A., Athanasiu, L., ... Havdahl, A. (2022). The norwegian mother, father, and child cohort study (MoBa) genotyping data resource: MoBaPsychGen pipeline v.1. *bioRxiv*, 2022.06.23.496289.
- Davies, P. S. W., Funder, J., Palmer, D. J., Sinn, J., Vickers, M. H., & Wall, C. R. (2016). Early life nutrition and the opportunity to influence long-term health: an Australasian perspective. *Journal of Developmental Origins of Health and Disease*, **7**, 440–448.
- Dharmage, S. C., Perret, J. L., & Custovic, A. (2019). Epidemiology of asthma in children and adults. *Frontiers in Pediatrics*, **7**, 246.
- DiLalla, L. F., Trask, M. M., Casher, G. A., & Long, S. S. (2020). Evidence for reliability and validity of parent reports of twin children's birth information. *Early Childhood Research Quarterly*, **51**, 267–274.
- Dudenev, J., Sharpe, L., Jaffe, A., Jones, E. B., & Hunt, C. (2017). Anxiety in youth with asthma: A meta-analysis. *Pediatric Pulmonology*, **52**, 1121–1129.
- Dyb, G., Stensland, S., & Zwart, J. A. (2015). Psychiatric comorbidity in childhood and adolescence headache. *Current Pain and Headache Reports*, **19**, 5.
- Ekinci, O., Titus, J. B., Rodopman, A. A., Berkem, M., & Trevathan, E. (2009). Depression and anxiety in children and adolescents with epilepsy: Prevalence, risk factors, and treatment. *Epilepsy & Behavior*, **14**, 8–18.
- Eyre, O., Bevan Jones, R., Agha, S. S., Wootton, R. E., Thapar, A. K., Stergiakouli, E., ... Riglin, L. (2021). Validation of the short mood and feelings questionnaire in young adulthood. *Journal of Affective Disorders*, **294**, 883–888.
- Fair, D. C., Rodriguez, M., Knight, A. M., & Rubinstein, T. B. (2019). Depression and anxiety in patients with juvenile idiopathic arthritis: Current insights and impact on quality of life A systematic review. *Open Access Rheumatology*, **11**, 237–252.
- Falla, K., Kuziek, J., Mahnaz, S. R., Noel, M., Ronksley, P. E., & Orr, S. L. (2022). Anxiety and depressive symptoms and disorders in children and adolescents with migraine: A systematic review and meta-analysis. *JAMA Pediatrics*, **176**, 1176–1187.
- Ferreira, M. A. R., Mathur, R., Vonk, J. M., Szwajda, A., Brumpton, B., Granell, R., ... Almqvist, C. (2019). Genetic architectures of childhood- and adult-onset asthma are partly distinct. *The American Journal of Human Genetics*, **104**, 665–684.
- Fewell, Z., Davey Smith, G., & Sterne, J. A. (2007). The impact of residual and unmeasured confounding in epidemiologic studies: A simulation study. *American Journal of Epidemiology*, **166**, 646–655.
- Finning, K., Neochoriti Varvarrigou, I., Ford, T., Panagi, L., & Ukoumunne, O. C. (2022). Mental health and school absenteeism in children with long-term physical conditions: A secondary analysis of the British Child and Adolescent Mental Health Surveys 2004 and 2007. *Child: Care, Health and Development*, **48**, 110–119.
- Fitzallen, G. C., Sagar, Y. K., Taylor, H. G., & Bora, S. (2021). Anxiety and depressive disorders in children born preterm: A meta-analysis. *Journal of Developmental & Behavioral Pediatrics*, **42**, 154–162.
- Forgetta, V., Manousaki, D., Istomine, R., Ross, S., Tessier, M. C., Marchand, L., ... Richards, J. B. (2020). Rare genetic variants of large effect influence risk of Type 1 diabetes. *Diabetes*, **69**, 784–795.
- Glazebrook, C., Hollis, C., Heussler, H., Goodman, R., & Coates, L. (2003). Detecting emotional and behavioural problems in paediatric clinics. *Child: Care, Health and Development*, **29**, 141–149.
- Harder, A., Nguyen, T. D., Pasma, J. A., Mosing, M. A., Hagg, S., & Lu, Y. (2022). Genetics of age-at-onset in major depression. *Translational Psychiatry*, **12**, 124.
- Hautakangas, H., Winsvold, B. S., Ruotsalainen, S. E., Bjornsdottir, G., Harder, A. V. E., Kogelman, L. J. A., ... Danish Blood Donor Study Genomic, C. (2022). Genome-wide analysis of 102,084 migraine cases identifies 123 risk loci and subtype-specific risk alleles. *Nature Genetics*, **54**, 152–160.
- Heikkila, K., Pulakka, A., Metsala, J., Alenius, S., Hovi, P., Gissler, M., ... Kajantie, E. (2021). Preterm birth and the risk of chronic disease multi-morbidity in adolescence and early adulthood: A population-based cohort study. *PLoS One*, **16**, e0261952.
- Hemani, G., Tilling, K., & Davey Smith, G. (2017). Orienting the causal relationship between imprecisely measured traits using GWAS summary data. *PLOS Genetics*, **13**, e1007081.
- Hemani, G., Zheng, J., Elsworth, B., Wade, K. H., Haberland, V., Baird, D., ... Haycock, P. C. (2018). The MR-Base platform supports systematic causal inference across the human phenome. *eLife*, **7**, e34408.
- Hinks, A., Cobb, J., Marion, M. C., Prahalad, S., Sudman, M., Bowes, J., ... Thompson, S. D. (2013). Dense genotyping of immune-related disease regions identifies 14 new susceptibility loci for juvenile idiopathic arthritis. *Nature Genetics*, **45**, 664–9.
- Hughes, A., Wade, K. H., Dickson, M., Rice, F., Davies, A., Davies, N. M., & Howe, L. D. (2021). Common health conditions in childhood and adolescence, school absence, and educational attainment: Mendelian randomization study. *npj Science of Learning*, **6**, 1.
- International League Against Epilepsy Consortium on Complex Epilepsies, Berkovic, S. F., Cavalleri, G. L., & Koeleman, B. P. (2022). Genome-wide meta-analysis of over 29,000 people with epilepsy reveals 26 loci and subtype-specific genetic architecture. *medRxiv*, 2022.06.08.22276120.
- Kaasboll, J., Skokauskas, N., Lydersen, S., & Sund, A. M. (2021). Parental Chronic Illness, Internalizing problems in young adulthood and the mediating role of adolescent attachment to parents: A prospective cohort study. *Frontiers in Psychiatry*, **12**, 807563.
- Kinnunen, L., Nordström, T., Niemelä, M., Räsänen, S., Whittle, S., & Miettinen, J. (2021). Parental physical illnesses and their association with subsequent externalizing and internalizing symptoms in children. *Journal of Child and Family Studies*, **30**, 2677–2689.
- Lai, E. T. C., Wickham, S., Law, C., Whitehead, M., Barr, B., & Taylor-Robinson, D. (2019). Poverty dynamics and health in late childhood in the UK: evidence from the Millennium Cohort Study. *Archives of Disease in Childhood*, **104**, 1049–1055.
- Lawlor, D. A., Tilling, K., & Davey Smith, G. (2016). Triangulation in aetiological epidemiology. *International Journal of Epidemiology*, **45**, 1866–1886.
- Lindberg, L., Hagman, E., Danielsson, P., Marcus, C., & Persson, M. (2020). Anxiety and depression in children and adolescents with obesity: A nationwide study in Sweden. *BMC Medicine*, **18**, 30.
- Liu, J., Ji, X., Pitt, S., Wang, G., Rovit, E., Lipman, T., & Jiang, F. (2022). Childhood sleep: Physical, cognitive, and behavioral consequences and implications. *World Journal of Pediatrics*, 1–11.
- Liu, S., Leone, M., Ludvigsson, J. F., Lichtenstein, P., D'Onofrio, B., Svensson, A. M., ... Butwicki, A. (2022). Association and familial coaggregation of childhood-onset Type 1 diabetes with depression, anxiety, and stress-related disorders: A population-based cohort study. *Diabetes Care*, **45**, 1987–1993.
- Loades, M. E., Read, R., Smith, L., Higson-Sweeney, N. T., Laffan, A., Stallard, P., ... Crawley, E. (2021). How common are depression and anxiety in adolescents with chronic fatigue syndrome (CFS) and how should we screen for these mental health co-morbidities? A clinical cohort study. *European Child & Adolescent Psychiatry*, **30**, 1733–1743.
- Lum, A., Wakefield, C. E., Donnan, B., Burns, M. A., Fardell, J. E., & Marshall, G. M. (2017). Understanding the school experiences of children and

- adolescents with serious chronic illness: a systematic meta-review. *Child: Care, Health and Development*, **43**, 645–662.
- Magnus, P., Birke, C., Vejrup, K., Haugan, A., Alsaker, E., Daltveit, A. K., ... Stoltenberg, C. (2016). Cohort profile update: The Norwegian mother and child cohort study (MoBa). *International Journal of Epidemiology*, **45**, 382–388.
- McGrath, J. J., Al-Hamzawi, A., Alonso, J., Altwajiri, Y., Andrade, L. H., Bromet, E. J., ... Collaborators, W. H. O. W. M. H. S. (2023). Age of onset and cumulative risk of mental disorders: a cross-national analysis of population surveys from 29 countries. *Lancet Psychiatry*, **10**(9), 668–681.
- Miller, L. E., Perkins, K. A., Dai, Y. G., & Fein, D. A. (2017). Comparison of parent report and direct assessment of Child skills in toddlers. *Research in Autism Spectrum Disorders*, **41–42**, 57–65.
- Mokkink, L. B., van der Lee, J. H., Grootenhuis, M. A., Offringa, M., Heymans, H. S., Dutch National Consensus Committee Chronic, D., & Health Conditions in, C. (2008). Defining chronic diseases and health conditions in childhood (0-18 years of age): national consensus in the Netherlands. *European Journal of Pediatrics*, **167**, 1441–1417.
- Najman, J. M., Hayatbakhsh, M. R., Clavarino, A., Bor, W., O'Callaghan, M. J., & Williams, G. M. (2010). Family poverty over the early life course and recurrent adolescent and young adult anxiety and depression: A longitudinal study. *American Journal of Public Health*, **100**, 1719–1723.
- Nguyen, T.-D., Kowalec, K., Pasma, J., Larsson, H., Lichtenstein, P., Dalman, C., ... Lu, Y. (2023). Genetic contribution to the heterogeneity of major depressive disorder: evidence from a sibling-based design using swedish national registers. *American Journal of Psychiatry*, **180**, 714–722.
- Orzan, E., Battelino, S., Ciciriello, E., Bonifacio, S., Pellizzoni, S., & Saksida, A. (2021). Reliability of parental assessment of auditory skills in young children: A cross-sectional study in Italian language. *BMJ Open*, **11**, e042297.
- Paltiel, L., Anita, H., Skjerdet, T., Harbak, K., Bækken, S., Nina Kristin, S., ... Magnus, P. (2014). The biobank of the Norwegian Mother and Child Cohort Study – present status. *Norsk Epidemiologi*, **24**(1), 29–35.
- Panagi, L., Newlove-Delgado, T., White, S. R., Bennett, S., Heyman, I., Shafraan, R., & Ford, T. (2022a). Trends in comorbid physical and mental health conditions in children from 1999 to 2017 in England. *European Child & Adolescent Psychiatry*.
- Panagi, L., White, S. R., Patel, S., Bennett, S., Shafraan, R., & Ford, T. (2022b). The importance of definitions in the measurement of long-term health conditions in childhood. Variations in prevalence of long-term health conditions in the UK using data from the Millennium Cohort Study, 2004–2015. *International Journal of Methods in Psychiatric Research*, **31**, e1926.
- Paternoster, L., Standl, M., Waage, J., Baurecht, H., Hotze, M., Strachan, D. P., ... Weidinger, S. (2015). Multi-ancestry genome-wide association study of 21,000 cases and 95,000 controls identifies new risk loci for atopic dermatitis. *Nature Genetics*, **47**, 1449–1456.
- Pedersen, C. B., Bybjerg-Grauholm, J., Pedersen, M. G., Grove, J., Agerbo, E., Bækvad-Hansen, M., ... Mortensen, P. B. (2018). The iPSYCH2012 case-cohort sample: New directions for unravelling genetic and environmental architectures of severe mental disorders. *Molecular Psychiatry*, **23**, 6–14.
- Pinquart, M. (2013a). Do the Parent–Child Relationship and parenting behaviors differ between families with a child with and without chronic illness? A Meta-analysis. *Journal of Pediatric Psychology*, **38**, 708–721.
- Pinquart, M. (2013b). Self-esteem of children and adolescents with chronic illness: A meta-analysis. *Child: Care, Health and Development*, **39**, 153–161.
- Pinquart, M. (2017). Systematic review: Bullying involvement of children with and without chronic physical illness and/or physical/sensory disability—a meta-analytic comparison with healthy/nondisabled peers. *Journal of Pediatric Psychology*, **42**, 245–259.
- Pinquart, M., & Shen, Y. (2011a). Anxiety in children and adolescents with chronic physical illnesses: A meta-analysis. *Acta Paediatrica*, **100**, 1069–1076.
- Pinquart, M., & Shen, Y. (2011b). Depressive symptoms in children and adolescents with chronic physical illness: An updated meta-analysis. *Journal of Pediatric Psychology*, **36**, 375–84.
- Pittet, I., Berchtold, A., Akre, C., Michaud, P. A., & Suris, J. C. (2010). Are adolescents with chronic conditions particularly at risk for bullying? *Archives of Disease in Childhood*, **95**, 711.
- Purves, K. L., Coleman, J. R. L., Meier, S. M., Rayner, C., Davis, K. A. S., Cheesman, R., ... Eley, T. C. (2020). A major role for common genetic variation in anxiety disorders. *Molecular Psychiatry*, **25**, 3292–3303.
- Qiu, Y., Xu, L., Pan, Y., He, C., Huang, Y., Xu, H., ... Dong, C. (2021). Family resilience, parenting styles and psychosocial adjustment of children with chronic illness: A cross-sectional study. *Frontiers in Psychiatry*, **12**.
- R Core Team (2022). *R: A language and environment for statistical Computing. R foundation for statistical computing*. Vienna, Austria.
- Royal College of Paediatrics and Child Health. (2020). *State of child health*. RCPCH: London.
- Sanderson, E., Glymour, M. M., Holmes, M. V., Kang, H., Morrison, J., Munafò, M. R., ... Davey Smith, G. (2022). Mendelian randomization. *Nature Reviews Methods Primers*, **2**.
- Shaikh, U., Nettiksimmons, J., Bell, R. A., Tancredi, D., & Romano, P. S. (2012). Accuracy of parental report and electronic health record documentation as measures of diet and physical activity counseling. *Academic Pediatrics*, **12**, 81–87.
- Sieh, D. S., Meijer, A. M., Oort, F. J., Visser-Meily, J. M., & Van der Leij, D. A. (2010). Problem behavior in children of chronically ill parents: a meta-analysis. *Clinical Child and Family Psychology Review*, **13**, 384–397.
- Skrivankova, V. W., Richmond, R. C., Woolf, B. A. R., Davies, N. M., Swanson, S. A., VanderWeele, T. J., ... Richards, J. B. (2021). Strengthening the reporting of observational studies in epidemiology using mendelian randomisation (STROBE-MR): Explanation and elaboration. *BMJ*, **375**, n2233.
- Slob, E. A. W., & Burgess, S. (2020). A comparison of robust Mendelian randomization methods using summary data. *Genetic Epidemiology*, **44**, 313–329.
- Smith, G. D., & Ebrahim, S. (2003). 'Mendelian randomization': Can genetic epidemiology contribute to understanding environmental determinants of disease? *International Journal of Epidemiology*, **32**, 1–22.
- Stevenson, J., Kreppner, J., Pimperton, H., Worsfold, S., & Kennedy, C. (2015). Emotional and behavioural difficulties in children and adolescents with hearing impairment: a systematic review and meta-analysis. *European Child & Adolescent Psychiatry European Child & Adolescent Psychiatry*, **24**, 477–496.
- Sundell, A. L., & Angelhoff, C. (2021). Sleep and its relation to health-related quality of life in 3–10-year-old children. *BMC Public Health*, **21**, 1043.
- Suryavanshi, M. S., & Yang, Y. (2016). Clinical and economic burden of mental disorders among children with chronic physical conditions, United States, 2008–2013. *Preventing Chronic Disease*, **13**:150535.
- Thapar, A., Eyre, O., Patel, V., & Brent, D. (2022). Depression in young people. *Lancet*, **400**, 617–631.
- Thapar, A., & Riglin, L. (2020). The importance of a developmental perspective in Psychiatry: What do recent genetic-epidemiological findings show? *Molecular Psychiatry*, **25**, 1631–1639.
- Thapar, A., & Rutter, M. (2019). Do natural experiments have an important future in the study of mental disorders? *Psychological Medicine*, **49**, 1079–1088.
- Van Cleave, J., Gortmaker, S. L., & Perrin, J. M. (2010). Dynamics of obesity and chronic health conditions among children and youth. *JAMA*, **303**, 623–630.
- Vejrup, K., Hillesund, E. R., Agnihotri, N., Helle, C., & Øverby, N. C. (2023). Diet in early life is related to child mental health and personality at 8 years: findings from the norwegian mother, father and child cohort study (MoBa). *Nutrients*, **15**.
- Verbanck, M., Chen, C.-Y., Neale, B. & Do, R. (2018). Detection of widespread horizontal pleiotropy in causal relationships inferred from Mendelian randomization between complex traits and diseases. *Nature Genetics*, **50**, 693–698.
- Wachowska, K., Bliźniewska-Kowalska, K., Sławek, J., Adamczyk-Sowa, M., Szulc, A., Maes, M., ... Gałeczki, P. (2023). Common pathomechanism of migraine and depression. *Psychiatria Polska*, **57**, 405–419.
- Zhu, Z., Zhu, X., Liu, C. L., Shi, H., Shen, S., Yang, Y., ... Liang, L. (2019). Shared genetics of asthma and mental health disorders: a large-scale genome-wide cross-trait analysis. *European Respiratory Journal*, **54**.