



Eigenvalue Optimisation on Flat Tori and Lattice Points in Anisotropically Expanding Domains

Jean Lagacé

Abstract. This paper is concerned with the maximisation of the k -th eigenvalue of the Laplacian amongst flat tori of unit volume in dimension d as k goes to infinity. We show that in any dimension maximisers exist for any given k , but that any sequence of maximisers degenerates as k goes to infinity when the dimension is at most 10. Furthermore, we obtain specific upper and lower bounds for the injectivity radius of any sequence of maximisers. We also prove that flat Klein bottles maximising the k -th eigenvalue of the Laplacian exhibit the same behaviour. These results contrast with those obtained recently by Gittins and Larson, stating that sequences of optimal cuboids for either Dirichlet or Neumann boundary conditions converge to the cube no matter the dimension. We obtain these results via Weyl asymptotics with explicit control of the remainder in terms of the injectivity radius. We reduce the problem at hand to counting lattice points inside anisotropically expanding domains, where we generalise methods of Yu. Kordyukov and A. Yakovlev by considering domains that expand at different rates in various directions.

1 Introduction and Main Results

Let (M, g) be a smooth closed Riemannian manifold of dimension d . We study the Laplace eigenvalue problem

$$\Delta u + \lambda u = 0.$$

The eigenvalues of the Laplacian form a discrete, nondecreasing sequence, repeating every eigenvalue according to multiplicity,

$$0 = \lambda_0(M, g) \leq \lambda_1(M, g) \leq \cdots \nearrow \infty,$$

accumulating only at infinity.

1.1 Asymptotic Eigenvalue Optimisation

In this paper, we study the maximisation problem

$$(1.1) \quad \Lambda_k^*(\mathcal{G}) := \sup_{g \in \mathcal{G}} \Lambda_k(M, g) := \sup_{g \in \mathcal{G}} \text{Vol}_g(M)^{2/d} \lambda_k(M, g),$$

where \mathcal{G} is a class of metrics on M . This problem has been studied extensively for $k = 1$ in many settings: closed manifolds, manifolds with Neumann boundary conditions,

Received by the editors September 14, 2018.

Published online on Cambridge Core March 7, 2019.

The research of the author was supported by NSERC's Alexander-Graham-Bell doctoral scholarship.

AMS subject classification: 35P20, 11H06, 52C07.

Keywords: spectral optimisation, Laplacian, eigenvalue, asymptotics.

and manifolds with Dirichlet boundary conditions, in which case one minimises Λ_k . Note that for closed manifolds it only makes sense to maximise Λ_k . Indeed, for any k one can find a sequence of metrics g_n of unit volume such that $\Lambda_k(M, g_n) \rightarrow 0$ as $n \rightarrow \infty$ by considering a sequence of metrics that degenerate to a disjoint union of $k + 1$ closed manifolds touching at a point.

An interesting feature is that the extremisers for low eigenvalues are in general very symmetric. Indeed, the Faber–Krahn inequality [12, 23, 24] and the Szegő–Weinberger inequality [27, 28] imply that the ball is the extremiser for Λ_1 with Dirichlet or Neumann boundary conditions in any dimension. In the case of closed surfaces, Hersch [15] has shown that the round sphere is the maximiser for Λ_1 amongst two-dimensional spheres, and Nadirashvili [26] has shown that the equilateral flat torus is the maximiser for Λ_1 amongst surfaces of genus one.

For higher eigenvalues on domains, one does not expect those symmetries to appear. Indeed, A. Berger [6] has shown that disks or unions of disks can minimise Λ_k on domains in the plane with Dirichlet boundary conditions only finitely many times. Furthermore, numerical experiments of Antunes and Freitas [1] suggest that optimal domains in \mathbb{R}^2 may not exhibit many symmetries for $k \geq 5$. However, the same authors investigated in [2] the behaviour of optimal domains as k goes to infinity. More specifically, they showed that amongst rectangles with Dirichlet boundary condition, the sequence of rectangles minimising Λ_k converges to the square in the Hausdorff metric. This has led to a series of papers [4, 5, 13] culminating in a proof by Gittins and Larson, who show that in any dimension and with either Neumann or Dirichlet boundary conditions, the sequence of optimal cuboids converges to the cube.

Without any restriction on the metric, one does not even have a maximiser amongst closed manifolds. Indeed, Colbois and Dodziuk [10] have shown that amongst all metrics of fixed volume on a manifold, one can make λ_1 as large as possible. For metrics on closed surfaces, one does not necessarily expect the sequence of maximising metrics to converge to a smooth metric. For instance, Karpukhin, Nadirashvili, Penskoi, and Polterovich [18] obtained in a recent preprint that the maximising metric on the two-dimensional sphere for the k -th Laplace eigenvalue degenerates to a union of k kissing round spheres.

We study the maximisation problem (1.1) for metrics on two classes of closed manifold. The first one is the class \mathcal{M} of flat metrics on tori in dimension d . Let $\mathcal{L} = \mathrm{GL}_d(\mathbb{R})/\mathrm{GL}_d(\mathbb{Z})$ be the set of lattices in \mathbb{R}^d equipped with the quotient topology. We identify \mathcal{M} with \mathcal{L} , since

$$\mathcal{M} = \{\mathbb{T}_\Gamma = \mathbb{R}^d/\Gamma : \Gamma \in \mathcal{L}\}.$$

As such, convergence in \mathcal{M} will be identified with convergence in \mathcal{L} . We study the properties of maximisers to (1.1) in \mathcal{L}_0 the subset of all lattices with unit determinant, which corresponds to subset \mathcal{M}_0 of flat tori with unit volume.

The second class that we study is the set \mathcal{E} of flat metrics on Klein bottles. Flat Klein bottles are quotients of two-dimensional flat rectangular tori, and as such are described by the two-parameter family

$$\mathcal{E} := \{K(a, b) := (\mathbb{R}^d/(a\mathbb{Z} \oplus b\mathbb{Z})) / \sim : (a, b) \in \mathbb{R}_+^2\},$$

where \sim is the relation $(x, y) \sim (x + \frac{a}{2}, b - y)$. Once again, we study the properties of maximisers of (1.1) in the class \mathcal{E}_0 of Klein bottles with unit volume, i.e., the family $K(a, b)$ where $ab = 2$.

Before discussing asymptotic properties of maximisers to the problem (1.1), we start by proving that such maximisers do exist.

Theorem 1.1 *For all $k \in \mathbb{N}$, there exist $\mathbb{T}_k^* \in \mathcal{M}_0$ and $K_k^* \in \mathcal{E}_0$ maximising the variational problems*

$$\Lambda_k^*(\mathcal{M}) = \sup_{\mathbb{T} \in \mathcal{M}} \Lambda_k(\mathbb{T}) \quad \text{and} \quad \Lambda_k^*(\mathcal{E}) = \sup_{K \in \mathcal{E}} \Lambda_k(K).$$

The behaviour of maximisers for tori and Klein bottles contrasts both with the results obtained for cuboids where the optimal cuboid converges to the cube and with the degeneracy results of [10, 18]. Indeed, we show that for tori of dimension $2 \leq d \leq 10$, the sequence of optimisers has no limit points in \mathcal{M}_0 . However, we also show that this degeneracy can happen without changing the curvature as was done in [18] or in [10].

Furthermore, we obtain a rate of degeneracy in terms of the injectivity radius. This is similar to the results in [13], where the rate of convergence to the cube is given. The range $2 \leq d \leq 10$ are the dimensions for which the volume of the unit ball ω_d is larger than $\omega_1 = 2$. In higher dimensions, the same type of result may hold, but the degeneracy certainly does not happen in the same way.

Theorem 1.2 *In dimension $2 \leq d \leq 10$, there are no accumulation points in \mathcal{M}_0 of any sequence $\{\mathbb{T}_k^*\}$. The injectivity radius of \mathbb{T}_k^* respects*

$$k^{-\frac{(1-d)^2}{d}} \ll \text{inj}(\mathbb{T}_k^*) \ll k^{-\frac{1}{d}}.$$

The lower bound is valid for all dimensions $d \in \mathbb{N}$.

Remark 1.3 In dimension 2, the lower bound and the upper bound are, at least to polynomial order, the same. In dimensions $3 \leq d \leq 10$ there is a discrepancy which is due to the fact that the lower bound is obtained in a less natural way (see Section 4.2 and the extra step involving Minkowski's successive minima theorem for the lower bound). We conjecture that any actual sequence of maximisers for $2 \leq d \leq 10$ respects $\text{inj}(\mathbb{T}_k^*) \asymp k^{-\frac{1}{d}}$.

In [17], Kao, Lai, and Osting conjectured that in dimension 2, the optimal flat torus was given by $\mathbb{T}_{2k}^* = \mathbb{R}^2 / \Gamma_{2k}$, where Γ_{2k} is the lattice spanned over \mathbb{Z} by the vectors

$$(1.2) \quad \gamma_1^{(2k)} = (k^2 - 1/4)^{-1/4} (1, 0) \quad \text{and} \quad \gamma_2^{(2k)} = (k^2 - 1/4)^{-1/4} (1/2, \sqrt{k^2 - 1/4}).$$

In dimension 2, flat tori of unit volume form a two-dimensional moduli space with parameters a, b , with $a \in (-1/2, 1/2]$, $b > 0$ such that $a^2 + b^2 \geq 1$. The associated lattices are spanned by

$$\gamma_1(a, b) = b^{-1/2} (1, 0) \quad \text{and} \quad \gamma_2(a, b) = b^{-1/2} (a, b).$$

It is shown in [17] that the flat torus in equation (1.2) is indeed maximal for Λ_{2k} amongst tori for which $a^2 + b^2 \geq (k-1)^2$. The upper bound on the injectivity radius in Theorem 1.2 yields that there exists a constant $c > 0$ such that the same torus has a higher Λ_{2k} than every flat tori such that $a^2 + b^2 \leq ck^2$.

Our methods also allow us to study sequences of optimisers in the moduli space \mathcal{E} of flat Klein bottles. Indeed, we also have degeneracy in this case, and we can also describe the rate of degeneracy.

Theorem 1.4 *There are no accumulation points in \mathcal{E}_0 of any sequence $\{K_k^*\}$. The injectivity radius of K_k^* respects $k^{-\frac{1}{2}} \ll \text{inj}(K_k^*) \ll k^{-\frac{1}{2}}$.*

1.2 Explicit Exponent for the Remainder in Weyl's Law

In the papers [2, 4, 5, 13] on optimal cuboids a prominent feature consisted in finding uniform bounds on the eigenvalue counting function

$$N(\lambda; M) = \#\{\lambda_k(M) < \lambda\}.$$

Weyl's law states that for any fixed (M, g) , the counting function $N(\lambda; M)$ enjoys the asymptotics

$$(1.3) \quad N(\lambda; M) = \frac{\omega_d}{(2\pi)^d} \lambda^{\frac{d}{2}} + R(\lambda; M),$$

where $R(\lambda; M) = o(\lambda^{\frac{d}{2}})$ and ω_d is the volume of a unit ball in dimension d . Under the hypothesis that periodic geodesics have measure 0 in the cosphere bundle of M , Duistermaat and Guillemin [11] have shown that the remainder in equation (1.3) satisfies $R(\lambda; M) = o(\lambda^{\frac{d-1}{2}})$. Note that the size of $R(\lambda; M)$ depends on the geometry of M in a non-trivial way. Indeed, for any fixed λ , one can find a sequence g_n of metrics on M such that $N(\lambda; (M, g_n)) \rightarrow \infty$ as $n \rightarrow \infty$ for the same reason one can make λ_k arbitrarily small. However, one can still ask under what geometric conditions on M does there exists a function $R(\lambda)$ such that

$$(1.4) \quad N(\lambda; M) = \frac{\omega_d}{(2\pi)^d} \lambda^{\frac{d}{2}} + R(\lambda)$$

with $R(\lambda) = O(\lambda^\tau)$ independent of M , with $\tau < d/2$. The search for this type of uniform bound was a prominent feature in the above-mentioned papers [2, 4, 5, 13]. The presence of the boundary allowed them to derive a two-term Weyl type bound; closed manifolds do not exhibit this behaviour.

In [8, Theorem 6.2], Buser has obtained bounds on the eigenvalue λ_k of a closed manifold, valid when k was large enough in terms of the injectivity radius; see also [14, equation 1.2.5] where this result is reformulated in terms of the counting function. The following theorem states that we can find explicit bounds on the remainder in (1.4) depending on the injectivity radius.

Theorem 1.5 *There is $C > 0$ such that for all $\lambda \geq 2\pi$ and all flat tori of unit volume, we have that*

$$\left| N(\lambda; \mathbb{T}_\Gamma) - \frac{\omega_d}{(2\pi)^d} \lambda^{d/2} \right| \leq C \lambda^{\frac{d}{2} - \frac{d}{d+1}} \text{inj}(\Gamma)^{-\frac{2d}{d+1}}.$$

Moreover, for any flat Klein bottle $K(a, b) \in \mathcal{E}_0$,

$$\left| N(\lambda; K(a, b)) - \frac{1}{4\pi} \lambda \right| \leq C \lambda^{\frac{1}{3}} \operatorname{inj}(K(a, b))^{-\frac{2}{3}}.$$

We make the following remarks as to the naturality and sharpness of those results.

Remark 1.6 The remainder in the previous theorem is natural in the following sense. If we take the normalisation

$$N_0(\lambda) = \lambda^{-\frac{d}{2}} N(\lambda)$$

such that $N_0(\lambda)$ has a limit as $\lambda \rightarrow \infty$, then the remainder obtained in Theorem 1.5 is invariant under homothetic rescaling of the metric.

Remark 1.7 If $\operatorname{inj}(\mathbb{T}_\Gamma)$ is of the order of $\lambda^{-1/2}$, the remainder in Theorem 1.5 is of the order of the principal term. This can indeed happen. As part of the proof of Theorem 1.2, we will construct an explicit sequence of flat tori $\mathbb{T}_k \in \mathcal{M}_0$ such that

$$\operatorname{inj}(\mathbb{T}_k) = \frac{\lambda_{2k}(\mathbb{T}_k)^{-1/2}}{2\pi},$$

whose eigenvalue counting functions satisfy

$$\left| N(\lambda_{2k}(\mathbb{T}_k); \mathbb{T}_k) - \frac{\omega_d}{(2\pi)^d} \lambda_{2k}(\mathbb{T}_k)^{d/2} \right| \gg \lambda^{d/2}.$$

In fact, one will be able to compute explicitly

$$\left| N(\lambda_{2k}(\mathbb{T}_k); \mathbb{T}_k) - \frac{\omega_1}{(2\pi)^d} \lambda_{2k}(\mathbb{T}_k)^{d/2} \right| = 2d - 1$$

and $\omega_1 \neq \omega_d$.

This also implies that one cannot improve the order of error term in the spectral parameter without making it worse in terms of the injectivity radius, and vice versa.

1.3 Lattice Points Inside Domains

We translate the problems at hand in the language of lattice point counting. The spectrum of the Laplacian on a flat torus is given by

$$(1.5) \quad \sigma(\mathbb{T}_\Gamma) = \{4\pi^2 |\gamma^*|^2 : \gamma^* \in \Gamma^*\},$$

where Γ^* is the lattice dual to Γ defined by

$$\Gamma^* := \{\gamma^* \in \mathbb{R}^d : (\gamma^*, \Gamma) \subset \mathbb{Z}\}.$$

Similarly, the spectrum of the Laplacian on a flat Klein bottle is given in [7] to be

$$(1.6) \quad \sigma(K(a, b)) := \left\{ 4\pi^2 \left(\frac{m^2}{a^2} + \frac{n^2}{b^2} \right) : (m, n) \in \mathbb{Z} \times \mathbb{N}_0, (m, n) \neq (2\ell + 1, 0) \right\}.$$

A classical problem in the geometry of numbers consists in counting the number of points of an isotropically shrinking lattice $\Gamma_\lambda := \lambda^{-1}\Gamma$ inside a domain Ω containing the origin as $\lambda \rightarrow \infty$. This dates back to the Gauss circle problem and has been studied in great details for various type of domains over the years. Denote

$$|\Omega| = \operatorname{Vol}_d(\Omega) \quad \text{and} \quad |\Gamma| = \det(A_\Gamma),$$

where A_Γ is any matrix such that $A_\Gamma \mathbb{Z}^d = \Gamma$. In general, one aims for asymptotics of the form

$$(1.7) \quad N(\Omega; \Gamma_\lambda) := \#(\Omega \cap \Gamma_\lambda) = \frac{|\Omega|}{|\Gamma_\lambda|} + R(\lambda; \Omega; \Gamma),$$

where

$$(1.8) \quad R(\lambda; \Omega; \Gamma) = O(|\Gamma_\lambda|^{-\eta})$$

with $\eta < 1$. The implicit constant on the right-hand side of equation (1.8) depends on the geometry of Ω , the geometry of its boundary, and on Γ . In general, given non-compact families of lattices or domains, the implicit constant is not uniform, and therefore formula (1.7) cannot be used directly to find extremisers to $N(\Omega; \Gamma_\lambda)$ for large λ . Note that maximising this counting function does not makes sense, even while keeping the lattice determinant and the volume of the domain fixed. Indeed, for a fixed Ω containing the origin and ε small enough, the lattice $\varepsilon^{d-1}\mathbb{Z} \oplus \varepsilon^{-1}\mathbb{Z}^{d-1}$ has arbitrarily many points in Ω and determinant 1.

We formulate the results of the two previous sections in terms of lattices. From the fact that

$$\#\{\mathbb{Z}^d \cap A_\Gamma^{-1}(B_1)\} = \#\{A_\Gamma \mathbb{Z}^d \cap B_1\},$$

the following two questions are equivalent.

- What's the largest lattice determinant of a lattice with at least k points in B_1 ?
- What's the smallest area of an ellipsoid enclosing at least k points of the lattice \mathbb{Z}^d ?

Symmetry of ellipsoids or lattices with respect to the transformation $x \mapsto -x$ means that no generality is lost by asking these questions for only even (or odd) k . Let us order elements of any lattice as

$$\Gamma = \{\gamma_k : k \in \mathbb{N}_0\}$$

with $\gamma_0 = 0$ and $\gamma < \tilde{\gamma}$ if $|\gamma| < |\tilde{\gamma}|$, and if their norms are equal by lexicographic order. The scaling invariance of the problem is made explicit by studying maximisers to the functional

$$\tilde{\Lambda}_k(\Gamma) = |\Gamma|^{-1/d} |\gamma_k|.$$

We obtain the following restatement of Theorem 1.1 in terms of lattices.

Theorem 1.8 (Lattice version of Theorem 1.1) *For every $k \in \mathbb{N}$, there exists $\Gamma_k^* \in \mathcal{L}$ maximising $\tilde{\Lambda}_k$.*

Remark 1.9 The maximiser in the previous theorem is not unique, in particular if Γ is a maximiser, then $\mu\Gamma$ is also one. We will, depending on what is pertinent at the right moment, either normalise them by determinant or by $|\gamma_k|$. Note that even within \mathcal{L}_0 , uniqueness is not guaranteed.

We now study properties of the maximisers Γ_k^* . The degeneracy of a sequence Γ_k^* is given in terms of their *successive minima*, the lattice invariants $\mu_j(\Gamma)$ defined for $1 \leq j \leq d$ by

$$\mu_j(\Gamma) = \inf \left\{ \mu : \dim(\text{span}(\Gamma \cap B_\mu)) \geq j \right\}.$$

We prove the following restatement of Theorem 1.2.

Theorem 1.10 (Lattice version of Theorem 1.2) *Let $\{\Gamma_k^*\} \subset \mathcal{L}_0$ be a sequence of maximisers of $\tilde{\Lambda}_k$ normalised by $|\Gamma_k^*| = 1$, in dimension $d \leq 10$. Then the following hold.*

- (i) *The sequence Γ_k^* has no accumulation points in \mathcal{L}_0 .*
- (ii) *The successive minima of the sequence Γ_k^* satisfy the asymptotic bounds*

$$\mu_1(\Gamma_k^*) \gg k^{-1+\frac{1}{d}} \quad \text{and} \quad \mu_d(\Gamma_k^*) \gg k^{\frac{1}{d}}.$$

This will be proved thanks to the following restatement of Theorem 1.5 in terms of lattices.

Theorem 1.11 (Lattice version of Theorem 1.5) *There exists a constant C such that for all lattices with $|\Gamma| \leq 1$*

$$\left| N(B_1; \Gamma) - \frac{\omega_d}{|\Gamma|} \right| \leq C |\Gamma|^{-1} \mu_d(\Gamma)^{\frac{2d}{d+1}}.$$

1.4 Plan of the Paper and Sketch of the Proofs

We start in Section 2 by exposing general facts about lattices that will be used in the sequel. More specifically, we describe the relevant lattice invariants and state theorems of Minkowski and Banaszczyk that are important later, for ease of reference.

In Section 3, we prove Theorems 1.8 and 1.10. Inspired by a construction of Kao, Lai, and Osting [17] in dimension 2, we produce in Section 3.2 in any dimension a sequence of lattices Θ_{2k} such that

$$|\theta_{2k-1}| = |\theta_{2k}| = \left(\frac{2k}{\omega_1} \right)^{1/d}.$$

However, Theorem 1.11 implies that for any lattice Γ of unit determinant whose successive minima satisfy $\mu_d(\Gamma) = o(k^{1/d})$, we have that

$$|\gamma_{2k-1}| = |\gamma_{2k}| \leq \left(\frac{2k}{\omega_d} \right)^{1/d} (1 + o(1)),$$

where ω_d is the volume of the unit ball. One can see that while the sequence ω_d converges to 0 as $d \rightarrow \infty$, it is initially increasing. Indeed, for all $2 \leq d \leq 10$, we have that $\omega_d > \omega_1$.

In Section 4, we will show that the spectral theoretic versions of Theorems 1.1, 1.2, and 1.5 are implied by Theorems 1.8, 1.10, and 1.11 using Banaszczyk's transference Theorem 2.2 and Minkowski's successive minima Theorem 2.1.

In Section 5, we switch gears and describe Theorem 1.11 in terms of points of \mathbb{Z}^d sitting inside anisotropically expanding domains. These were studied by Yu. Kordyukov and A. Yakovlev in a series of papers [19–22], and we generalise their results and methods to our setting.

In Section 6, we prove the theorems about the number of points of a lattice sitting inside anisotropically expanding domains using the Poisson summation formula method. In the classical version of this problem, one uses global estimates on the Fourier transform of the indicator of a convex set to obtain bounds on the counting function of lattice points inside an expanding domain. It is, however, not possible

to make these kinds of computations uniformly when the expansion is anisotropic. The main idea, inspired by [21], is to only use Fourier transform estimates along the subspace where the expansion is the fastest and to use trivial L^∞ estimates in the orthogonal complement.

2 Some Facts about Lattices in \mathbb{R}^d

For most standard results on lattices, one can see [9]. The set of all full-rank lattices in \mathbb{R}^d can be realised as $\mathcal{L} = \text{GL}_d(\mathbb{R})/\text{GL}_d(\mathbb{Z})$, equipped with the quotient topology. A lattice $\Gamma \in \mathcal{L}$ is identified with its generator matrix A_Γ , the matrix such that $A_\Gamma \mathbb{Z}^d = \Gamma$. Every lattice determines uniquely a flat torus $\mathbb{T}_\Gamma = \mathbb{R}^d/\Gamma$.

Two relevant lattice invariants that are of interest in this paper are the determinant (or volume) and the successive minima. The determinant is defined as

$$|\Gamma| := \det A_\Gamma = \text{Vol}_d(\mathbb{T}_\Gamma).$$

By convention, we assign a volume of 1 to the trivial lattice. The successive minima $\mu_j(\Gamma)$ are defined for $1 \leq j \leq d$ as

$$\mu_j(\Gamma) := \inf\{\mu : \dim(\text{span}(\Gamma \cap B_\mu)) \geq j\}.$$

Note that μ_j is always attained, *i.e.*, there is always $\gamma \in \Gamma$ such that $\mu_j(\Gamma) = |\gamma|$. Furthermore, the first successive minimum gives the injectivity radius of the associated torus, *i.e.*, $\mu_1(\Gamma) = \text{inj}(\mathbb{T}_\Gamma)$.

The successive minima of a lattice and the determinant are related through a theorem of Minkowski.

Theorem 2.1 (Minkowski's successive minima theorem) *Let μ_1, \dots, μ_d be the successive minima of a lattice Γ . Then there exists constants $c, C > 0$ such that*

$$c|\Gamma| \leq \prod_{j=1}^d \mu_j \leq C|\Gamma|.$$

With any lattice Γ we associate the dual lattice

$$\Gamma^* = \{\gamma^* \in \mathbb{R}^d, (\gamma^*, \Gamma) \subset \mathbb{Z}\}.$$

The operation $*$ is a continuous involution on \mathcal{L} ; hence, a set $\mathcal{K} \subset \mathcal{L}$ is compact if and only if \mathcal{K}^* is. Let A_Γ be the generating matrix for Γ ; then $A_{\Gamma^*} = (A_\Gamma^*)^{-1}$. From this we infer that $|\Gamma^*| = |\Gamma|^{-1}$.

The following theorem from Banaszczyk [3] is also useful in the sequel and relates the successive minima of Γ to those of Γ^* .

Theorem 2.2 (Banaszczyk's transference theorem) *For any $1 \leq j \leq d$, the following inequalities hold between the successive minima of the lattices Γ and Γ^* :*

$$1 \leq \mu_j(\Gamma) \mu_{d-j+1}(\Gamma^*) \leq d.$$

The lattice invariants can be used to characterise compactness in \mathcal{L} , by Mahler's selection theorem [9, Theorems 5.3, 5.4 and Lemma 8.3]. This theorem states that a set $\mathcal{K} \subset \mathcal{L}$ is compact if and only if the determinant is bounded and the first

minimum μ_1 is bounded away from zero on \mathcal{K} . Equivalently, it is compact if and only if the determinant is bounded away from zero and μ_d is bounded on \mathcal{K} . Compactness in the moduli space of all flat tori is obtained by identifying a torus with its lattice.

Definition 2.3 A sequence of lattices $\{\Gamma_k\}$ is said to *degenerate* if either $|\Gamma_k| \rightarrow \infty$ or if $\mu_1(\Gamma_k) \rightarrow 0$. In other words, it degenerates if it is not contained in some compact set in \mathcal{L} .

We will be interested in the number of lattice points inside the unit ball B_1 ; denote this quantity by $N(\Gamma; 1)$. Denoting by $\mathbf{1}_S$ the indicator function of a set S , we have that

$$N(\Gamma; 1) = \sum_{\gamma \in \Gamma} \mathbf{1}_{B_1}(\gamma).$$

Finally, we say that a subspace $V \subset \mathbb{R}^d$ is a Γ -subspace if it is spanned by a subset of Γ . The set $\Gamma(V) := \Gamma \cap V$ is a lattice in V .

3 Optimal Lattices

In this section, we prove Theorems 1.8 and 1.10, assuming Theorem 1.11. Order elements of a lattice Γ with respect to their norms and by lexicographic order whenever the norms are equal. We write $\Gamma = \{\gamma_k : k \in \mathbb{N}_0\}$. We study sequences of lattices maximising the functionals

$$\tilde{\Lambda}_k(\Gamma) = |\Gamma|^{-1/d} |\gamma_k|.$$

Note that for any lattice Γ and $m \geq 1$ we have that $\tilde{\Lambda}_{2m-1}(\Gamma) = \tilde{\Lambda}_{2m}(\Gamma)$; we will therefore only consider maximisers for even k .

3.1 Proof of Theorem 1.8

Consider a maximising sequence $\{\Gamma_n\}$ for $\tilde{\Lambda}_k$. Without loss of generality, from the definition of $\tilde{\Lambda}_k$ we can suppose that $|\Gamma_n| = 1$ for all n . Suppose that $\mu_1(\Gamma_n) \rightarrow 0$. Then for some n , we have that $\mu_1(\Gamma_n) < 1/k$. Let $\gamma \in \Gamma_n$ be a lattice point realising $\mu_1(\Gamma_n)$. Then $1 > |k\gamma| > |\gamma_k|$. However, the k -th element of \mathbb{Z}^d has norm greater than 1, contradicting that $\{\Gamma_n\}$ was a maximising sequence. By Mahler's selection theorem, $\{\Gamma_n\}$ has a convergent subsequence, and by continuity of the norm and the determinant, it converges to a maximiser for $\tilde{\Lambda}_k$. ■

3.2 Lattices with Large $\tilde{\Lambda}_k$

In this section we study a specific sequence of lattices that we will use as a measuring stick for other sequences of lattices. Note that we make no claim of these lattices being the optimisers. Consider the lattices

$$\Theta_{2k} = k^{-1+\frac{1}{d}} \mathbb{Z} \oplus k^{\frac{1}{d}} \mathbb{Z}^{d-1}.$$

Then we have

$$|\theta_{2k-1}| = |\theta_{2k}| = k^{1/d} \quad \text{and} \quad |\Theta_{2k}| = 1.$$

In particular, we have that

$$\tilde{\Lambda}_{2k}(\Theta_{2k}) = k^{1/d},$$

which will be the quantity to beat. Observe that the sequence Θ_{2k} degenerates and that $\mu_d(\Theta_{2k}) = k^{1/d}$.

3.3 Proof of Theorem 1.10

Let $\{\Gamma_k\}$ be a sequence of lattices of unit volume such that $\mu_d(\Gamma_k) = o(k^{1/d})$. We will show that under such conditions, Γ_k cannot be a maximiser for $\tilde{\Lambda}_k$ infinitely often. This is done by showing that for large k and any fixed $t > 0$,

$$\#(B_{k^{1/d-t}} \cap \Gamma_{2k}) > 2k,$$

implying that

$$\tilde{\Lambda}_{2k}(\Gamma_{2k}) \leq k^{1/d} - t < \tilde{\Lambda}_{2k}(\Theta_{2k}).$$

We have that

$$\begin{aligned} \#(B_{k^{1/d-t}} \cap \Gamma_{2k}) &= \#(B_1 \cap \frac{1}{k^{1/d-t}} \Gamma_{2k}), \\ \mu_d\left(\frac{1}{k^{1/d-t}} \Gamma_{2k}\right) &= o(1), \\ \left|\frac{1}{k^{1/d-t}} \Gamma_{2k}\right| &= k^{-1}(1 - tk^{-1/d})^{-d}. \end{aligned}$$

Thus, we satisfy the hypotheses of Theorem 1.11 and therefore get

$$\#(B_{k^{1/d-t}} \cap \Gamma_{2k}) = \omega_d(1 - tk^{-1/d})^d k(1 + o(1)).$$

For $2 \leq d \leq 10$, we have that $\omega_d > \omega_1 = 2$. Hence, there is K such that for $k > K$,

$$\#(B_{k^{1/d-t}} \cap \Gamma_{2k}) > 2k,$$

proving that there is a finite number of maximisers in the sequence $\{\Gamma_k\}$. This implies that there is constant c such that any sequence of normalised maximisers respects $\mu_d(\Gamma_k) \geq ck^{1/d}$, also implying that the sequence degenerates.

For the lower bound on $\mu_1(\Gamma_k)$, any sequence Γ_k normalised by determinants such that $\mu_1(\Gamma_{2k}) < k^{-1+1/d}$ has that

$$\tilde{\Lambda}_{2k}(\Gamma_{2k}) \leq k\mu_1(\Gamma_{2k}) < \tilde{\Lambda}_{2k}(\Theta_{2k});$$

hence, this is not a sequence of maximisers. ■

4 From Lattices to Tori

In this section we prove the spectral theoretic versions of Theorems 1.1, 1.2, and 1.5, as well as Theorem 1.4. For any lattice Γ we denote by γ_k^* the k -th ordered element of the dual lattice Γ^* . Since $\lambda_k(\mathbb{T}_\Gamma) = 4\pi^2|\gamma_k^*|^2$ and $\text{Vol}(\mathbb{T}_\Gamma) = |\Gamma^*|^{-1}$, we have that

$$\Lambda_k(\mathbb{T}_\Gamma) = (2\pi\tilde{\Lambda}_k(\Gamma^*))^2.$$

Since these quantities are positive, the problem of maximising Λ_k on flat tori is the same as the problem of maximising $\tilde{\Lambda}_k$ on the dual lattices of those tori.

4.1 Proof of Theorem 1.1

By Theorem 1.8, there exists a lattice Γ_k^* maximising $\tilde{\Lambda}_k$. The torus with lattice $\Gamma = (\Gamma_k^*)^*$ is therefore a maximiser for Λ_k .

For flat Klein bottles, we have from equation (1.6) that the eigenvalues of $K(a, b)$ are continuous in the parameters a and b . Normalising by $ab = 2$, it is easy to see that for any k , $\lambda_k(K(a, b))$ goes to 0 when either a or b goes to zero. Hence, for any fixed k we can restrict ourselves to a compact subset of the parameters a, b , and the maximiser exists. ■

4.2 Proof of Theorem 1.2

Denote by Γ_k^* a sequence of optimal lattices with unit determinant for Λ_k and denote by \mathbb{T}_k^* the corresponding optimal torus $T_k^* = \mathbb{R}^d / (\Gamma_k^*)^*$. Since compactness of a set $\mathcal{K} \subset \mathcal{L}_0$ is equivalent to compactness of the set of duals \mathcal{K}^* , we have that the sequence of optimal tori degenerates.

We now turn to the geometric constraints. Recall that $\text{inj}(\mathbb{T}_k^*) = \mu_1((\Gamma_k^*)^*)$. By Banaszczyk's transference theorem, we have that

$$\mu_1((\Gamma_k^*)^*) \leq \frac{d}{\mu_d(\Gamma_k^*)}.$$

Hence, from the lower bound for $\mu_d(\Gamma_k^*)$ in Theorem 1.10, we have that

$$\text{inj}(\mathbb{T}_k^*) = \mu_1((\Gamma_k^*)^*) \ll k^{-1/d}.$$

On the other hand, by Minkowski's successive minima theorem, there is a constant C such that

$$\mu_d(\Gamma_k^*) \leq C\mu_1(\Gamma_k^*)^{1-d} \leq k^{\frac{(1-d)^2}{d}}.$$

Once again, Banaszczyk's transference theorem yields

$$\text{inj}(\mathbb{T}_k^*) \geq k^{-\frac{(1-d)^2}{d}},$$

finishing the proof. ■

4.3 Proof of Theorem 1.4

For flat Klein bottles, observe that the injectivity radius of $K(a, b)$ is given by

$$\text{inj}(K(a, b)) = \min(a, b/2).$$

Let $\Gamma(a, b)$ be the lattice defined by

$$\Gamma(a, b) := \frac{2\pi}{a}\mathbb{Z} \oplus \frac{2\pi}{b}\mathbb{Z}.$$

It is not hard to see that $\Gamma(a, b)$ has the property

$$N(\lambda; K(a, b)) = \frac{1}{2} \#(\Gamma(a, b) \cap B_{\sqrt{\lambda}}) + O(1).$$

Indeed, let $\Xi(a, b)$ be the set

$$\Xi(a, b) := \left(\frac{2\pi}{a}\mathbb{Z} \oplus \frac{2\pi}{b}N_0 \right) \setminus \left\{ \frac{2\pi}{a}(2\ell + 1, 0) : \ell \in \mathbb{Z} \right\}.$$

Then the spectrum of $K(a, b)$ is the same as the square of the norm of elements of $\Xi(a, b)$. However, it is easy to see that if we take the union of $\Xi(a, b)$ and $-\Xi(a, b)$, we recover $\Gamma(a, b)$ except for points of the form $(2(2\ell + 1)\pi/a, 0)$, but the elements of the form $(4\pi\ell/a, 0)$ are added twice. Hence, we have that

$$\left| \#(\Gamma(a, b) \cap B_{\sqrt{\lambda}}) - \#(\Xi(a, b) \cap B_{\sqrt{\lambda}}) - \#(-\Xi(a, b) \cap B_{\sqrt{\lambda}}) \right| \leq 3.$$

Now, for rectangular lattices, we have that

$$\mu_1(\Gamma(a, b)) = 2\pi \min(a^{-1}, b^{-1}) \quad \text{and} \quad \mu_2(\Gamma(a, b)) = 2\pi \max(a^{-1}, b^{-1}).$$

The rest of the analysis is performed exactly in the same way as for flat tori. ■

4.4 Proof of Theorem 1.5

Let \mathbb{T}_Γ be any flat torus of unit volume. Observe that, by Banaszczyk's transference theorem, we have that $\text{inj}(\mathbb{T}_\Gamma) \asymp \mu_d(\Gamma^*)^{-1}$. We have from equation (1.5) that

$$N(\lambda; T_\Gamma) = \#(2\pi\lambda^{-1/2}\Gamma^* \cap B_1).$$

Denote by Γ_λ^* the rescaled lattice $2\pi\lambda^{-1/2}\Gamma^*$. By Theorem 1.11, we have that

$$(4.1) \quad N(\lambda; T_\Gamma) = \frac{\omega_d}{|\Gamma_\lambda^*|} + O(|\Gamma_\lambda^*|^{-1} \mu_d(\Gamma_\lambda^*)^{\frac{2d}{d+1}}).$$

We have that

$$\mu_d(\Gamma_\lambda^*) = 2\pi\lambda^{-1/2} \mu_d(\Gamma^*) \quad \text{and} \quad |\Gamma_\lambda^*| = \frac{(2\pi)^d}{\lambda^{d/2}}.$$

Inserting those values into equation (4.1) yields the desired asymptotic in Theorem 1.5. ■

5 Anisotropically Expanding Domains

We now ground the statement of Theorem 1.11 in terms of the counting of lattice points sitting inside anisotropically expanding domains developed by Yu. Kordyukov and A. Yakovlev in [19–22]. Consider the decomposition of \mathbb{R}^d as

$$\mathbb{R}^d := E := \bigoplus_{j=1}^d V_j.$$

We will use E to refer to a specific decomposition for \mathbb{R}^d . For $\varepsilon = (\varepsilon_1, \dots, \varepsilon_d)$ consider the linear transformation T_ε given by

$$T_\varepsilon = \sum_{j=1}^d \varepsilon_j^{-1} \mathbf{x}_j,$$

with $\mathbf{x}_j \in V_j$. Without loss of generality, we suppose that $\varepsilon_1 \leq \dots \leq \varepsilon_d$. We denote the set of all such transformations by \mathcal{T}_E and the union of all such transformations over decompositions E by \mathcal{T} , and we say that T_ε is anisotropic whenever not all ε_j are equal.

For Ω a bounded subset of Euclidean space and $\Gamma \in \mathcal{L}$, let

$$n_\varepsilon(\Omega; \Gamma; \mathbf{y}) := \#(\Gamma \cap (T_\varepsilon \Omega + \mathbf{y})) = \#(T_\varepsilon^{-1}(\Gamma - \mathbf{y}) \cap \Omega).$$

Kordyukov and Yakovlev have studied asymptotics for n_ε in the specific case where a subspace V of \mathbb{R}^d is fixed, and Ω is stretched along its orthogonal complement. In our notation, this corresponds to $E = V_1 \oplus V_2$ with $\varepsilon_1 \rightarrow 0$ and $\varepsilon_2 = 1$.

In our case, the expansion is happening at different rates along different subspaces. We split the remainder of this section in three parts. First, we describe asymptotics for n_ε in terms of the decomposition E with an explicit dependence on the ε_j . Then we show that from the perspective of the counting function, we can describe any lattice using the transformations T_ε . Finally, we derive Theorem 1.11 from Theorem 5.1.

5.1 Lattice Points Inside Anisotropically Expanding Domains

We start by fixing some notation. Denote $|\varepsilon| = \det T_\varepsilon = \prod_j \varepsilon_j$; asymptotic results will be given in terms of $|\varepsilon|$ as it goes to zero, and in terms of how fast the ε_j 's goes to zero in relation to $|\varepsilon|$. Let us split the decomposition E into three parts. We first write

$$V_0 = \bigoplus_{j: \varepsilon_j = 0} V_j,$$

and let W be the maximal Γ^* -subspace in V_0 , and write $d_W = \dim(W)$ and $\Gamma_W = \Gamma^*(V)^*$. We further decompose E as

$$E = V \oplus V' \oplus W$$

in such a way that $\Gamma^* \cap V' = \{0\}$. We set $d_V = \dim(V)$ and $d_{V'}$ analogously. Finally, denote by

$$\delta_V = \|T_\varepsilon^{-1}\|_V$$

the norm of $\mathbb{T}_\varepsilon^{-1}$ restricted to V .¹ We obtain the following theorem.

Theorem 5.1 Suppose that Ω is a bounded open subset of \mathbb{R}^d with smooth boundary such that for all $\gamma \in \Gamma_W$, $\Omega \cap (\gamma + W^\perp)$ is strictly convex. Then,

$$(5.1) \quad n_\varepsilon(\Omega; \Gamma; \mathbf{y}) = \frac{|\varepsilon|^{-1} |\Gamma_W|}{|\Gamma|} \sum_{\gamma \in \Gamma_W} \text{Vol}(\Omega \cap (\gamma + W^\perp)) + \left(|\varepsilon|^{-1} \delta_V^{\frac{2d_V}{1+d_V+2d_{V'}}} \right),$$

with the implicit constant only dependant on Γ_W , V' and Ω .

Remark 5.2 If $W = \Gamma_W = \{0\}$, the condition on Ω becomes strict convexity, and the asymptotic formula becomes

$$n_\varepsilon(\Omega; \Gamma; \mathbf{y}) = \frac{|\Omega|}{|\Gamma|} |\varepsilon|^{-1} + \left(|\varepsilon|^{-1} \delta_V^{\frac{2d_V}{1+d_V+2d_{V'}}} \right)$$

with the implicit constants dependant on V' and Ω .

Furthermore, from [22, Section 3.2], this is the only case we need to prove. Indeed, they show that there are \mathbf{y}_γ such that

$$n_\varepsilon(\Omega; \Gamma; \mathbf{y}) = \sum_{\gamma \in \Gamma_W} n_\varepsilon(\Omega \cap (\gamma + W^\perp); \Gamma(W^\perp); \mathbf{y}_\gamma).$$

¹If $V = V_{j_1} \oplus \dots \oplus V_{j_m}$ with the indices in increasing order, this is equal to ε_{j_m} .

They then show in [22, Lemma 3.3] that $\Gamma(W^\perp)^* \cap (V' \cap W^\perp) = \{0\}$. Since the sum in equation (5.1) is finite, we obtain the desired result by applying Theorem 5.1 with $W = \{0\}$ term by term.

5.2 From \mathcal{T} to Lattices

We start by showing that we can restrict ourselves to lattices of the form $T_\varepsilon^{-1}\mathbb{Z}^d$ in our investigation of Theorem 1.11.

Lemma 5.3 *For every $\Gamma \in \mathcal{L}$, there exists a decomposition*

$$\mathbb{R}^d = E = \bigoplus_{j=1}^d V_j$$

and $T_\varepsilon \in \mathcal{T}_E$ such that

$$(5.2) \quad N(\Gamma; B_1) = n_\varepsilon(B_1; \mathbb{Z}^d; 0).$$

For every $T_\varepsilon \in \mathcal{T}$, there exists Γ , such that equation (5.2) holds.

Proof Let $A_\Gamma \in \text{GL}_d(\mathbb{R})$ be such that $A_\Gamma \mathbb{Z}^d = \Gamma$. Then

$$\sum_{\gamma \in \Gamma} \mathbf{1}_{B_1}(\gamma) = \sum_{\gamma \in \Gamma} \mathbf{1}_{A_\Gamma(B_1)}(A_\Gamma^{-1}\gamma) = \sum_{\mathbf{n} \in \mathbb{Z}^d} \mathbf{1}_{A_\Gamma(B_1)}(\mathbf{n})$$

Observe now that since $B_1 = \{\mathbf{x} \in \mathbb{R}^d : \mathbf{x}^* \mathbf{x} \leq 1\}$,

$$A_\Gamma(B_1) = \{\mathbf{x} \in \mathbb{R}^d : \mathbf{x}^* (A_\Gamma^*)^{-1} A_\Gamma^{-1} \mathbf{x} \leq 1\}.$$

Since $(A_\Gamma A_\Gamma^*)^{-1}$ is symmetric definite positive, it can be diagonalised as

$$(A_\Gamma A_\Gamma^*)^{-1} = U^* D^{1/2} D^{1/2} U$$

with U orthogonal. Let $\varepsilon = \text{diag}(D^{1/2})$ and V_j be eigenspaces of $(A_\Gamma A_\Gamma^*)^{-1}$. Since $N(\Gamma; B_1)$ is invariant under orthogonal transformations of Γ , we have that

$$N(\Gamma; 1) = N(UT; 1) = \#\{A_\Gamma^{-1}UT \cap A_\Gamma^{-1}B_1\} = n_\varepsilon(B_1).$$

On the other hand, this process can be inverted. Given T_ε , we take Γ to be the lattice with generating matrix T_ε^{-1} . ■

The previous lemma allows us to consider only the lattices of the form $\Gamma = T_\varepsilon^{-1}\mathbb{Z}^d$. The following lemma relates the lattice invariants to the associated transformation T_ε .

Lemma 5.4 *Let Γ be a lattice in \mathcal{L} . Then for any $T_\varepsilon \in \mathcal{T}$ such that $\Gamma = T_\varepsilon^{-1}U\mathbb{Z}^d$ for some orthogonal transformation U , we have that*

$$|\Gamma| = \det(T_\varepsilon^{-1}) = |\varepsilon|.$$

and that the following bounds hold for the successive minima $\mu_1(\Gamma)$ and $\mu_d(\Gamma)$:

$$\varepsilon_1 \leq \mu_1(\Gamma) \leq \mu_d(\Gamma) \leq \varepsilon_d.$$

Furthermore, one can choose T_ε such that $\Gamma = T_\varepsilon^{-1}U\mathbb{Z}^d$ and

$$\mu_1(\Gamma) \leq \frac{d^{5/2}}{2} \varepsilon_1 \quad \text{and} \quad \mu_d(\Gamma) \geq \frac{2}{d^{3/2}} \varepsilon_d.$$

Proof Without loss of generality, since the determinant and successive minima are invariant under orthogonal transformations, we suppose that $U = I$. The assertion on determinants holds by definition. Let \mathbf{n} be any non-zero element of \mathbb{Z}^d , and write $\mathbf{n} = \mathbf{n}_1 + \cdots + \mathbf{n}_d$ with $\mathbf{n}_j \in V_j$. Then

$$|T_\epsilon^{-1}\mathbf{n}|^2 = \sum_{j=1}^d \epsilon_j^2 |\mathbf{n}_j|^2 \geq \epsilon_1^2 \sum_{j=1}^d |\mathbf{n}_j|^2 = \epsilon_1^2 |\mathbf{n}|^2.$$

Since $\mathbf{n} \neq 0$, we have that $\mu_1(\Gamma) \geq \epsilon_1$. For the upper bound on μ_d , observe that any T_ϵ sends bases of \mathbb{R}^d to bases of \mathbb{R}^d . As such, from the definition of μ_d , we have that

$$\mu_d(\Gamma) \leq \sup_{\mathbf{n} \in \mathbb{Z}^d \setminus \{0\}} \frac{|T_\epsilon^{-1}\mathbf{n}|}{|\mathbf{n}|} \leq \epsilon_d.$$

We now obtain the lower bound on μ_d for a specific T_ϵ . There is a basis of Γ whose elements all have norm smaller than $d\mu_d(\Gamma)/2$ [9, Lemma V.8]. Let $T_{\epsilon, \Gamma}^{-1}$ be the square root of the diagonalised Gram matrix G_Γ associated with that basis. By Cauchy–Schwartz, the entries of the Gram matrices G_Γ all bounded by $\frac{d^2 \mu_d(\Gamma)^2}{4}$. Let $\nu_d(G_\Gamma)$ be the largest eigenvalue of G_Γ . It satisfies the bound

$$\nu_d(G_\Gamma) \leq \sqrt{\text{tr}(G_\Gamma^* G_\Gamma)} \leq \frac{d^3 \mu_d(\Gamma)^2}{4}.$$

Note that the eigenvalues of G_Γ are the same as those of $T_{\epsilon, \Gamma}^{-2}$, hence we have that

$$\epsilon_d \leq \frac{d^{3/2}}{2} \mu_d(\Gamma),$$

yielding the desired result. For the upper bound on μ_1 , observe that a generating matrix for Γ^* is T_ϵ . Hence, by the previous argument we have that

$$\mu_d(\Gamma^*) \geq \frac{2}{d^{3/2}} \epsilon_1^{-1}.$$

From Banaszczyk's transference theorem, we can then infer that

$$\mu_1(\Gamma) \leq d\mu_d(\Gamma^*)^{-1} \leq \frac{d^{5/2}}{2} \epsilon_1,$$

finishing the proof. ■

5.3 Proof of Theorem 1.11

Given lattice Γ with $|\Gamma| < 1$, we know from Lemma 5.3 that one can find a decomposition E of \mathbb{R}^d and a transformation $T_\epsilon \in \mathcal{T}_E$ such that

$$N(\Gamma; B_1) = n_\epsilon(B_1; \mathbb{Z}^d; 0).$$

Furthermore, from Lemma 5.4 we get that one can choose T_ϵ in such a way that

$$\epsilon_d \leq \frac{d^{3/2}}{2} \mu_d(\Gamma).$$

We therefore satisfy the hypotheses of Proposition 5.1 with $V = E$, $V' = W = \{0\}$, and $\delta_E = \varepsilon_d$, and we deduce that

$$N(\Gamma; B_1) = \omega_d |\varepsilon|^{-1} + O(\varepsilon^{-1} \varepsilon_d^{\frac{2d}{1+d}}).$$

Plugging in $|\Gamma| = |\varepsilon|$ and $\mu_d \asymp \varepsilon_d$ gives the desired asymptotics. ■

6 Asymptotic Estimates

In this section, we prove Theorem 5.1 using the Poisson summation formula. We follow the structure set out by the author and Parnowski [25]. The first thing we have to do is a mollification of $\mathbf{1}_\Omega$ so that it is smooth enough for the Poisson summation formula to be used, and we will get estimates from above and below using the mollified functions. In the second part, we obtain estimates on partial Fourier transforms of such functions. Finally, we use the Poisson summation formula to obtain asymptotics for the counting function.

6.1 Mollification

Let $\rho \in C_c^\infty(\mathbb{R}^d)$ be a non-negative bump function supported in the unit ball and such that

$$\int_{\mathbb{R}^d} \rho(\mathbf{x}) \, d\mathbf{x} = 1.$$

We also let $\mathbf{h} = (h_1, \dots, h_d)$ be a set of parameters to be fixed later, and we set

$$\rho_{\mathbf{h}}(\mathbf{x}) = \frac{1}{h_1 \cdots h_d} \rho(T_{\mathbf{h}}(\mathbf{x})).$$

Note that $\rho_{\mathbf{h}}$ is supported in the ellipsoid

$$E_{\mathbf{h}} = \{\mathbf{x} \in V : \|T_{\mathbf{h}}\mathbf{x}\| < 1\}.$$

For any function $f : \mathbb{R}^d \rightarrow \mathbb{R}$, let $f^{(\mathbf{h})}$ be the mollification of f by $\rho_{\mathbf{h}}$, that is,

$$f^{(\mathbf{h})}(\mathbf{x}) = [f * \rho_{\mathbf{h}}](\mathbf{x}) = \int_{\mathbb{R}^d} f(\mathbf{x} - \mathbf{y}) \rho_{\mathbf{h}}(\mathbf{y}) \, d\mathbf{y}.$$

Let us now approximate $\mathbf{1}_\Omega$ by smooth functions. For any set B , define the sets

$$B_{\mathbf{h}} = \bigcup_{\mathbf{x} \in B} (\mathbf{x} + E_{\mathbf{h}}) \quad \text{and} \quad B_{-\mathbf{h}} = \mathbb{R}^d \setminus (\mathbb{R}^d \setminus B)_{\mathbf{h}}.$$

The following lemma will be needed regarding these sets.

Lemma 6.1 *Let $\varepsilon \mathbf{h} = (\varepsilon_1 h_1, \dots, \varepsilon_d h_d)$ and $B \subset V$. Then*

$$T_{\varepsilon}(B)_{\pm \mathbf{h}} = T_{\varepsilon}(B_{\pm \varepsilon \mathbf{h}}).$$

Proof This follows simply from linearity of T_{ε} and the fact that $T_{\varepsilon} E_{\mathbf{h}} = E_{\varepsilon \mathbf{h}}$. ■

We now prove that $\mathbf{1}^{(\mathbf{h})}$ provides a good approximation to $\mathbf{1}$.

Lemma 6.2 *Let $\Omega \subset \mathbb{R}^d$ and $\mathbf{x} \in \mathbb{R}^d$. Then*

$$(6.1) \quad \mathbf{1}_{T_{\varepsilon}(\Omega)_{-\mathbf{h}}}^{(\mathbf{h})}(\mathbf{x}) \leq \mathbf{1}_{T_{\varepsilon}(\Omega)}(\mathbf{x}) \leq \mathbf{1}_{T_{\varepsilon}(\Omega)_{\mathbf{h}}}^{(\mathbf{h})}(\mathbf{x}).$$

Proof For any set B , we have that $0 \leq \mathbf{1}_B^{(\mathbf{h})} \leq 1$. Hence, to show the rightmost inequality in (6.1) it suffices to show that for any $\mathbf{x} \in T_\varepsilon(\Omega)$, we have that $\mathbf{1}_{T_\varepsilon(\Omega)_\mathbf{h}}^{(\mathbf{h})}(\mathbf{x}) = 1$. By definition $\mathbf{x} + E_\mathbf{h} \subset T_\varepsilon(\Omega)_\mathbf{h}$, hence

$$\mathbf{1}_{(T_\varepsilon(\Omega)_\mathbf{h})}^{(\mathbf{h})}(\mathbf{x}) = \int_{E_\mathbf{h}} \rho_\mathbf{h}(\mathbf{y}) \, d\mathbf{y} = 1.$$

To prove the leftmost inequality in (6.1), it suffices to show that for any $\mathbf{x} \in E \setminus T_\varepsilon(\Omega)$, we have that $\mathbf{1}_{(T_\varepsilon(\Omega))_\mathbf{h}}^{(\mathbf{h})}(\mathbf{x}) = 0$. We have that

$$\mathbf{x} + E_\mathbf{h} \subset (\mathbb{R}^d \setminus T_\varepsilon(\Omega))_\mathbf{h},$$

and $\mathbf{1}_{(T_\varepsilon(\Omega))_\mathbf{h}}$ is supported in the complement of that set. Hence,

$$\mathbf{1}_{(T_\varepsilon(\Omega))_\mathbf{h}}^{(\mathbf{h})}(\mathbf{x}) = \int_{E_\mathbf{h}} \mathbf{1}_{T_\varepsilon(\Omega)_\mathbf{h}}(\mathbf{x} - \mathbf{y}) \rho_\mathbf{h}(\mathbf{y}) \, d\mathbf{y} = 0,$$

finishing the proof. ■

The following corollary follows directly from the previous lemma.

Corollary 6.3 *Defining*

$$n_\varepsilon^\pm(\Omega) = \sum_{\gamma \in \Gamma} \mathbf{1}_{T_\varepsilon(\Omega)_{\pm \mathbf{h}}}^{(\mathbf{h})}(\gamma),$$

the inequalities $n_\varepsilon^-(\Omega) \leq n_\varepsilon(\Omega) \leq n_\varepsilon^+(\Omega)$ hold for all ε .

6.2 Fourier Transform Estimates

Let V be a subspace of \mathbb{R}^d and write $\mathbf{x} = \mathbf{x}_V + \mathbf{x}'$ for any $\mathbf{x} \in \mathbb{R}^d$. We define the V -Fourier transform of a sufficiently rapidly decaying function f as

$$[\mathcal{F}_V f](\xi_V, \mathbf{x}') = \int_V e^{-2\pi i \mathbf{x}_V \cdot \xi_V} f(\mathbf{x}_V, \mathbf{x}') \, d\mathbf{x}_V.$$

When $V = \mathbb{R}^d$, we will write $[\mathcal{F}f] := [\mathcal{F}_{\mathbb{R}^d} f]$. We obtain estimates for the decay of $[\mathcal{F}f](\mathbf{x})$ in terms of $[\mathcal{F}_V f]$. Observe that

$$\begin{aligned} (6.2) \quad |[\mathcal{F}f](\xi)| &= \left| \int_{\mathbb{R}^d} e^{-2\pi i \mathbf{x} \cdot \xi} f(\mathbf{x}) \, d\mathbf{x} \right| \\ &= \left\{ \int_{V^\perp} e^{-2\pi i \mathbf{x}' \cdot \xi'} \int_V e^{-2\pi i \mathbf{x}_V \cdot \xi_V} f(\mathbf{x}) \, d\mathbf{x}_V \, d\mathbf{x}' \right\} \\ &= \left\{ \int_{V^\perp} e^{-2\pi i \mathbf{x}' \cdot \xi'} [\mathcal{F}_V f](\xi_V, \mathbf{x}') \, d\mathbf{x}' \right\} \\ &\leq \int_{V^\perp} |[\mathcal{F}_V f](\xi_V, \mathbf{x}')| \, d\mathbf{x}'. \end{aligned}$$

From this we get the following lemma.

Lemma 6.4 *Let Ω be a bounded domain, and let V be a subspace of dimension d_V such that the intersection $\Omega \cap V$ is strictly convex. Then*

$$[\mathcal{F}\mathbf{1}_\Omega](\xi) = O(|\xi_V|^{-\frac{d_V+1}{2}}).$$

Proof Standard results about the Fourier transform of the indicator of a strictly convex set (see, e.g., [16, Theorem 2.29]) tell us that

$$[\mathcal{F}_V \mathbf{1}_\Omega](\xi_V, \mathbf{x}') = O_{\mathbf{x}'}(|\xi_V|^{-\frac{d_V+1}{2}}).$$

From equation (6.2), we have that

$$|[\mathcal{F} \mathbf{1}]_\Omega(\xi)| \leq \int_{V^\perp} |[\mathcal{F}_V \mathbf{1}]_\Omega(\xi_V, \mathbf{x}')| d\mathbf{x}'.$$

Since $[\mathcal{F}_V \mathbf{1}_\Omega](\xi_V, \mathbf{x}')$ is compactly supported in \mathbf{x}' , we obtain the desired result, finishing the proof. ■

6.3 Poisson Summation Formula

Let us apply the Poisson summation formula to the smoothed sums $n_\varepsilon^\pm(\Omega; \Gamma; \mathbf{y})$. Denote $\Gamma' = \Gamma^* \setminus \{0\}$ to obtain

$$\begin{aligned} (6.3) \quad n_\varepsilon^\pm(\Omega, \mathbf{y}; 0) &= \sum_{\gamma \in \Gamma} \mathbf{1}_{T_\varepsilon(\Omega)_{\pm \mathbf{h}} + \mathbf{y}}^{(\mathbf{h})}(\gamma) = \frac{1}{|\Gamma|} \sum_{\gamma^* \in \Gamma^*} [\mathbf{1}_{T_\varepsilon(\Omega)_{\pm \mathbf{h}} + \mathbf{y}}^{(\mathbf{h})}](\gamma^*); \\ &= \frac{1}{|\Gamma|} [\mathbf{1}_{T_\varepsilon(\Omega)_{\pm \mathbf{h}} + \mathbf{y}}^{(\mathbf{h})}](0) + \Sigma(\varepsilon, \mathbf{h}, \mathbf{y}). \end{aligned}$$

Observe that

$$(6.4) \quad [\mathbf{1}_{T_\varepsilon(\Omega)_{\pm \mathbf{h}} + \mathbf{y}}^{(\mathbf{h})}](\xi) = e^{i\mathbf{y} \cdot \xi} [\mathbf{1}_{T_\varepsilon(\Omega)_{\pm \mathbf{h}}}^{(\mathbf{h})}](\xi) [\rho_{\mathbf{h}}](\xi - \mathbf{y}).$$

Since we will only find bounds using the absolute values of the terms in the previous equation, and since

$$[\mathcal{F} \mathbf{1}_{T_\varepsilon(\Omega)_{\pm \mathbf{h}} + \mathbf{y}}^{(\mathbf{h})}](0) = [\mathbf{1}_{T_\varepsilon(\Omega)_{\pm \mathbf{h}}}^{(\mathbf{h})}](0),$$

we suppose without loss of generality that $\mathbf{y} = 0$.

We first turn our attention to $[\mathbf{1}_{T_\varepsilon(\Omega)_{\pm \mathbf{h}}}^{(\mathbf{h})}](0)$. Using properties of the Fourier transform and Lemma 6.1, we have that

$$[\mathbf{1}_{T_\varepsilon(\Omega)_{\pm \mathbf{h}}}^{(\mathbf{h})}](\xi) = |\varepsilon|^{-1} [\mathbf{1}_{\Omega_{\pm \mathbf{h}}}](T_\varepsilon(\xi))$$

and from equation (6.1) that

$$(6.5) \quad [\rho_{\mathbf{h}}](\xi) = [\rho](T_{\mathbf{h}}^{-1}(\xi)).$$

Hence, the first term in equation (6.3) is given by

$$[\mathbf{1}_{T_\varepsilon(\Omega)_{\pm \mathbf{h}}}^{(\mathbf{h})}](0) = \frac{|\varepsilon|^{-1}}{|\Gamma|} [\mathcal{F} \mathbf{1}_{\Omega_{\pm \mathbf{h}}}](T_\varepsilon(0)) [\mathcal{F} \rho](T_{\mathbf{h}}^{-1}(0)) = \frac{|\varepsilon|^{-1}}{|\Gamma|} \text{Vol}(\Omega_{\pm \mathbf{h}})$$

As long as all the $\varepsilon_j h_j$ remain bounded, we have that there exists a constant C such that

$$\text{Vol}(\Omega_{\varepsilon \mathbf{h}} \setminus \Omega) \leq C \left(\sum_{j=1}^d \varepsilon_j h_j \right), \quad \text{and} \quad \text{Vol}(\Omega \setminus \Omega_{-\varepsilon \mathbf{h}}) \leq C \left(\sum_{j=1}^d \varepsilon_j h_j \right),$$

hence, writing $\Omega_{\varepsilon \mathbf{h}} = \Omega \cup (\Omega_{\varepsilon \mathbf{h}} \setminus \Omega)$ and $\Omega = \Omega_{-\varepsilon \mathbf{h}} \cup (\Omega \setminus \Omega_{-\varepsilon \mathbf{h}})$, we have

$$[\mathbf{1}_{T_\varepsilon(\Omega)_{\pm \mathbf{h}}}^{(\mathbf{h})}](0) = \frac{|\varepsilon|^{-1}}{|\Gamma|} \text{Vol}(\Omega) + \left(|\varepsilon|^{-1} \sum_{k=1}^d \varepsilon_k h_k \right).$$

Let us now study $\Sigma(\boldsymbol{\varepsilon}, \mathbf{h}, 0)$ in equation (6.3). Using equations (6.4) and (6.5) we deduce that

$$\Sigma(\boldsymbol{\varepsilon}, \mathbf{h}, 0) = |\boldsymbol{\varepsilon}|^{-1} \sum_{\gamma^* \in \Gamma'} [\mathcal{P}\mathbf{I}_{\Omega_{\text{eh}}}] (T_{\boldsymbol{\varepsilon}}(\gamma^*)) [\mathcal{P}\rho] (T_{\mathbf{h}}^{-1}(\gamma^*)).$$

We have that $[\mathcal{P}\rho_{\mathbf{h}}]$ is a Schwartz function; i.e., for any N

$$[\mathcal{P}\rho_{\mathbf{h}}](\boldsymbol{\xi}) = O((1 + |\boldsymbol{\xi}|)^{-N}).$$

Hence, we have that

$$\Sigma(\boldsymbol{\varepsilon}, \mathbf{h}, 0) = |\boldsymbol{\varepsilon}|^{-1} \sum_{\gamma^* \in \Gamma'} \frac{[\mathcal{P}\mathbf{I}_{\Omega_{\text{eh}}}] (T_{\boldsymbol{\varepsilon}}(\gamma^*))}{(1 + (h_1 \mathbf{x}_1)^N + \dots + (h_d \mathbf{x}_d)^N)}.$$

6.4 Proof of Proposition 5.1

All that remains is to bound $\Sigma(\boldsymbol{\varepsilon}, \mathbf{h}, 0)$ and to balance it with the error term coming from the Fourier transform evaluated at 0. Let

$$j_V = \operatorname{argmax}(j : V_j \subset V),$$

and set $h_V = h_{j_V}$. Choose

$$h_k = \delta_V^{\frac{2d_V}{1+d_V+2d_{V'}}} \varepsilon_k^{-1},$$

hence,

$$|\mathbf{h}| = \delta_V^{\frac{2d_V(d_V+d_{V'})}{1+d_V+2d_{V'}}} |\boldsymbol{\varepsilon}|^{-1}.$$

For all $\gamma^* \in \Gamma'$, we have that $\gamma_V^* \neq 0$. From Lemma 6.4, we obtain the bound

$$\begin{aligned} \Sigma(\boldsymbol{\varepsilon}, \mathbf{h}, 0) &\ll |\boldsymbol{\varepsilon}|^{-1} \sum_{\gamma^* \in \Gamma'} \frac{\delta_V^{\frac{d_V+1}{2}}}{|\gamma_V^*|^{\frac{m+1}{2}} (1 + (h_1 \gamma_1^*)^N + \dots + (h_d \gamma_d^*)^N)} \\ &\ll \delta_V^{\frac{d_V+1}{2}} |\boldsymbol{\varepsilon}|^{-1} \int_{\mathbb{R}^d} \frac{|\mathbf{x}_V|^{-\frac{d_V+1}{2}}}{(1 + (h_1 \mathbf{x}_1)^N + \dots + (h_d \mathbf{x}_d)^N)} d\mathbf{x} \\ &\ll (\delta_V h_V)^{-\frac{d_V+1}{2}} |\boldsymbol{\varepsilon}|^{-1} |\mathbf{h}|^{-1}. \end{aligned}$$

Combining with the estimate on $[\mathcal{P}\mathbf{I}_{T_{\boldsymbol{\varepsilon}}(\Omega)_{\pm\mathbf{h}}}^{(\mathbf{h})}](0)$, we have that

$$(6.6) \quad n_{\boldsymbol{\varepsilon}}^{\pm}(\Omega) = \frac{|\boldsymbol{\varepsilon}|^{-1}}{|\Gamma|} \operatorname{Vol}(\Omega) + \left(|\boldsymbol{\varepsilon}|^{-1} \sum_{k=1}^d \varepsilon_k h_k \right) + \left((\delta_V h_V)^{-\frac{d_V+1}{2}} |\boldsymbol{\varepsilon}|^{-1} |\mathbf{h}|^{-1} \right).$$

Using the fact that $|\boldsymbol{\varepsilon}| \leq \delta_V^{d_V}$, we obtain that equation (6.6) reduces to

$$n_{\boldsymbol{\varepsilon}}^{\pm}(\Omega) = \frac{|\boldsymbol{\varepsilon}|^{-1}}{|\Gamma|} \operatorname{Vol}(\Omega) + \left(|\boldsymbol{\varepsilon}|^{-1} \delta_V^{\frac{2d_V}{1+d_V+2d_{V'}}} \right). \quad \blacksquare$$

Acknowledgments This work is part of the author's doctoral studies at Université de Montréal under the supervision of Iosif Polterovich. We thank Pedro Freitas, Katie Gittins, Corentin Léna, and Braxton Osting for useful discussions.

References

- [1] P. R. S. Antunes and P. Freitas, *Numerical optimization of low eigenvalues of the Dirichlet and Neumann Laplacians*. J. Optim. Theory Appl. 154(2012), 235–257. <https://doi.org/10.1007/s10957-011-9983-3>
- [2] P. R. S. Antunes and P. Freitas, *Optimal spectral rectangles and lattice ellipses*. Proc. R. Soc. Lond. Ser. A Math. Phys. Eng. Sci. 469(2013), no. 2015, 20120492. <https://doi.org/10.1098/rspa.2012.0492>
- [3] W. Banaszczyk, *New bounds in some transference theorems in the geometry of numbers*. Math. Ann. 296(1993), 625–635. <https://doi.org/10.1007/BF01445125>
- [4] M. van den Berg, D. Bucur, and K. Gittins, *Maximising Neumann eigenvalues on rectangles*. Bull. Lond. Math. Soc. 48(2016), 877–894. <https://doi.org/10.1112/blms/bdw049>
- [5] M. van den Berg and K. Gittins, *Minimizing Dirichlet eigenvalues on cuboids of unit measure*. Mathematika 63(2017), 469–482. <https://doi.org/10.1112/S0025579316000413>
- [6] A. Berger, *The eigenvalues of the Laplacian with Dirichlet boundary condition in \mathbb{R}^2 are almost never minimized by disks*. Ann. Global Anal. Geom. 47(2015), 285–304. <https://doi.org/10.1007/s10455-014-9446-9>
- [7] M. Berger, P. Gauduchon, and E. Mazet, *Le spectre d'une variété riemannienne*. Lecture Notes in Mathematics, 194, Springer-Verlag, Berlin-New York, 1971.
- [8] P. Buser, *A note on the isoperimetric constant*. Ann. Sci. École Norm. Sup. (4) 15(1982), 213–230.
- [9] J. W. S. Cassels, *An introduction to the geometry of numbers*. Die Grundlehren der mathematischen Wissenschaften, Band 99, Springer-Verlag, Berlin-New York, 1971.
- [10] B. Colbois and J. Dodziuk, *Riemannian metrics with large λ_1* . Proc. Amer. Math. Soc. 122(1994), 905–906. <https://doi.org/10.2307/2160770>
- [11] J. J. Duistermaat and V. W. Guillemin, *The spectrum of positive elliptic operators and periodic bicharacteristics*. Invent. Math. 29(1975), 39–79. <https://doi.org/10.1007/BF01405172>
- [12] G. Faber, *Beweis, dass unter allen homogenen Membranen von gleicher Fläche und gleicher Spannung die kreisförmige den tiefsten Grundton gibt*. Sitzungsber. Bayer. Akad. Wiss. München, Math.-Phys. Kl., 1923, pp. 169–172.
- [13] K. Gittins and S. Larson, *Asymptotic behaviour of cuboids optimising Laplacian eigenvalues*. Integral Equations Operator Theory 89(2017), 607–629. <https://doi.org/10.1007/s00020-017-2407-5>
- [14] A. Hassannezhad, G. Kokarev, and I. Polterovich, *Eigenvalue inequalities on Riemannian manifolds with a lower Ricci curvature bound*. J. Spectr. Theory 6(2016), 807–835. <https://doi.org/10.4171/JST/143>
- [15] J. Hersch, *Quatre propriétés isopérimétriques de membranes sphériques homogènes*. C. R. Acad. Sci. Paris Sér. A-B 270(1970), A1645–A1648.
- [16] A. Iosevich and E. Liflyand, *Decay of the Fourier transform, analytic and geometric aspects*. Birkhäuser/Springer, Basel, 2014. <https://doi.org/10.1007/978-3-0348-0625-1>
- [17] C.-Y. Kao, R. Lai, and B. Osting, *Maximization of Laplace-Beltrami eigenvalues on closed Riemannian surfaces*. ESAIM Control Optim. Calc. Var. 23(2017), 685–720. <https://doi.org/10.1051/cocv/2016008>
- [18] M. Karpukhin, N. Nadirashvili, A. V. Penskoi, and I. Polterovich, *An isoperimetric inequality for Laplace eigenvalues on the sphere*. J. Diff. Geom., to appear.
- [19] Yu. A. Kordyukov and A. A. Yakovlev, *Lattice points in domains and adiabatic limits*. (Russian) Algebra i Analiz 23(2011), 80–95. <https://doi.org/10.1090/S1061-0022-2012-01225-2>
- [20] Yu. A. Kordyukov and A. A. Yakovlev, *The problem of the number of integer points in families of anisotropically expanding domains, with applications to spectral theory*. Mat. Zametki 92(2012); trans. in Math. Notes 92(2012), no. 3–4, 574–576. <https://doi.org/10.1134/S0001434612090295>
- [21] Yu. A. Kordyukov and A. A. Yakovlev, *The number of integer points in a family of anisotropically expanding domains*. Monatsh. Math. 178(2015), 97–111. <https://doi.org/10.1007/s00605-015-0787-7>
- [22] Yu. A. Kordyukov and A. A. Yakovlev, *On a problem in geometry of numbers arising in spectral theory*. Russ. J. Math. Phys. 22(2015), 473–482. <https://doi.org/10.1134/S106192081504007X>
- [23] E. Krahn, *Über eine von Rayleigh formulierte Minimaleigenschaft des Kreises*. Math. Ann. 94(1925), 97–100. <https://doi.org/10.1007/BF01208645>
- [24] E. Krahn, *Über Minimaleigenschaften der Kugel in drei und mehr Dimensionen*. Acta Comm. Univ. Tartu (Dorpat) A9(1926), 1–44.
- [25] J. Lagacé and L. Parnowski, *A generalised Gauss circle problem and integrated density of states*. J. Spectr. Theory 6(2016), 859–879. <https://doi.org/10.4171/JST/145>
- [26] N. Nadirashvili, *Berger's isoperimetric problem and minimal immersions of surfaces*. Geom. Funct. Anal. 6(1996), 877–897. <https://doi.org/10.1007/BF02246788>

- [27] G. Szegő, *Inequalities for certain eigenvalues of a membrane of given area*. J. Rational Mech. Anal. 3(1954), 343–356. <https://doi.org/10.1512/iumj.1954.3.53017>
- [28] H. F. Weinberger, *An isoperimetric inequality for the N -dimensional free membrane problem*. J. Rational Mech. Anal. 5(1956), 633–636. <https://doi.org/10.1512/iumj.1956.5.55021>

Department of Mathematics, University College London, Gower Street, London, WC1E 6BT, United Kingdom
e-mail: j.lagace@ucl.ac.uk