

## Better dietary adherence and weight maintenance achieved by a long-term moderate-fat diet

Leila Azadbakht, Parvin Mirmiran, Ahmad Esmailzadeh and Fereidoun Azizi\*

Endocrine Research Center, Shaheed Beheshti University of Medical Sciences, PO Box 19395-4763, Tehran, Iran

(Received 24 January 2006 – Revised 31 July 2006 – Accepted 24 August 2006)

The objective of the present study was to determine the effects of a long-term moderate-fat diet (30% energy from fat) v. a low-fat one (20% energy from fat) on metabolic risks. The study was a randomised, prospective 14-month trial on overweight and obese patients (eighty-nine overweight and obese men and women). The intervention was a moderate-fat diet (30% energy) or a low-fat diet (20% energy). The main outcome measurements were change in body weight, waist circumference, LDL-cholesterol, HDL-cholesterol, total cholesterol, TAG, and systolic and diastolic blood pressure. Forty-five subjects on the moderate-fat diet and forty-four subjects on the low-fat one were studied. Characteristics of all randomised participants were similar in both groups. After 7 months, the moderate- and low-fat diets had similar effects on cardiovascular risks. The moderate-fat diet was more successful after 14 months in reducing weight ( $-5.0$  (SD 2.5) kg in the moderate-fat group v.  $-1.2$  (SD 1.1) kg in the low-fat one;  $P < 0.0001$ ), waist circumference ( $-5.5$  (SD 2.4) cm in the moderate-fat group v.  $-2.3$  (SD 1.3) cm in the low-fat one;  $P < 0.0001$ ), and other cardiovascular risk factors as well (LDL, TAG, total cholesterol and systolic blood pressure). In conclusion, a moderate-fat energy-restricted diet in the long term might have more beneficial effects on weight maintenance and cardiovascular risk factors compared with a low-fat diet. Better dietary adherence with the moderate-fat diet may be the reason for its successful effects.

**Moderate-fat diet: Low-fat diet: Cardiovascular risk factors: Long-term diet: Weight reducing**

Today, obesity is a major health problem commonly faced by developed and developing countries alike. Several studies indicate that obese individuals are more susceptible to chronic non-communicable disease than their normal-weight counterparts (Higgins *et al.* 1988; Pi-Sunyer, 1993; Bray, 1998; Must *et al.* 1999). Overweight and obesity are related with major chronic conditions, including hypertension, coronary artery disease, diabetes, arthritis, cancer and many causes of mortality (National Institutes of Health; Anonymous, 1998). Weight loss could help control other diseases and related risk factors (US Department of Health & Human Services, 2005).

Current dietary guidelines for weight-loss promotion recommend a reduction in fat intake (Pi-Sunyer, 1993). The standard approach for weight loss is a low-fat, high-carbohydrate diet, with an energy level below that required for weight maintenance (Bray, 1998). During the past few years there have been reports on the favourable effect of a restricted carbohydrate diet on weight loss and CVD (Bouche *et al.* 2003; Brynes *et al.* 2003; Ebbeling *et al.* 2003, 2005; Pereira *et al.* 2004; Stern *et al.* 2004). Ebbeling *et al.* (2005) showed that a low-fat diet significantly worsened thrombogenicity. There are no obvious suggestions, however, on the appropriate amount of fat in the diet to promote long-term weight loss and maintenance and decrease cardiovascular risk factors. Most previous studies have demonstrated the short-term success of low-fat diets (Sheppard *et al.* 1991; Powell *et al.* 1994; Jeffery *et al.* 1995; McManus *et al.* 2001). Possibly the limited long-term success of some low-fat diets

is associated with the fact that in the long term these diets are less appetising, making their long-term compliance difficult. A successful study on weight loss was done by McManus *et al.* (2001), which showed the priority of a moderate-fat diet on weight loss, participation and adherence. But, there are no reports on the effect on a moderate-fat diet on metabolic risks. We hypothesised that by using a moderate-fat diet, we could use higher amounts of unsaturated fat, such as  $\omega_3$ , which is more beneficial for the prevention of CVD. Furthermore, dietary adherence to a moderate-fat diet may be better than to a low-fat one. Therefore, the present study was conducted to determine the effects of a long-term moderate-fat diet (30% energy from fat) v. a low-fat one (20% energy from fat) on metabolic risk factors.

### Subjects and methods

Eighty-nine overweight and obese subjects (twenty-five men and sixty-four women) were recruited for a 14-month randomised trial from among participants of the Tehran Lipid and Glucose Study attending the diet therapy clinic at the Lipid and Glucose Unit of the Endocrine Research Center. They had not participated in weight-reduction programmes during the previous 6 months and had maintained a stable weight ( $\pm 1$  kg); none of the study participants had previously participated in dietary studies. They were non-smokers, free of chronic disease and readily participated in the monthly visits. Each patient was asked to complete personal health

\* Corresponding author: Dr Fereidoun Azizi, fax +98 21 2402463, email azizi@erc.ac.ir

and medical history questionnaires that were used as a screening tool. Patients were excluded if they were taking any medications affecting nutrient metabolism, blood lipids and blood pressure, vitamin and mineral supplements and antacids containing Mg or Ca. The proposal of the present study was approved by the research council of the Endocrine Research Center of Shaheed Beheshti University of Medical Sciences (Tehran, Iran) and informed written consent was obtained from each subject. Forty-five subjects were randomised using a program generated by a random number table to an energy-controlled diet containing 30% energy as fat, and forty-four subjects to an energy-controlled diet containing 20% energy as fat (Fig. 1). The nutritionist who prescribed the diets had to be aware of the group assignment. Laboratory staff were not aware to which group the patients had been assigned. All subjects were given general oral and written information about healthy food choices and a diet 2090 kJ (500 kcal) below their energy needs according to their weight, which was offered according to specific individualised programmes at baseline and at subsequent visits. Nutrient goals for the intervention diets are shown in Table 1. Under- and over-reporters were defined as reported daily energy intakes less than 3350 kJ/d (800 kcal/d) and over 17 570 kJ/d (4200 kcal/d) respectively (Fung *et al.* 2002).

#### Study procedures

During the 3 weeks of run-in and 14 months of intervention feeding, we followed patients every month and checked their food records and current use of medications and measured their anthropometric indices. Habitual physical activity levels were ascertained by the Lipid Research Clinic questionnaire (Ainsworth *et al.* 1993) at baseline and after 7 and 14 months of intervention.

#### Measurements

Weight was measured while the subjects were minimally clothed without shoes using digital scales and recorded to the nearest 0.1 kg. Height was measured in a standing position, without shoes, using a tape meter while the shoulders were in a normal state. BMI was calculated as weight in kg divided by height in metres squared. Waist circumference was measured at the narrowest level over light clothing, using an unstretched tape meter, without any pressure to the body surface (Esmailzadeh *et al.* 2004). Blood samples (12 h fasting) were collected into tubes containing 0.1% EDTA and centrifuged at 4°C and 500 g for 10 min to separate the plasma (Azizi *et al.* 2003b). Serum total cholesterol, HDL and TAG concentrations were measured by commercially available enzymic reagents (Pars Azmoon, Iran) adapted to the Selectra Autoanalyzer (Vital Scientific, Spankeren, Netherlands). HDL-cholesterol was measured after precipitation of the apo B-containing lipoproteins with phosphotungstic acid. LDL-cholesterol was calculated according to the Friedewald method (Friedewald *et al.* 1972). All samples were analysed when internal quality control met the acceptable criteria. Blood pressure was measured twice after the participants sat for 10 min (Azizi *et al.* 2002). Additional covariate information regarding age, smoking habits (Azizi *et al.* 2003a), medical history and current use of medication (Mirmiran *et al.* 2003) was obtained using validated questionnaires, as reported earlier, completed during the screening and every month. Measurements were obtained at baseline, and after 7 and 14 months of the dietary programme.

#### Statistical analysis

Data are presented as means and standard deviations unless stated otherwise. The sample of eighty subjects was



**Fig. 1.** Design of the trial. Among 121 subjects, 100 individuals that met the study criteria were randomly assigned to a moderate-fat diet and a low-fat diet (fifty on the moderate-fat diet and fifty on the low-fat one). Forty-five subjects on the moderate-fat diet and forty-four subjects on the low-fat diet followed-up the study for 14 months.

**Table 1.** Nutrient goals (% energy) for the intervention diets

	Moderate-fat diet	Low-fat diet
Fat	30	20
SFA	5	5–6
MUFA	15	7
PUFA	10	7
Protein	15	15
Carbohydrate	55	65
Cholesterol (mg)	< 200	< 200
Dietary fibre (g)	25	25

determined to provide at least an 80% power to detect a difference in weight loss of 2.1 kg between the groups, allowing for a dropout rate of 30%. The primary outcome variable was change in body weight. Other outcome variables were serum LDL, HDL, TAG, total cholesterol, systolic and diastolic blood pressure, from baseline to 14 months. The changes in outcome variables were compared between the diet groups by unpaired *t* test and within each group by paired *t* test. *P* values less than 0.05, two sided, were considered statistically significant. All analyses were conducted using SPSS version 11.1 (SPSS Inc., Chicago, IL, USA).

## Results

Baseline characteristics for all subjects are shown in Table 2. The mean age of the two groups was similar (45 (SD 5) years in the moderate-fat-diet group and 46 (SD 6) years in the low-fat-diet group). Mean weight was 80 (SD 11) kg in the

**Table 2.** Baseline characteristics of study participants of both groups (Mean values and standard deviations)

Characteristics	Moderate-fat diet (n 45)		Low-fat diet (n 44)	
	Mean	SD	Mean	SD
Age (years)	45	5	46	6
BMI (kg/m <sup>2</sup> )	29.0	10	29.2	11
Weight (kg)	80	11	79	12
Waist circumference (cm)	99	19	101	21
Medications (%)				
LDL (mg/dl)	120	36	117	38
HDL (mg/dl)	36.5	15	35.3	14
TAG (mg/dl)	196	28	203	29
Total cholesterol (mg/dl)	211	32	210	31
Systolic blood pressure (mmHg)	131	24	133	25
Diastolic blood pressure (mmHg)	80	11	81	11
Female				
<i>n</i>	31		33	
%	68		75	
Male				
<i>n</i>	14		11	
%	32		25	
Physical activity (%)				
Very light	61		60	
Light	36		38	
Moderate	3		2	
Medications (%)				
Oestrogen-replacement therapy	5		5	
Oral contraception	6		5	

moderate-fat group and 79 (SD 12) kg in the low-fat group. The mean of the serum lipid profiles and blood pressure were not different in the two groups at baseline. Table 3 shows the reported nutrient intake at baseline and after 7 and 14 months' intervention, in both the moderate- and low-fat-diet groups. Reported nutrient intake at baseline was similar in the two groups. The reported total fat intake decreased from 29 to 20% at 7 and 14 months in the low-fat-diet group. In the moderate-fat group, total fat intake was similar at baseline, and after 7 and 14 months. There was no significant difference between SFA intake in the two groups any time after 7 and 14 months' intervention. The MUFA and PUFA intake was higher in the moderate-fat-diet group than in the low-fat one (15.1 (SD 4.1) % energy for the moderate-fat and 9.1 (SD 4.2) % energy for the low-fat one at 7 months for MUFA; 15.0 (SD 4.9) % energy for the moderate-fat and 9.9 (SD 5.3) % energy for the low-fat one at 14 months for MUFA; 9.9 (SD 3.9) % energy for the moderate-fat and 6.9 (SD 4.2) % energy for the low-fat one at 7 months for PUFA; 10.1 (SD 4.8) % energy for the moderate-fat diet and 6.9 (SD 5.1) % for the low-fat one at 14 months for PUFA). The reported carbohydrate intake for both groups was similar at baseline but it increased for the low-fat group at 7 and 14 months. Protein, fibre and cholesterol intakes of the two groups were similar.

Changes in cardiovascular risk factors after 7 months are shown in Table 4. The highest reduction in body weight was seen after the first 7 months both in the moderate- and low-fat groups. Weight reduction was similar and significant after 7 months in both groups;  $-5.3$  (SD 1.3) kg in the moderate-fat group ( $P < 0.001$ ), and  $-5.2$  (SD 1.9) kg in the low-fat group ( $P < 0.001$ ). Changes in waist circumference were also similar to those seen in weight. The effects of low-fat and moderate-fat diets on cardiovascular risk factors were similar after 7 months.

Table 5 shows the changes in cardiovascular risk factors after 14 months' intervention. The moderate-fat diet was more successful after 14 months in reducing weight, waist circumference, LDL-cholesterol, TAG, and systolic and diastolic blood pressure. Although only the moderate-fat diet could significantly increase HDL after 14 months, there was no significant difference between the two groups at any time.

## Discussion

The results of the present study testing two approaches to weight loss, a low-fat diet and a moderate-fat one, both having the same goals for energy, showed that the moderate-fat diet had more favourable effects on cardiovascular risks in the long term. The present results showed that after 14 months' intervention the changes in reduction of weight, waist circumference, LDL, TAG, total cholesterol, and systolic and diastolic blood pressure in the moderate-fat-diet group were significantly greater than for the low-fat one. However, after 7 months of intervention there were no significant differences between the two groups in decreasing the cardiovascular risks. Therefore, we can conclude that the moderate-fat diet is more suitable in the long term. As a low-fat diet is more restrictive to follow, it seems that dietary adherence to a moderate-fat diet is better than to low-fat ones. Therefore, weight maintenance, a key issue in long-term weight-reducing programmes, is achieved in energy-restricted moderate-fat diets.

**Table 3.** Reported nutrient intake among subjects at time points during the study (Mean values and standard deviations)

Variable	Baseline				<i>P</i> *	7 months				<i>P</i> *	14 months				<i>P</i> *
	Moderate-fat (n 50)		Low-fat (n 50)			Moderate-fat (n 47)		Low-fat (n 45)			Moderate-fat (n 45)		Low-fat (n 44)		
	Mean	SD	Mean	SD		Mean	SD	Mean	SD		Mean	SD	Mean	SD	
Total energy (kJ/d)	10611	172	10506	167	0.81	8368	126	8389	130	0.89	8745	155	9205	163	0.79
Total fat (% energy)	29	6.1	29	6.0	0.90	30	7.0	20	6.5	0.01	30	7.2	20	10	0.04
SFA (% energy)	7.3	2.3	7.5	2.4	0.68	7.1	2.1	7.0	2.0	0.84	7.0	3.6	6.8	4.2	0.85
MUFA (% energy)	11.1	3.9	10.6	3.7	0.32	15.1	4.1	7.1	4.2	0.01	15.0	4.9	7.9	5.3	0.01
PUFA (% energy)	5.9	2.6	6.9	2.9	0.09	9.9	3.9	6.9	4.2	0.01	10.1	4.8	6.9	5.1	0.02
Carbohydrate (% energy)	55	11	58	12	0.22	55	13	64	16	0.01	55	14	65	19	0.03
Protein (% energy)	15	6.1	15	6.0	0.99	15	8.0	15	10.1	0.99	15	9.6	15	11	0.99
Dietary fibre (g/1000 kJ)	2.5	1.4	2.6	1.4	0.68	2.5	1.5	2.6	1.6	0.71	2.5	1.7	2.6	2.1	0.89
Cholesterol (mg/1000 kJ)	55	5	56	5	0.52	49	7	49	9	0.51	49	9	49	11	0.52

\* *P* values for difference between the moderate-fat and low-fat groups by *t* test.

**Table 4.** Changes in cardiovascular risk factors assessed after 7 months of moderate- and low-fat diets (Mean values and standard deviations)

Variable	Moderate-fat diet (n 45)				<i>P</i> *	Low-fat diet (n 44)				<i>P</i> *	Between-group change		<i>P</i> †
	7 months		Change			7 months		Change			Difference (95% CI)		
	Mean	SD	Mean	SD		Mean	SD	Mean	SD				
Weight (kg)	74.9	10	-5.1	1.3	0.02	73.7	9	-5.3	1.9	0.02	-5.2, 2.8	0.51	
Waist circumference (cm)	91.4	16	-7.6	2.1	0.04	93.0	19	-7.4	2.1	0.04	-5.5, 9.9	0.57	
HDL (mg/dl)	36.9	16	+3.1	3.0	0.34	41.9	17	+6.6	3.2	0.04	-4.6, 9.2	0.51	
LDL (mg/dl)	110.1	32	-10.5	3.9	0.14	103.7	30	-6.9	4.6	0.05	-18.5, 10.9	0.60	
TAG (mg/dl)	183.2	23	-7.5	3.3	0.16	191.5	27	-11.3	5.6	0.81	-4.2, 16.8	0.23	
Total cholesterol (mg/dl)	203.0	22	-8.3	3.7	0.11	201.0	26	-8.7	5.6	0.15	-12.1, 8.1	0.69	
Systolic blood pressure (mmHg)	128.9	20	-7.1	2.6	0.05	126.1	21	-6.9	2.3	0.05	-11.4, 5.8	0.52	
Diastolic blood pressure (mmHg)	82.8	11	-2.2	0.9	0.45	78.1	10	-2.9	1.1	0.48	-9.1, -0.2	0.03	

\* *P* values for difference from baseline by paired *t* test.

† *P* values for difference between moderate-fat and low-fat group by *t* test.

**Table 5.** Changes in cardiovascular risk factors assessed after 14 months of moderate- and low-fat diets (Mean values and standard deviations)

Variable	Moderate-fat diet (n 45)				Low-fat diet (n 44)				Between-group change		
	14 months		Change		14 months		Change		P*	Difference (95% CI)	P†
	Mean	SD	Mean	SD	Mean	SD	Mean	SD			
Weight (kg)	75.0	10	-5.0	2.5	77.8	11	-1.2	1.1	0.51	-2.9, 4.6	0.0001
Waist circumference (cm)	93.5	18	-5.5	2.4	98.7	20	-2.3	1.3	0.61	2.3, 4.0	0.0001
HDL (mg/dl)	43.1	17	+6.6	3.2	39.1	16	3.8	3.6	0.56	-10.9, 2.9	0.25
LDL (mg/dl)	113	37	-6.9	4.6	112	38	-3.9	5.9	0.49	0.77, 5.2	0.008
TAG (mg/dl)	186	26	-10.1	5.1	200	24	-2.3	1.6	0.53	6.2, 9.4	0.0001
Total cholesterol (mg/dl)	201	22	-10.3	4.1	203	26	-5.9	3.3	<0.05	2.8, 5.9	0.0001
Systolic blood pressure (mmHg)	128	2	-7.4	2.3	129	23	-3.3	1.2	0.31	3.3, 4.8	0.0001
Diastolic blood pressure (mmHg)	82.1	11	-2.9	1.2	79.7	10	-1.3	1.1	0.62	-6.8, 2.0	0.28

\* P values for difference from baseline by paired t test.

† P values for difference between moderate-fat and low-fat group by t test.

A meta-analysis (Pirozzo *et al.* 2002) showed that low-fat diets were not better than energy-restricted diets in achieving long-term weight loss. In their study, there were four studies at the 6-month follow-up, five studies at the 12-month follow-up and three studies at the 18-month follow-up. In most of the mentioned studies, there were small, non-significant differences in weight loss between the low-fat-diet groups and the comparison groups. The changes in metabolic risks also were not clinically significant in the low-fat-diet groups compared with other groups. However, in the present study the weight loss in the low-fat-diet group was modest and it was less than observed in other studies.

Regarding weight reduction, McManus *et al.* (2001) also showed that a moderate-fat Mediterranean diet, controlled in energy, was more successful in long-term weight loss as compared with a low-fat one. In the present study, the results at 7 months showed similar amounts of weight loss in the two groups – about 5 kg. However, while the moderate-fat-diet group maintained their lower average weight through the 14-month study period, the low-fat group regained the weight lost. The moderate-fat-diet group was more likely to adhere to the diet; it seems that low-fat diets are less appetising, which makes long-term compliance difficult.

Esposito *et al.* (2004) also showed that consumption of a Mediterranean diet is more effective than a low-fat one in changing serum lipid concentrations.

Jeffery *et al.* (1995) showed that a low-fat diet caused more weight loss early in the programme as compared with a higher-fat diet, but after 2 years both groups gained back all the weight lost. Toubro & Astrup (1997) showed a non-significant difference ( $P=0.08$ ) in weight regained after a 1-year weight-maintenance programme between an *ad libitum* low-fat (25% energy) diet and a higher fat (unspecified fat content) one. Bahadori *et al.* (2005) in a 24-week study showed that a low-fat, low-glycaemic-index diet induced a significant reduction of fat mass; adherence to the diet was very good.

As has been reported in other studies, many obese individuals under-report their food intake (Lichtman *et al.* 1992; Johnson *et al.* 1994). Therefore, we tested under- and over-reporting and there were no individuals who had overestimated or underestimated their energy intakes in our sample. It seems that because individuals have been well educated on all portion sizes and the food amounts that should be consumed, they reported their energy intake correctly.

The results of the limited success of the low-fat diet would not have been predicted by experiments on metabolism. After consuming a meal, carbohydrate is used preferentially for oxidation compared with fat, which is stored (McManus *et al.* 2001). Therefore satiety might be more stimulated by carbohydrates rather than fat. Of course, motivation and adherence are very difficult to sustain in a weight-loss programme. Fat may enhance the palatability of foods, especially vegetables, in the moderate-fat diet.

In the present study, while outlining the specified diet, we did not actually provide the foods mentioned; this is one of our limitations. Diet adherence was, however, checked by food records. The strengths of the present study are its randomised design, long intervention period and frequent follow-up of patients. The present study provides the opportunity to assess the change in the serum lipid profile and blood pressure, which have been little emphasised in previous



studies with the aim of comparing the long-term effects of moderate- and low-fat diets.

According to the results of the present study, we can conclude that the long-term effects of a moderate-fat diet, predominantly with unsaturated fatty acids, are more favourable than those of a low-fat one. Actually, the better adherence of subjects in the modified fat group may result in weight maintenance in the long term. Further studies need to be conducted on the use of this diet to prevent obesity. Also, the effect of this diet on other specific cardiovascular risks and inflammatory markers should be examined.

## References

- Ainsworth BE, Jacobs DR Jr & Leon AS (1993) Validity and reliability of self-reported physical activity status: the Lipid Research Clinics questionnaire. *Med Sci Sports Exerc* **25**, 92–98.
- Anonymous (1998) Clinical guidelines on the identification, evaluation, and treatment of overweight and obesity in adults – the Evidence Report. National Institutes of Health. *Obesity Res* **6**, Suppl. 2, 51S–209S.
- Azizi F, Emami H, Salehi P, Ghanbarian A, Mirmiran P, Mirbolooki M & Azizi T (2003a) Cardiovascular risk factors in the elderly: the Tehran Lipid and Glucose Study. *J Cardiovasc Risk* **10**, 65–73.
- Azizi F, Ghanbarian A, Madjid M & Rahmani M (2002) Distribution of blood pressure and prevalence of hypertension in Tehran adult population: Tehran Lipid and Glucose Study (TLGS), 1999–2000. *J Hum Hypertens* **16**, 305–312.
- Azizi F, Rahmani M, Ghanbarian A, Emami H, Salehi P, Mirmiran P & Sarbazi N (2003b) Serum lipid levels in an Iranian adults population: Tehran Lipid and Glucose Study. *Eur J Epidemiol* **18**, 311–319.
- Bahadori B, Yazdani-Biuki B, Krippel P, Brath H, Uitz E & Wascher TC (2005) Low-fat, high-carbohydrate (low-glycaemic index) diet induces weight loss and preserves lean body mass in obese healthy subjects: results of a 24-week study. *Diabetes Obes Metab* **7**, 290–293.
- Bouche C, Rizkalla S, Luo J, Vidal H, Veronese A, Pacher N, Fouquet C, Lang V & Slama G (2003) Five-week, low-glycemic-index diet decreases total fat mass and improves plasma lipid profile in moderately overweight nondiabetic men. *Diabetes Care* **2**, 822–828.
- Bray GA (1998) Obesity: a time bomb to be defused. (Comment). *Lancet* **352**, 160–161.
- Brynes AE, Edwards CM, Ghatei M, Dornhorst A, Morgan LM, Bloom SR & Frost GS (2003) A randomised four-intervention crossover study investigating the effect of carbohydrates on day-time profiles of insulin, glucose, non-esterified fatty acids and triacylglycerols in middle-aged men. *Br J Nutr* **89**, 207–218.
- Ebbeling C, Leidig M, Sinclair K, Hangen J & Ludwig D (2003) A reduced-glycemic load diet in the treatment of adolescent obesity. *Arch Pediatr Adolesc Med* **157**, 773–779.
- Ebbeling CB, Leidig MM, Sinclair KB, Seger-Shippe LG, Feldman HA & Ludwig DS (2005) Effects of an *ad libitum* low-glycemic load diet on cardiovascular disease risk factors in obese young adults. *Am J Clin Nutr* **81**, 976–982.
- Esmailzadeh A, Mirmiran P & Azizi F (2004) Waist-to-hip ratio is a better screening measure for cardiovascular risk factors than other anthropometric indicators in Tehranian adult men. *Int J Obes Relat Metab Disord* **28**, 1325–1332.
- Esposito K, Marfella R, Ciotola M, Di Palo C, Giugliano F, Giugliano G, D'Armiento M, D'Andrea F & Giugliano D (2004) Effect of a Mediterranean-style diet on endothelial dysfunction and markers of vascular inflammation in the metabolic syndrome: a randomized trial. *JAMA* **292**, 1490.
- Friedewald WT, Levy RI & Fredrickson DS (1972) Estimation of the concentration of low-density lipoprotein cholesterol in plasma, without use of the preparative ultracentrifuge. *Clin Chem* **18**, 499–502.
- Fung TT, Hu FB, Pereira MA, Liu S, Stampfer MJ, Colditz GA & Willett WG (2002) Whole-grain intake and the risk of type 2 diabetes: a prospective study in men. *Am J Clin Nutr* **76**, 35–40.
- Higgins M, Kannel W, Garrison R, Pinsky J & Stokes J III (1988) Hazards of obesity – the Framingham experience. *Acta Med Scand* **723**, S23–S36.
- Jeffery RW, Hellerstedt WL, French SA & Baxter JE (1995) A randomized trial of counseling for fat restriction versus calorie restriction in the treatment of obesity. *Int J Obes Relat Metab Disord* **19**, 132–137.
- Johnson RK, Goran MI & Poehlman ET (1994) Correlates of over- and underreporting of energy intake in healthy older men and women. *Am J Clin Nutr* **59**, 1286–1290.
- Lichtman SW, Pisarska K, Berman ER, Pestone M, Dowling H, Offenbacher E, Weisel H, Heshka S, Matthews DE & Heymsfield SB (1992) Discrepancy between self-reported and actual caloric intake and exercise in obese subjects. *N Engl J Med* **327**, 1893–1898.
- McManus K, Antinoro L & Sacks F (2001) A randomized controlled trial of a moderate-fat, low-energy diet compared with a low fat, low-energy diet for weight loss in overweight adults. *Int J Obes Relat Metab Disord* **25**, 1503–1511.
- Mirmiran P, Mohammadi F, Allahverdi S & Azizi F (2003) Estimation of energy requirements for adults: Tehran lipid and glucose study. *Int J Vitam Nutr Res* **73**, 193–200.
- Must A, Spadano J, Coakley EH, Field AE, Colditz G & Dietz WH (1999) The disease burden associated with overweight and obesity. *JAMA* **282**, 1523–1529.
- Pereira MA, Swain J, Goldfine AB, Rifai N & Ludwig DS (2004) Effects of a low-glycemic load diet on resting energy expenditure and heart disease risk factors during weight loss. *JAMA* **292**, 2482–2490.
- Pirozzo S, Summerbell C, Cameron C & Glasziou P (2002) Advice on low-fat diets for obesity. *Cochrane Database Syst Rev* **2**, CD003640.
- Pi-Sunyer FX (1993) Medical hazards of obesity. *Ann Intern Med* **119**, 655–660.
- Powell JJ, Tucker T, Fisher AG & Wilcox K (1994) The effects of different percentage of dietary fat intake, exercise, and calorie restriction of body composition and body weight in obese females. *Am J Health Prom* **8**, 443–448.
- Sheppard L, Kristal AR & Kushi LH (1991) Weight loss in women participating in a randomized trial of low-fat diets. *Am J Clin Nutr* **54**, 821–828.
- Stern L, Iqbal N, Seshadri P, Chicano KL, Daily DA, McGrory J, Williams M, Gracely EJ & Samaha FF (2004) The effects of low-carbohydrate versus conventional weight loss diets in severely obese adults: one-year follow-up of a randomized trial. *Ann Intern Med* **140**, 778–785.
- Toubro S & Astrup A (1997) Randomised comparison of diets for maintaining obese subjects' weight after major weight loss: ad lib, low fat, high carbohydrate diet v fixed energy intake. *BMJ* **314**, 29–34.
- US Department of Health and Human Services (2005) *Dietary Guidelines for Americans 2005*. Washington, DC: Government Printing Office.