

Food and Waterborne Disease in the Greater New York City Area Following Hurricane Sandy in 2012

Michael S. Bloom, PhD; Jillian Palumbo, MPH; Nazia Saiyed, MPH;
 Ursula Lauper, MA MPH; Shao Lin, MD, PhD

ABSTRACT

Objective: We aimed to evaluate residence in evacuation areas (storm areas) as a risk factor for food and waterborne disease (FWBD) associated with Hurricane Sandy flooding.

Methods: We captured 9601 incident outpatient and inpatient FWBD hospital discharge diagnoses for residents of the greater New York City area. We used Poisson or negative binomial regression models to compare the covariate-adjusted risk for a FWBD diagnosis, pre-Sandy (10/28-11/09, 2001-2011) vs. post-Sandy (10/28-11/09, 2012), for residents of “storm” and “non-storm” areas.

Results: Outpatient FWBD risk was lower for storm area residents after Hurricane Sandy (risk ratio [RR] = 0.58, 95% confidence interval [CI]: 0.46-0.74), and varied by age, sex, and county. However, storm area residents 65 years of age or older experienced higher risk after Hurricane Sandy (RR = 2.16, 95% CI: 1.11-4.19), albeit based on few cases. Inpatient FWBD risk was lower for non-storm area residents after Hurricane Sandy (RR = 0.79, 95% CI: 0.66-0.95), and varied by age, race, and county, although there was no significant change for storm area residents (RR = 0.86, 95% CI: 0.69-1.08). Those ≥ 65 years of age were also at lower risk for inpatient FWBD diagnosis, yet the effect was weaker for storm area (RR = 0.89, 95% CI: 0.67-1.18) than for non-storm area residents (RR = 0.68, 95% CI: 0.52-0.89).

Conclusions: Hurricane preparation, mitigation, and response activities in the greater New York City area may have led to “protective” effects for FWBD. (*Disaster Med Public Health Preparedness*. 2016;10:503-511)

Key Words: hurricane, infectious disease transmission, public health, floods

Hurricane Sandy made landfall in New York State (NYS) on October 29, 2012, battering parts of New York City (NYC), Long Island, and Westchester County with high winds and strong rains through October 30.¹ More than 12 million area residents, making up over half of NYS’s population, were affected by the storm to some degree.² Extensive flooding damaged property and critical infrastructure, around 7 to 8 million customers lost power, thousands of injuries were reported, and 117 deaths were attributed to the storm.^{1,3} The Office of the Mayor of New York City activated a Coastal Storm Plan on October 28, 2012, implementing mandatory evacuations for 375,000 residents of low-lying coastal zones.¹ Still, the flooding overwhelmed the NYC evacuation zones in Bronx, Kings (Brooklyn), New York (Manhattan), Queens, and Richmond (Staten Island) counties, as well as hurricane storm surge zones in nearby Nassau, Suffolk, and Westchester counties. The inundation disrupted power supplies and disabled sewage and water treatment facilities, possibly cross-contaminating municipal water supplies, and placing the population at increased risk for contracting food and waterborne disease (FWBD).^{4,5}

Extreme weather events, including hurricane-instigated flooding, can lead to direct contamination of food and drinking water sources by pathogenic microorganisms or to indirect contamination following refrigeration-compromising power outages and disrupted food distribution networks.^{6,7} Power was lost by 94.4% of Nassau, 77.8% of Suffolk, 69.5% of Richmond, and 40.5% of New York counties; the number of blackouts peaked on October 30, 2012.⁸ FWBD in association with flooding is a primary concern of health authorities and the public following extreme weather events.⁹ Crowding and inadequate hygiene, in association with temporary shelters for displaced persons, can introduce additional hazards.^{10,11} For example, epidemics of diarrheal illness have been reported after heavy rainfall and flooding in Austria,¹² Bangladesh,^{13,14} China,¹⁵ India,¹⁶ Mozambique,¹⁷ Pakistan,¹⁸ and the United States,¹⁹⁻²¹ most often following displacement of evacuees to inadequate and overcrowded facilities.²² The elderly, infants, and children are likely to be especially vulnerable to FWBD following extreme weather events.^{7,23,24} In urban areas, low-income, inner-city populations, and those without access to

motor vehicles, appear to be more vulnerable to adverse health impacts from extreme weather events as well.^{25,26}

Our aim in this analysis was to evaluate the impact of residence in low-lying, coastal storm-surge areas of NYC, and nearby Nassau, Suffolk, and Westchester counties, as a risk factor for FWBD associated with Hurricane Sandy and the mandatory evacuation order. In doing so, our goal was to offer recommendations for preventive measures to reduce adverse health impacts following future extreme weather events.

METHODS

Study Population

The study population comprised individuals residing in NYC (Bronx, Kings, New York, Queens, and Richmond counties), Long Island (Nassau and Suffolk counties), and Westchester County in NYS, between October 28 and November 9 during 2001 to 2012. The risk period was defined according to the emergency declaration and evacuation order issued in advance of Hurricane Sandy's landfall on October 28, 2012, and the end of the US Federal Emergency Management Agency (FEMA) emergency incident period on November 9, 2012. We geocoded individuals living in the study catchment area as residents of either the "storm area" or the "non-storm area" by using street address, based on emergency management agency-established coastal storm plan evacuation zones in NYC, or hurricane storm surge zones in Nassau, Suffolk, and Westchester counties (Supplementary Figure 1 in the online data supplement).

Study Outcome

We retrospectively captured incident outpatient (ie, emergency department) and inpatient FWBD diagnoses by using the NYS Department of Health Statewide Planning and Research Cooperative System (SPARCS). SPARCS is a legislatively mandated database capturing discharge diagnoses for approximately 95% of NYS hospitals (ie, excludes only federal and psychiatric facilities). Diagnoses are indexed by billing address, rather than treatment site, allowing us to identify the likely location of cases' residence. Using Map Marker v.22 (Pitney Bowes Software Inc, Troy, NY) and SAS v. 9.3 (SAS Institute, Inc, Cary, NC) most cases (98%) were geocoded to street-level resolution. Outpatient SPARCS records were available to us beginning in 2005, while inpatient records were available beginning in 2001. We captured all NYS outpatient and inpatient FWBD discharges between October 28 and November 9, 2001-2012, by including the 9th Revision of the International Classification of Diseases (ICD-9) discharge codes 003-009 (see Supplementary Table 1 in the online data supplement). Data access and use was approved by the NYS Department of Health Institutional Review Board and the SPARCS Data Protection Review Board.

Statistical Analysis

We characterized the number and proportion of outpatient and inpatient FWBD diagnoses made in the 13 days between October 28 and November 9 (ie, the "study interval"), in 2005 to 2012 or 2001 to 2012, respectively. Using the chi-square test, we assessed unadjusted differences by study area (storm and non-storm), for age, sex, race, ethnicity, county of residence, year of diagnosis, and source of payment for health care services as an indicator of socioeconomic status. We used multivariable Poisson or negative binomial (ie, to accommodate overdispersion) regression models to calculate risk ratios (RRs) and 95% confidence intervals (CIs) for FWBD diagnoses made during the "post-Sandy" period, defined as the study interval in 2012, relative to the average "pre-Sandy" baseline period, defined as the study interval in 2005-2011 (outpatient) or 2001-2011 (inpatient). We used generalized estimating equations (GEEs) to accommodate observations correlated by county of residence, 2010 US census population totals as the offset based on zip code centroids (census tracts did not conform to storm and non-storm areas),² and included variables representing the pre-/post-Sandy interval, study area (storm or non-storm), the cross product of pre-/post-Sandy interval \times study area to assess the interaction, day of the week (ie, short-term trend), and year (ie, long-term trend). Models were implemented in proc genmod by using the slice statement to assess overall effects. We constructed individual regression models for outpatient and inpatient diagnoses, confirmed the plausibility of model assumptions, and stratified by sex, age, race, and county of residence. We further assessed associations by ICD-9 code, to evaluate heterogeneity across FWBD diagnoses. In a sensitivity analysis, we also evaluated the impact of a 28-day risk period on outpatient FWBD diagnosis. SAS v. 9.3 (SAS Institute, Inc) was used for all analysis and statistical significance was defined as $P < 0.05$ for a two-tailed test.

RESULTS

Descriptive Analysis

Table 1 characterizes 2372 total outpatient FWBD events reported by hospitals in the study area between October 28 and November 9, from 2005-2012 (characterized graphically by Figure 1). Most cases lived in the non-storm area (75.0%), and approximately 25% of cases resided in the storm area, primarily in NYC counties (83.7%). The age distribution differed somewhat for outpatient diagnoses in the storm and non-storm areas ($P = 0.04$). A higher proportion of diagnoses in persons < 19 years of age was made in the non-storm area (63.3%) than in the storm area (57.8%), whereas a higher proportion of diagnoses among persons ≥ 20 years of age was made in the storm area (42.2%) than in the non-storm area (36.6%). Blacks (37.0% vs. 24.9%, $P < 0.01$) and persons of non-Hispanic ethnicity (63.5% vs. 49.7%, $P < 0.01$) made up larger proportions of outpatient FWBD cases in the storm area than in the non-storm area. There also existed

TABLE 1

Demographics to Characterize Incident Food and Waterborne Disease Outpatient Visits Among Residents of Hurricane Sandy Storm and Non-Storm Areas, 2005-2012^a				
Demographic Factor	Overall, No. (%)	Storm Area, No. (%)	Non-Storm Area, No. (%)	P value^b
Total	2372 (100.0)	592 (25.0)	1780 (75.0)	-
Age, years				
< 5	887 (37.4)	200 (33.8)	687 (38.6)	0.04
5-19	582 (24.5)	142 (24.0)	440 (24.7)	
20-64	806 (34.0)	218 (36.8)	588 (33.0)	
≥ 65	97 (4.1)	32 (5.4)	65 (3.6)	
Sex				
Male	1144 (48.2)	277 (46.8)	867 (48.7)	0.42
Female	1228 (51.8)	315 (53.2)	913 (51.3)	
Race				
White	632 (26.6)	145 (24.5)	487 (27.4)	<0.01
Black	663 (28.0)	219 (37.0)	444 (24.9)	
Other/Unknown	1077 (45.4)	228 (38.5)	849 (47.7)	
Ethnicity				
Spanish/Hispanic	748 (31.5)	156 (26.4)	592 (33.3)	<0.01
Non-Hispanic	1260 (53.1)	376 (63.5)	884 (49.7)	
Unknown	364 (15.3)	60 (10.1)	304 (17.1)	
County of residence				
Nassau	97 (4.1)	33 (5.6)	64 (3.6)	<0.01
Suffolk	184 (7.8)	27 (4.6)	157 (8.8)	
Westchester	105 (4.4)	1 (0.2)	104 (5.8)	
Bronx	376 (15.9)	43 (7.3)	333 (18.7)	
Kings (Brooklyn)	591 (24.9)	204 (34.5)	387 (21.7)	
New York (Manhattan)	379 (16.0)	169 (28.5)	210 (11.8)	
Queens	608 (25.6)	110 (18.6)	498 (28.0)	
Richmond (Staten Island)	32 (1.3)	5 (0.8)	27 (1.5)	
Year				
2005	155 (6.5)	27 (4.6)	128 (7.2)	0.25
2006	257 (10.8)	71 (12.0)	186 (10.4)	
2007	210 (8.9)	50 (8.4)	160 (9.0)	
2008	412 (17.4)	114 (19.3)	298 (16.7)	
2009	349 (14.7)	84 (14.2)	265 (14.9)	
2010	370 (15.6)	99 (16.7)	271 (15.2)	
2011	320 (13.5)	79 (13.3)	241 (13.5)	
2012	299 (12.6)	68 (11.5)	231 (13.0)	
Source of payment				
Self-pay	233 (9.8)	68 (11.5)	165 (9.3)	0.10
Medicare	92 (3.9)	31 (5.2)	61 (3.4)	
Medicaid	223 (9.4)	60 (10.1)	163 (9.2)	
Insurance company	1339 (56.4)	318 (53.7)	1021 (57.4)	
Other/unknown	485 (20.4)	115 (19.4)	370 (20.8)	

^aStorm areas defined as New York City Evacuation Zones (Bronx, Queens, New York, Kings, and Richmond Counties) or as Hurricane Storm Surge Zones in Nassau, Suffolk, and Westchester Counties.


^bP Value for difference between storm and non-storm areas by chi-square test.

substantial heterogeneity by county in terms of the proportion of outpatient FWBD diagnoses among residents of the storm and non-storm areas ($P < 0.01$). There was no difference in the distribution of outpatient FWBD diagnoses and sex ($P = 0.42$), year of diagnosis ($P = 0.25$), or source of payment for medical services ($P = 0.10$). Most outpatient FWBD diagnoses (69.2%) were attributed to ICD-9 008 (intestinal infections due to other organisms; Supplementary Table 1 in the online data supplement).

We also captured 7229 total inpatient FWBD diagnoses reported by hospitals in the study area, between October 28 and November 9, from 2001-2012 (Supplementary Table 2 in the online data supplement and Figure 1). Most inpatient FWBD diagnoses were made in the non-storm area, with the exception of participants 65 years of age and older, for whom more diagnoses were from the storm area (62.9% vs. 59.7%, $P = 0.01$). Inpatient FWBD diagnoses were also more common among blacks ($P < 0.01$) in the storm area (23.1%)

FIGURE 1

Total Number of Outpatient and Inpatient Food and Waterborne Disease Diagnoses in the Study Catchment Area, From October 28 to November 9, 2005-2012 and 2001-2012, Respectively.



relative to the non-storm area (16.5%), whereas diagnoses among “other/unknown” race was more likely in the non-storm area (23.6%) than in the storm area (20.0%). Again, there was substantial heterogeneity by county in terms of the proportion of FWBD diagnoses among residents of the storm and non-storm areas ($P < 0.01$). However, we did not detect differences by sex or year of diagnosis for inpatient FWBD diagnosis. Storm area residents receiving an inpatient FWBD diagnosis were less likely than non-storm area residents ($P < 0.01$) to self-pay (1.6% vs. 3.3%), to use Medicaid (9.8% vs. 11.5%), or to use private insurance (30.2% vs. 32.4%). The majority of inpatient FWBD diagnoses (86.8%) were again attributed to ICD-9 008 (intestinal infections due to other organisms; Supplementary Table 1 in the online data supplement).

Multivariable Analysis

Table 2 describes covariate-adjusted RRs and 95% CIs for outpatient FWBD diagnosis in the storm and non-storm areas, during the post-Sandy period relative to the 2005-2011 pre-Sandy period. Risk was lower among storm area residents during the post-Sandy period relative to the pre-Sandy period (RR = 0.58, 95% CI: 0.46-0.74), with a similar trend, albeit not statistically significant, for residents of non-storm areas (RR = 0.63, 95% CI: 0.38-1.04). The effect appeared somewhat stronger for males (RR = 0.53, 95% CI: 0.38-0.75) than for females (RR = 0.65, 95% CI: 0.45-0.93). In contrast, outpatient FWBD risk was higher in the post-Sandy period for storm area residents 65 years of age and older, relative to the pre-Sandy period (RR = 2.16, 95% CI: 1.11-4.19), adjusted for the impact in the non-storm area (RR = 1.45, 95% CI: 0.64-3.25). This manifested as 5.36 additional outpatient FWBD diagnoses per 100,000 residents aged ≥ 65 years in 2012 relative to 2005-2011. Yet, we captured

only $n = 24$ outpatient FWBD cases aged ≥ 65 years in 2012 and so the effect estimates were imprecise (ie, ICD-9 codes 009 [$n = 9$], 008 [$n = 8$], and 005 [$n = 7$], distributed uniformly across the risk interval, and with greatest frequencies from Nassau [29.2%] and Queens [29.2%] counties). Younger storm area residents were, however, at lower risk post-Sandy (RR = 0.34, 95% CI: 0.18-0.64 for 5-19 years; RR = 0.61, 95% CI: 0.48-0.76 for 20-64 years). In both storm and non-storm areas, residents of New York County (RR = 0.50, 95% CI: 0.29-0.87 and RR = 0.37, 95% CI: 0.21-0.64, respectively) and Queens County (RR = 0.40, 95% CI: 0.20-0.77 and RR = 0.26, 95% CI: 0.18-0.39, respectively) experienced lower risks for outpatient FWBD diagnosis in the post-Sandy period than were experienced in the pre-Sandy period. With few exceptions, other effect estimates, although not statistically significant, suggested reduced risks. Owing to small cell sizes ($n = 1$ case in the storm area), we did not calculate separate estimates for Westchester County. There were no meaningful differences in outpatient FWBD diagnosis effects using a 28-day risk interval in lieu of the 13-day risk interval (Supplementary Table 3 in the online data supplement).

Table 3 describes covariate-adjusted RRs and 95% CIs for inpatient FWBD diagnosis in the storm and non-storm areas, during the pre-Sandy period relative to the 2001-2011 post-Sandy period. We did not detect significant main effects for individuals residing in the storm area. However, non-storm area females were again at a reduced risk (RR = 0.76, 95% CI: 0.66-0.95), as were whites (RR = 0.78, 95% CI: 0.62-0.98), other races (RR = 0.75, 95% CI: 0.63-0.89), and Queens County residents (RR = 0.62, 95% CI: 0.41-0.94). In contrast to outpatient FWBD risks, the risk for inpatient FWBD was lower in the post-Sandy period than in the pre-Sandy period for individuals aged ≥ 65 years, and the effect

TABLE 2

Risk Ratios (95% Confidence Intervals) for Incident Food and Waterborne Disease Outpatient Diagnoses, From October 28 to November 9, 2012, Relative to October 28 to November 9, 2005-2011, Among Residents of Hurricane Sandy Storm Areas and Non-Storm Areas^a

Demographic factor	Storm Area ^b	Non-Storm Area ^c
Overall	0.58 (0.46-0.74)	0.63 (0.38-1.04)
Sex		
Male	0.53 (0.38-0.75)	0.64 (0.38-1.06)
Female	0.65 (0.45-0.93)	0.63 (0.38-1.07)
Age, years		
0-4	0.61 (0.36-1.03)	0.61 (0.30-1.23)
5-19	0.34 (0.18-0.64)	0.46 (0.15-1.47)
20-64	0.61 (0.48-0.76)	0.73 (0.55-0.96)
≥65	2.16 (1.11-4.19)	1.45 (0.64-3.25)
Race		
White	0.67 (0.50-0.89)	0.64 (0.35-1.18)
Black	0.64 (0.42-0.96)	0.62 (0.44-0.88)
Other	0.52 (0.35-0.76)	0.65 (0.30-1.39)
County of residence		
Nassau	0.90 (0.29-2.85)	2.07 (0.98-4.36)
Suffolk	0.97 (0.31-2.98)	0.79 (0.45-1.39)
Bronx	0.51 (0.15-1.71)	1.04 (0.69-1.55)
Kings (Brooklyn)	0.73 (0.45-1.19)	0.72 (0.50-1.04)
New York (Manhattan)	0.50 (0.29-0.87)	0.37 (0.21-0.64)
Queens	0.40 (0.20-0.77)	0.26 (0.18-0.39)
Richmond (Staten Island)	0.66 (0.07-6.65)	1.76 (0.59-5.22)

^aPoisson or negative binomial regression models adjusted for day of the week (short-term trend), year (long-term trend), including a cross product term between pre-/post-Sandy interval and study area, and offset by the 2010 US Census population total. Statistically significant results ($P < 0.05$) in boldface; time-trend effects in storm and non-storm areas not statistically different ($P > 0.05$).

^bStorm area includes those categorized as evacuation zones A, B, and C in Bronx, Kings, New York, Queens, and Richmond counties, and areas categorized as storm surge zones categories 1, 2, 3, and 4 in Nassau and Suffolk Counties.

^cNon-storm areas include those not categorized as storm areas.

was significantly stronger for non-storm areas (RR = 0.68, 95% CI: 0.52-0.89) than for storm areas (RR = 0.89, 95% CI: 0.67-1.18), as assessed by the interaction. We also detected an interaction for residents of Nassau County, in which storm area residents experienced little change for inpatient FWBD risk (RR = 1.06, 95% CI: 0.64-1.75), while non-storm area residents experienced approximately half the risk (RR = 0.53, 95% CI: 0.32-0.89); again, the effect in non-storm area residents was significantly larger than the effect in storm area residents. As for outpatient FWBD, with few exceptions, the remaining effect estimates suggested reduced, although not statistically significant, risks for inpatient FWBD. We were unable to generate stratified estimates for Westchester County, given $n = 1$ case in the storm area.

We also evaluated covariate-adjusted risk ratios for specific FWBD diagnoses during the post-Sandy period relative to the

TABLE 3

Risk Ratios (95% Confidence Intervals) for Incident Food and Waterborne Disease Inpatient Diagnoses, From October 28 to November 9, 2012, Relative to October 28 to November 9, 2001-2011, Among Residents of Hurricane Sandy Storm Areas and Non-Storm Areas^a

Demographic Factor	Storm Area ^b	Non-Storm Area ^c
Overall	0.86 (0.69-1.08)	0.79 (0.66-0.95)
Sex		
Male	0.88 (0.67-1.16)	0.91 (0.74-1.11)
Female	0.85 (0.62-1.16)	0.76 (0.61-0.95)
Age, years		
0-4	1.71 (0.67-4.38)	1.02 (0.57-1.81)
5-19	0.47 (0.12-1.82)	1.17 (0.54-2.53)
20-64	0.80 (0.55-1.17)	0.95 (0.77-1.17)
≥65 ^d	0.89 (0.67-1.18)	0.68 (0.52-0.89)
Race		
White	0.90 (0.67-1.20)	0.78 (0.62-0.98)
Black	0.88 (0.75-1.04)	0.98 (0.72-1.32)
Other	0.73 (0.39-1.39)	0.75 (0.63-0.89)
County of residence		
Nassau ^d	1.06 (0.64-1.75)	0.53 (0.32-0.89)
Suffolk	1.77 (0.86-3.64)	0.95 (0.67-1.36)
Bronx	0.54 (0.23-1.27)	0.77 (0.54-1.11)
Kings (Brooklyn)	0.65 (0.40-1.06)	0.78 (0.52-1.18)
New York (Manhattan)	0.89 (0.51-1.54)	0.61 (0.36-1.04)
Queens	0.69 (0.34-1.39)	0.62 (0.41-0.94)
Richmond (Staten Island)	2.31 (0.72-7.45)	0.62 (0.27-1.44)

^aPoisson or negative binomial regression models adjusted for day of the week (short-term trend), year (long-term trend), including a cross product term between pre-/post-Sandy interval and study area, and offset by the 2010 US Census population total. Statistically significant results ($P < 0.05$) in boldface.

^bStorm area includes those categorized as evacuation zones A, B, and C in Bronx, Kings, New York, Queens, and Richmond counties, and areas categorized as storm surge zones categories 1, 2, 3, and 4 in Nassau, Suffolk, and Westchester Counties.

^cNon-storm areas include those not categorized as storm areas.

^dTime trend effects in the storm and non-storm areas significantly different ($P < 0.05$).

pre-Sandy period (Table 4). We detected a lower risk for outpatient diagnosis of ICD-9 008 (intestinal infections due to other organisms) in storm areas (RR = 0.50, 95% CI: 0.34-0.74) and for inpatient diagnosis in non-storm areas (RR = 0.73, 95% CI: 0.57-0.94). Owing to small numbers of cases, we were unable to generate stratified effect estimates for ICD-9 003 (other Salmonella infections, outpatient) or for ICD-9 004 (Shigellosis), ICD-9 006 (Amebiasis), or ICD-9 007 (other protozoal intestinal diseases).

DISCUSSION

We conducted a retrospective time-trend analysis of incident FWBD diagnoses to assess the health impact of Hurricane Sandy-associated flooding in the greater NYC area and compared effects in storm area residents to non-storm area residents. Overall, our data suggest lower risks for FWBD in the post-Sandy period relative to the pre-Sandy period, both

TABLE 4

Risk Ratios (95% Confidence Intervals) for Specific Food and Waterborne Disease Outpatient and Inpatient Diagnoses, From October 28 to November 9, 2012, Relative to October 28 to November 9, 2005-2011 or 2001-2011, Respectively, Among Residents of Hurricane Sandy Storm Areas and Non-Storm Areas^a

Case Diagnosis	Storm Area ^b	Non-Storm Area ^c
Outpatient (ICD-9 code)		
Other food poisoning (bacterial) (005)	1.01 (0.54-1.89)	1.02 (0.63-1.66)
Intestinal infections due to other organisms (008)	0.50 (0.34-0.74)	0.54 (0.29-1.01)
Ill-defined intestinal infections (009)	0.70 (0.29-1.72)	0.88 (0.47-1.64)
Inpatient (ICD-9 code)		
Other Salmonella infections (002-003)	2.15 (0.54-8.53)	0.93 (0.49-1.78)
Other food poisoning (bacterial) (005)	0.90 (0.23-3.61)	0.85 (0.54-1.35)
Intestinal infections due to other organisms (008)	0.89 (0.69-1.14)	0.73 (0.57-0.94)
Ill-defined intestinal infections (009)	0.59 (0.30-1.18)	0.98 (0.74-1.29)

^aAbbreviation: ICD-9, International Classification of Disease, 9th Revision. Poisson or negative binomial regression models adjusted for day of the week (short-term trend), year (long-term trend), including a cross product term between pre-/post-Sandy interval and study area, and offset by the 2010 US Census population total. Statistically significant results ($P < 0.05$) in boldface. There were an insufficient number of case diagnoses to assess outpatient "other Salmonella infections" (ICD-9 002-003), or for "shigellosis" (ICD-9 004), "amebiasis" (ICD-9 006), and "other protozoal intestinal diseases" (ICD-9 007). Time-trend effects in storm and non-storm areas not statistically different ($P > 0.05$).

^bStorm area includes those categorized as evacuation zones A, B, and C in Bronx, Kings, New York, Queens, and Richmond counties, and areas categorized as storm surge zones categories 1, 2, 3, and 4 in Nassau, Suffolk, and Westchester Counties.

^cNon-storm areas include those not categorized as storm areas.

for residents of storm areas and for residents of non-storm areas. This pattern was generally robust to stratification by sex, race, and county of residence. The change appears to have been due in large part to reduced risks for diagnosis of ICD-9 008 (intestinal infections due to other organisms), without concomitant increases in related diagnoses. Still, the post-Sandy outpatient FWBD risk for storm area residents aged ≥ 65 years was more than double the pre-Sandy risk, consistent with reported vulnerabilities among the aging.²³ However, the effect estimate was based on very few exposed cases and so coupled to multiple conducted statistical tests it may reflect a chance observation and should be interpreted with caution. In contrast, inpatient FWBD risk was decreased during the post-Sandy period for age ≥ 65 years and with a stronger effect for non-storm area residents. It seems unlikely for a "competing risk" to account for this pattern, in that persons who would be hospitalized as inpatients in the pre-Sandy period were treated as outpatient in the post-Sandy period. Rather, it is tempting to speculate that risk among older residents increased for more moderate FWBD (ie, not severe enough to merit hospitalization), whereas risks for more severe FWBD (ie, requiring hospitalization) decreased in the post-Sandy period.

Our overall results are consistent with a building consensus that the risk for infectious disease is typically low following hydro-meteorological disasters in highly developed, high-income

nations; in particular, in the absence of large-scale population displacement.^{22,27} A previous NYC study of 42 reportable diseases, including 16 gastrointestinal and related illnesses after Hurricane Sandy also reported that there was no increased risk for hurricane-affected relative to unaffected areas.²⁸ Furthermore, no increased risk for gastrointestinal distress was reported following earlier hurricane events in Connecticut, Mississippi, and Rhode Island.²⁹ However, a large outbreak of norovirus-associated gastroenteritis was reported from 2 refugee facilities, housing more than 125,000 displaced New Orleans residents in Houston, Texas, possibly due to sewage exposure from flooding instigated by Hurricane Katrina.^{21,30} Although NYC opened 73 evacuation shelters in preparation for Hurricane Sandy, most closed in less than 48 hours, and only 12 continued to operate for 7 or more days.³¹ Still, an emergency shelter-based outbreak of gastrointestinal disease, likely norovirus, was reported 5 to 6 days after the storm,³¹ underscoring the vulnerability of displaced persons. Illnesses such as this shelter-based outbreak are unlikely to be captured in the SPARCS dataset. The overall limited and mostly short-term population displacement associated with Hurricane Sandy in the greater NYC area may also have mitigated the risk for FWBD during the post-Sandy interval.^{31,32}

The counterintuitively lower post-Sandy FWBD risks detected in our study might reflect the impact of public health measures and increased vigilance on the part of catchment

area residents. For example, on October 29, 2012, NYC raised chlorine levels and implemented ultraviolet disinfection in response to high turbidity detected in a nearby Westchester County drinking water reservoir; the turbid source was quickly replaced.³³ In collaboration with private suppliers, NYC also provided more than 2.1 million meals ready to eat, 1,000,000 bottles of water, and 719,000 hot lunches and dinners to affected residents, between November 1, 2012, and January 31, 2013.¹ NYC is home to more than 460 combined sewage outflows (CSOs), which discharge raw domestic sewage into the Hudson River and the New York Harbor during heavy precipitation events.^{7,34} In Massachusetts, even small-scale flooding is associated with a higher rate of emergency room gastrointestinal complaints, up to 4 days following heavy rainfall events in areas with CSOs.³⁵ Heavy rainfall was also correlated to diminished water quality in urban areas serviced by CSOs in a recent Canadian study.³⁶ Despite the high density of CSOs in the study catchment areas, and substantial flooding associated with the Hurricane Sandy storm surge, we detected increased FWBD risk for only older residents of storm areas, intimating the general effectiveness of disaster management during this event.

A recent systematic review concluded that little evidence exists to support the effectiveness of basic water, sanitation, and hygiene interventions (“WASH”) following humanitarian crises in poorly developed, or in low- and middle-income nations.³⁷ However, such interventions appear to be more effective in well-developed and high-income nations, including in the United States, possibly related to a lower likelihood for population displacement following an event, more extensive and widespread interventions, more effective population uptake, and more intensive surveillance activities.³⁸ In fact, trauma-related morbidity, rather than illness-related morbidity, is reported most frequently in association with extreme weather events occurring in high-income nations such as the United States.³⁹ Our study results reinforce this perspective, and furthermore underscore the likely value of public health planning activities in advance of extreme weather events.

Our study results may have been impacted by a dynamic study population, given evacuations and hospital closings in the catchment area during the storm. Approximately 375,000 individuals resided in mandatory NYC evacuation zones, and 5 hospital facilities evacuated.¹ Yet, only 37% of the residential population appear to have complied with the evacuation order, half of them having left before Hurricane Sandy made landfall.³² It is possible that our study results reflect diminished medical services utilization or case under-ascertainment in the post-Sandy interval, with only the most serious cases visiting hospitals for treatment, or use of more distant treatment facilities by evacuees or due to closings. Still, we identified similar decreased FWBD risks for residents of storm areas and nearby non-storm areas, and we did not detect increased risks for inpatient FWBD. It is also unlikely

that differential use of health care facilities outside of NYS, during the pre-Sandy and post-Sandy periods, by both storm area and non-storm area residents, accounts for our results. Admission of NYS residents to hospitals outside of NYS is reported to and captured by the SPARCS database. Still, the total number of outpatient FWBD diagnoses in the post-Sandy period was consistent with a decreasing trend for the preceding years and with similar numbers for inpatient diagnosis and so any impact was likely to have been modest.

Notwithstanding a large number of incident cases overall, including the pre-Sandy and post-Sandy study intervals, and linkage to residential street billing address, several important factors limit our study results. Our risk period was defined as the 13 FEMA emergency incident period days, ending on November 9, 2012, to capture case diagnosis we believe most likely to have been associated with Sandy and the mandatory evacuation order issued on October 28, 2012. We compared this interval with the same interval, for the 7 (outpatient) or 11 (inpatient) preceding years in a time-trend analysis,⁴⁰ adjusting for year to accommodate long-term trends, and using 2010 Census data for the denominator.² We also compared the time-trend to nearby “non-storm” areas to accommodate year-to-year variability in seasonal factors or admission policies. We may have missed cases with incubation periods longer than the 13-day risk window; however, we had similar outpatient results using a 28-day risk period. With few exceptions (eg, amebiasis), the incubation period for most FWBD ranges from <2 to 8 days.^{41,42} We captured few cases of protozoal infection, and so were unable to assess the time trend. A future investigation focused on protozoal infection is needed to more definitively assess the risk. Furthermore, to reduce outcome misclassification we captured only clinically confirmed FWBD diagnoses as study outcomes, rather than use a syndromic case definition. Finally, our ecological exposure assessment strategy, in which residents were classified as “storm area” vs. “non-storm area” may have introduced misclassification, such as individuals living in non-storm areas in close proximity to and affected similarly to those in storm areas, or individuals living in unaffected subsections of storm areas. Still, we would expect such misclassification to be unrelated to diagnosis and to have biased the results towards the null hypothesis.

CONCLUSIONS

Overall, our study results suggest that most FWBD risks decreased in the post-Sandy period relative to the pre-Sandy period, for residents of storm-affected and non-storm affected areas. In the absence of bias, these results suggest that disaster management planning and mitigation, and public health interventions, were effective in preventing FWBD associated with flooding due to the storm surge. However, the risk for outpatient FWBD diagnosis may have increased for residents aged ≥ 65 years in the post-Sandy period, raising the possibility for a vulnerable population to which targeted strategies

should be developed for mitigating risk during future extreme weather events.

About the Authors

Departments of Environmental Health Sciences (Dr Bloom, Ms Palumbo, Ms Lauper, Dr Lin) and Epidemiology and Biostatistics (Dr Bloom, Ms Saiyed, Dr Lin), University at Albany, State University of New York, Rensselaer, New York and Sinai Urban Health Institute, Chicago, Illinois (Ms Saiyed).

Correspondence and reprint requests to Michael S. Bloom, Department of Environmental Health Sciences, School of Public Health Rm #157, One University Place, Rensselaer, NY 12144 (e-mail: mbloom@albany.edu).

Acknowledgments

The authors thank Mr Cris Pantea and Dr Seema Nayak from the Center for Environmental Health, New York State Department of Health; Dr Srishti Shrestha from the National Institute of Environmental Health Sciences; and Mr Ziqiang Lin and Mr Wangjian Zhang from the University at Albany, SUNY, for their support on data management, analysis, and results table preparation. We also thank the staff and administrators from Bureau of Environmental and Occupational Epidemiology, Center for Environmental Health, for their support on this project.

Funding

This work was supported by Grant #1U01 TP000566-01 from the Centers for Disease Control and Prevention (CDC). The content is solely the responsibility of the authors and does not necessarily represent the official views of the CDC.

Supplementary Material

To view supplementary material for this article, please visit <http://dx.doi.org/10.1017/dmp.2016.85>

Published online: May 16, 2016.

REFERENCES

- Gibbs LI, Holloway CFH. Hurricane Sandy After Action: Report and Recommendations to Mayor Michael S. Bloomberg. http://www.nyc.gov/html/recovery/downloads/pdf/sandy_aar_5.2.13.pdf 2013. Published May 2013. Accessed April 29, 2016.
- US Census Bureau. 2010 Census Data. <http://www.census.gov/2010census/data/>. Published 2012. Accessed May 1, 2015.
- Casey-Lockyer M, Heick RJ, Mertzluft CE, et al. Deaths associated with Hurricane Sandy - October-November 2012. *MMWR Morb Mortal Wkly Rep.* 2013;62:393-397.
- Rose JB, Epstein PR, Lipp EK, et al. Climate variability and change in the United States: potential impacts on water- and foodborne diseases caused by microbiologic agents. *Environ Health Perspect.* 2001;109:211-221. <http://dx.doi.org/10.2307/3435011>.
- McMichael AJ. Extreme weather events and infectious disease outbreaks. *Violence.* 2015;6(6):543-547. <http://dx.doi.org/10.4161/21505594.2014.975022>.
- Centers for Disease Control and Prevention. Keep Food and Water Safe After a Disaster or Emergency. <http://www.bt.cdc.gov/disasters/foodwater/facts.asp>. Published 2014. Accessed October 16, 2015.
- Patz JA, Vavrus SJ, Uejio CK, et al. Climate change and waterborne disease risk in the Great Lakes Region of the US. *Am J Prev Med.* 2008;35(5):451-458. <http://dx.doi.org/10.1016/j.amepre.2008.08.026>.
- Lin S, Lu Y, Justino J, et al. What happened to our environment and mental health as a result of Hurricane Sandy [published online ahead of print]? *Disaster Med Public Health Prep.* <http://dx.doi.org/10.1017/dmp.2016.51>.
- Epidemiologic assessment of the impact of four hurricanes - Florida, 2004 (Reprinted from MMWR, vol 54, pg 693-697, 2005). *JAMA.* 2005; 294(12):1484-1485. <http://dx.doi.org/10.1001/jama.294.12.1484>.
- Watson JT, Gayer M, Connolly MA. Epidemics after natural disasters. *Emerg Infect Dis.* 2007;13(1):1-5. <http://dx.doi.org/10.3201/eid1301.060779>.
- Alderman K, Turner LR, Tong S. Floods and human health: a systematic review. *Environ Int.* 2012;47:37-47. <http://dx.doi.org/10.1016/j.envint.2012.06.003>.
- Schmid D, Lederer I, Much P, et al. Outbreak of norovirus infection associated with contaminated flood water, Salzburg, 2005. *Euro Surveill.* 2005;10:E050616.050613-E050616.050613.
- Qadri F, Khan AI, Faruque ASG, et al. Enterotoxigenic Escherichia coli and Vibrio cholerae diarrhea, Bangladesh, 2004. *Emerg Infect Dis.* 2005;11(7):1104-1107. <http://dx.doi.org/10.3201/eid1107.041266>.
- Schwartz BS, Harris JB, Khan AI, et al. Diarrheal epidemics in Dhaka, Bangladesh, during three consecutive floods: 1988, 1998, and 2004. *Am J Trop Med Hyg.* 2006;74:1067-1073.
- Liu ZD, Ding GY, Zhang Y, et al. Analysis of risk and burden of dysentery associated with floods from 2004 to 2010 in Nanning, China. *Am J Trop Med Hyg.* 2015;93(5):925-930. <http://dx.doi.org/10.4269/ajtmh.14-0825>.
- Fredrick T, Ponnaiah M, Murhekar MV, et al. Cholera outbreak linked with lack of safe water supply following a tropical cyclone in Pondicherry, India, 2012. *J Health Popul Nutr.* 2015;33:31-38.
- Kondo H, Seo N, Yasuda T, et al. Post-flood-infectious diseases in Mozambique. *Prehosp Disaster Med.* 2002;17(03):126-133. <http://dx.doi.org/10.1017/S1049023X0000340>.
- Ahmed Z, Khan AA, Nisar N. Frequency of infectious diseases among flood affected people at district Rajanpur, Pakistan. *Pak J Med Sci.* 2011;27:866-869.
- Wade TJ, Sandhu SK, Levy D, et al. Did a severe flood in the midwest cause an increase in the incidence of gastrointestinal symptoms? *Am J Epidemiol.* 2004;159(4):398-405. <http://dx.doi.org/10.1093/aje/kwh050>.
- Wade TJ, Lin CJ, Jagai JS, et al. Flooding and emergency room visits for gastrointestinal illness in Massachusetts: a case-crossover study. *PLoS One.* 2014;9(10):e110474. <http://dx.doi.org/10.1371/journal.pone.0110474>.
- Yee EL, Palacio H, Atmar RL, et al. Widespread outbreak of norovirus gastroenteritis among evacuees of Hurricane Katrina residing in a large "megashelter" in Houston, Texas: lessons learned for prevention. *Clin Infect Dis.* 2007;44(8):1032-1039. <http://dx.doi.org/10.1086/512195>.
- Kouadio IK, Aljunid S, Kamigaki T, et al. Infectious diseases following natural disasters: prevention and control measures. *Expert Rev Anti Infect Ther.* 2012;10(1):95-104. <http://dx.doi.org/10.1586/eri.11.155>.
- Wu J, Xiao J, Li T, et al. A cross-sectional survey on the health status and the health-related quality of life of the elderly after flood disaster in Bazhong city, Sichuan, China. *BMC Public Health.* 2015;15(1):163. <http://dx.doi.org/10.1186/s12889-015-1402-5>.
- Callaghan WM, Rasmussen SA, Jamieson DJ, et al. Health concerns of women and infants in times of natural disasters: lessons learned from Hurricane Katrina. *Matern Child Health J.* 2007;11(4):307-311. <http://dx.doi.org/10.1007/s10995-007-0177-4>.
- Renne JL, Sanchez TW, Litman T. Carless and special needs evacuation planning: a literature review. *J Plann Lit.* 2011;26(4):420-431. <http://dx.doi.org/10.1177/0885412211412315>.
- Flanagan BE, Gregory EW, Hallisey EJ, et al. A social vulnerability index for disaster management. *J Homel Secur Emerg Manage.* 2011;8(1):3.
- Ivers LC, Ryan ET. Infectious diseases of severe weather-related and flood-related natural disasters. *Curr Opin Infect Dis.* 2006;19(5):408-414. <http://dx.doi.org/10.1097/01.qco.0000244044.85393.9e>.
- Greene SK, Wilson EL, Konty KJ, et al. Assessment of reportable disease incidence after Hurricane Sandy, New York City, 2012. *Disaster Med Public Health Prep.* 2013;7(05):513-521. <http://dx.doi.org/10.1017/dmp.2013.98>.
- CDC. Epidemiologic notes and reports hurricanes and hospital emergency-room visits - Mississippi, Rhode Island, Connecticut. *MMWR Morb Mortal Wkly Rep.* 1986;34:765-770.
- Murray KO, Kilborn C, DesVignes-Kendrick M, et al. Emerging disease syndromic surveillance for Hurricane Katrina evacuees seeking shelter in

- Houston's Astrodome and Reliant Park complex. *Public Health Rep.* 2009;124:364-371.
31. Ridpath AD, Bregman B, Jones L, et al. Challenges to implementing communicable disease surveillance in New York City evacuation shelters after Hurricane Sandy, November 2012. *Public Health Rep.* 2015;130:48-53.
 32. Brown S, Parton H. *Evacuation in New York City During Hurricanes Irene and Sandy*. New York: New York City Department of Health and Mental Hygiene; 2014.
 33. New York City Department of Environmental Protection (DEP). New York City 2012 Drinking Water Supply and Quality Report. <http://www.nyc.gov/html/dep/pdf/wsstate12.pdf>. Published 2013. Accessed April 29, 2016.
 34. Riverkeeper. Combined Sewage Overflows (CSOs). <http://www.riverkeeper.org/campaigns/stop-polluters/sewage-contamination/cso/>. Published 2015. Accessed October 19, 2015.
 35. Jagai JS, Li Q, Wang S, et al. Extreme precipitation and emergency room visits for gastrointestinal illness in areas with and without combined sewer systems: an analysis of Massachusetts data, 2003-2007. *Environ Health Perspect.* 2015;123:873-879.
 36. Gooré Bi E, Monette F, Gasperi J. Analysis of the influence of rainfall variables on urban effluents concentrations and fluxes in wet weather. *J Hydrol (Amst)*. 2015;523:320-332. <http://dx.doi.org/10.1016/j.jhydrol.2015.01.017>.
 37. Ramesh A, Blanchet K, Ensink JHJ, Roberts B. Evidence on the effectiveness of water, sanitation, and hygiene (WASH) interventions on health outcomes in humanitarian crises: a systematic review. *PLoS One*. 2015;10(9):e0124688. <http://dx.doi.org/10.1371/journal.pone.0124688>.
 38. Brown L, Murray V. Examining the relationship between infectious diseases and flooding in Europe. *Disaster Health.* 2013;1(2):117-127. <http://dx.doi.org/10.4161/dish.25216>.
 39. Doocy S, Dick A, Daniels A, et al. The human impact of tropical cyclones: a historical review of events 1980-2009 and systematic literature review. *PLoS Curr.* 2013 Apr 16. doi: 10.1371/currents.dis.2664354a5571512063ed29d25ffbc74.
 40. Stroup DF, Wharton M, Kafadar K, et al. Evaluation of a method for detecting aberrations in public-health surveillance data. *Am J Epidemiol.* 1993;137:373-380.
 41. Lee RM, Lessler J, Lee RA, et al. Incubation periods of viral gastroenteritis: a systematic review. *BMC Infect Dis.* 2013;13:446. doi: 10.1186/1471-2334-13-446.
 42. Heymann D. ed. *Control of Communicable Diseases Manual*. 19th ed. Washington, DC: American Public Health Association; 2009.