

**Introduction** Further research is still needed to demonstrate the benefits of animal-assisted therapy (AAT) for specific participant profiles, such as children with behavioural disorders.

**Objectives** We wanted to find out if AAT could be considered an efficient therapeutic strategy for the treatment of children with behavioural disorders.

**Aims** We wanted to study the effects of a preestablished AAT program on the behaviour of children with emotional and behavioural issues in 6 different reception centres for children under government guardianship.

**Methods** Forty-five children (12 to 17 years old) with emotional and behavioural issues participated in a 14-session AAT program. Behavioural measures were those routinely scored as part of therapy; an observational report of 3 different problematic behaviours (such as impulsivity, lack of social skills or lack of personal recognition) was made twice a week for each child (with a score of frequency and intensity). A pre- and post-treatment "global behaviour score" was calculated for each child, as an average value of the 3 problematic behaviours measured during the month pre-treatment and the month post-treatment.

**Results** The 45 participants attended, on average, 72.8% of AAT sessions. Independent behaviour scores differed between the pre- and post-intervention evaluations ( $n = 135$  behaviours) (Wilcoxon test;  $P < 0.0001$ ). Based on the global behaviour score for each child ( $n = 45$ ), significant change was found between pre- and post-intervention evaluations (Wilcoxon test;  $P = 0.0011$ ).

**Conclusions** Our results suggest AAT could be a beneficial intervention for children with behavioural issues in terms of program adherence and behaviour improvement.

**Disclosure of interest** The authors have not supplied their declaration of competing interest.

<http://dx.doi.org/10.1016/j.eurpsy.2017.01.418>

#### EV0090

### Exercise for depression in adolescents: A meta-analysis of randomised controlled trials

T. Carter<sup>1,\*</sup>, I. Morres<sup>2</sup>, O. Meade<sup>3</sup>, C. Patrick<sup>3</sup>

<sup>1</sup> The University of Nottingham, School of Health Sciences, Nottingham, United Kingdom

<sup>2</sup> University of Thessaly, School of Physical Education and Sport Science, School of Physical Education and Sport Science, Greece

<sup>3</sup> University of Nottingham, School of Health Sciences, Nottingham, United Kingdom

\* Corresponding author.

**Background** The reported lifetime prevalence of depression in adolescents has increased drastically in recent decades. Depression in this population can be particularly damaging and can frequently have a long-lasting adverse impact. The common treatment approach includes psychological therapy and/or anti-depressant medication. However, the efficacy of these approaches, both singularly and combined, is far from conclusive. Recently, there have been an increased number of studies investigating the effect of exercise and physical activity on adolescent depression, however, despite this increased attention, there are no recent reviews and meta-analyses synthesising such studies.

**Aim** The review examines the treatment effect of exercise on depression symptoms for adolescents aged 13–17 years of age.

**Methods** A systematic search of seven electronic databases identified relevant randomised controlled trials. Screening, data extraction and trial methodological quality were undertaken by two independent researchers. Standardized mean differences were used for pooling post-intervention depressive symptom scores.

**Results** Eleven trials met the inclusion criteria, eight of which provided the necessary data for calculation of standardized effect size. Exercise showed a statistically significant moderate overall

effect on depressive symptom reduction. Sensitivity analyses were also conducted and will be presented.

**Discussion** Exercise and physical activity appear to improve depression symptoms in adolescents, especially in clinical samples, suggesting that exercise may be a useful treatment strategy for adolescents with depression.

**Conclusion** Despite the positive findings, large clinical trials that adequately minimise bias are required for firmer conclusions on the effectiveness of exercise as an antidepressant treatment.

**Disclosure of interest** The authors have not supplied their declaration of competing interest.

<http://dx.doi.org/10.1016/j.eurpsy.2017.01.419>

#### EV0091

### Sixteen-year population-based cohort study of main class International Classification of Diseases associated with psychiatric disorders in a sample under the age of two years

D. Cawthorpe

The University of Calgary, Psychiatry/Community Health, Calgary, Canada

**Introduction** This paper illustrates the use of cohort data from a population to describe the early life prevalence and odds ratios (ORs) of the main classes of International Classification of Diseases (ICD) associated with any mental disorder arising at any time during the 16 year study period.

**Objectives** The main ICD disorder classes were examined in relation to psychiatric disorders over 16 years in a cohort under the age of two years between April 1st, 1993, and January 1st, 1995.

**Aims** To demonstrate the utility of studying the complete profile of associated diagnoses over time in a population cohort.

**Methods** The total number of individuals under the age of two years before 1995 ( $n = 17,603$ ) were tallied within each main class of ICD disorder by year and expressed as ORs of those with and without any 16-year psychiatric disorder.

**Results** The greatest annual rates observed in the early years of life were for the following main ICD classes of disease: respiratory system, sense organs, symptoms signs ill-defined conditions, no diagnosis, injury poisoning, and skin subcutaneous tissue disorders. These disorders also had the highest ORs in early life given the presence of a mental disorder at any time during the study period.

**Discussion** Knowing the early life main class diagnoses associated with psychiatric disorders could guide both basic science research as well as early intervention social and health investment policies.

**Disclosure of interest** The authors have not supplied their declaration of competing interest.

<http://dx.doi.org/10.1016/j.eurpsy.2017.01.420>

#### EV0092

### Drawing in autistic spectrum disorder children

S. Cepedello\*, A.S. Adrian, G.B. LAura, M.T. Eduardo, D.U.V. Nieves, D.L.R. Henar, Á.A. Aldara

Hospital Clinico Universitario, Psiquiatria, Valladolid, Spain

\* Corresponding author.

**Introduction** Drawing can be a tool to complete the psychodiagnose process, especially in children with expression and verbal problems.

The autistic spectrum disorder (ASD) children have problems to describe their feelings and emotions, they can provide us many information drawing their own world.

**Objective** The purpose of this study was to discover the psychopathology of children with ASD through their drawings.