


First identification of SARS-CoV-2 lambda (C.37) variant in Southern Brazil

Priscila Lamb Wink PhD^{1,2,3} , Fabiana Caroline Zempulski Volpato MSc^{1,2,4}, Francielle Liz Monteiro PhD^{1,2,5}, Julia Biz Willig MSc^{1,2}, Alexandre Prehn Zavascki MD, PhD^{1,6,7}, Afonso Luís Barth PhD^{1,2,3,4} and Andreza Francisco Martins PhD^{1,3,8,9}

¹LABRESIS – Laboratório de Pesquisa em Resistência Bacteriana, Hospital de Clínicas de Porto Alegre, Porto Alegre, Rio Grande do Sul, Brazil, ²Laboratório de Diagnóstico de SARS-CoV-2, Hospital de Clínicas de Porto Alegre, Rio Grande do Sul, Brazil, ³Programa de Pós-Graduação em Ciências Farmacêuticas, Universidade Federal do Rio Grande do Sul, Porto Alegre, Rio Grande do Sul, Brazil, ⁴Programa de Pós-Graduação em Ciências Médicas, Universidade Federal do Rio Grande do Sul, Porto Alegre, Rio Grande do Sul, Brazil, ⁵Laboratório de Diagnóstico Molecular, Universidade Franciscana, Santa Maria, Rio Grande do Sul, Brazil, ⁶Infectious Diseases Service, Hospital de Clínicas de Porto Alegre, Porto Alegre, Rio Grande do Sul, Brazil, ⁷Internal Medicine Department, Universidade Federal do Rio Grande do Sul, Porto Alegre, Rio Grande do Sul, Brazil, ⁸Bioinformatic Core, Hospital de Clínicas de Porto Alegre, Rio Grande do Sul, Brazil and ⁹Departamento de Microbiologia, Imunologia e Parasitologia, Instituto de Ciências Básicas, Universidade Federal do Rio Grande do Sul, Brazil

To the Editor—Since the emergence of the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) in December 2019, new lineages of this virus have been described. The recent emergence of SARS-CoV-2 variants of interest (VOIs) and variants of concern (VOCs) with potentially increased transmissibility and reduced sensitivity to antibody neutralization may affect the efficacy of strategies to control the coronavirus 2019 (COVID-19) pandemic.¹ A novel VOI within the B.1.1.1 lineage (termed C.37), which was designated by the World Health Organization (WHO) as “the lambda variant” in June 15, 2021, was detected in Peru in August 2020. It has been identified in 26 countries in America, Europe, and Oceania, drawing international attention for its rapid spread.^{2–5} Despite the global dissemination of the lambda variant, in Brazil this lineage was reported only in São Paulo state in February 2021.⁵ Here, we describe the first report of the SARS-CoV-2 lambda variant in Southern Brazil. This study was approved by the Ethics Committees from Hospital de Clínicas de Porto Alegre (CAAE: 30767420.2.0000.5327).

A young male who had been in Argentina initiated respiratory symptoms compatible with viral infection while returning to his town, which borders Argentina, in Rio Grande do Sul, the southernmost Brazilian state. On day 4 after the beginning of the symptoms, he tested positive for SARS-CoV-2 by a point-of-care test. On the following day, he was admitted to a local hospital. Due to the worsening of symptoms, he was transferred to the intensive care unit of Hospital de Clínicas de Porto Alegre, a tertiary-care, COVID-19 referral hospital in the state, 2 days after hospital admission. Oro/nasopharyngeal swabs were collected, and real-time reverse transcription-PCR testing for 2 genes of the nucleocapsid protein (N1 and N2) for SARS-CoV-2 was performed,⁶ confirming the SARS-CoV-2 infection. The specimen (203_LABRESIS) was submitted for whole-genome sequencing (WGS) as part of an epidemiological study. Sequencing libraries were prepared using the CleanPlex SARS-CoV-2 panel protocol (Paragon Genomics, Hayward, CA) for target enrichment and library preparation, following the manufacturer’s instructions ([https://www.paragongenomics.com/wp-content/uploads/2020/](https://www.paragongenomics.com/wp-content/uploads/2020/03/UG4001-01_-CleanPlex-SARS-CoV-2-Panel-User-Guide.pdf)

[03/UG4001-01_-CleanPlex-SARS-CoV-2-Panel-User-Guide.pdf](https://www.paragongenomics.com/wp-content/uploads/2020/03/UG4001-01_-CleanPlex-SARS-CoV-2-Panel-User-Guide.pdf)). The resulting libraries were sequenced using MiSeq (Illumina, San Diego, CA). Consensus sequences were generated by the QIASeq SARS-CoV-2 pipeline (QIAGEN CLC Genomics Workbench 21, Germantown, PA) with high-quality whole-genome sequence (average coverage 3,142; <2% Ns; 29,873 Kb). The specimen 203_LABRESIS was classified as the C.37 variant using the Phylogenetic Assignment of Named Global Outbreak Lineages (Pangolin) software tool (version 3.0.2),³ and the sequence was deposited in the GISAID database (no. EPI_ISL_2617911).⁵

The lambda variant (C.37) is defined by a deletion in the ORF1a gene (Δ 3675–3677), which is also present in the alpha (B.1.1.7, United Kingdom), beta (B.1.351, South African), and gamma (P.1, Brazil) VOCs. In addition, the lambda variant displays a novel deletion and multiple nonsynonymous mutations in the spike gene (Δ 246–252, G75V, T76I, L452Q, F490S, D614G, and T859N).^{2,4} The mutations L452Q and F490S are mapped in the receptor-binding domain (RBD) region and the F490S has been associated with reduced susceptibility to antibody neutralization.¹ The isolate 203_LABRESIS presented all 8 of the C.37-defining lineage mutations described above in addition to other 19 mutations that have already been described in members of this lineage (Supplementary Material online).^{2,4,7} The 203_LABRESIS sequence was aligned with high quality and coverage-complete genome sequences from Brazil (n = 1), Argentina (n = 16), Chile (n = 24), and Peru (n = 26) that were available in the EpiCoV database in GISAID. We used them to perform a comparative phylogenetic tree. The 203_LABRESIS sequence clustered together with another C.37 variant sequence from Argentina that was also identified in February 2021 (Supplementary Material online).

The high prevalence of the lambda variant has been described particularly in South American countries such as Chile, Peru, Ecuador, and Argentina, where this new VOI is associated with substantial rates of community transmission.² The critical health-care system situation and recent reports of increased deaths in these countries are thought to be associated with the increasing prevalence of the lambda variant.^{8–10}

Notably, on June 15, 2021, the lambda variant was deemed a VOI by the World Health Organization.² Whether the lambda variant is more transmissible or more pathogenic than other variants or whether it is able to evade the effects of available vaccines

Author for correspondence: Priscila Lamb Wink, E-mail: pris_farma@yahoo.com.br
Cite this article: Wink PL, *et al.* (2022). First identification of SARS-CoV-2 lambda (C.37) variant in Southern Brazil. *Infection Control & Hospital Epidemiology*, 43: 1996–1997, <https://doi.org/10.1017/ice.2021.390>

remains unknown. The novel S: Δ 246-252 deletion and additional mutations in the spike protein should be considered to understand their effects on viral fitness and host interaction. This first report of the SARS-CoV-2 lambda variant in Southern Brazil raises concern regarding a possible dissemination of this lineage in the region. Moreover, considering that this VOI has rapidly spread in Peru, Ecuador, Chile, and Argentina, we believe that this lambda variant has considerable potential to become a VOC.

Supplementary material. To view supplementary material for this article, please visit <https://doi.org/10.1017/ice.2021.390>

Acknowledgments. We thank the staff of Laboratório de Diagnóstico de SARS-CoV-2 as well as the staff of the Hospital Infection Committee of our institution (Hospital de Clínicas de Porto Alegre) for providing data used in this study. This research was conducted using the Hospital de Clínicas de Porto Alegre Biobank resources (application no. GPPG 2020-0163).


Financial support. This study was funded by FAPERGS (20/2551-0000265-9), National Institute of Antimicrobial Resistance Research - INPRA (MCTI/CNPq/CAPEF/FAPs nº 16/2014) and by Fundo de Incentivo a Pesquisa e Eventos do Hospital de Clínicas de Porto Alegre (FIPE/HCPA) (Project no. 2020-0163). P.L.W. and F.C.Z.V. were supported by a grant from the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES).

Conflicts of interest. A.P.Z., A.L.B., and A.F.M. are research fellows of the National Council for Scientific and Technological Development (CNPq), Ministry of Science and Technology, Brazil. A.P.Z. has received a research grant from Pfizer unrelated to this work. All other authors have no conflict to declare.

References

- Liu Z, Van Blargan LA, Louis-Marie Bloyet, *et al.* Identification of SARS-CoV-2 spike mutations that attenuate monoclonal and serum antibody neutralization. *Cell Host Microbe* 2021;29:477–488.
- Technical report. COVID-19 weekly epidemiological update, edition 44. World Health Organization website. https://reliefweb.int/sites/reliefweb.int/files/resources/20210615_Weekly_Epi_Update_44.pdf. Published 2021. Accessed June 15, 2021.
- Pangolin: lineage assignment in an emerging pandemic as an epidemiological tool. [v1.1.21]. Pangolin website. <http://pangolin.cog-uk.io/>. Published 2021. Accessed June 15, 2021.
- Latif AA, Mullen JL, Alkuzweny M, *et al.* C.37 lineage report. Outbreak Info website. <https://outbreak.info/situation-reports?pango=C.37>. Published 2021. Published June 2021. Accessed June 15, 2021.
- Tracking of variants. GISAID website. www.gisaid.org/hcov19-variants. Accessed June 15, 2021.
- Wink PL, Volpato F, de Lima-Morales, *et al.* RT-qPCR half reaction optimization for the detection of SARS-CoV-2. 1. medRxiv 2021. doi: [10.1101/2021.05.19.21257470](https://doi.org/10.1101/2021.05.19.21257470).
- Romero PE, Dávila-Barclay A, Gonzáles L, *et al.* C.37: Novel lineage expanding in Peru and Chile, with a convergent deletion in the ORF1a gene (Δ 3675-3677) and a novel deletion in the Spike gene (Δ 246-252, G75V, T76I, L452Q, F490S, T859N). <https://virological.org/t/novel-sublineage-within-b-1-1-1-currently-expanding-in-peru-and-chile-with-a-convergent-deletion-in-the-orf1a-gene-3675-3677-and-a-novel-deletion-in-the-spike-gene-246-252-g75v-t76i-l452q-f490s-t859n/685>. Published April 2021. Accessed October 20, 2021.
- INS confirma presencia de variante C-37 del coronavirus en Perú, 25 Mayo 2021. Peru Ministerio de Salud: Instituto Nacional de Salud website. <https://web.ins.gob.pe/index.php/es/prensa/noticia/minsa-ins-confirma-presencia-de-variante-c-37-del-coronavirus-en-peru> Published 2021. Published 2021. Accessed June 15, 2021.
- Vigilancia de variantes de SARS-CoV-2 en CABA, Provincia de Buenos Aires, Córdoba, Entre Ríos, Neuquén y Santa Fe. Argentina government website. <https://www.argentina.gob.ar/noticias/vigilancia-de-variantes-de-sars-cov-2-en-caba-provincia-de-buenos-aires-cordoba-entre-rios>. Published 2021. Accessed June 15, 2021.
- Latif AA, Mullen JL, Alkuzweny M, *et al.* Chile mutation report. Outbreak Info website. <https://outbreak.info/location-reports?loc=CHL&pango=C.37>. Published 2021. Accessed June 15, 2021.

Early detection of the SARS-CoV-2 P.1 variant in Rio Grande do Sul, Brazil: a case report

Fabiana Caroline Zempulski Volpato MSc^{1,2,3} , Priscila Lamb Wink PhD^{1,2,4}, Francielle Liz Monteiro PhD^{1,2,5}, Julia Biz Willig MSc^{1,2}, Tarsila Vieceli MD⁶, Andreza Francisco Martins Ph.D.,^{1,4,7,8} Alexandre Prehn Zavascki MD, PhD^{1,6,9} and Afonso Luís Barth Ph.D.^{1,2,3,4}

¹LABRESIS – Laboratório de Pesquisa em Resistência Bacteriana, Hospital de Clínicas de Porto Alegre, Porto Alegre, Rio Grande do Sul, Brasil, ²Laboratório de Diagnóstico de SARS-CoV-2, Hospital de Clínicas de Porto Alegre, Rio Grande do Sul, Brasil, ³Programa de Pós-Graduação em Ciências Médicas, Universidade Federal do Rio Grande do Sul, Porto Alegre, Rio Grande do Sul, Brasil, ⁴Programa de Pós-Graduação em Ciências Farmacêuticas, Universidade Federal do Rio Grande do Sul, Porto Alegre, Rio Grande do Sul, Brasil, ⁵Laboratório de Diagnóstico Molecular, Universidade Franciscana, Santa Maria, Rio Grande do Sul, Brasil, ⁶Infectious Diseases Service, Hospital de Clínicas de Porto Alegre, Porto Alegre, Rio Grande do Sul, Brasil, ⁷Bioinformatic Core, Hospital de Clínicas de Porto Alegre, Rio Grande do Sul, Brasil, ⁸Departamento de Microbiologia, Imunologia e Parasitologia, Instituto de Ciências Básicas, Universidade Federal do Rio Grande do Sul, Brasil and ⁹Internal Medicine Department, Universidade Federal do Rio Grande do Sul, Porto Alegre, Rio Grande do Sul, Brazil

Author for correspondence: Fabiana Caroline Zempulski Volpato, E-mail: fabiana_volpato@yahoo.com.br

Cite this article: Volpato FCZ, *et al.* (2022). Early detection of the SARS-CoV-2 P.1 variant in Rio Grande do Sul, Brazil: a case report. *Infection Control & Hospital Epidemiology*, 43: 1997–1999, <https://doi.org/10.1017/ice.2021.387>

© The Author(s), 2021. Published by Cambridge University Press on behalf of The Society for Healthcare Epidemiology of America.

To the Editor—In December 2019, a new coronavirus, severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), emerged in China and was associated with coronavirus disease 2019 (COVID-19). Since it was first described, multiple lineages of SARS-CoV-2 have been identified worldwide. Due to the increased transmissibility of the new variants, the initially