

ARTICLE

A rainbow Dirac theorem for loose Hamilton cycles in hypergraphs

Amarja Kathapurkar^{1*}, Patrick Morris^{2†} and Guillem Perarnau^{2,3‡}

¹University of Birmingham, Birmingham, UK, ²Universitat Politècnica de Catalunya (UPC), Barcelona, Spain, and ³Centre de Recerca Matemàtica, Bellaterra, Spain

Corresponding author: Patrick Morris; Email: pmorrismaths@gmail.com

(Received 13 January 2025; revised 15 October 2025; accepted 15 October 2025)

Abstract

A meta-conjecture of Coulson, Keevash, Perarnau, and Yepremyan [12] states that above the extremal threshold for a given spanning structure in a (hyper-)graph, one can find a rainbow version of that spanning structure in any suitably bounded colouring of the host (hyper-)graph. We solve one of the most pertinent outstanding cases of this conjecture by showing that for any $1 \leq j \leq k - 1$, if G is a k -uniform hypergraph above the j -degree threshold for a loose Hamilton cycle, then any globally bounded colouring of G contains a rainbow loose Hamilton cycle.

Keywords: Spanning structures; Hypergraphs; extremal graph theory; local lemma

2020 MSC Codes: Primary: 05C35, 05D40; Secondary: 05C65

1. Introduction

A seminal theorem of Dirac [16] states that any n -vertex graph G with $\delta(G) \geq n/2$ contains a Hamilton cycle. This inspired many further results exploring the optimal minimum degree conditions for certain spanning structures in a host (hyper-)graph, and this area, sometimes referred to as ‘Dirac theory’, is a cornerstone of modern extremal combinatorics. The field has flourished in recent decades due to powerful tools being developed to tackle these questions, such as the regularity method and absorption. In graphs, this has led to a deep understanding of the full picture.

In hypergraphs, the situation is considerably more complex. This is, in part, due to the various ways in which one can generalise the graph case. For example, when generalising Dirac’s theorem to hypergraphs, one has a range of choices as to which minimum degree condition is considered and what type of Hamilton cycle is desired. Indeed, for a k -uniform hypergraph G (k -graph for short), one can consider

$$\delta_j(G) := \min \left\{ |\{e \in E(G) : T \subset e\}| : T \in \binom{V(G)}{j} \right\},$$

*Research supported by EPSRC Research grant EP/R034389/1.

†Research supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) Walter Benjamin programme - project number 504502205 and by the European Union’s Horizon Europe Marie Skłodowska-Curie grant RAND-COMB-DESIGN - project number 101106032 .

‡Research supported by the grants RED2022-134947-T, PID2023-147202NB-I00, PCI2024-155080-2 and the Programme Severo Ochoa y María de Maeztu por Centros y Unidades de Excelencia en I&D (CEX2020-001084-M), all of them funded by MICIU/AEI/10.13039/501100011033.

for $1 \leq j \leq k-1$. Likewise with Hamilton cycles, one can consider a cyclic ordering of the vertices of G and require that each edge of the Hamilton cycle occupies k consecutive vertices in the ordering and every pair of consecutive edges intersect in precisely ℓ vertices for some $1 \leq \ell \leq k-1$. Such a Hamilton cycle is called a *Hamilton ℓ -cycle*, and when $\ell = 1$, we refer to it as a *loose cycle*, whilst the case $\ell = k-1$ is referred to as a *tight cycle*. Note that if an n -vertex k -graph G has a Hamilton ℓ -cycle, then one necessarily has that $(k-\ell)|n$.

In hypergraphs, our understanding of minimum degree thresholds is far from complete, even in simple cases such as perfect matchings (see e.g. [38]). For Hamilton cycles, we make the following definition to capture the relevant extremal minimum degree conditions.

Definition 1.1 (Hypergraph Hamilton cycle thresholds). *For $1 \leq j, \ell \leq k-1$, we denote by $\delta_j^k(\ell)$ the smallest real number $\delta \geq 0$ such that for any $\varepsilon > 0$ and $n \in (k-\ell)\mathbb{N}$ sufficiently large, any n -vertex k -graph G with $\delta_j(G) \geq (\delta + \varepsilon)n^{k-j}$ contains a Hamilton ℓ -cycle.*

Determining these threshold values has proven to be a considerable challenge. For example, in the case of loose Hamilton cycles (when $\ell = 1$), a case that has been of particular interest, we only know the location of $\delta_j^k(1)$ when $j = k-2$ or $j = k-1$. For $j = k-1$, the so-called *codegree threshold*, this was established by Kühn and Osthus [28] when $k = 3$ and in general by Keevash, Kühn, Mycroft, and Osthus [26] and, independently, Hàn and Schacht [23]. For $j = k-2$, this was again solved first in the 3-uniform case, by Buß, Hàn, and Schacht [8], and then for general k by Bastos, Mota, Schacht, Schnitzer, and Schenburger [15]. For other values of ℓ , the results are also sporadic; we refer the reader to the survey [38] for a comprehensive review.

Beyond determining extremal degree thresholds, another fruitful line of research in Dirac theory has been to show *robustness* for these thresholds. The aim here is to show that when a (hyper-)graph is above the minimum degree threshold (we will informally refer to such (hyper-)graphs as being ‘Dirac’) with respect to a given spanning structure, the Dirac (hyper-)graph is in fact *robust* with respect to containing that spanning structure. This includes considerations such as the existence of many spanning structures and resilience to random edge deletions. We refer to the nice survey of Sudakov [35] and the references therein for an introduction to these results in the context of Dirac’s condition for Hamilton cycles. In this paper, we consider a notion of robustness related to finding *rainbow* spanning structures in any bounded edge colouring of the Dirac (hyper-)graph. This is motivated by the classical study of rainbow spanning structures in certain colourings of graphs.

1.1 Rainbow-spanning structures

A subgraph H of an edge-coloured graph G is *rainbow* if each of the edges of H is a different colour. Rainbow subgraphs appeared early on in combinatorics via connections to design theory with several beautiful conjectures being posed. Perhaps the most famous such conjecture, known as the Ryser-Brualdi-Stein conjecture [7, 33, 34] states that every $n \times n$ Latin square has a transversal of size at least $n-1$ and one of size n when n is odd. The first part of this (establishing the existence of transversals of size $n-1$) has only recently been solved by Montgomery [30]. Translating to colourings of graphs, the Ryser-Brualdi-Stein conjecture is equivalent to the assertion that one can always find a rainbow matching of size $n-1$ (or n when n is odd) in any properly coloured $K_{n,n}$ with n colours.

From a graph-theoretic perspective one can ask more generally what conditions on a colouring of a host graph guarantee the existence of a rainbow (almost) spanning structure of interest. A colouring being proper is equivalent to saying that the colouring is *t -locally bounded* (at each vertex we see every colour at most t times) with $t = 1$. One can also then consider *globally bounded* conditions where we instead bound the size of each colour class in the whole (hyper-)graph. The first result of this kind was due to Erdős and Spencer [19], who showed that any colouring of $K_{n,n}$ with at most $\frac{n}{16}$ edges of each colour contains a rainbow perfect matching. Another early example of interest was due to Erdős and Stein (see [18]) who asked whether there is some constant $c > 0$

such that any colouring of K_n with at most cn edges of each colour contains a rainbow Hamilton cycle. This was then explicitly conjectured by Hahn and Thomassen [22] and, after several results towards the conjecture, was solved by Albert, Frieze and Reed [1]. A generalisation to hypergraph Hamilton cycles was then given by Dudek, Frieze and Ruciński [17]. There has been a wealth of similar results studying different spanning structures.

1.2 Rainbow structures in Dirac (hyper-)graphs

The majority of results concerning rainbow spanning substructures in bounded (and proper) colourings have focused on the case where the host graph is a complete (hyper-)graph or complete bipartite graph. When considering other possible host graphs, Dirac graphs arise naturally. This perspective was first considered by Cano, Perarnau, and Serra [9] who showed that one can find a rainbow Hamilton cycle in any globally $o(n)$ -bounded colouring of G when G is either an n -vertex graph or a balanced bipartite graph with n vertices in each part, and such that G has minimum degree $\delta(G) \geq (1 + o(1))\frac{n}{2}$. The asymptotic minimum degree condition was then replaced to give an exact minimum degree condition $\delta(G) \geq \frac{n}{2}$ by Coulson and Perarnau, first in the bipartite case [13] and then in the non-bipartite case [14] as in Dirac's original theorem. These results thus give evidence of robustness for the extremal thresholds for Hamilton cycles. Note also that in the bipartite case, these results can be seen as a direct strengthening of the result of Erdős and Spencer [19], allowing for host graphs that are not complete (at the expense of a worse constant for the boundedness).

Further examples of these types of results came from Coulson, Keevash, Perarnau, and Yepremyan [12] who proved that (asymptotically) above the minimum degree for a given (hyper-)graph F -factor, one finds a rainbow F -factor in any suitably bounded colouring, and from Glock and Joos [21] who gave a rainbow version of the famous blow-up lemma [27], which allowed them to give results of this flavour in considerable generality for graphs, in particular providing a rainbow version of the bandwidth theorem [6]. We remark that a nice feature of the work of [12] is that they could establish such a result, even in cases where the minimum degree threshold has not yet been determined.

All of these results provide evidence of a general phenomenon and caused Coulson, Keevash, Perarnau, and Yepremyan [12] to explicitly give the “meta-conjecture” that once one is above the extremal threshold for a given spanning structure, rainbow copies of that structure can be found in any suitably bounded colouring of the Dirac graph. Our main result provides further evidence for this conjecture by establishing that this is the case for loose Hamilton cycles in hypergraphs.

Theorem 1.2 (Main Theorem). *Let $1 \leq j < k \in \mathbb{N}$ and let $\varepsilon > 0$ be arbitrary. Then there exists $\mu > 0$ such that for any sufficiently large $n \in (k-1)\mathbb{N}$, we have that if G is an n -vertex k -graph with $\delta_j(G) \geq (\delta_j^k(1) + \varepsilon)n^{k-j}$, the following holds. For any colouring $\chi: E(G) \rightarrow \mathbb{N}$ of G such that*

$$(B) \quad |\chi^{-1}(c)| \leq \mu n^{k-1} \text{ for all } c \in \mathbb{N},$$

we have that there is a rainbow loose Hamilton cycle in G coloured by χ .

Theorem 1.2 provides a first generalisation of the result of Coulson and Perarnau [14] to the hypergraph setting and strengthens the previously mentioned work of Dudek, Frieze, and Ruciński [17] who proved the statement in the case that the host hypergraph G is complete. Note that the bound in (B) is tight up to the choice of μ as some bound of the order of n^{k-1} is required in order to have sufficiently many colours available for a rainbow loose Hamilton cycle. Note also that, as with the previous results on F -factors in hypergraphs due to Coulson, Keevash, Perarnau, and Yepremyan [12], our result works even for the values of j where the threshold value $\delta_j^k(1)$ is not yet known.

We remark that Theorem 1.2 and the aforementioned result [12] on F -factors are the only results of this kind, proving robustness via rainbow copies in bounded colourings, in the realm of hypergraphs. It is not surprising that progress has been much slower when dealing with higher

uniformities as many of the tools used in the graph setting [13, 21], in particular the regularity method, face barriers for hypergraphs in Dirac theory, as discussed above. Our proof of Theorem 1.2 follows the same scheme as that of [12] for F -factors, appealing to the local lemma and using random samples to perform switches. The main barrier in applying these methods in the setting of Hamilton cycles is that the structure we are trying to find is *connected*. This prevents us from being able to locally adapt a copy of our desired spanning structure independently from the structure as a whole. These local adaptations are at the core of the switching method used in [12] and, at a high level, we overcome this obstacle by using absorption techniques to piece back together loose Hamilton cycles after local alterations. Making this work requires new ideas to overcome technical hurdles at many parts of the proof and we also require recent work of Alvarado, Kohayakawa, Lang, Mota, and Stagni [3] who showed the existence of an absorption scheme when the location of the threshold is not known. Another technical contribution of our work is to remove the need for a local bound on the colouring. Indeed, the result of Coulson, Keevash, Perarnau, and Yepremyan [12] found F -factors under the assumption (B) as well as an additional constraint that every $(k-1)$ -set lies in at most μn edges of each colour. This local boundedness condition is superfluous when $k=2$ (the graph case), but it is not implied by the global condition (B) when $k \geq 3$. The authors of [12] justify this extra condition by showing that for certain F , for example when F is a clique with more than one edge, the condition is in fact *necessary*. However, in the setting of loose Hamilton cycles, Theorem 1.2 shows that the global condition (B) suffices. Within the proof scheme of [12] and our work here, removing the need for the local condition requires new ideas to maintain the existence of a rainbow structure. We develop a novel strategy using random partitions and deterministic adjustments to the partition and we remark that these ideas can be used to strengthen the result of [12] to be able to drop the local boundedness condition on the colouring to get F -factors when F is *linear*, that is, when any pair of edges in F intersect in at most one vertex. We discuss this further in the concluding remarks, Section 7.

1.3 Organisation

In the next section, we collect some notation and terminology as well as some probabilistic tools that we will use throughout. In Section 3, we then provide a proof overview and reduce the proof of Theorem 1.2 to three main lemmas which are proven in Sections 4, 5 and 6. Finally in Section 7, we discuss potential future directions of research.

2. Preliminaries

2.1 Notation and terminology

For k -graphs H and G , we use the notation $H \subset G$ to indicate that H is a *subhypergraph* (or *subgraph* for short) of G . A *loose path* (or simply path, for short) of *length* t in a k -graph G is a subgraph P of G defined by a collection of t edges $e_1, \dots, e_t \in E(G)$ such that there is some ordering $v_1, \dots, v_{t(k-1)+1}$ of $V(P) \subset V(G)$ with $e_i = \{v_{1+(i-1)(k-1)}, \dots, v_{k+(i-1)(k-1)}\}$ for $i = 1, \dots, t$. The vertices $V^E(P) := \{v_1, v_{t(k-1)+1}\}$ are referred to as the *endvertices* of the path.

A loose Hamilton cycle H comes with a cyclic order on $E(H)$ such that the only intersecting edges of H are pairs of consecutive edges in the ordering (which intersect in exactly one vertex). For each loose Hamilton cycle H of G , we will act under the convention that there is a fixed orientation of this order of $E(H)$ and will refer to an *increasing* path, of length t say, from an edge $e \in E(H)$ to be the path defined by taking e and the $t-1$ edges that follow e when traversing the loose Hamilton cycle according to the orientated order on $E(H)$.

For a k -graph G and $S \subset V(G)$ with $|S| \leq k$, we define the set $E_G(S) \subseteq E(G)$ of edges *incident to* S to be

$$E_G(S) := \{e \in E(G) : S \subseteq e\},$$

and we define the *degree* of S in G to be

$$\deg_G(S) = |E_G(S)|.$$

We write simply $E(S)$ and $\deg(S)$ if the k -graph G is clear from context. We will also need neighbourhoods and degrees defined relative to vertex subsets. For a k -graph G and vertex subsets $S, W \subset V(G)$ with $|S| \leq k$, we define the set $E_G(S; W) \subseteq E(G)$ to be

$$E_G(S; W) := \{e \in E(G) : S \subseteq e, e \setminus S \subset W\},$$

and we define the *degree* of S in G relative to W to be

$$\deg_G(S; W) = |E_G(S; W)|.$$

We write simply $E(S; W)$ and $\deg(S; W)$ if the k -graph G is clear from context.

2.2 Probabilistic tools

In this section we collect various probabilistic tools that will be used in the proof of Theorem 1.2. We start with a general version of the (lopsided) local lemma, see [2, Lemma 5.1.1] and the remarks following the lemma.

Lemma 2.1 (The lopsided local lemma). *Suppose \mathcal{A} is a set of events in a finite probability space, Γ is a graph with $V(\Gamma) = \mathcal{A}$ and there are real numbers $\{x_A : A \in \mathcal{A}\}$ with $0 \leq x_A < 1$ for all $A \in \mathcal{A}$. Suppose further that for any $A \in \mathcal{A}$ and subset $\mathcal{A}' \subseteq \mathcal{A} \setminus \{A\}$ such that $AA' \notin E(\Gamma)$ for all $A' \in \mathcal{A}'$ and $\mathbb{P}[\cap_{A' \in \mathcal{A}'} \overline{A'}] > 0$, we have that*

$$\mathbb{P}[A | \cap_{A' \in \mathcal{A}'} \overline{A'}] \leq x_A \prod_{B: AB \in E(\Gamma)} (1 - x_B). \quad (2.1)$$

Then with positive probability none of the events $A \in \mathcal{A}$ occur.

Remark 2.2. *The fact that we only have a condition for families \mathcal{A}' for which $\mathbb{P}[\cap_{A' \in \mathcal{A}'} \overline{A'}] > 0$ is not mentioned in [2, Section 5.1] (or elsewhere in the literature). However such a condition is necessary for the probability in (2.1) to be well-defined. This omission presumably stems from the fact that Lemma 2.1 is in fact proven in [2] under the stronger assumption that A is mutually independent from all of the events $\{A' : AA' \in E(\Gamma)\}$ and $\mathbb{P}[A]$ satisfies the inequality in (2.1). Therefore one can bypass the issue of having $\mathbb{P}[\cap_{A' \in \mathcal{A}'} \overline{A'}] = 0$ by using the mutual independence to deduce that $\mathbb{P}[A | \cap_{A' \in \mathcal{A}'} \overline{A'}] = \mathbb{P}[A]$ and bounding $\mathbb{P}[A]$. Strictly speaking, there is still a potential issue here with $\mathbb{P}[A | \cap_{A' \in \mathcal{A}'} \overline{A'}]$ being well-defined but an inspection of the proof of [2, Lemma 5.1.1] shows that one can add an extra induction over the size of $|\mathcal{A}'|$ which will show that $\mathbb{P}[\cap_{A' \in \mathcal{A}'} \overline{A'}] > 0$ for all $\mathcal{A}' \subseteq \mathcal{A}$ (and ultimately \mathcal{A} itself) and so imposing (2.1) only for \mathcal{A}' with $\mathbb{P}[\cap_{A' \in \mathcal{A}'} \overline{A'}] > 0$ is justified.*

In fact, it will be easier for us to work with a slight modification of Lemma 2.1, for which we make the following definition.

Definition 2.3. *Let \mathcal{A} be a set of events in a finite probability space. Suppose Γ is a graph with $V(\Gamma) = \mathcal{A}$ and $\mathbf{p} \in [0, 1/2]^{\mathcal{A}}$. We call Γ a \mathbf{p} -dependency graph for \mathcal{A} if for every $A \in \mathcal{A}$ and $\mathcal{A}' \subseteq \mathcal{A} \setminus \{A\}$ such that $AA' \notin E(\Gamma)$ for all $A' \in \mathcal{A}'$ and $\mathbb{P}[\cap_{A' \in \mathcal{A}'} \overline{A'}] > 0$, we have $\mathbb{P}[A | \cap_{A' \in \mathcal{A}'} \overline{A'}] \leq p_A$.*

We now deduce the following corollary of Lemma 2.1 which we will use in applications. This result is already known in the literature but we add its proof for completeness.

Corollary 2.4. *Under the setting of Definition 2.3, if $\sum \{p_B : AB \in E(\Gamma)\} \leq 1/4$ for all $A \in \mathcal{A}$ then with positive probability none of the events in \mathcal{A} occur.*

Proof. For each $A \in \mathcal{A}$, we set $x_A = 2p_A$ and need to show that the condition (2.1) holds for such values. So fix some $A \in \mathcal{A}$ and some $\mathcal{A}' \subseteq \mathcal{A}$ with $\mathbb{P}[\cap_{A' \in \mathcal{A}'} \overline{A'}] > 0$ and such that $AA' \notin E(\Gamma)$ for all $A' \in \mathcal{A}'$. Then

$$x_A \prod_{B: AB \in E(\Gamma)} (1 - x_B) \geq x_A \left(1 - \sum_{B: AB \in E(\Gamma)} x_B\right) = 2p_A \left(1 - 2 \sum_{B: AB \in E(\Gamma)} p_B\right) \geq p_A \geq \mathbb{P}[A | \cap_{A' \in \mathcal{A}'} \overline{A'}],$$

as required. \square

We will frequently use concentration inequalities for random variables. The first such inequality, Chernoff's inequality [11] (see also [24, Corollaries 2.3 and 2.4]), deals with the case of binomial random variables.

Theorem 2.5 (Chernoff inequality). *Let X be the sum of a set of mutually independent Bernoulli random variables and let $\lambda = \mathbb{E}[X]$. Then for any $0 < \delta < \frac{3}{2}$, we have that*

$$\mathbb{P}[X \geq (1 + \delta)\lambda] \leq e^{-\delta^2 \lambda / 3} \quad \text{and} \quad \mathbb{P}[X \leq (1 - \delta)\lambda] \leq e^{-\delta^2 \lambda / 2}.$$

Moreover, if $x \geq 7\lambda$, then $\mathbb{P}[X \geq x] \leq e^{-x}$.

The following well known bound gives a lower bound on the probability that a binomial random variable hits its expected value. The form we give here can be found in [12, Lemma 3.6].

Lemma 2.6. *Let X be binomially distributed with parameters (n, p) and suppose that $\lambda := \mathbb{E}[X] = np \in \mathbb{N}$ is an integer with $\lambda = o(\sqrt{n})$. Then $\mathbb{P}[X = \lambda] \geq 1/(4\sqrt{\lambda})$.*

We will also be interested in concentration inequalities when dealing with random variables that are not mutually independent. Let $\{X_i\}_{i \in \mathcal{I}}$ be a family of random variables on a common probability space. A (strong) *dependency graph* for $\{X_i\}_{i \in \mathcal{I}}$ is any (2-uniform) graph Γ with $V(\Gamma) = \mathcal{I}$ and such that if A and B are disjoint subsets of \mathcal{I} and $\{e \in E(\Gamma) : e \cap A, e \cap B \neq \emptyset\} = \emptyset$, then the families $\{X_i\}_{i \in A}$ and $\{X_i\}_{i \in B}$ are independent of each other. The following inequality is a version of Suen's inequality [36] given by Janson [25] (see also [24, Theorem 2.23]) and it provides a bound for the lower tail in the case that there is a strong dependency graph which is sufficiently sparse.

Lemma 2.7 (Suen's inequality). *Let $\{I_i\}_{i \in \mathcal{I}}$ be a finite family of indicator random variables with some strong dependency graph Γ . Let $X := \sum_{i \in \mathcal{I}} I_i$, $\lambda := \mathbb{E}(X)$ and write $i \sim j$ if $ij \in E(\Gamma)$. Moreover, let $\Delta_X := \sum_{(i,j) \in \mathcal{I}^2: i \sim j} \mathbb{E}[I_i I_j]$ and $\delta_X = \max_{i \in \mathcal{I}} \sum_{k \in \mathcal{I}: k \sim i} \mathbb{P}[I_k = 1]$. Then for any $0 \leq t \leq \lambda$,*

$$\mathbb{P}[X \leq \lambda - t] \leq \exp \left(- \min \left(\frac{t^2}{4(\lambda + \Delta_X)}, \frac{t}{6\delta_X} \right) \right). \quad (2.2)$$

Finally we need the following inequality of Talagrand [37], which we give in the following form, see [29, Section 10.1].

Lemma 2.8 (Talagrand's Inequality). *Let $c, r > 0$ and X be a non-negative random variable determined by n independent trials satisfying the following conditions:*

- (*c*-Lipschitz) *changing the outcome of one trial can affect the value of X by at most c ; and*
- (*r*-certifiable) *for any $s > 0$, if $X \geq s$ then there is a set of at most rs trials whose outcomes certify that $X \geq s$.*

Then for any $0 \leq t \leq \lambda := \mathbb{E}[X]$, we have that

$$\mathbb{P} \left[|X - \lambda| > t + 60c\sqrt{r\lambda} \right] \leq 4e^{-\frac{t^2}{8c^2 r \lambda}}.$$

3. Proof overview

In the remainder of the paper (with the exclusion of the concluding remarks in Section 7), we fix $1 \leq j < k \in \mathbb{N}$. For convenience, we also collect the following hierarchy of constants which is adhered to throughout;

$$1/n \ll \mu \ll \gamma \ll 1/t \ll \beta \ll \varepsilon, 1/k. \quad (3.1)$$

By this we mean that given any $\varepsilon > 0$ and $2 \leq k \in \mathbb{N}$, we can choose constants from right to left so that all required relations with these constants in our proof are satisfied. Indeed any statement concerning these parameters in our proofs will be true under conditions of the form $c' \leq f(c_1, \dots, c_\ell)$ where c' is below c_1, \dots, c_ℓ in the hierarchy. In order to highlight when we are appealing to this hierarchy, we will state that we are using that $c' \ll c_1, \dots, c_\ell$. We will also fix the following key constants

$$T := (t(k-1) + 1), \quad \tilde{m} := t^{100k}, \quad m := T \cdot \tilde{m} \quad \text{and} \quad M := T \cdot m = T^2 \cdot \tilde{m}. \quad (3.2)$$

throughout. The definition of \tilde{m} has some flexibility with the important thing being that it is much larger than t but still bounded by some polynomial in t . The definitions of T , m and M are more rigid with it being crucial that $m/\tilde{m} = M/m = T$. Note that due to how the hierarchy is defined, we can assume that $\gamma \ll 1/M, 1/m, 1/\tilde{m}, 1/T$ and likewise with all constants to the left of γ in the hierarchy. Finally, we will assume that $n \in (k-1)\mathbb{N}$ and fix some n -vertex k -graph G and a colouring $\chi : E(G) \rightarrow \mathbb{N}$ satisfying the assumptions of Theorem 1.2.

Our proof will work through an application of the (lopsided) Lovász local lemma (Lemma 2.1/Corollary 2.4), which is a powerful tool for finding rainbow spanning structures in (hyper)graphs [1, 12–14, 17, 19]. Our probability space will be a choice of a uniformly random loose Hamilton cycle H_* in G , noting that the existence of one such cycle is guaranteed by the fact that G has minimum degree above the extremal threshold. Our bad events \mathcal{A} will encode the possibility of having two edges of the same colour in H_* and thus our application of the local lemma will imply that with positive probability, the random loose Hamilton cycle H_* is rainbow.

In order to appeal to the local lemma, we need to bound from above probabilities of the form $\mathbb{P}[A \mid \cap_{A' \in \mathcal{A}'} A']$ for certain $\mathcal{A}' \subset \mathcal{A}$. Ignoring the conditioning of these events momentarily, we need an upper bound on the probability of some bad event A occurring, where A encodes the appearance of two edges of the same colour appearing in H_* . The key idea to estimate such probabilities is to use *switchings*. Broadly speaking, we do a double-counting argument of the pairs of loose Hamilton cycles (H, H') of G such that H satisfies A , H' does not satisfy it, and H' can be obtained from H by reshuffling a small number of disjoint subpaths. Exhibiting many such pairs for each H in which the bad event A occurs will allow us to conclude an upper bound on $\mathbb{P}[A]$. The collection of subpaths that we will use to reshuffle the loose Hamilton cycle are what we will call a *splitting* of the Hamilton cycle, as in the following definition.

Definition 3.1 (A Hamilton cycle splitting). *Given a loose Hamilton cycle H in G , an m -splitting of H is a collection of vertex-disjoint loose paths $\mathcal{P} = \{P_1, \dots, P_m\}$ in G such that each path P_i is a subgraph of H . We say the splitting is t -balanced if each P_i has length t and we say that the splitting is t -bounded if each P_i has length at most t .*

We define $V(\mathcal{P}) = \cup\{V(P) : P \in \mathcal{P}\}$ to be vertices of \mathcal{P} , $V^E(\mathcal{P}) := \cup\{V^E(P) : P \in \mathcal{P}\}$ to be the endvertices of \mathcal{P} and $V^I(\mathcal{P}) := V(\mathcal{P}) \setminus V^E(\mathcal{P}) = \cup\{V(P) \setminus V^E(P) : P \in \mathcal{P}\}$ to be the interior vertices of \mathcal{P} .

Given a splitting \mathcal{P} of a loose Hamilton cycle H satisfying a bad event A , we will choose new edges of G to piece back together a new loose Hamilton cycle H' which avoids A . The edges that we use to do this will lie on the vertices $V(\mathcal{P})$ of the splitting and furthermore, they will be *transverse*, that is, each new edge will contain at most one vertex from each of the paths in the splitting \mathcal{P} . Using transverse edges in the new loose Hamilton cycle H' will give us control over the colours that feature on new edges which, as we will see shortly, is essential to our approach. In general, we make the following definition of transverse sets and partitions.

Definition 3.2 (Transverse with respect to a splitting). *Given a loose Hamilton cycle H in G and an m -splitting $\mathcal{P} = \{P_1, \dots, P_m\}$ of H , a set of vertices $S \subset V(\mathcal{P})$ is transverse (with respect to \mathcal{P}) if $|S \cap V(P)| \leq 1$ for all $P \in \mathcal{P}$. A partition of $V(\mathcal{P})$ is transverse (with respect to \mathcal{P}) if each part of the partition is transverse.*

Next we come to the notion of a *switching*, which encapsulates how we reshuffle a loose Hamilton cycle H on the vertices of some splitting \mathcal{P} to arrive at a new loose Hamilton cycle H' . The switching will depend on a path P_0 which is a subpath of the original loose Hamilton cycle H that can be thought of as a “troublesome” part of H . In applications, this path P_0 will contain (at least one of) the pair of monochromatic edges described by some bad event A that occurs in H .

Definition 3.3 (Switchings). *Given a loose Hamilton cycle H in G and a loose Hamilton path P_0 of length t with $P_0 \subset H$, we define a (P_0, H) -switching of size m to be a triple $(\mathcal{P}, H', \mathcal{P}')$ such that \mathcal{P} is a t -balanced m -splitting of H with $P_0 \in \mathcal{P}$, H' is a loose Hamilton cycle of G , \mathcal{P}' is a $2t$ -bounded m -splitting of H' , and the following conditions are satisfied.*

- (I) $H[V(G) \setminus V^I(\mathcal{P})] = H'[V(G) \setminus V^I(\mathcal{P}')]$;
- (II) $V(P_0)$ is transverse with respect to \mathcal{P}' .

Condition (I) essentially says that when we switch from H to H' , all of the alteration takes place on the vertices of a splitting \mathcal{P} . Note also that condition (I) necessarily implies that $V^E(\mathcal{P}) = V^E(\mathcal{P}')$. Condition (II) will imply that in the new loose Hamilton cycle H' , the vertices of $V(P_0)$ are far apart. In particular, this will imply that our bad event A in H will no longer occur in the new loose Hamilton cycle H' as at least one of the pair of monochromatic edges described by A will no longer feature in H' .

As previously mentioned, we will use switchings to double count pairs (H, H') of loose Hamilton cycles in G in which some bad event A occurs in H but not in H' . This in turn will lead to upper bounds on the probability $\mathbb{P}[A]$ of A occurring. However, in order to appeal to the lopsided Lovász local lemma (Lemma 2.1), what we actually want to bound is probabilities of the form $\mathbb{P}[A \mid \cap_{A' \in \mathcal{A}'} \overline{A'}]$ for any collection of events \mathcal{A}' that do not lie in the neighbourhood of A in the dependency graph Γ (see Definition 2.3). As usual in the local lemma, in order to be effective, we will define Γ *locally* and so bad events B that are adjacent to A in Γ will be those in which the edges involved in the event B intersect the edges that are involved in the event A . In practice this will translate to having bad events B that are adjacent to A being those that involve edges intersecting the troublesome path P_0 and all other bad events might feature in the set \mathcal{A}' . We will condition on some such collection of events \mathcal{A}' *not* happening and in order to estimate $\mathbb{P}[A \mid \cap_{A' \in \mathcal{A}'} \overline{A'}]$, our double counting will focus on loose Hamilton cycles such that no event in \mathcal{A}' occurs. Thus we assume that no event in \mathcal{A}' occurs in H and our switching will also need to produce some loose Hamilton cycle H' in which no event in \mathcal{A}' occurs. This will be achieved by enforcing that the switching is *feasible* as in the following definition.

Definition 3.4 (Feasible switchings). *Given a loose Hamilton cycle H in G and a loose Hamilton path P_0 of length t with $P_0 \subset H$, we say that a (P_0, H) -switching $(\mathcal{P}, H', \mathcal{P}')$ is feasible if $F' := H'[V(\mathcal{P}') \setminus V(P_0)]$ is rainbow and no edge of F' shares a colour with $H[V(G) \setminus V(P_0)]$.*

Note that in order for a switching to be feasible we do *not* require that H' is fully rainbow outside of $V(P_0)$. Indeed, it could be that there are edges of the same colour in $H'[V(G) \setminus V^I(\mathcal{P}')] = H[V(G) \setminus V^I(\mathcal{P})]$. The key point is that in switching from H to H' , we are not introducing any *new* pairs of monochromatic edges, apart from possibly some that include vertices of $V(P_0)$. As we assume that no event in our collection \mathcal{A}' occurs in H , this will guarantee that none of the events in \mathcal{A}' will occur in H' neither. There is the exception of pairs of monochromatic edges that intersect $V(P_0)$ and might appear in H' (and not in H) but, as alluded to above, we will define our

dependency graph Γ in such a way that these events will be adjacent to A in Γ and thus will not feature in \mathcal{A}' .

We are now in a position to state our first main lemma which precisely shows that if we can find many switchings for any loose Hamilton cycle H and troublesome subpath P_0 , then we will be able to prove the existence of a rainbow loose Hamilton cycle. The proof of Lemma 3.5, which we give in Section 4, will follow from applying the local lemma (Corollary 2.4) and the double counting argument sketched above.

Lemma 3.5 (Many feasible switchings implies a rainbow Hamilton cycle). *Suppose that for every loose Hamilton cycle H in G and loose path P_0 of length t with $P_0 \subset H$, there are at least γn^{m-1} feasible (P_0, H) -switchings of size m . Then G contains a rainbow loose Hamilton cycle in G .*

With Lemma 3.5 in hand, it remains to find many feasible switchings for any loose Hamilton cycle H and subpath P_0 . To find a feasible switching, we first need to find a splitting \mathcal{P} of H that contains P_0 . The next definition captures the desired properties of such a splitting, that will allow us to use it in a feasible switching.

Definition 3.6 (Suitable splittings). *We say an m -splitting \mathcal{P} of a loose Hamilton cycle H is suitable for $P_0 \in \mathcal{P}$ if*

- (i) *for any transverse $S \subset V(\mathcal{P}) \setminus V(P_0)$ with $|S| = k - 1$, there are at most $\varepsilon m/4$ edges in $E_{G[V(\mathcal{P})]}(S)$ that share a colour with an edge in H ;*
- (ii) *for any transverse $X \subset V(\mathcal{P}) \setminus V(P_0)$, there is no pair of edges $e, f \in E(G[X])$ with $\chi(e) = \chi(f)$ and $|e \cap f| = 1$; and*
- (iii) *for any pair of disjoint transverse edges $e, f \in E(G[V(\mathcal{P}) \setminus V(P_0)])$ with $\chi(e) = \chi(f)$, we have that there are two distinct $P_i, P_{i'} \in \mathcal{P}$ such that $P_x \cap e, P_x \cap f \neq \emptyset$ for $x = i, i'$.*

Remark 3.7. *Note that condition (iii) of Definition 3.6 in particular implies that for any transverse $X \subset V(\mathcal{P}) \setminus V(P_0)$, there is no pair of disjoint edges $e, f \in E(G[X])$ with $\chi(e) = \chi(f)$.*

Whilst the conditions for a splitting being suitable are somewhat technical, suffice it to say here that these conditions will guarantee that we can choose new edges (all of which will be transverse with respect to the splitting \mathcal{P}) to contribute to the new loose Hamilton cycle H' in our switching, in such a way that the switching will be feasible (Definition 3.4) and so not create any new unwanted pairs of monochromatic edges.

On top of finding a suitable splitting, we also need to specify how we build our new H' for our splitting and in particular, the endpoints of the new paths in the splitting \mathcal{P}' of H' . This will be given by a rerouting.

Definition 3.8 (Rerouting). *Given a loose Hamilton cycle H in G and an m -splitting $\mathcal{P} = \{P_1, \dots, P_m\}$ of H , a rerouting of \mathcal{P} is a partition of $V^E(\mathcal{P})$ into pairs $\{\{v_a^i, v_b^i\} : i = 1, \dots, m\}$ such that connecting the paths in $H[V(G) \setminus V^I(\mathcal{P})]$ by identifying v_a^i and v_b^i for $i = 1, \dots, m$ results in a (single) loose cycle.*

Recall the definition of a partition being transverse with respect to an m -splitting (Definition 3.2) and note that if we have a t -balanced m -splitting \mathcal{P} and $\mathcal{X} = \{X_1, \dots, X_T\}$ a partition of $V(\mathcal{P})$ that is transverse with respect to \mathcal{P} , then necessarily we have that each part in \mathcal{X} has size exactly $m = T \cdot \tilde{m}$. Given a (suitable) splitting \mathcal{P} of a loose Hamilton cycle H , in order to find a feasible switching $(\mathcal{P}, H', \mathcal{P}')$ which uses \mathcal{P} , we will first find transverse partition \mathcal{X} of $V(\mathcal{P})$ and then define our new loose Hamilton cycle H' and corresponding splitting \mathcal{P}' in such a way that each path in \mathcal{P}' belongs to one part of the partition \mathcal{X} . In particular, this will guarantee that each of the edges that features in H' and not in H will be transverse with respect to \mathcal{P} . The next definition captures what properties we require from the partition of $V(\mathcal{P})$.

Definition 3.9 (Viable partition). *For a t -balanced m -splitting \mathcal{P} of a loose Hamilton cycle H , we say a partition $\mathcal{X} = \{X_1, \dots, X_T\}$ of $V(\mathcal{P})$ is viable if it is transverse with respect to \mathcal{P} and*

- (a) *there exists a rerouting $\{\{v_a^i, v_b^i\} : i = 1, \dots, m\}$ of \mathcal{P} such that for each $h \in [T]$ there are (exactly) \tilde{m} indices $i \in [m]$ such that $\{v_a^i, v_b^i\} \subset X_h$; and*
- (b) $\delta_j(G[X_h]) \geq (\delta_j^k(1) + \frac{\varepsilon}{2})m^{k-j}$ for each $h \in [T]$.

Condition (a) is necessary because we wish to build the paths in \mathcal{P}' so that each path is contained in one part of \mathcal{X} and so we certainly need a rerouting (which dictates the endpoints of the paths in \mathcal{P}') in which pairs of vertices that feature in the rerouting lie in one part of the partition \mathcal{X} . The condition that there are *exactly* \tilde{m} pairs in each part will imply that necessary divisibility conditions are adhered to so that we can cover the vertices of each part with disjoint paths between pairs of the rerouting. The condition (b) will be useful also to imply the existence of these disjoint paths. Indeed, without a minimum degree condition we will have no guarantee that $G[X_h]$ for $h \in [T]$ contains any sort of useful spanning structure for creating H' .

Our next major lemma, proved in Section 5, shows that it in order to find a feasible switching, it is enough to find a suitable switching that admits a viable partition.

Lemma 3.10 (Suitable splittings with viable partitions give feasible switchings). *For a loose Hamilton cycle H in G and a path P_0 of length t with $P_0 \subset H$, if \mathcal{P} is a t -balanced m -splitting of H that is suitable for P_0 and such that $V(\mathcal{P})$ admits a viable partition, then there is some (size m) feasible (P_0, H) -switching $(\mathcal{P}, H', \mathcal{P}')$.*

To prove this lemma, we will build the paths in \mathcal{P}' which define the new loose Hamilton cycle H' , by building paths in each part of the viable partition \mathcal{X} , one at a time. Within each part X_h , at a high level, we use absorption techniques to find the relevant collection of disjoint paths that cover the vertices X_h . In fact, we are able to use a recent result of Alvarado, Kohayakawa, Lang, Mota, and Stagni [3] (which is proven using absorption techniques) as a black box to find our desired paths. Still though, there is some technical work needed to get in to a position where we can apply their result, and we also need to prove that we can build our paths in such a way that the resulting switching is feasible, which we will be able to achieve by appealing to the conditions of \mathcal{P} to be suitable.

Finally then, to complete the proof of Theorem 1.2, it will remain to prove that we can indeed find suitable splittings that admit feasible partitions. This is the content of our last main lemma, Lemma 3.11, which is proven in Section 6 by sampling both a random splitting and a random partition and showing that with positive probability, all the required conditions will hold in these random samples.

Lemma 3.11 (Many suitable splittings with viable partitions). *For a loose Hamilton cycle H in G and a path P_0 of length t with $P_0 \subset H$, there exists γn^{m-1} distinct choices of t -balanced m -splittings \mathcal{P} of H such that \mathcal{P} is suitable for P_0 and $V(\mathcal{P})$ admits a viable partition.*

With these three main lemmas in hand, the proof of Theorem 1.2 is immediate. We conclude this section by spelling it out in detail.

Proof of Theorem 1.2. By Lemma 3.5, it suffices to show that for any loose Hamilton cycle H in G and loose path $P_0 \subset H$ of length t , there are at least γn^{m-1} feasible (P_0, H) -switchings of size m . Lemma 3.10 then implies that it is sufficient to find γn^{m-1} t -balanced m -splittings \mathcal{P} of H with $P_0 \in \mathcal{P}$ and such that \mathcal{P} is suitable for P_0 and admits a viable partition. The existence of these splittings is then given by Lemma 3.11. \square

4. Many feasible switchings imply a rainbow Hamilton cycle

In this section we prove Lemma 3.5, which has a similar proof to that of [12, Lemma 2.2].

Proof of 3.5. Let H_* be a loose Hamilton cycle of G chosen uniformly at random. Recall the hierarchy of the parameters: $1/n \ll \mu \ll \gamma \ll 1/m \ll 1/t$ in (3.1).

For every P_0 loose path of length t in G and every $e_0, f_0 \in E(P_0)$ with $|e_0 \cap f_0| \leq 1$ and $\chi(e) = \chi(f)$ define the events $A(e_0, f_0; P_0) = \{P_0 \subset H_*\}$. Similarly, for every pair of vertex-disjoint loose paths P_0, Q_0 of length t in G and every $e_0 \in E(P_0)$ and $f_0 \in E(Q_0)$ with $\chi(e) = \chi(f)$, define $B(e_0, f_0; P_0, Q_0) = \{P_0, Q_0 \subseteq H_*\}$. Then we define

$$\begin{aligned}\mathcal{A} &= \{A(e_0, f_0; P_0) : \mathbb{P}[A(e_0, f_0; P_0)] > 0\}, \\ \mathcal{B} &= \{B(e_0, f_0; P_0, Q_0) : \mathbb{P}[B(e_0, f_0; P_0, Q_0)] > 0\}, \\ \mathcal{E} &= \mathcal{A} \cup \mathcal{B}.\end{aligned}$$

Note that if none of the events in \mathcal{E} hold, then H_* is rainbow. Indeed, for every pair $e_0, f_0 \in E(H_*)$ we have $|e_0 \cap f_0| \leq 1$ and by avoiding the events in \mathcal{E} , any such pair satisfies $\chi(e) \neq \chi(f)$.

For each $A = A(e_0, f_0; P_0) \in \mathcal{A}$, we write $V(A) := V(P_0)$ and for each $B = B(e_0, f_0; P_0, Q_0)$, we write $V(B) := V(P_0) \cup V(Q_0)$. Consider the graph Γ with $V(\Gamma) = \mathcal{E}$ where for every $E_1, E_2 \in \mathcal{E}$ we have $E_1 E_2 \in E(\Gamma)$ if and only if $V(E_1) \cap V(E_2) \neq \emptyset$. We will choose values $p_A, p_B \in [0, 1/2]$ such that Γ is a \mathbf{p} -dependency graph for \mathcal{E} , where $p_A = p_{\mathcal{A}}$ for all $A \in \mathcal{A}$ and $p_B = p_{\mathcal{B}}$ for all $B \in \mathcal{B}$. For $\mathcal{X} \in \{\mathcal{A}, \mathcal{B}\}$, define

$$d_{\mathcal{X}} = \max_{E \in \mathcal{E}} |\{X \in \mathcal{X} : EX \in E(\Gamma)\}|.$$

To apply Corollary 2.4, it suffices to check that $p_{\mathcal{A}} d_{\mathcal{A}} + p_{\mathcal{B}} d_{\mathcal{B}} \leq 1/4$.

To bound the degrees in Γ , we bound the number of events whose vertex set intersects a given vertex $v \in V(G)$. Recall that $T = t(k-1) + 1$ is the number of vertices of a loose path of length t .

For every $v \in V(G)$, we claim that

$$|A \in \mathcal{A} : v \in V(A)| \leq 2(k+1)T!\mu n^{T-1}. \quad (4.1)$$

Indeed, suppose that $A = A(e_0, f_0; P_0) \in \mathcal{A}$ with $v \in V(A)$. There are at most $(k+1)\mu n^{2k-1}$ pairs of edges $e_0, f_0 \in E(H_*)$ of the same colour with $|e_0 \cap f_0| \leq 1$; indeed, there are at most n^k choices for e_0 , at most $(k+1)$ for $e_0 \cap f_0$ and at most μn^{k-1} for f_0 , by (B). From them, at most $(k+1)\mu n^{2k-2}$ pairs satisfy $v \in e_0 \cup f_0$ (as one can assume that $v \in e_0$ which gives at most n^{k-1} choices for e_0). If $v \notin (e_0 \cup f_0)$, then there are at most $T!n^{T-2k}$ ways to complete the path P_0 containing both edges, and if $v \in (e_0 \cup f_0)$ there are at most $T!n^{T-2k+1}$ ways to do it. In total, v intersects with at most $2(k+1)T!\mu n^{T-1}$ events in \mathcal{A} , proving (4.1).

For every $v \in V(G)$, we claim that

$$|B \in \mathcal{B} : v \in V(B)| \leq 2(T!)^2 \mu n^{2T-2}. \quad (4.2)$$

Indeed, suppose that $B = B(e_0, f_0; P_0, Q_0) \in \mathcal{B}$ with $v \in V(B)$. If $v \in (e_0 \cup f_0)$, we may assume that $v \in e_0$ and there are at most n^{k-1} choices for e_0 and at most μn^{k-1} for f_0 of the same colour, again by (B). Then there are at most $(T!)^2 n^{2T-2k}$ choices for P_0 and Q_0 containing e_0 and f_0 respectively. If $v \notin (e_0 \cup f_0)$, then there are at most μn^{2k-1} pairs e_0, f_0 of the same colour, and at most $(T!)^2 n^{2T-2k-1}$ choices for P_0 and Q_0 containing e_0 and f_0 respectively, such that $v \in P_0 \cup Q_0$. In total, v intersects at most $2(T!)^2 \mu n^{2T-2}$ events in \mathcal{B} , proving (4.2).

By the definition of Γ and since $|V(E)| \leq 2T$ for all $E \in \mathcal{E}$, there exists $C = C(k, t)$ such that

$$d_{\mathcal{A}} < C\mu n^{T-1} \quad \text{and} \quad d_{\mathcal{B}} < C\mu n^{2T-2}. \quad (4.3)$$

In order to bound p_A and p_B we will use switchings. Recall that p_A is an upper bound on $\mathbb{P}[A \mid \cap_{E \in \mathcal{E}'} \bar{E}]$ where $A \in \mathcal{A}$ and

$$\mathcal{E}' \subseteq \{E \in \mathcal{E} \setminus \{A\} : AE \notin E(\Gamma)\}. \quad (4.4)$$

Fix $A = A(e_0, f_0; P_0) \in \mathcal{A}$ and \mathcal{E}' as in (4.4). Let $\mathcal{H} = \mathcal{H}(\mathcal{E}')$ be the collection of loose Hamilton cycles of G that satisfy $\cap_{E \in \mathcal{E}'} \bar{E}$, which we may assume is non-empty. Let $\mathcal{H}_0 = \{H \in \mathcal{H} : P_0 \subset H\}$. Consider the bipartite multigraph \mathcal{F}_A with parts \mathcal{H}_0 and \mathcal{H} and where an edge is added between $H \in \mathcal{H}_0$ and $H' \in \mathcal{H}$ for each feasible (P_0, H) -switching $Y = (\mathcal{P}, H', \mathcal{P}')$ of size m .

In fact, for every feasible (P_0, H) -switching $Y = (\mathcal{P}, H', \mathcal{P}')$, we have $H' \in \mathcal{H} \setminus \mathcal{H}_0$. Indeed, by Definition 3.3, since $V(P_0)$ is transverse with respect to \mathcal{P}' , P_0 is not in H' . Moreover, by Definition 3.4, all edges in H' that are not in H either intersect P_0 or have a unique colour in H' . However, by the definition of Γ and (4.4), all events that intersect P_0 are not in \mathcal{E}' , thus $H' \in \mathcal{H}$.

Let δ_A be the minimum degree in \mathcal{F}_A of the vertices in \mathcal{H}_0 and Δ_A be the maximum degree in \mathcal{F}_A of the vertices in \mathcal{H} . Double-counting the edges of \mathcal{F}_A , we obtain

$$\mathbb{P}[A \mid \cap_{E \in \mathcal{E}'} \bar{E}] = \frac{|\mathcal{H}_0|}{|\mathcal{H}|} \leq \frac{\Delta_A}{\delta_A}. \quad (4.5)$$

It suffices to upper bound Δ_A and lower bound δ_A . The hypothesis of the lemma and the fact that every feasible switching produces a loose Hamilton cycle in \mathcal{H} , imply that $\delta_A \geq \gamma n^{m-1}$. To bound Δ_A , we fix any $H' \in \mathcal{H}$ and bound the number of pairs (H, Y) where $H \in \mathcal{H}_0$ and $Y = (\mathcal{P}, H', \mathcal{P}')$ is a feasible (P_0, H) -switching of size m . By the previous observation, it suffices to check it for $H' \in \mathcal{H} \setminus \mathcal{H}_0$. By condition (II) in Definition 3.3, all vertices in $V(P_0)$ are in different parts of \mathcal{P}' . We first construct \mathcal{P}' . As P_0 is fixed by A , the number of ways to choose T paths in H' of length at most $2t$ and each one containing exactly one vertex in $V(P_0)$ is a constant only depending on t, k . Additionally, there are at most n^{m-T} choices to complete \mathcal{P}' with $m - T$ paths in H' (each factor of n corresponding to a choice of starting point in H' for one of the remaining paths in \mathcal{P}'). Once \mathcal{P}' has been fixed, the number of ways to construct \mathcal{P} and H is a constant only depending on t, k, m . So $\Delta_A \leq C' n^{m-T}$, for $C' = C'(k, t, m)$. We obtain

$$\mathbb{P}[A \mid \cap_{E \in \mathcal{E}'} \bar{E}] \leq C' \gamma^{-1} n^{1-T} =: p_A. \quad (4.6)$$

Similarly, we choose p_B to be an upper bound on $\mathbb{P}[B \mid \cap_{E \in \mathcal{E}'} \bar{E}]$ where $B = B(e_0, f_0; P_0, Q_0) \in \mathcal{B}$ and \mathcal{E}' satisfying (4.4) where A is replaced by B . Define $\mathcal{H} = \mathcal{H}(\mathcal{E}')$ as before and now let $\mathcal{H}_0 = \{H \in \mathcal{H} : P_0, Q_0 \subset H\}$ and construct the bipartite multigraph \mathcal{F}_B with parts \mathcal{H}_0 and \mathcal{H} , where an edge is added between $H \in \mathcal{H}_0$ and $H' \in \mathcal{H}$ for each pair (Y, Z) where

- $Y = (\mathcal{P}, H_Y, \mathcal{P}_Y)$ is a feasible (P_0, H) -switching of size m , such that $V(\mathcal{P}) \cap V(Q_0) = \emptyset$, and
- $Z = (\mathcal{Q}, H', \mathcal{Q}')$ is a feasible (Q_0, H_Y) -switching of size m , such that $V(\mathcal{Q}) \cap V(P_0) = \emptyset$.

As in the previous observation, if there is an edge between H and H' , then $H' \in \mathcal{H} \setminus \mathcal{H}_0$. Define Δ_B and δ_B analogously to Δ_A and δ_A for the multigraph \mathcal{F}_B . To compute δ_B , note that some switchings are invalid as they violate the disjointness conditions. However, given P_0 and Q_0 , there is some constant $\tilde{C} > 0$ depending only on t and k such that there are at most $\tilde{C} n^{m-2}$ many (P_0, H) -switchings $Y = (\mathcal{P}, H_Y, \mathcal{P}_Y)$ such that $V(\mathcal{P}) \cap V(Q_0) \neq \emptyset$ and at most $\tilde{C} n^{m-2}$ many (Q_0, H_Y) -switchings $Z = (\mathcal{Q}, H', \mathcal{Q}')$ such that $V(\mathcal{Q}) \cap V(P_0) \neq \emptyset$. Therefore a negligible proportion of the pairs of switchings are disallowed and by the hypothesis of the lemma, $\delta_B \geq \frac{1}{2} \gamma^2 n^{2m-2}$. Similarly as before, we have that $\Delta_B \leq C' n^{2(m-T)}$. It follows that

$$\mathbb{P}[B \mid \cap_{E \in \mathcal{E}'} \bar{E}] \leq 2C' \gamma^{-2} n^{2-2T} =: p_B. \quad (4.7)$$

Combining equation (4.3), (4.6) and (4.7), and since $\mu \ll \gamma, 1/m, 1/t, 1/k$, we have that $p_A d_A + p_B d_B \leq 1/4$. The lemma now follows by applying Corollary 2.4. \square

5. Suitable splittings with viable partitions give feasible switchings

In this section, we will prove Lemma 3.10 showing that for a loose Hamilton cycle H containing some length t path P_0 , if we have a t -balanced m -splitting \mathcal{P} that is suitable for P_0 and such that $V(\mathcal{P})$ admits a viable partition, then we can find some feasible (P_0, H) -switching $(\mathcal{P}, H', \mathcal{P}')$. We therefore need to find a new loose Hamilton cycle H' which is the same as H outside of the vertex set $V^I(\mathcal{P})$. Recall that given the viable partition $\mathcal{X} = \{X_1, \dots, X_T\}$ of $V(\mathcal{P})$, there is some rerouting $\mathcal{R} = \{\{v_a^i, v_b^i\} : i = 1, \dots, m\}$ of \mathcal{P} such that each X_h contains exactly \tilde{m} pairs in \mathcal{R} . We will define H' by considering each X_h in the viable partition \mathcal{X} individually and finding some collection \mathcal{C}_h of \tilde{m} vertex-disjoint paths in X_h that cover the vertex set of X_h and such that each path $P \in \mathcal{C}_h$ has as endpoints one of the pairs in \mathcal{R} that lies in X_h . Note that, due to the fact that \mathcal{R} is a rerouting (Definition 3.8) and our requirement that the collections \mathcal{C}_h will cover all the vertices in $V(\mathcal{P})$, taking the paths in the collections along with the paths in $H[V(G) \setminus V^I(\mathcal{P})]$ will indeed define a new loose Hamilton cycle H' .

For each $h \in [T]$, in order to find the required path collection \mathcal{C}_h covering X_h , we will use the following lemma.

Lemma 5.1 (Path tilings). *Suppose that $G' \subset G$ is a k -graph with $|V(G')| = m$ and $\delta_j(G') \geq (\delta_j^k(1) + \frac{\varepsilon}{4})m^{k-j}$. Suppose also that there is some collection $\mathcal{R}' = \{\{u_a^i, u_b^i\} : i \in [\tilde{m}]\} \subset \binom{V(G')}{2}$ of \tilde{m} disjoint pairs of vertices in G' . Then there is a collection $\mathcal{C}' = \{P_i : i \in [\tilde{m}]\}$ of \tilde{m} vertex-disjoint (loose) paths in G' each of length at most $2t$ and such that $V(G') = \cup_{i \in [\tilde{m}]} V(P_i)$ and $V^E(P_i) = \{u_a^i, u_b^i\}$ for each $i \in [\tilde{m}]$. Moreover, for any $(2$ -uniform) graph B with $V(B) = V(G')$ and $\Delta(B) \leq 2tk^2$, we can find a collection $\mathcal{C}' = \{P_i : i \in [\tilde{m}]\}$ as above such that no edge of any path P_i contains an edge of B .*

We now show how to use Lemma 5.1 to establish Lemma 3.10.

Proof of Lemma 3.10. We fix some path P_0 of length t in G , some loose Hamilton cycle H containing P_0 and some t -balanced m -splitting $\mathcal{P} = \{P_0, \dots, P_{m-1}\}$ of H , that is suitable for P_0 and such that $V(\mathcal{P})$ admits a viable partition $\mathcal{X} = \{X_1, \dots, X_T\}$. By the definition of \mathcal{X} being viable (Definition 3.9), we have that there exists a rerouting $\mathcal{R} = \{\{v_a^i, v_b^i\} : i = 1, \dots, m\}$ of \mathcal{P} and some partition of \mathcal{R} as $\mathcal{R} = \mathcal{R}_1 \cup \dots \cup \mathcal{R}_T$ with $|\mathcal{R}_h| = \tilde{m}$ and $\mathcal{R}_h \subset \binom{X_h}{2}$ for each $h \in [T]$. Moreover, as \mathcal{X} is a viable (and hence transverse) partition of $V(\mathcal{P})$ of size T we necessarily have that for each $h \in [T]$ and each $i = 0, \dots, m-1$, the intersection $X_h \cap V(P_i)$ contains a single vertex. We label the vertices of each X_h as $X_h = \{u_{0,h}, \dots, u_{m-1,h}\}$ so that $u_{i,h} \in V(P_i)$ for $i = 0, \dots, m-1$. We further set $\hat{X}_h := X_h \setminus \{u_{0,h}\} = \{u_{i,h} : i \in [m-1]\}$ to be the vertex set obtained from X_h by removing $u_{0,h} \in V(P_0)$.

Our aim is to show that there is some feasible (P_0, H) -switching $(\mathcal{P}, H', \mathcal{P}')$. We will do this by finding a collection \mathcal{C}_h of paths in X_h , for $h \in [T]$ in succession. For each h , we will guarantee the following properties of \mathcal{C}_h :

- (1) \mathcal{C}_h is a *path tiling* in $G[X_h]$, that is, the collection of paths in \mathcal{C}_h are vertex-disjoint and cover the vertex set X_h ;
- (2) The set of pairs of endpoints $\{V^E(P) : P \in \mathcal{C}_h\}$ is precisely set \mathcal{R}_h and each path in \mathcal{C}_h has length at most $2t$.

For the next conditions we define $\hat{\mathcal{C}}_h := \{P[\hat{X}_h] : P \in \mathcal{C}_h\}$ to be the collection of paths obtained from \mathcal{C}_h after deleting the vertex $u_{0,h} \in X_h$ and the edges that contain it. Note that by removing $u_{0,h}$, one of the paths in \mathcal{C}_h might split into two paths in $\hat{\mathcal{C}}_h$. When we refer to edges in $\hat{\mathcal{C}}_h$, we mean the collection of edges that feature on some subgraph belonging to the set $\hat{\mathcal{C}}_h$.

- (3) No edge in \hat{C}_h shares a colour with an edge in $H[V(G) \setminus V(\mathcal{P}^I)]$;
- (4) No edge in \hat{C}_h shares a colour with an edge in $\cup_{1 \leq h' < h} \hat{C}_{h'}$;
- (5) The subgraph of G obtained by taking all edges in \hat{C}_h is rainbow.

We claim that after finding such C_h , for all $h \in [T]$, we will be done. Indeed, note that the graph obtained by taking $H[V(G) \setminus V(\mathcal{P})]$ and all the paths in $\mathcal{P}' := \cup_{h=1}^T \{P : P \in C_h\}$, defines a new loose Hamilton cycle H' in G , using here the conditions (1) and (2) above, and the fact that \mathcal{R} is a rerouting as in Definition 3.8. Moreover, by (2), note that \mathcal{P}' is an m -splitting of H' which is $2t$ -bounded. Note also that $H'[V(G) \setminus V^I(\mathcal{P}')] is precisely the graph $H[V(G) \setminus V^I(\mathcal{P})]$ and that $V(P_0)$ is transverse with respect to \mathcal{P}' as the partition \mathcal{X} was transverse with respect to \mathcal{P} and all the paths in \mathcal{P}' lie in one part of \mathcal{X} . Thus the triple $(\mathcal{P}, H', \mathcal{P}')$ defines a (P_0, H) -switching of size m (as defined in Definition 3.3). Furthermore, the graph $F' := H'[V(\mathcal{P}') \setminus V(P_0)]$ is rainbow due to the conditions (4) and (5) and no edge in F' shares a colour with an edge in $H[V(G) \setminus V(\mathcal{P}^I)]$ due to the conditions given by (3). Hence $(\mathcal{P}, H', \mathcal{P}')$ is a feasible (P_0, H) -switching, as required.$

It remains to prove that we can find the collections C_h as claimed and as previously mentioned, we will show how to construct these sequentially for $h = 1, \dots, T$. So suppose that we are at the point of constructing C_h for some $h \in [T]$ and that all $C_{h'}$ with $1 \leq h' < h$ have already been constructed. Firstly, we let $G' \subset G[X_h]$ be the k -graph obtained from $G[X_h]$ by deleting any edge $e \subset \hat{X}_h = X_h \setminus \{u_{0,h}\}$ that shares a colour with some edge in H . Note that as \mathcal{P} is a suitable splitting of H and \mathcal{X} is a viable partition, we have that $\delta_j(G') \geq (\delta_j^k(1) + \frac{\varepsilon}{4})m^{k-j}$. Indeed, for any subset $S \subset X_h$ with $|S| = j$, property (b) of Definition 3.9 implies that S is contained in at least $(\delta_j^k(1) + \frac{\varepsilon}{2})m^{k-j}$ edges in $G[X_h]$ and property (i) of Definition 3.6 implies that at most $\frac{\varepsilon}{4}m^{k-j}$ of these share a colour with some edge in H . We will find our paths in G' and thus ensure that property (3) holds.

Next we define a (2-uniform) graph B_h with vertex set $V(B_h) = X_h = \{u_{0,h}, \dots, u_{m-1,h}\}$ and a pair of vertices $\{u_{i,h}, u_{i',h}\} \in \binom{X_h}{2}$ being an edge of B_h if and only if $0 \notin \{i, i'\}$ and there is some $1 \leq h' < h$ such that the pair $\{u_{i,h'}, u_{i',h'}\}$ lies in an edge $e \in E(G[\hat{X}_{h'}])$ that features in $\hat{C}_{h'}$. We claim that $\Delta(B_h) \leq 2tk^2$. Indeed, for a vertex $u = u_{i,h} \in X_h$ we have that $\deg_{B_h}(u) = 0$ if $i = 0$ and otherwise we bound the number of choices for i' with $\{u, u_{i',h}\} \in E(B_h)$ by a choice of $h' < h$ (at most $t(k-1) \leq tk$ choices), a choice of edge $e \in G[\hat{X}_{h'}]$ that lies in $\hat{C}_{h'} \subset C_{h'}$ and contains $u_{i,h'}$ (at most 2 choices) and a choice of vertex $u_{i',h'}$ that lies in e (at most $k-1 \leq k$ choices).

By the ‘moreover’ part of Lemma 5.1, we can obtain a collection C_h of \tilde{m} paths in $G'[X_h]$ that satisfy the conditions (1) and (2) above and such that no edge in a path in C_h contains an edge of B_h . It remains to establish properties (4) and (5). For (5), note that as \hat{C}_h is a collection of vertex-disjoint loose paths, any pair of edges $e, f \subseteq \hat{X}_h \subset V(\mathcal{P}) \setminus V(P_0)$ that feature in \hat{C}_h necessarily have that $|e \cap f| \leq 1$. Thus, as \hat{X}_h is transverse with respect to \mathcal{P} , we have that $\chi(e) \neq \chi(f)$ due to conditions (ii) (for $|e \cap f| = 1$) and (iii) (for $|e \cap f| = 0$, see Remark 3.7) of Definition 3.6 and the fact that \mathcal{P} is suitable for P_0 . This establishes (5). Finally, for (4) suppose for a contradiction that there is some edge e in \hat{C}_h that shares a colour with some edge f in $\hat{C}_{h'}$ for some $1 \leq h' < h$. As \mathcal{P} is suitable for P_0 and $e, f \subset V(\mathcal{P}) \setminus V(P_0)$ are disjoint edges that are both transverse with respect to \mathcal{P} (as they lie in different parts of the transverse partition \mathcal{X}), condition (iii) of Definition 3.6 implies that there exists distinct $i, i' \in [m-1]$ such that $u_{i,h'}, u_{i',h'} \subset f$ and $u_{i,h}, u_{i',h} \subset e$. However, this implies that $\{u_{i,h}, u_{i',h}\} \in E(B_h)$ contradicting the fact that no edge on a path in C_h contains an edge of B_h . This concludes the proof. \square

5.1 Path tilings

It remains to prove Lemma 5.1. For this, we will use the following recent theorem due to Alvarado, Kohayakawa, Lang, Mota, and Stagni [3, Theorem 9.3] which shows that the j -degree threshold for a loose Hamilton cycle coincides with the threshold for a loose Hamilton path between any pair of vertices.

Theorem 5.2 (Hamilton paths). *For any $n_0 \geq t/2$ such that $n_0 - 1 \in (k-1)\mathbb{N}$, if G_0 is an n_0 -vertex k -graph with $\delta_j(G_0) \geq (\delta_j^k(1) + \frac{\varepsilon}{8})n_0^{k-j}$, then for any pair of vertices $w^1, w^2 \in V(G_0)$, there is a loose Hamilton path in G_0 with endpoints w^1, w^2 .*

With Theorem 5.2 in hand, the simple idea behind proving Lemma 5.1 is to partition $V(G')$ into \tilde{m} sets $V_1, \dots, V_{\tilde{m}}$ of (roughly) equal sizes such that the collection of pairs $\mathcal{R}' = \{\{u_a^i, u_b^i\} : i \in [\tilde{m}]\}$ is distributed amongst the vertex sets with $\{u_a^i, u_b^i\} \subset V_i$ for each $i \in [\tilde{m}]$. We can then hope to apply Theorem 5.2 to each $G'[V_i]$ individually to get a path P_i in $G'[V_i]$ between u_a^i and u_b^i and thus obtain the desired path tiling $\mathcal{C}' = \{P_i : i \in [\tilde{m}]\}$. In order to be able to apply Theorem 5.2 in each of the vertex sets V_i , we need the graphs $G'[V_i]$ to inherit the j -degree condition of G' . This is achieved by taking a random partition (apart from pairs in \mathcal{R}' which are placed in the desired set V_i). We also need the divisibility conditions that $|V_i| - 1 \in (k-1)\mathbb{N}$ for each $i \in [\tilde{m}]$. When randomly distributing the vertices amongst the V_i , we will not be able to ensure such a divisibility condition but we overcome this by setting aside some vertex sets W_i with $|W_i| = k-2$ for $i \in [\tilde{m}]$ which will be used to adjust the sets V_i and establish the required divisibility conditions. In order to prove the ‘moreover’ statement of Lemma 5.1, we show that with high probability the random sets V_i induce almost independent sets in B . If each V_i was truly an independent set in B then the loose Hamilton paths output by Theorem 5.2 would not use hyperedges that contain edges of B . To turn the almost independent sets into truly independent sets in B , we will need to adjust our sets slightly by moving some limited number of vertices between parts of the partition. Moreover, by choosing our sets W_i carefully to avoid edges of B , we can also maintain that the $B[V_i]$ are empty graphs after adding vertices from W_i to amend the divisibility constraints. We now give the details.

Proof of Lemma 5.1. We will prove the whole statement including the ‘moreover’ part and so we fix some G' with vertex set $V = V(G') \subset V(G)$, some $(2$ -uniform) graph B with $V(B) = V$ and some collection of pairs of vertices $\mathcal{R}' = \{\{u_a^i, u_b^i\} : i \in [\tilde{m}]\}$ as in the statement of the lemma. We let $R := \bigcup_{i=1}^{\tilde{m}} \{u_a^i, u_b^i\}$ be the set of vertices that appear in a pair in \mathcal{R}' . We will now define vertex sets $W_i, Y_i \subset V$ for $i = 1, \dots, \tilde{m}-1$ sequentially, and initiate the process by defining $W_0 = \emptyset$ and $Y_0 = V \setminus R$. For each $i \in [\tilde{m}-1]$ we will define $W_i \subset Y_{i-1}$ to have $|W_i| = k-2$ and Y_i will be defined as $Y_i = Y_{i-1} \setminus W_{i-1}$. This will guarantee that the sets W_i are disjoint from each other and for each $i \in [\tilde{m}-1]$ we will have that $|Y_i| \geq m - 2\tilde{m} - (k-2)(i-1) \geq (t-1)\tilde{m}$. We will also impose that $W_i \subset Y_{i-1}$ has the property that the only possible edges in $B[\{u_a^i, u_b^i\} \cup W_{i-1} \cup W_i]$ and $B[\{u_a^{i+1}, u_b^{i+1}\} \cup W_i]$ are $\{u_a^i, u_b^i\}$ and $\{u_a^{i+1}, u_b^{i+1}\}$ respectively. Let us see that such a vertex set W_i indeed exists. We start with Y_{i-1} which has size at least $(t-1)\tilde{m}$ and remove all the vertices that are B -neighbours of a vertex in $\{u_a^i, u_b^i, u_a^{i+1}, u_b^{i+1}\} \cup W_{i-1}$ to get some vertex set $Y'_{i-1} \subset Y_{i-1}$ which has size at least $(t-1)\tilde{m} - (4 + (k-2))2tk^2 \geq (t-2)\tilde{m}$ as $1/t \ll 1/k$ and $\tilde{m} = t^{100k}$ by (3.1) and (3.2). As $\Delta(B) \leq 2tk^2$ there is some vertex set $Y''_{i-1} \subset Y'_{i-1}$ with $|Y''_{i-1}| \geq |Y'_{i-1}|/(2tk^2 + 1) \geq \tilde{m}/2tk^3 \geq k-1$ such that $B[Y''_{i-1}]$ is empty. Choosing W_i arbitrarily from this vertex set Y''_{i-1} gives the desired properties, noting here that we have previously chosen W_{i-1} so that the only possible edge in $B[\{u_a^i, u_b^i\} \cup W_{i-1}]$ is $\{u_a^i, u_b^i\}$. After choosing W_i , we define $Y_i = Y_{i-1} \setminus W_i$ by removing the vertices of W_i from Y_{i-1} and move onto the next step $i+1$ (or finish, if $i = \tilde{m}-1$).

At the end of the process above we have defined W_i for $i = 0, 1, \dots, \tilde{m}-1$ and the vertex set $V' := Y_{\tilde{m}-1}$ which is the vertex set obtained from V by removing R and all of the sets W_i .

We further define $W_{\tilde{m}} = \emptyset$. Note that $|V'| = T \cdot \tilde{m} - 2\tilde{m} - (k-2)(\tilde{m}-1) = \tilde{m}(t-1)(k-1) + k-2$. We proceed with the following claim, which will find a partition $V' = U_1 \cup \dots \cup U_{\tilde{m}}$ of the vertices V' which have certain properties. For each $i \in [\tilde{m}]$ we will define $U_i^+ := U_i \cup \{u_a^i, u_b^i\} \cup W_i \cup W_{i-1}$. Recall that by construction, there are no edges in $B[U_i^+ \setminus U_i]$ with the possible exception of the edge $\{u_a^i, u_b^i\}$. We say that U_i is *good* if there is no edge in $B[U_i^+]$ different from $\{u_a^i, u_b^i\}$, *bad* if there is at least one edge in $B[U_i^+]$ different from $\{u_a^i, u_b^i\}$. Moreover we say U_i is *very bad* if there is no choice of a single vertex $u \in U_i$ that is incident to all edges in $B[U_i^+]$ other than $\{u_a^i, u_b^i\}$. Recall that we have chosen β such that $1/t \ll \beta \ll \varepsilon$.

Claim 5.3. *There is a partition $V' = U_1 \cup \dots \cup U_{\tilde{m}}$ of the vertices V' such that the following holds:*

- (A) $(t-1)(k-1) - \beta t \leq |U_i| \leq (t-1)(k-1) + \beta t$ for all $i \in [\tilde{m}]$;
- (B) for any $S \in \binom{V'}{j}$ and $i \in [\tilde{m}]$, we have that $\delta_{G'}(S; U_i^+) \geq (\delta_j^k(1) + \frac{3\varepsilon}{16})|U_i^+|^{k-j}$;
- (C) none of the U_i with $i \in [\tilde{m}]$ are very bad; and
- (D) at most $t^3 k^3$ of the sets U_i with $i \in [\tilde{m}]$ are bad.

Proof of Claim: We will consider a random partition of the vertices of V' and show that all the properties (A), (B), (C) and (D) hold with probability at least $4/5$. Thus with probability at least $1/5$ all four conditions will be satisfied and so a partition as claimed does indeed exist. In more detail, each vertex $v \in V'$ is assigned to U_i with probability $1/\tilde{m}$ for each $i \in [\tilde{m}]$ and the allocations of different vertices is done independently. Note that the expected size of each U_i is $\lambda := (t-1)(k-1) + \frac{(k-2)}{\tilde{m}} \leq (t-1)(k-1) + 1$.

For part (A), note that for a fixed $i \in [\tilde{m}]$ we have that $|U_i|$ is binomially distributed and so by Chernoff bounds (Theorem 2.5),

$$\mathbb{P}[|U_i| \notin [\lambda - \beta t, \lambda + \beta t]] \leq 2e^{-\frac{\beta^2 t}{6k^2}} \leq e^{-\sqrt{t}}.$$

Thus the fact that with probability at least $9/10 \geq 4/5$, (A) holds follows from a union bound over all $\tilde{m} = t^{100k}$ choices of $i \in [\tilde{m}]$, using here that t is sufficiently large.

For condition (B), we fix some $i \in [\tilde{m}]$ and some j -set $S \subset V = V(G')$. We define $N(S) := \{A \subset V' : A \cup S \in E(G')\}$ and note that $|N(S)| \geq (\delta_j^k(1) + \frac{\varepsilon}{4} - \beta)m^{k-j}$. Indeed, we have that the number of edges in G' containing S is at least $(\delta_j^k(1) + \frac{\varepsilon}{4})m^{k-j}$ due to the minimum degree condition on G' and at most $|V \setminus V'|m^{k-j-1} \leq k\tilde{m}m^{k-j-1} \leq \beta m^{k-j}$ of the sets $A \subset V(G')$ with $A \cup S \in E(G')$ are such that $A \not\subseteq V'$. Now let $X_i(S) = |\{A \in N(S) : A \subset U_i\}|$ be the random variable that counts the number of sets in $N(S)$ that end up in U_i . Each set in $N(S)$ is in U_i with probability $(1/\tilde{m})^{k-j}$ and so $\lambda_i(S) := \mathbb{E}[X_i(S)] \geq (\delta_j^k(1) + \frac{\varepsilon}{4} - \beta)(t(k-1))^{k-j}$. We will use Suen's inequality (Lemma 2.7) to prove concentration for $\lambda_i(S)$ and so for each $A \in N(S)$ we let I_A be indicator random variable for the event that $A \subset U_i$ and define a graph Γ with vertex set $V(\Gamma) = N(S)$ such that $A \sim A'$ if and only if $A \cap A' \neq \emptyset$. Then Γ is a strong dependency graph for the family of indicator random variables $\{I_A\}_{A \in N(S)}$ and following the notation of Lemma 2.7, we have that

$$\Delta_{X_i(S)} = \sum_{\ell=1}^{k-j-1} \sum_{\substack{(A, A') \in N(S)^2 \\ |A \cap A'| = \ell}} \mathbb{E}[I_A I_{A'}] \leq \sum_{\ell=1}^{k-j-1} \binom{2(k-j)-\ell}{\ell} m^{2(k-j)-\ell} \left(\frac{1}{\tilde{m}}\right)^{2(k-j)-\ell} \leq (2k)^{3k} t^{2(k-j)-1},$$

and similarly

$$\delta_{X_i(S)} = \max_{A \in N(S)} \sum_{A' : |A' \cap A| \geq 1} \mathbb{E}[I_{A'}] \leq \sum_{\ell=1}^{k-j-1} \binom{k-j}{\ell} m^{k-j-\ell} \left(\frac{1}{\tilde{m}}\right)^{k-j} \leq (2tk)^{3k} m^{-1}.$$

By Suen's inequality (Lemma 2.7), we have that $\mathbb{P}[X_i(S) \leq \lambda_i(S) - \beta(t(k-1))^{k-j}] \leq e^{-\sqrt{t}}$ and so a union bound over all $m^j = (t^{100k}T)^j$ choices of $S \subseteq V$ and all $\tilde{m} = t^{100k}$ choices of U_i gives that with probability at least 9/10 we have that $X_i(S) \geq \lambda_i(S) - \beta(t(k-1))^{k-j}$ for each $i \in [\tilde{m}]$ and $S \in \binom{V}{j}$, using here that t is sufficiently large. Moreover, we showed that condition (A) holds with probability at least 9/10 and so with probability at least 4/5, both (A) and the lower bounds on all $X_i(S)$ hold. This establishes the degree conditions (B) by using our lower bound on the values $\lambda_i(S)$, the upper bound on the sizes of $|U_i|$ (and the resulting upper bounds on $|U_i^+|$) from (A) and the fact that $\beta \ll \varepsilon$ in our hierarchy (3.1).

It remains to establish conditions (C) and (D). Again, note that we defined the sets W_i in such a way that the only possible edge in $B[U_i^+ \setminus U_i]$ is $\{u_a^i, u_b^i\}$ for all $i \in [\tilde{m}]$. For each $i \in [\tilde{m}]$, it suffices to consider edges in $B[U_i]$ and edges in B between U_i and $U_i^+ \setminus U_i$. Fixing some $i \in [\tilde{m}]$ and considering edges in $B[U_i]$, we note that by the maximum degree condition on B , we have that $B[V']$ contains at most $2tk^2m$ edges and each edge lands in U_i with probability $1/\tilde{m}^2$. Similarly for edges between $U_i^+ \setminus U_i$ and U_i , we have that there are at most $(2k-2)2tk^2$ edges between $U_i^+ \setminus U_i = \{u_a^i, u_b^i\} \cup W_i \cup W_{i-1}$ and V' and each such edge lies in $B[U_i^+]$ with probability $1/\tilde{m}$. Therefore the expected number of edges with at least one endpoint in U_i is at most $2tk^2m \cdot (1/\tilde{m})^2 + (2k-2)2tk^2 \cdot (1/\tilde{m}) \leq 6t^2k^3/\tilde{m}$ and by Markov's inequality, we have that $\mathbb{P}[U_i \text{ is bad}] \leq 6t^2k^3/\tilde{m}$ for each $i \in [\tilde{m}]$. Similarly, the probability that U_i is very bad can be upper bounded by the expected number of pairs of edges in $B[U_i^+]$, each having an endpoint in U_i using Markov's inequality. We get that

$$\mathbb{P}[U_i \text{ is very bad}] \leq \frac{(2tk^2m)^2}{\tilde{m}^4} + \frac{2tk^2m \cdot (2k-2)2tk^2}{\tilde{m}^3} + \frac{((2k-2)2tk^2)^2}{\tilde{m}^2} \leq \frac{40k^6t^4}{\tilde{m}^2},$$

where the first summand accounts for vertex-disjoint pairs of edges in $B[U_i]$, the second one accounts for an edge in $B[U_i]$ paired with a disjoint edge between U_i and $U_i^+ \setminus U_i$ and the last one counts pairs of edges which are both between U_i and $U_i^+ \setminus U_i$, and which by definition of very bad, do not intersect in U_i . Note that one of these three scenarios must necessarily occur for U_i to be very bad as otherwise there would be one vertex in U_i incident to all edges in $B[U_i^+]$ other than $\{u_a^i, u_b^i\}$. Thus, by linearity of expectation, the expected number of very bad sets U_i is less than $40k^6t^4/\tilde{m} < 1/5$ and the fact that 3 holds with probability at least 4/5 follows from Markov's inequality. Similarly, the expected number of bad U_i is at most $6t^2k^3$ by linearity of expectation, and so $\mathbb{P}[\neg(D)] \leq 6/t \leq 1/5$ as required. \square

We now fix some partition $V' = U_1 \cup \dots \cup U_{\tilde{m}}$ as in Claim 5.3 and we will first slightly adjust the partition so that all the sets become good. By condition (D) of Claim 5.3 there is some set of indices $I_B \subset [\tilde{m}]$ with $|I_B| \leq t^3k^3$ and such that $U_{i'}$ is good for all $i' \in [\tilde{m}] \setminus I_B$. For each bad index $i \in I_B$ as U_i is not very bad due to (C), we have that there is some vertex $z_i \in U_i$ such that $U'_i := U_i \setminus \{z_i\}$ is good, that is, such that the only possible edge in $B[U'_i \cup \{u_a^i, u_b^i\} \cup W_i \cup W_{i-1}]$ is $\{u_a^i, u_b^i\}$. Moreover, there are at least $\tilde{m} - t^3k^3 - 4tk^2 \geq \tilde{m}/2$ indices $i' \in [\tilde{m}] \setminus I_B$ such that adding z_i to $U_{i'}$ results in a set $U'_{i'} = U_{i'} \cup \{z_i\}$ which is good. Indeed, z_i has at most $2tk^2$ neighbours in B and each neighbour of z_i rules out at most 2 possible choices for such an i' (the 2 here is necessary in case the neighbour lies in some set $W_{i''}$). Therefore for each $i \in I_B$ corresponding to a bad set U_i we can find a $i' = i'(i)$ such that adding z_i to $U_{i'}$ results in both $U'_i := U_i \setminus \{z_i\}$ and $U'_{i'} := U_{i'} \cup \{z_i\}$ being good and we can do this for all $i \in I_B$ in such a way that $i'(i_1) \neq i'(i_2)$ for distinct $i_1, i_2 \in I_B$. Let $V' = U'_1 \cup \dots \cup U'_{\tilde{m}}$ be the resulting partition after switching all the z_i . Here, for any set $i_0 \in [\tilde{m}] \setminus I_B$ such that i_0 is not chosen as $i'(i)$ for some $i \in I_B$, we simply leave U_{i_0} unchanged and define $U'_{i_0} := U_{i_0}$.

Next we will distribute vertices in the sets W_i to get a partition of $V = V(G')$ as $V_1 \cup \dots \cup V_{\tilde{m}}$ with $U'_i \cup \{u_a^i, u_b^i\} \subset V_i$ and $|V_i| - 1 \in (k-1)\mathbb{N}$ for each $i \in [\tilde{m}]$. We define V_i for $i = 1, \dots, \tilde{m}$ sequentially via the following process, initiating with $W_0^- := W_0 = \emptyset$ and $V_0 = \emptyset$. Suppose we

are at step $i \in [\tilde{m}]$ and that $V_{i'}$ and $W_{i'}^-$ are defined for all $0 \leq i' < i$. Then we let $k_i := (|U_i'| + |W_{i-1}^-| + 1) \bmod (k-1)$ and note that $k_i \in \{0, \dots, k-2\}$. We define $W_i^+ \subset W_i$ to be an empty set if $k_i = 0$ and some arbitrary set of size $k-1-k_i$ if $k_i \neq 0$. Furthermore, we define V_i to be $V_i := U_i' \cup W_{i-1}^- \cup \{u_a^i, u_b^i\} \cup W_i^+$ and $W_i^- := W_i \setminus W_i^+$. We then continue to step $i+1$ or finish if $i = \tilde{m}$. Note that, for all $i \in [\tilde{m}-1]$ this process will succeed at step i and result in a set V_i such that $|V_i| \bmod (k-1) = |U_i'| + |W_{i-1}^-| + 2 + (k-1-k_i) \bmod (k-1)$ and so we indeed have that $|V_i| - 1 \in (k-1)\mathbb{N}$. For $i = \tilde{m}$ we claim that $k_i = 0$ and so the algorithm also succeeds and $|V_{\tilde{m}}| - 1 \in (k-1)\mathbb{N}$ (if $k_{\tilde{m}} \neq 0$ then the algorithm would stall as $W_{\tilde{m}} = \emptyset$ and so an appropriate $W_{\tilde{m}}^+$ would not exist). To see it, observe that $U_{\tilde{m}}' \cup W_{\tilde{m}-1}^- \cup \{u_a^{\tilde{m}}, u_b^{\tilde{m}}\} = V \setminus (\bigcup_{i=1}^{\tilde{m}-1} V_i)$ and for each $i \in [\tilde{m}-1]$ we have that $|V_i| = 1 \bmod (k-1)$ from above. Thus as $|V| = T \cdot \tilde{m} = (t(k-1) + 1)\tilde{m} = \tilde{m} \bmod (k-1)$, we get that $|U_{\tilde{m}}'| + |W_{\tilde{m}-1}^-| + 2 = \tilde{m} - (\tilde{m}-1) = 1 \bmod (k-1)$ and indeed $k_{\tilde{m}} = 0$ as required.

Note that for each $i \in [\tilde{m}]$, the vertex set V_i and the set U_i^+ defined via Claim 5.3 differ by at most $2(k-1) + 1$ vertices, and so condition (B) from Claim 5.3 implies that $\delta_j(G'[V_i]) \geq (\delta_j^k(1) + \frac{\varepsilon}{8})|V_i|^{k-j}$. Note also that $|V_i| \geq |U_i| \geq (t-1)(k-1) - \beta t \geq t/2$ by part (A) of Claim 5.3. We are therefore in a position to apply Theorem 5.2 in $G'[V_i]$ to obtain some loose Hamilton path P_i in $G'[V_i]$ between u_a^i and u_b^i . Doing this for each $i \in [\tilde{m}]$ results in some collection $\mathcal{C}' := \{P_i; i \in [\tilde{m}]\}$ which we claim satisfies the required conditions. Indeed, part (A) of Claim 5.3 and how we formed V_i implies that we certainly have $|V_i| \leq tk$ and so P_i will have length less than $2t$, for each $i \in [\tilde{m}]$. Moreover, for each such $i \in [\tilde{m}]$, due to how we defined V_i (and U_i'), we have that the only possible edge in $B[V_i]$ is $\{u_a^i, u_b^i\}$ and the pair $\{u_a^i, u_b^i\}$ does not lie in any edge of P_i as they are the endpoints of the path. This concludes the proof. \square

6. Many suitable splittings with viable partitions

In this section, we prove Lemma 3.11 showing that for a fixed loose Hamilton cycle H and loose path $P_0 \subset H$, there are many suitable splittings containing P_0 that admit viable partitions. We encourage the reader to remind themselves of the definitions of suitable (Definition 3.6) and viable (Definition 3.9) in order to follow the contents of this section. Lemma 3.11 will be achieved by random sampling, showing that with probability bounded away from 0, a randomly sampled m -splitting will suffice. As there are many m -splittings containing P_0 , indeed there are $\Omega(n^{m-1})$, we thus obtain the desired lower bound. In fact, instead of sampling an m -splitting, we will use a slightly more general model to sample, with more independence which will be useful in the analysis. Moreover, we will achieve our aim with two rounds of randomness, first showing that our random sampling results in a suitable m -splitting with probability bounded from 0 and then showing that such an m -splitting admits a viable partition which will be found again by appealing to a random choice of partition. In order to set ourselves up for this second round of randomness, we need slightly more from our random m -splitting than it just being suitable, as the following lemma details (see (3.2) for the definition of M).

Lemma 6.1. *Let G be as described in Theorem 1.2. For a loose Hamilton cycle H in G and a path $P_0 \subset H$ of length t , let $E_p \subset E(H)$ be a random subset of $E(H)$ obtained by sampling each edge independently with probability $p := \frac{(m-1)(k-1)}{n}$ and \mathcal{P}_p be the collection of length t increasing paths in H starting from an edge in E_p . Then with probability at least $1/m$, we have that $P_0 \notin \mathcal{P}_p$, the collection of paths $\mathcal{P}_* := \mathcal{P}_p \cup \{P_0\}$ is a t -balanced m -splitting¹ of H suitable for P_0 and $\delta_j(G[V(\mathcal{P}_*)]) \geq (\delta_j^k(1) + \frac{3\varepsilon}{4})M^{k-j}$.*

¹In particular the paths in \mathcal{P}_p are disjoint from P_0 .

Proof. Before embarking on the proof, we make some useful definitions and observations. We say that a pair of vertices $u, v \in V(G)$ are *close* if there are edges $e, f \in E(H)$ with $u \in e, v \in f$ and such that there is a path of length at most $2t + 1$ in H containing both e and f . We say that $u, v \in V(G)$ are *spread* if they are not close and we say that a set $S \subset V(G)$ is spread if there are no pairs of close vertices contained in S . We say that a subset $S \subset V(G)$ is *almost spread* if there is *exactly one* pair of vertices $u, v \in S$ that are close but all other pairs of vertices in S are spread.

Now note that for a vertex $v \in V(G)$, the probability that v belongs to the set $V(\mathcal{P}_p)$ is $1 - (1 - p)^{t+1}$ if v is a vertex of degree 2 in H and $1 - (1 - p)^t$ if v is not a vertex of degree 1 in H . In either case we have that

$$\frac{(1 - \beta)M}{n} \leq \mathbb{P}[v \in V(\mathcal{P}_p)] \leq \frac{2tmk}{n}, \quad (6.1)$$

recalling that β is fixed so that $1/t \ll \beta \ll \varepsilon$ in our hierarchy (3.1) and m is much larger than t ; see (3.2). For a spread set of vertices $S \subset V(G)$ note that the events that each $v \in V(\mathcal{P}_p)$ are independent for $v \in S$ and so $\mathbb{P}[S \subset V(\mathcal{P}_p)] = \prod_{v \in S} \mathbb{P}[v \in V(\mathcal{P}_p)]$.

We collect the following events:

- $\mathcal{E}_1 := \{V(\mathcal{P}_p) \text{ contains a spread } S \text{ with } |S| = k - 1 \text{ and there are at least } \varepsilon m/4 \text{ edges in } E_{G[V(\mathcal{P}_*)]}(S) \text{ that share a colour with an edge in } H\},$
- $\mathcal{E}_2 := \{V(\mathcal{P}_p) \text{ contains a pair } e, f \in E(G) \text{ with } \chi(e) = \chi(f), |e \cap f| \leq 1 \text{ and with } e \cup f \text{ spread}\},$
- $\mathcal{E}_3 := \{V(\mathcal{P}_p) \text{ contains a pair } e, f \in E(G) \text{ with } \chi(e) = \chi(f), e \cap f = \emptyset \text{ and } e \cup f \text{ almost spread}\},$
- $\mathcal{E}_4 := \{\delta_j(G[V(\mathcal{P}_*)]) < (\delta_j^k(1) + \frac{3\varepsilon}{4})M^{k-j}\}, \text{ and}$
- $\mathcal{E}_5 := \{V(\mathcal{P}_*) \text{ contains a pair of close vertices in distinct paths in } \mathcal{P}_*\}.$

The proof of the lemma will be deduced from the following claim.

Claim 6.2. *For each $i \in [5]$, we have that $\mathbb{P}[\mathcal{E}_i] \leq 1/m$.*

To see that Claim 6.2 implies the lemma, first note that by Lemma 2.6 with probability at least $1/(4\sqrt{m}) \geq 6/m$, we have that $|E_p| = \mathbb{E}[|E_p|] = m - 1$ and hence $|\mathcal{P}_*| = m$. Thus, with probability at least $6/m - 5 \cdot (1/m) = 1/m$, we have that $|\mathcal{P}_*| = m$ and none of the events $\mathcal{E}_1, \dots, \mathcal{E}_5$ hold, by Claim 6.2. In particular, event \mathcal{E}_5 not holding implies that there are no pairs of intersecting paths in \mathcal{P}_* and so \mathcal{P}_* is indeed a t -balanced m -splitting of H . The degree condition in the statement of the lemma is given directly by event \mathcal{E}_4 not holding and it remains to establish that \mathcal{P}_* is suitable for P_0 . For this, note that any transverse $Y \subset V(\mathcal{P}) \setminus V(P_0)$ is necessarily spread by event \mathcal{E}_5 not happening and so condition (i) of being suitable (Definition 3.6) follows from the absence of event \mathcal{E}_1 , whilst condition (ii) follows from the absence of event \mathcal{E}_2 . Finally, for condition (iii) consider two disjoint edges $e, f \in G[V(\mathcal{P}_*) \setminus V(P_0)]$ with $\chi(e) = \chi(f)$ and suppose that there are *not* two distinct $P_i, P_{i'} \in \mathcal{P}_* \setminus P_0$ with $P_x \cap e, P_x \cap f \neq \emptyset$ for $x = i, i'$. Then $e \cup f$ intersects at least $2k - 1$ members of \mathcal{P}_* and thus $e \cup f$ intersects at most one $P_i \in \mathcal{P}_*$ in two vertices (and no P_i in more than 2 vertices). Due to the absence of event \mathcal{E}_5 , there is no pair of close vertices in distinct paths in \mathcal{P}_* . Hence we have that $e \cup f$ can contain at most one pair of close vertices (coming from the at most one path that it intersects in two vertices), that is, we must have that $e \cup f$ is spread or almost spread. If the former, then we get a contradiction due to \mathcal{E}_2 not happening whilst if the latter is the case, then we contradict that \mathcal{E}_3 did not happen. Thus (iii) indeed occurs and \mathcal{P}_* is suitable for P_0 , as required.

It remains to establish Claim 6.2 which we now do by considering each event separately.

Bounding $\mathbb{P}[\mathcal{E}_1]$. Let $\chi(H)$ denote the colours featuring on edges in H and $E_{\chi(H)} := \{e \in E(G) : \chi(e) \in \chi(H)\}$ be the set of edges that share a colour with an edge of H . Let us fix $\alpha := \varepsilon/(32tk)$ and say that a $(k - 1)$ -set $S \in \binom{V(G)}{k-1}$ of vertices in G is *bad* if $|E_G(S) \cap E_{\chi(H)}| \geq \alpha n$ and *good* otherwise. We begin by showing that the probability that there exists some bad spread set

$S \in \binom{V(G)}{k-1}$ such that $S \subset V(\mathcal{P}_p)$ is at most $1/2m$. Indeed, we have that $|E_{\chi(H)}| \leq \frac{n}{k-1} \cdot \mu n^{k-1} \leq \mu n^k$ using that H has $n/(k-1)$ edges (and so at most this many colours featuring in $\chi(H)$) and the global bound (B) of Theorem 1.2. Hence the number of bad spread $(k-1)$ -sets in G is at most $(\mu k n^{k-1})/\alpha$. Each such set S appears in $V(\mathcal{P}_p)$ with probability at most $(2tmk/n)^{k-1}$ using the upper bound in (6.1) and the fact that S is spread and so each vertex of S appears independently. Hence, by Markov's inequality the probability that there exists a bad spread set in $V(\mathcal{P}_p)$ is at most

$$\frac{\mu k n^{k-1}}{\alpha} \cdot \frac{(2tmk)^{k-1}}{n^{k-1}} \leq \frac{1}{2m},$$

using that $\mu \ll 1/m, 1/t, 1/k, \varepsilon$ and $\alpha = \varepsilon/(32tk)$.

Now for a good $(k-1)$ -set $S \in \binom{V(G)}{k-1}$, let $Y(S) := |E_{G[V(\mathcal{P}_*)]}(S) \cap E_{\chi(H)}|$ be the random variable which counts the number of edges in $G[V(\mathcal{P}_*)]$ containing S that share a colour with an edge of H . We will show that the probability that there is some good spread S that appears in $V(\mathcal{P}_p)$ with $Y(S) \geq \varepsilon m/4$ is at most $1/(2m)$. Hence with probability at least $1 - 1/m$ all the spread $(k-1)$ -sets in $V(\mathcal{P}_p)$ are good and have a bounded $Y(S)$, establishing Claim 6.2 for \mathcal{E}_1 .

In order to measure the probability of some good S having large $Y(S)$, consider the subset $N(S)$ obtained from $\{e \setminus S : e \in E_G(S) \cap E_{\chi(H)}\} \subset V(G)$ by removing all vertices that are close to a vertex in $S \cup V(P_0)$. Then consider the random variable $Z(S) := |N(S) \cap V(\mathcal{P}_p)|$. As $Y(S) \leq Z(S) + (2t+1) \cdot 2k^2 \leq Z(S) + \varepsilon m/8 =: Z'(S)$ for any good S , we will instead bound the probability that the random variable $Z'(S)$ is larger than $\varepsilon m/4$. For bounding $Z'(S)$, note that for any good spread S , we have that $Z'(S)$ is tk -Lipschitz and 1-certifiable. Moreover, using (6.1) and the definition of α we have that $\lambda := \mathbb{E}[Z'(S)] \leq \alpha n \cdot (2tmk)/n + \varepsilon m/8 \leq 3\varepsilon m/16$. Hence by Talagrand's Inequality (Lemma 2.8),

$$\mathbb{P}[Z'(S) \geq \varepsilon m/4] \leq \mathbb{P}[Z'(S) \geq \varepsilon m/8 + 60tk\sqrt{\lambda}] \leq 4e^{-\frac{\varepsilon^2 m^2}{512t^2 k^2 \lambda}} \leq m^{-(k+2)},$$

as $1/m \ll 1/t, 1/k, \varepsilon$; see (3.1). Moreover, due to the fact that we removed vertices close to S in order to get $N(S)$, the events $\{S \subset V(\mathcal{P}_p)\}$ and $\{Z'(S) \geq \varepsilon m/4\}$ are independent and so occur concurrently with probability at most $(2tmk/n)^{k-1} \cdot (1/m)^{k+2}$. The required bound on the probability of some good spread $S \subset V(\mathcal{P}_p)$ having $Y(S) \geq \varepsilon m/4$ thus follows from a union bound over the at most n^{k-1} choices for good spread S in $V(G)$.

Bounding $\mathbb{P}[\mathcal{E}_2]$. For a pair of edges $e, f \in E(G)$ with $e \cup f$ being spread and $|e \cap f| \leq 1$ we have that $\mathbb{P}[e \cup f \subset V(\mathcal{P}_p)] \leq (2tmk/n)^{2k-1}$ by (6.1). The number of such pairs e, f with $\chi(e) = \chi(f)$ is at most μn^{2k-1} due to the bound (B) on the colouring in Theorem 1.2. Hence, by a union bound, $\mathbb{P}[\mathcal{E}_2] \leq \mu n^{2k-1} \cdot \left(\frac{2tmk}{n}\right)^{2k-1} \leq \frac{1}{m}$, using here that $\mu \ll 1/m, 1/t, 1/k$.

Bounding $\mathbb{P}[\mathcal{E}_3]$. The number of pairs of disjoint edges $e, f \in E(G)$ with $\chi(e) = \chi(f)$ is at most μn^{2k-1} due to the global bound 1 on the colouring in Theorem 1.2. If such a pair is almost spread, then $\mathbb{P}[e \cup f \subset V(\mathcal{P}_p)] \leq (2tmk/n)^{2k-1}$. Indeed this follows from (6.1) and the fact that we can remove one vertex from $e \cup f$ to get a spread set. Hence by a union bound, $\mathbb{P}[\mathcal{E}_3] \leq \mu n^{2k-1} \cdot \left(\frac{2tmk}{n}\right)^{2k-1} \leq \frac{1}{m}$, as required.

Bounding $\mathbb{P}[\mathcal{E}_4]$. Let $S \in \binom{V(G)}{j}$ be some j -set in $V(G)$ and define $E_G(S) - S := \{e \setminus S : e \in E_G(S)\} \subset \binom{V(G)}{k-j}$ to be the collection of subsets that form an edge with S . Moreover, let $N'(S) \subseteq E_G(S) - S$ be the subcollection of sets A such that A is spread and no vertex in A is close to a vertex in $S \cup P_0$. Note that $N'(S)$ can be obtained from $E_G(S) - S$ by deleting at most $2tk^2 n^{k-1-j} + n^{k-1-j}$.

$2tk \leq \beta n^{k-j}$ sets and so $|N'(S)| \geq (\delta_j^k(1) + \varepsilon - \beta)n^{k-j}$ by our minimum j -degree condition on G in the statement of the lemma.

Let now $X(S) := |\{A \in N'(S) : A \subset V(\mathcal{P}_p)\}|$ be the random variable that counts the number of sets in $N'(S)$ whose vertices are contained in the random set $V(\mathcal{P}_p)$. The fact that $N'(S)$ does not contain any close pairs implies that vertices of sets in $N'(S)$ appear in $V(\mathcal{P}_p)$ independently. Due to this as well as our lower bound on $N'(S)$ and the lower bound in (6.1), we have that

$$\lambda' := \mathbb{E}[X(S)] \geq (\delta_j^k(1) + \varepsilon - \beta)n^{k-j} \cdot \left(\frac{(1-\beta)M}{n}\right)^{k-j} = (\delta_j^k(1) + 7\varepsilon/8)M^{k-j}, \quad (6.2)$$

using here that $\beta \ll \varepsilon$. Let Γ be the auxiliary graph with vertex set $N'(S)$ such that two sets $A, A' \in N'(S)$ are adjacent in Γ whenever there is a pair of close vertices u, u' with $u \in A$ and $u' \in A'$. Then Γ is a strong dependency graph for the indicator random variables $\{A \subset V(\mathcal{P}_p)\}_{A \in N'(S)}$ and we are in the setting of Suen's inequality (Lemma 2.7). For $A, A' \in N(G')$ let $s(A, A')$ denote the size of a largest spread subset of $A \cup A'$ and note that when $AA' \in E(\Gamma)$ (equivalently, $A \sim A'$) one necessarily has $k-j \leq s(A, A') \leq 2(k-j) - 1$. Then in the notation of Lemma 2.7, we have that

$$\Delta_{X(S)} \leq \sum_{s=k-j}^{2(k-j)-1} \sum_{A \sim A' : s(A, A')=s} \mathbb{P}[A \cup A' \subset V(\mathcal{P}_p)] \leq \sum_{s=k-j}^{2(k-j)-1} n^s \cdot (4tk)^{2k} \left(\frac{2tmk}{n}\right)^s \leq (4tk)^{6k} m^{2(k-j)-1},$$

using (6.1). Similarly we have that $\delta_{X(S)} \leq 2 \cdot (4tk) \cdot n^{k-1-j} (2tmk/n)^{k-j} \leq 1$, using that sets in $N'(S)$ do not contain pairs of close vertices and the fact that $1/n \ll 1/m, 1/t, 1/k$ (3.1). Therefore appealing to Lemma 2.7, and using (6.2) we get that

$$\begin{aligned} \mathbb{P}\left[X(S) < \left(\delta_j^k(1) + \frac{3\varepsilon}{4}\right)M^{k-j}\right] &\leq \mathbb{P}\left[X(S) \leq \lambda' - \frac{\varepsilon}{8}M^{k-j}\right] \\ &\leq \exp\left(-\frac{\varepsilon^2 M^{2(k-j)}}{512(4tk)^{6k} m^{2(k-j)-1}}\right) \leq \exp(-\sqrt{m}), \end{aligned} \quad (6.3)$$

where in the last inequality we used $1/m \ll 1/t, \varepsilon, 1/k$.

Finally then we need to sum over all possible $S \in \binom{V(G)}{j}$. For any such S , let $s(S)$ be the maximal size of a subset of S which is spread and contains no vertices close to P_0 . Then $\mathbb{P}[S \subset V(\mathcal{P}_p)] \leq (2tmk/n)^{s(S)}$ using (6.1) and the fact that vertices in such a maximal subset appear independently. Moreover, as sets in $N'(S)$ do not contain vertices close to S , we have that the events $\{S \subset V(\mathcal{P}_p)\}$ and $\{X(S) < (\delta_j^k(1) + 3\varepsilon/4)M^{k-j}\}$ are independent. Thus by a union bound

$$\begin{aligned} \mathbb{P}[\mathcal{E}_4] &\leq \sum_{S \in \binom{V(G)}{j}} \mathbb{P}[S \subset V(\mathcal{P}_p)] \cdot \mathbb{P}\left[X(S) < \left(\delta_j^k(1) + 3\varepsilon/4\right)M^{k-j}\right] \\ &\leq \sum_{s=0}^j \sum_{\{S: s(S)=s\}} \left(\frac{2tmk}{n}\right)^s \exp(-\sqrt{m}) \\ &\leq \sum_{s=0}^j n^s \cdot ((s+tk)(4tk))^{j-s} \cdot \left(\frac{2tmk}{n}\right)^s \cdot \exp(-\sqrt{m}) \\ &\leq \frac{1}{m}, \end{aligned}$$

where the second summand in the penultimate line accounts for choosing vertices in S that are close to a vertex in P_0 or a vertex in the maximal spread subset of S with size $s(S)$.

Bounding $\mathbb{P}[\mathcal{E}_5]$. The expected number of vertices in \mathcal{P}_p that are close to some vertex in $V(P_0)$ is at most

$$(3t+1)k \cdot \left(\frac{2tmk}{n} \right) \leq \frac{1}{2m},$$

by (6.1). Moreover, again using the upper bound of (6.1), we have that the expected number of pairs of vertices that are close and appear in distinct paths in \mathcal{P}_p is at most

$$n \cdot 4tk \cdot \left(\frac{2tmk}{n} \right)^2 \leq \frac{1}{2m}.$$

Therefore by Markov's inequality, the probability that there is some pair of close vertices that appear in distinct paths in \mathcal{P}_* is at most $1/m$, as required.

Next we show that any splitting as in Lemma 6.1 admits a viable partition. We will first consider a random partition and show that it has some of the desired properties, for example satisfying the degree condition 2 of Definition 3.9. We will then utilise the directed version of Dirac's theorem for Hamilton cycles in digraphs [20] (see Theorem 6.3 below) to dictate some small adjustments to this random partition that will result in a viable partition.

Let us first recall some relevant notation and terminology for digraphs. A *digraph* \vec{D} has a vertex set $V(\vec{D})$ and edge set $\vec{E}(\vec{D})$ consisting of ordered pairs \vec{uv} of vertices $u \neq v \in V(\vec{D})$. The *out-degree* of a vertex $u \in V(\vec{D})$ is $d^+(u) := |\{\vec{uv} \in \vec{E}(\vec{D}) : v \in V(\vec{D})\}|$ and the *in-degree* of $u \in V(\vec{D})$ is $d^-(u) := |\{\vec{vu} \in \vec{E}(\vec{D}) : v \in V(\vec{D})\}|$. The *minimum out-degree* is $\delta^+(\vec{D}) := \min\{d^+(u) : u \in V(\vec{D})\}$ and the *minimum in-degree* $\delta^-(\vec{D})$ is defined analogously. A *Hamilton dicycle* \vec{H} in an n' -vertex digraph \vec{D} is a subdigraph of \vec{D} defined by the edge set $E(\vec{H}) = \{\vec{u_i u_{i+1}} : i = 0, \dots, n' - 1\}$ for some labelling of the vertices of \vec{D} as $V(\vec{D}) = \{u_0, \dots, u_{n'-1}\}$ (and taking $u_{n'} := u_0$). Ghouila-Houri [20] proved the following natural generalisation of Dirac's theorem to digraphs.

Theorem 6.3 (Dirac's theorem for digraphs [20]). *If \vec{D} is an n' -vertex digraph with $\delta^-(\vec{D}), \delta^+(\vec{D}) \geq n'/2$, then \vec{D} contains a Hamilton dicycle.*

We are now in a position to prove the second part of Lemma 3.11, by finding a viable partition of our suitable splitting.

Lemma 6.4. *For a loose Hamilton cycle H in G and a path $P_0 \subset H$ of length t , suppose that \mathcal{P} is a t -balanced m -splitting of H suitable for P_0 and $\delta_j(G[V(\mathcal{P})]) \geq (\delta_j^k(1) + \frac{3\varepsilon}{4})M^{k-j}$. Then $V(\mathcal{P})$ admits a viable partition.*

Proof. Let $\mathcal{P} = \{P_0, \dots, P_{m-1}\}$ and for each $i = 0, \dots, m-1$, label the endvertices of the path P_i as $V^E(P_i) := \{u_a^i, u_b^i\}$ in such a way that if we start at some edge $e \in E(H) \setminus E(P_i)$ and traverse H according to the implicit orientated cyclic order on $E(H)$, we meet the edge of P_i containing u_a^i before meeting the edge of P_i containing u_b^i . Further, we fix $U_a := \{u_a^i : i = 0, \dots, m-1\}$ and $U_b := \{u_b^i : i = 0, \dots, m-1\}$ so that $V^E(\mathcal{P}) = U_a \cup U_b$. Finally we follow the convention that $u_a^m := u_a^0, u_b^m := u_b^0, u_a^{-1} := u_a^{m-1}$ and $u_b^{-1} := u_b^{m-1}$. \square

We proceed with the following claim.

Claim 6.5. *There exists a transverse partition $\mathcal{Y} = \{Y_1, \dots, Y_T\}$ of $V(\mathcal{P})$ with respect to \mathcal{P} (with parts of size m) such that for each $h \in [T]$ we have that*

- (A) $|Y_h \cap U_b| = \tilde{m}$;
- (B) $|Y_h \cap U_a| \leq \beta m$; and
- (C) $\deg_G(S; Y_h) \geq (\delta_j^k(1) + \frac{5\varepsilon}{8})m^{k-j}$ for all $S \in \binom{V(\mathcal{P})}{j}$.

Proof of Claim.: The claim is proven by considering a uniformly random transverse partition. We will show that with probability at least $1/(4\sqrt{\tilde{m}})^{t(k-1)}$ property (A) holds for all Y_h . On the other hand, we will show that the probability that (B) or (C) fails is at most $e^{-\sqrt{m}}$. Hence using that $\tilde{m} = t^{100k}$ and $m \geq \tilde{m}$ by (3.2), with positive probability the conditions are all satisfied and such a transverse partition does indeed exist.

First let us lower bound the probability of property (A) occurring for all Y_h . Note that we can generate the uniformly random partition \mathcal{Y} via the following random process. We first choose Y_1 by taking a uniformly random vertex from each of the $V(P_i)$ with $i = 0, \dots, m-1$. Next, we choose Y_2 by taking a uniformly random vertex from each of the remaining sets $V(P_i) \setminus Y_1$ with $i = 0, \dots, m-1$ and so on. At the last step our random choice for $Y_{t(k-1)}$ also defines Y_T as the only remaining vertex from each of the $V(P_i)$ with $i = 0, \dots, m-1$. Now we have that

$$\mathbb{P}[(A)] = \prod_{h=1}^{t(k-1)} \mathbb{P}[|Y_h \cap U_b| = \tilde{m} \mid |Y_{h'} \cap U_b| = \tilde{m} \text{ for all } h' < h], \quad (6.4)$$

noting again that Y_T is fully determined by the previous choices of $Y_{h'}$ and if each of the choices satisfy $|Y_{h'} \cap U_b| = \tilde{m}$ then certainly we will have $|Y_T \cap U_b| = \tilde{m}$. Now for a given $h \in [t(k-1)]$, given that $|Y_{h'} \cap U_b| = \tilde{m}$ for all $h' < h$, we have that there exactly $m - (h-1)\tilde{m} = \tilde{m} \cdot (t(k-1) + 2 - h)$ indices $i \in \{0, \dots, m-1\}$ such that u_b^i is available for Y_h and not used in previous $Y_{h'}$. Each of these vertices is chosen for Y_h with probability exactly $1/(t(k-1) + 2 - h)$. Thus in expectation we have that $|Y_h \cap U_b| = \tilde{m}$ and as the choices are independent for the different $i \in \{0, \dots, m-1\}$ we have that $|Y_h \cap U_b|$ is binomially distributed. Using Lemma 2.6 to lower bound each of the terms in the product in (6.4) thus gives our required lower bound on the probability.

We now turn to (B). Fixing a $h \in [T]$ we have that each vertex in U_a lies in Y_h with probability $1/T$. Hence $\mathbb{E}[|U_a \cap Y_h|] = m/T = \tilde{m}$. Moreover the events $\{u \in Y_h\}$ are independent for different $u \in U_a$. Therefore we are in the setting of Chernoff's inequality (Theorem 2.5) and as $1/t \ll \beta$, we get that $\mathbb{P}[|U_a \cap Y_h| \geq \beta m/4] \leq \exp(-\beta m/4)$. Taking a union bound over the choices of $h \in [T]$, we get that with probability at least $1 - \exp(-\sqrt{m})$ all the constraints on the sizes of these intersections are satisfied.

Finally, we address the degree condition (C). For $S \in \binom{V(\mathcal{P})}{j}$, define

$$\hat{N}(S) := \{A = e \setminus S : e \in E_{G[V(\mathcal{P})]}(S) \text{ and } A \text{ is transverse with respect to } \mathcal{P}\}$$

As $\hat{N}(S)$ is obtained from the collection of all sets of the form $e \setminus S$ with $e \in E_{G[V(\mathcal{P})]}(S)$ by deleting at most $M^{k-j-1} \cdot (tk) \leq \beta M^{k-j}$ sets, we have that $|\hat{N}(S)| \geq (\delta_j^k(1) + \frac{3\varepsilon}{4} - \beta)M^{k-j}$ for all S , using the minimum degree condition in the statement of Lemma 6.4. As there are at most M^j j -sets in $V(\mathcal{P})$ and T different Y_h , using a union bound, it suffices to show that for any fixed $S \in \binom{V(\mathcal{P})}{j}$ and any $h \in [T]$, we have that $\mathbb{P}[\hat{Z}_h(S) \leq (\delta_j^k(1) + \frac{5\varepsilon}{8})m^{k-j}] \leq \exp(-m^{3/4})$, where $\hat{Z}_h(S) = |\{A \in \hat{N}(S) : A \subseteq Y_h\}|$ counts the number of sets in $\hat{N}(S)$ that fall into Y_h . Now for any such S and $h \in [T]$ we have that $A \subset Y_h$ with probability $(1/T)^{k-j} = (m/M)^{k-j}$, as it is transverse, and so using our lower bound on $|\hat{N}(S)|$, we get that $\hat{\lambda} := \mathbb{E}[\hat{Z}_h(S)] \geq (\delta_j^k(1) + \frac{3\varepsilon}{4} - \beta)m^{k-j}$. An application of Suen's inequality (Lemma 2.7) will then give exponential concentration to this mean and imply that indeed $\mathbb{P}[\hat{Z}_h(S) \leq (\delta_j^k(1) + \frac{5\varepsilon}{8})m^{k-j}] \leq \exp(-m^{3/4})$, as required. As the calculations for $\Delta_{\hat{Z}(S)}$ and $\delta_{\hat{Z}(S)}$ are very similar to those done in bounding $\mathbb{P}[\mathcal{E}_4]$ in the proof of Lemma 6.1, we skip the details here. \square

Claim 6.5 puts us in good stead to prove the lemma. Indeed, the partition \mathcal{Y} is transverse with respect to \mathcal{P} and condition (b) of Definition 3.9 is already satisfied (with some room to spare) due to condition (C). However, we are missing condition (a) of being viable and need to give a




Figure 1. On the left, an example of a transverse partition $\mathcal{Y} = \{Y_1, Y_2, Y_3, Y_4\}$ with paths P_0, \dots, P_7 (which are just single edges of a 4-uniform hypergraph in this case) such that $V(P_i) = \{u_a^i, u_b^i, u_c^i, u_d^i\}$ (denoted in the figure simply as a, b, c, d in each row) and u_a^i and u_b^i are the endpoints of P_i . On the right, the digraph $\vec{K}_8 \setminus \vec{D}$ where \vec{K}_8 is the complete digraph on 8 vertices with all edges $\vec{ii'}$ with $0 \leq i < i' \leq 8$, and \vec{D} is the auxiliary digraph defined by \mathcal{Y} .




Figure 2. An example of Hamilton dicycle \vec{H} in the digraph \vec{D} given by the partition \mathcal{Y} given in Fig. 1.

rerouting which is compatible with the partition. Claim 6.5 gives us some help to do this as we already have that each part Y_h contains the correct number of vertices from U_b that we need for a rerouting. Hence we will leave these U_b vertices fixed and move the U_a vertices, slightly modifying the partition \mathcal{Y} in the process, in order to get some viable partition \mathcal{X} .

Now to find the reshuffling, we appeal to an auxiliary digraph \vec{D} with vertex set $V(\vec{D}) = \{0, \dots, m-1\}$ and for $i \neq i'$ we have $\vec{ii'} \in \vec{E}(\vec{D})$ if and only if u_b^i and $u_b^{i'-1}$ lie in different sets Y_h . An example of (the complement of) this digraph is given on the right side of Fig. 1. Note that, by definition, $\vec{ii_+} \notin \vec{E}(\vec{D})$ where $i_+ := i+1 \pmod m$, for all $i = 0, \dots, m-1$. Due to property (A) given by Claim 6.5, we have that $\delta^-(\vec{D}), \delta^+(\vec{D}) \geq m-1-\tilde{m} \geq m/2$ and therefore by Theorem 6.3, \vec{D} contains some Hamilton dicycle \vec{H} . See Fig. 2 for an example of such a Hamilton dicycle for the partition from Fig. 1.

Letting $\mathcal{R} := \{\{u_a^i, u_b^{i'-1}\} : \vec{ii'} \in \vec{E}(\vec{H})\}$, we have that \mathcal{R} is a *rerouting* (Definition 3.8). Indeed, any path in $H[V(G) \setminus V^I(P)]$ has endpoints u_b^{i-1} and u_a^i for some $i \in \{0, \dots, m-1\}$ and identifying a pair of vertices in \mathcal{R} results in joining two of these paths by identifying an endpoint from each. The fact that \vec{H} is a Hamilton dicycle guarantees that identifying all the pairs in \mathcal{R} results in a single loose cycle. See Fig. 3 for the pairs \mathcal{R} defined by the Hamilton dicycle in Fig. 2 for the partition \mathcal{Y} in Fig. 1.

\mathcal{Y}	Y_1	Y_2	Y_3	Y_4		\mathcal{X}	X_1	X_2	X_3	X_4
$V(P_0)$	\bullet^c	\times^a	\blacksquare^b	\bullet^d	$h_0 = 2, w_0 = u_a^0$	$V(P_0)$	\bullet^c	\times^a	\blacksquare^b	\bullet^d
$V(P_1)$	\circ^a	\times^b	\bullet^d	\bullet^c	$h_1 = 1, w_1 = u_a^1$	$V(P_1)$	\circ^a	\times^b	\bullet^d	\bullet^c
$V(P_2)$	\circ^b	\bullet^c	\bullet^d	\circ^a	$h_2 = 2, w_2 = u_c^2$	$V(P_2)$	\circ^b	\circ^a	\bullet^d	\bullet^c
$V(P_3)$	\bullet^d	\circ^b	\square^a	\bullet^c	$h_3 = 4, w_3 = u_c^3$	$V(P_3)$	\bullet^d	\circ^b	\bullet^c	\square^a
$V(P_4)$	\square^b	\blacksquare^a	\bullet^c	\bullet^d	$h_4 = 3, w_4 = u_c^4$	$V(P_4)$	\square^b	\bullet^c	\blacksquare^a	\bullet^d
$V(P_5)$	\bullet^d	\bullet^a	\blacksquare^b	\bullet^c	$h_5 = 4, w_5 = u_c^5$	$V(P_5)$	\bullet^d	\bullet^c	\blacksquare^b	\bullet^a
$V(P_6)$	\bullet^c	\blacksquare^a	\bullet^d	\square^b	$h_6 = 3, w_6 = u_d^6$	$V(P_6)$	\bullet^c	\bullet^d	\blacksquare^a	\square^b
$V(P_7)$	\square^a	\bullet^d	\bullet^c	\bullet^b	$h_7 = 1, w_7 = u_a^7$	$V(P_7)$	\square^a	\bullet^d	\bullet^c	\bullet^b

Figure 3. On the left, the pairs \mathcal{R} are identified for the partition \mathcal{Y} according to the Hamilton dicycle \vec{H} from Fig. 2. The values of h_i and the vertices w_i are then given and the resulting partition \mathcal{X} .

Now for $i \in \{0, \dots, m-1\}$, let $h_i \in [T]$ be the index such that $u_b^{i'-1} \in Y_{h_i}$ where $i' \in \{0, \dots, m-1\}$ is such that $i\vec{i}' \in \vec{E}(\vec{H})$. Further, define $w_i \in V(\mathcal{P})$ to be the unique vertex lying in the intersection of $V(P_i)$ and Y_{h_i} . Note that due to how we defined the auxiliary digraph \vec{D} , we have that $w_i \neq u_i^b$ as we guaranteed that for each edge $i\vec{i}' \in \vec{E}(\vec{H}) \subset \vec{E}(\vec{D})$ we have that u_b^i and $u_b^{i'-1}$ lie in different parts of \mathcal{Y} . It may be the case however that $u_a^i \in Y_{h_i}$ and thus $w_i = u_a^i$. Examples are given in Fig. 3.

We now define $\mathcal{X} := \{X_1, \dots, X_T\}$ to be the partition of $V(\mathcal{P})$ obtained from \mathcal{Y} by swapping all the pairs $\{u_a^i, w_i\} : i = 0, \dots, m-1\}$. Formally, we have that for $h \in [T]$,

$$X_h := (Y_h \setminus (U_a \cup \{w_{h_i} : i \in I_h\})) \cup \{w_i : i \in \{0, \dots, m-1\} \text{ with } u_a^i \in Y_h\} \cup \{u_a^i : i \in I_h\},$$

where $I_h := \{i \in \{0, \dots, m-1\} : h_i = h\}$. See Fig. 3 for an example.

Note that whenever we remove a vertex in $Y_h \cap V(P_i)$ for some $i \in \{0, \dots, m-1\}$ it is replaced by some other vertex in $Y_h \cap V(P_i)$ and so \mathcal{X} is again a transverse partition of $V(\mathcal{P})$ into parts of size m . Moreover, due to condition (A) of Claim 6.5, the fact that $w_i \notin U_b$ for all $i \in \{0, \dots, m-1\}$ and how we defined the partition \mathcal{X} , we have that each X_h contains exactly \tilde{m} pairs of the rerouting \mathcal{R} . Thus condition (a) of being viable (Definition 3.9) is satisfied. Note also that for each $h \in [T]$, we have that

$$|Y_h \setminus X_h| \leq |Y_h \cap U_a| + |I_h| \leq \beta m + \tilde{m} \leq \varepsilon m/8,$$

using properties (A) and (B) of Claim 6.5 and the fact that any index in I_h corresponds to a vertex in $U_b \cap Y_h$. Thus for any $S \subset X_h$ with $|S| = j$, there are at most $\varepsilon m^{k-j}/8$ sets in $\binom{Y_h}{k-j} \setminus \binom{X_h}{k-j}$. Hence the j -degree of S relative to Y_h is close to its j -degree in X_h . Thus, due to property (C) of Claim 6.5, we get that for each $h \in [T]$, we have $\delta_j(G[X_h]) \geq (\delta_j^k(1) + \frac{\varepsilon}{2})m^{k-j}$, verifying property (b) of being viable (Definition 3.9) and completing the proof of the lemma.

With Lemmas 6.1 and 6.4, Lemma 3.11 follows easily.

Proof of Lemma 3.11. Recall that H is a loose Hamilton cycle and $P_0 \subset H$. Let \mathcal{P} denote the collection of t -balanced m -splittings \mathcal{P} of H that are suitable for $P_0 \in \mathcal{P}$ and such that $V(\mathcal{P})$ admits a viable partition. Moreover, let \mathcal{P}_* be the output by the random procedure in Lemma 6.1. Then by Lemmas 6.1 and 6.4, we have that

$$1/m \leq \mathbb{P}[\mathcal{P}_* \in \mathcal{P}] = \sum_{i=1}^{n+1} \mathbb{P}[\mathcal{P}_* \in \mathcal{P} \mid |\mathcal{P}_*| = i] \mathbb{P}[|\mathcal{P}_*| = i] \leq \mathbb{P}[\mathcal{P}_* \in \mathcal{P} \mid |\mathcal{P}_*| = m],$$

using that $\mathbb{P}[\mathcal{P}_* \in \mathcal{P} \mid |\mathcal{P}_*| = i] = 0$ for all $i \neq m$ in the last inequality. Given that $|\mathcal{P}_*| = m$, each of the possible $\binom{n/(k-1)}{m-1}$ outcomes of \mathcal{P}_* (given by a choice of $E_p \subset E(H)$ with $|E_p| = m - 1$), are equally likely. Hence

$$|\mathcal{P}| = \mathbb{P}[\mathcal{P}_* \in \mathcal{P} \mid |\mathcal{P}_*| = m] \binom{\frac{n}{k-1}}{m-1} \geq \frac{1}{m} \binom{\frac{n}{k-1}}{m-1} \geq \gamma n^{m-1},$$

as required, using here that $\gamma \ll 1/m, 1/k$. \square

7. Concluding remarks

In this paper, we made progress towards the meta-conjecture of Coulson, Keevash, Perarnau, and Yepremyan [12] that whenever one is above the extremal threshold for a given spanning structure, one can find a rainbow copy of that spanning structure in any suitably bounded colouring. Indeed, Theorem 1.2 confirms this phenomenon for loose Hamilton cycles in hypergraphs. We also elucidate what constitutes a *suitable* bound on the colouring, with the global bound (B) in Theorem 1.2 capturing the optimal qualitative conditions in the setting of loose Hamilton cycles. As mentioned in the introduction, for F -factors in hypergraphs, the theorem of Coulson, Keevash, Perarnau, and Yepremyan [12] requires an extra *local* bound that every $(k - 1)$ -set is in $o(n)$ edges of the same colour. This is necessary in some cases. Indeed, with $k \geq 3$ consider G to be the complete k -graph and assign a unique colour to each $(k - 1)$ -set of vertices of G . If we then colour the edges of G by ordering its vertices and colouring each edge by the colour corresponding to the first $(k - 1)$ -set of vertices in the edge, we have that this colouring is globally but not locally bounded. Moreover, if F is such that every $(k - 1)$ -set of vertices of F is contained in at least two edges of F (for example, if F is a clique on at least $k + 1$ vertices), then one can see that any F -factor in G is *not* rainbow. This justifies the necessity of the local bound in general. However, as mentioned in the introduction, one can use our proof, in particular our method in Lemma 5.1 using a partition to avoid edges of the bad graph B , to remove the need for a local bound when F is *linear*, that is, when no pair of edges of F intersect in more than one vertex. It remains to fully determine optimal “suitable” conditions on bounds for *all* F -factors.

Even more pertinent is the meta-conjecture for other types of Hamilton cycles in hypergraphs, where we lack any sort of positive result. Focusing in on the other extreme to loose Hamilton cycles in hypergraphs, for tight Hamilton cycles the only result known is due to Dudek, Frieze, and Ruciński [17] who showed the existence of rainbow tight cycles in the complete k -uniform hypergraph and under a strong global bounded condition of having $o(n)$ edges of each colour. It would be interesting to generalise this result to Dirac hypergraphs and establish an analogue of Theorem 1.2 in the context of tight cycles (or indeed any Hamilton ℓ -cycle with $\ell \geq 2$). We remark that the proof scheme used here which appeals to switchings and double counting, seems to break down for these cases as tight Hamilton cycles are, in a sense, *more* connected and so there is less flexibility to locally alter a Hamilton cycle to obtain (enough) new ones. Ultimately, it would be interesting to establish optimal conditions on the boundedness needed to guarantee the existence of tight Hamilton cycles (and the other ℓ -cycles) in Dirac graphs, as we have for loose cycles. This was already raised explicitly in [4]. Even in the complete case, this remains mysterious and Dudek, Frieze and Ruciński [17] asked if some strong global bound (or a local bound as above) is in fact necessary. That is, they asked if there is some colouring of the complete k -graph with $\Theta(n)$ edges of each colour and no rainbow tight Hamilton cycle. They also suggested that they believe the answer should be positive. Alternatively, it could be that a weak global bound of $o(n^{k-1})$ edges of each colour (as in Theorem 1.2) already suffices to force a rainbow tight Hamilton cycle. This remains open but the following example shows that, at least in some range of minimum degree, bounding the colouring further than the weak global bound *is* necessary.




Figure 4. The construction for Example 7.1 in the case that $n \in 3\mathbb{N}$.

Example 7.1. For any $0 < \alpha < 2/3$ and n sufficiently large, there exists an n -vertex 3-graph G with $\delta_2(G) \geq \alpha n$ and a colouring of G with no colour appearing more than n times such that there is no rainbow tight Hamilton cycle.

Proof. We first define the graph G and the colouring, see Fig. 4. Partition the vertex set V of G into three vertex subsets V_1, V_2, V_3 such that $|V_i| \in \{\lfloor n/3 \rfloor, \lceil n/3 \rceil\}$ for each $i \in [3]$. For every pair of vertices $\{u, v\} \in \binom{V}{2}$ such that $\{u, v\} \subset V_i$ for some $i \in [3]$, define a unique colour $c(u, v)$. Now the edges of G are precisely the sets $S = \{u, v, w\}$ such that $|S \cap V_{i'}| = 2$ for some $i' \in [3]$ and we define the colouring of $S \in E(G)$ to be $c(S \cap V_{i'})$. The fact that $\delta_2(G) \geq \alpha n$ follows from the fact that $\alpha < 2/3$ and n is sufficiently large. It is also clear that each colour appears at most $2\lceil n/3 \rceil \leq n$ times. To see that G does not contain a rainbow tight Hamilton cycle, suppose for a contradiction that it does and let v_1, v_2, \dots, v_n be a cyclic ordering of V corresponding to the cycle. Moreover, without loss of generality, suppose that $v_1 \in V_1$ and $v_2 \in V_2$. Let $t \in \mathbb{N}$ be the least integer such that $v_t \in V_3$ and consider $v_{t-2}, v_{t-1} \in V_1 \cup V_2$. We must have that there is some $i_* \in [2]$ such that $v_{t-2}, v_{t-1} \in V_{i_*}$. Indeed otherwise, as the ordering corresponds to a tight Hamilton cycle, there would be an edge $\{v_{t-2}, v_{t-1}, v_t\} \in E(G)$ with one vertex in each part, which is not possible. Now $v_{t-3} \notin V_{i_*}$ as no edge is completely contained in V_{i_*} and so we must have that $v_{t-3} \in V_{3-i_*}$. Therefore both $\{v_{t-3}, v_{t-2}, v_{t-1}\}$ and $\{v_{t-2}, v_{t-1}, v_t\}$ lie on the Hamilton cycle and have colour $c(\{v_{t-2}, v_{t-1}\})$, meaning that the Hamilton cycle is not rainbow, a contradiction. \square

It was proven by Rödl, Ruciński and Szemerédi [32] that the codegree threshold for tight Hamilton cycles $\delta_2^3(2)$ (see Definition 1.1) is located at $1/2$. Therefore Example 7.1 shows that in the minimum degree range $1/2 < \delta(G) < 2/3$, there are 3-graphs above the extremal threshold for which an $o(n^2)$ global bound on the colouring does not force a rainbow Hamilton cycle and some extra local bound is necessary. In general, we believe that the optimal bounds on colourings for tight Hamilton cycles (and other Hamilton ℓ -cycles also) may depend on the minimum degree of the Dirac hypergraph in question and it would be interesting to explore this further.

Finally we remark that we have only focused on qualitative features of bounding colourings and one may also ask for the best possible quantitative bounds, for example trying to optimise the constant $\mu > 0$ in the global bound (B) in Theorem 1.2. This seems like a very challenging problem in general, even for the simplest case of perfect matchings in globally bounded colourings of complete bipartite graphs. Here, the constant of Erdős and Spencer [19] has been improved slightly by Bissacot, Fernández, Procacci and Scoppola [5] who showed that if no colour appears more than $27n/256$ times, then there is a rainbow perfect matching. From above, it is clear that a global bound of n is necessary in order to have enough colours. It follows from a recent construction of Pokrovskiy and Sudakov [31] (see also the very recent improvement [10]) that being globally n -bounded is not enough to guarantee a rainbow perfect matching, but it is still possible that a bound of the form $n - o(n)$ suffices.

References

- [1] Albert, M., Frieze, A. and Reed, B. (1995) Multicoloured Hamilton cycles. *Electron. J. Comb.* **2**(1) R10.
- [2] Alon, N. and Spencer, J. (2016) *The probabilistic method*. Fourth ed. Wiley Series in Discrete Mathematics and Optimization. John Wiley & Sons, Inc, Hoboken, NJ.
- [3] Alvarado, J. D., Kohayakawa, Y., Lang, R., Mota, G. O. and Stagni, H. (2023) *Resilience for loose Hamilton cycles*. arXiv preprint arXiv: [2309.14197](https://arxiv.org/abs/2309.14197). to appear in Forum of Mathematics, Sigma.
- [4] Antoniuk, S., Kamčev, N. and Ruciński, A. (2023) Properly colored Hamilton cycles in Dirac-type hypergraphs. *Electron. J. Comb.* **30** P1–44.
- [5] Bissacot, R., Fernández, R., Procacci, A. and Scoppola, B. (2011) An improvement of the Lovász local lemma via cluster expansion. *Comb. Prob. Comp.* **20**(5) 709–719.
- [6] Böttcher, J., Schacht, M. and Taraz, A. (2009) *Proof of the bandwidth conjecture of Bollobás and Komlós*. *Math. Ann.* **343**(1) 175–205.
- [7] Brualdi, R. A. and Ryser, H.J. (1991) *Combinatorial matrix theory*, Vol. 39, Springer.
- [8] Buß, E., Hàn, H. and Schacht, M. (2013) *Minimum vertex degree conditions for loose Hamilton cycles in 3-uniform hypergraphs*. *J. Comb. Theory, Ser. B* **103**(6) 658–678.
- [9] Cano, P., Perarnau, G. and Serra, O. (2017) Rainbow spanning subgraphs in bounded edge-colourings of graphs with large minimum degree. *Electron. Notes Discrete Math.* **61** 199–205.
- [10] Chakraborti, D., Christoph, M., Hunter, Z., Montgomery, R. and Petrov, T. (2024) Almost-full transversals in equi- n -squares. arXiv preprint arXiv: [2412.07733](https://arxiv.org/abs/2412.07733).
- [11] Chernoff, H. (1952) A measure of asymptotic efficiency for tests of a hypothesis based on the sum of observations. *Ann. Math. Stat.* **23** 493–507.
- [12] Coulson, M., Keevash, P., Perarnau, G. and Yepremyan, L. (2020) Rainbow factors in hypergraphs. *J. Comb. Theory, Ser. A* **172** 105184.
- [13] Coulson, M. and Perarnau, G. (2019) Rainbow matchings in Dirac bipartite graphs. *Random Struct. Algor.* **55**(2) 271–289.
- [14] Coulson, M. (2020) A rainbow Dirac’s theorem. *SIAM J. Discrete. Math.* **34**(3) 1670–1692.
- [15] de Oliveira Bastos, J., Mota, G. O., Schacht, M., Schnitzer, J. and Schulerburg, F. (2017) Loose Hamiltonian cycles forced by large $(k - 2)$ -degree—approximate version. *SIAM J. Discrete Math.* **31**(4) 2328–2347.
- [16] Dirac, G. A. (1952) Some theorems on abstract graphs. *Proc. Lond. Math. Soc.* **3**(1) 69–81.
- [17] Dudek, A., Frieze, A. and Ruciński, A. (2012) *Rainbow Hamilton cycles in uniform hypergraphs*. *Electron. J. Comb.* **19**(1) P46.
- [18] Erdős, P., Nešetřil, J. and Rödl, V. (1983) *Some problems related to partitions of edges of a graph*, Graphs and other combinatorial topics, Teubner, Leipzig 5463.
- [19] Erdős, P. and Spencer, J. (1991) Lopsided Lovász local lemma and Latin transversals. *Discrete Appl. Math.* **30**(2-3) 151–154.
- [20] Ghouila-Houri, A. (1960) Une condition suffisante d’existence d’un circuit hamiltonien. *C. R. Acad. Sci. Paris* **251** 495–497.
- [21] Glock, S. and Joos, F. (2020) A rainbow blow-up lemma. *Random Struct. Algor.* **56**(4) 1031–1069.
- [22] Hahn, G. and Thomassen, C. (1986) Path and cycle sub-Ramsey numbers and an edge-colouring conjecture. *Discrete Math.* **62**(1) 29–33.
- [23] Ha’n, H. and Schacht, M. (2010) Dirac-type results for loose Hamilton cycles in uniform hypergraphs. *J. Comb. Theory, Ser. B* **100**(3) 332–346.
- [24] Janson, S., uczak, T. L.- and Ruciński, A. (2000) *Random graphs*, Wiley-Interscience Series in Discrete Mathematics and Optimization, Wiley-Interscience, New York.
- [25] Janson, S. (1998) New versions of Suen’s correlation inequality. *Random Struct. Algor.* **13**(3-4) 467–483.
- [26] Keevash, P., Kühn, D., Mycroft, R. and Osthus, D. (2011) Loose Hamilton cycles in hypergraphs. *Discrete Math.* **311**(7) 544–559.
- [27] Komlós, J., Sárközy, G. N. and Szemerédi, E. (1997) Blow-up lemma. *Combinatorica* **17** 109–123.
- [28] Kühn, D. and Osthus, D. (2006) Loose Hamilton cycles in 3-uniform hypergraphs of high minimum degree. *J. Comb. Theory, Ser. B* **96**(6) 767–821.
- [29] Molloy, M. and Reed, B. (2002) *Graph colouring and the probabilistic method*, Vol. 23, Springer Science & Business Media.
- [30] Montgomery, R. (2023) A proof of the Ryser-Brualdi-Stein conjecture for large even n , arXiv preprint arXiv: [2310.19779](https://arxiv.org/abs/2310.19779).
- [31] Pokrovskiy, A. and Sudakov, B. (2019) *A counterexample to Stein’s Equi- n -square conjecture*. *Proc. Am. Math. Soc.* **147**(6) 2281–2287.
- [32] Rödl, V., Ruciński, A. and Szemerédi, E. (2006) A Dirac-type theorem for, 3-uniform hypergraphs. *Comb. Prob. Comp.* **15**(1-2) 229–251.
- [33] Ryser, H. J. (1967) Neuere probleme der kombinatorik. In *Vorträge über Kombinatorik*, Oberwolfach, Vol. 69, 35. no. 91.
- [34] Stein, S. K. (1975) Transversals of Latin squares and their generalizations. *Pac. J. Math.* **59**(2) 567–575.
- [35] Sudakov, B. (2017) Robustness of graph properties. In *Surveys in Combinatorics 2017*, Vol. 440 (A. Claesson, M. Dukes, S. Kitaev, D. Manlove, K. Meeks. eds), London Mathematical Society Lecture Note Series. Illustrated Publisher, Cambridge University Press, pp. 372–408.

- [36] Suen, W. C. S. (1990) A correlation inequality and a poisson limit theorem for nonoverlapping balanced subgraphs of a random graph. *Random Struct. Algor.* **1**(2) 231–242.
- [37] Talagrand, M. (1995) *Concentration of measure and isoperimetric inequalities in product spaces*, Publications Mathématiques de l’Institut des Hautes Etudes Scientifiques 81, 73–205.
- [38] Zhao, Y. (2016) Recent advances on Dirac-type problems for hypergraphs. In *Recent trends in combinatorics*, Vol. 159 (A. Beveridge, J. Griggs, L. Hogben, G. Musiker, P. Tetali, eds), The IMA Volumes in Mathematics and its Applications, Springer, Cham, pp. 145–165.