

KULIKOV SURFACES FORM A CONNECTED COMPONENT OF THE MODULI SPACE

TSZ ON MARIO CHAN AND STEPHEN COUGHLAN

Abstract. We show that the Kulikov surfaces form a connected component of the moduli space of surfaces of general type with $p_g = 0$ and $K^2 = 6$. We also give a new description for these surfaces, extending ideas of Inoue. Finally, we calculate the bicanonical degree of Kulikov surfaces and prove that they verify the Bloch conjecture.

§1. Introduction

In 1966, Burniat [Bu] constructed some new examples of surfaces of general type with $p_g = 0$ as $(\mathbb{Z}/2)^2$ -covers of the plane branched in a certain configuration of lines. Then in 1994, Inoue [I] constructed examples with the same numerical invariants as Burniat's examples by taking a $(\mathbb{Z}/2)^3$ -quotient of a hypersurface inside a product of three elliptic curves. It seemed to be well known that the two constructions were equivalent, and a proof was written down in [BC3]. The Inoue construction was then used in [BC3] to explicitly calculate the fundamental group and homology of the Burniat–Inoue surfaces. This enabled a proof that primary Burniat–Inoue surfaces (those with $K^2 = 6$) form a connected component of the moduli space of surfaces of general type, and further, that this component is actually closed under homotopy equivalence (see below for an explanation of this terminology). The first proof that Burniat–Inoue surfaces form a connected component of the moduli space appeared in [MP1].

Kulikov [K] exhibited a new example of a surface of general type with $p_g = 0$ and $K^2 = 6$ as a $(\mathbb{Z}/3)^2$ -cover of the plane branched in a different configuration of lines (see Figure 1). In this paper we show how the Kulikov

Received December 15, 2010. Revised December 23, 2011. Accepted February 15, 2012.
2010 Mathematics Subject Classification. Primary 14J29; Secondary 14J10, 14J25.

Chan's work was partially supported by the Deutscher Akademischer Austauschdienst (DAAD) Forschungsstipendien für Doktoranden.

Coughlan's work was partially supported by Deutsche Forschungsgemeinschaft (DFG) Forschergruppe 790 "Classification of algebraic surfaces and compact complex manifolds."

© 2013 by The Editorial Board of the *Nagoya Mathematical Journal*




Figure 1: The Kulikov configuration.

surfaces fit into the same framework as Burniat–Inoue surfaces by exhibiting an Inoue-type construction. We go on to prove that Kulikov surfaces form a 1-dimensional connected component of the moduli space, which is closed under homotopy equivalence.

The layout and main results of this paper are as follows. In Section 2 we discuss preliminary results on abelian covers of the plane branched in line configurations, and we use these to define the Kulikov surface. In Section 3 we prove that there is an Inoue-type construction for Kulikov surfaces and that they form a 1-dimensional irreducible component of the moduli space of surfaces of general type. We also describe three interesting degenerate Kulikov surfaces, which can be viewed as the boundary points of the compactified moduli space.

In Section 4, following ideas employed in [BC3], we use the Inoue-type construction to explicitly calculate the fundamental groups and homology of our surfaces.

THEOREM 1.1. *If X is a Kulikov surface, then $\pi_1(X) = \Gamma$ and $H_1(X, \mathbb{Z}) = (\mathbb{Z}/3)^3$.*

Here Γ is a certain infinite subgroup of the affine group $\mathbb{A}(3, \mathbb{C})$, described in terms of generators in Section 4. Having determined the fundamental group, we are able to prove our second main result.

THEOREM 1.2. *The moduli space of Kulikov surfaces is closed under homotopy equivalence.*

In other words, any compact complex surface which is homotopy equivalent to a Kulikov surface is given by our construction. We say that the moduli space is *closed under homotopy equivalence*. This implies that the pencil of Kulikov surfaces forms a connected component of the moduli space of surfaces of general type with $p_g = 0$ and $K^2 = 6$. One might conjecture that any surface with the same π_1 is given by our construction. We have not been able to prove this, because the proof of Proposition 4.5(2) relies on homotopy equivalence.

Section 5 contains a calculation of the degree of the bicanonical map of the Kulikov surfaces.

THEOREM 1.3. *The bicanonical morphism of a Kulikov surface is birational.*

This is the expected result; by [MP2] and [MP1], the possible values are 1 or 2, but it would be surprising if the degree did not divide 3. In the course of calculating the degree of the bicanonical map, we prove Proposition 5.1, which gives a formula for the eigenspace decomposition of the bicanonical sheaf ω_X^2 for an abelian cover. This complements the standard formulas of [P, Proposition 4.1], and we believe that it is of independent interest.

In our last section, we verify the Bloch conjecture on 0-cycles for Kulikov surfaces. Let $A_0^0(X)$ denote the group of 0-cycles of degree 0 on X , modulo rational equivalence.

THEOREM 1.4. *The Kulikov surface X has $A_0^0(X) = 0$.*

The theorem is proved using a well-known method of [IM], which involves a careful examination of various quotient surfaces associated to the maximal abelian cover $\widehat{X} \rightarrow X$. Unfortunately, the final step in our proof is done by computer algebra.

During the final preparation of the manuscript, we were informed by V. Alexeev that he and R. Pardini also computed the degenerate Kulikov surfaces of Section 3.4, using methods similar to those of [AP].

§2. Surfaces as abelian covers

2.1. Abelian covers

We briefly review the preliminary results needed for our calculations. More details can be found in several articles, including [BC1], [C], and [P]. Assume that Y is nonsingular, and let $\varphi: X \rightarrow Y$ be a finite morphism with a faithful action of the abelian group G on X so that Y is the quotient. Let

Δ be the branch divisor of φ ; then φ is determined by the surjective homomorphism $\Phi: \pi_1(Y \setminus \Delta) \rightarrow G$, which factors through $\Phi: H_1(Y \setminus \Delta, \mathbb{Z}) \rightarrow G$ since G is abelian. In this paper, Y is always \mathbb{P}^1 , \mathbb{P}^2 , or a nonsingular del Pezzo surface, and Δ is a configuration of points or lines; hence,

$$H_1(Y \setminus \Delta, \mathbb{Z}) \cong \frac{\bigoplus_{i=1}^n \mathbb{Z}\langle \Delta_i \rangle}{\Delta},$$

where Δ_i are the irreducible components of Δ so that $\Delta = \sum_{i=1}^n \Delta_i$.

Now suppose that P is a point of intersection of two or more branch lines l_1, \dots, l_k . Then X is nonsingular over P if and only if there are exactly two lines intersecting transversely at P , and

$$(1) \quad \langle \Phi(l_1), \Phi(l_2) \rangle = \langle \Phi(l_1) \rangle \oplus \langle \Phi(l_2) \rangle \subset G.$$

If we blow up P to obtain the exceptional curve E , then there is an induced group homomorphism on the blowup, which we still call Φ , defined by

$$(2) \quad \Phi(E) = \sum_{i=1}^k \Phi(l_i).$$

Note that the induced cover on the blowup may be unramified over E and that the blowup may or may not resolve the singularity on X over P .

We have

$$\varphi_* \mathcal{O}_X = \bigoplus_{\chi \in G^*} \mathcal{L}_\chi^{-1},$$

where $G^* = \text{Hom}(G, \mathbb{C}^*)$ is the group of characters of G , and G acts on \mathcal{L}_χ^{-1} with character χ . Fix $\chi \in G^*$, and choose a generator for the image of χ in \mathbb{C}^* , so that we view $\chi: G \rightarrow \mathbb{Z}/d$, where d is the order of χ . Then the following formula determines the eigensheaf $\mathcal{L}_\chi = \mathcal{O}_Y(L_\chi)$:

$$(3) \quad dL_\chi = \sum_{i=1}^n (\chi \circ \Phi(\Delta_i)) \Delta_i.$$

If X is nonsingular, the ramification formula for K_X gives

$$(4) \quad K_X = \varphi^* \left(K_Y + \sum_{i=1}^n \left(1 - \frac{1}{d_i} \right) \Delta_i \right),$$

where d_i is the order of $\Phi(\Delta_i)$ in G , and

$$(5) \quad \varphi_* \omega_X = \bigoplus_{\chi \in G^*} \omega_Y(L_\chi).$$

2.2. The Kulikov surface

In this section we construct the Kulikov surface. We also define the groups G^0, G^1, G^2 which appear throughout this article. Let $\varphi: \widehat{X} \rightarrow \mathbb{P}^2$ be the $(\mathbb{Z}/3)^5$ -cover of \mathbb{P}^2 branched in the Kulikov configuration $\Delta = \sum_{i=1}^6 \Delta_i$ of Figure 1. Then φ is determined by the group homomorphism $\Phi: H_1(\mathbb{P}^2 \setminus \Delta) \rightarrow (\mathbb{Z}/3)^6$, represented by the 6×6 matrix

$$(6) \quad \Phi = \begin{pmatrix} 0 & 1 & 0 & 2 & 0 & 0 \\ 0 & 0 & 1 & 2 & 0 & 0 \\ \hline 0 & 0 & 1 & 0 & 2 & 0 \\ 1 & 0 & 0 & 0 & 2 & 0 \\ \hline 1 & 0 & 0 & 0 & 0 & 2 \\ 0 & 1 & 0 & 0 & 0 & 2 \end{pmatrix}$$

of rank 5. Since Δ is supported on six lines, Φ is unique up to choice of generators for $(\mathbb{Z}/3)^6$. Thus, \widehat{X} is maximal, which means that any other $(\mathbb{Z}/3)^k$ -cover of \mathbb{P}^2 branched in Δ factorizes φ .

DEFINITION 2.1. Let $G^1 \cong (\mathbb{Z}/3)^5$ denote the image of Φ in $(\mathbb{Z}/3)^6$. Then G^1 is generated by the columns of Φ , and we label these $\delta_1, \delta_2, \delta_3, \omega_1, \omega_2$, and ω_3 , respectively. Thus, \widehat{X} is a G^1 -cover of \mathbb{P}^2 branched in the Kulikov configuration.

The Kulikov surface $\psi: X \rightarrow \mathbb{P}^2$ is a $(\mathbb{Z}/3)^2$ -cover of \mathbb{P}^2 branched in the Kulikov line configuration and first appeared in [K]. The cover is determined by the homomorphism $\Psi: H_1(\mathbb{P}^2 \setminus \Delta) \rightarrow (\mathbb{Z}/3)^2$, given by the matrix

$$(7) \quad \Psi = \begin{pmatrix} 1 & 1 & 1 & 0 & 1 & 2 \\ 0 & 0 & 0 & 1 & 1 & 1 \end{pmatrix}.$$

Now $\varphi: \widehat{X} \rightarrow \mathbb{P}^2$ factors through the Kulikov surface, giving commutative diagrams

$$(8) \quad \begin{array}{ccc} \widehat{X} & \xrightarrow{\varphi} & \mathbb{P}^2 \\ \downarrow \tilde{\psi} & \nearrow \psi & \\ X & & \end{array} \quad \begin{array}{ccc} (\mathbb{Z}/3)^6 & \xleftarrow{\Phi} & H_1(\mathbb{P}^2 \setminus \Delta, \mathbb{Z}) \\ \downarrow \tilde{\Psi} & \swarrow \Psi & \\ (\mathbb{Z}/3)^2 & & \end{array}$$

where

$$\tilde{\Psi} = \begin{pmatrix} 0 & 0 & 1 & 1 & 0 & 1 \\ 1 & 1 & 2 & 0 & 0 & 2 \end{pmatrix}$$

and where $\tilde{\psi}: \widehat{X} \rightarrow X$ is Galois étale with group $G^2 = \text{Ker}(\tilde{\Psi}) \cap \text{Im}(\Phi)$.

DEFINITION 2.2. We define $G^2 = \text{Ker}(\tilde{\Psi}) \cap G^1$, so that $G^2 = \langle g_1, g_2, g_3 \rangle \cong (\mathbb{Z}/3)^3$, where

$$(9) \quad \begin{aligned} g_1 &= {}^t(0, 0, 1, 0, 1, 2), \\ g_2 &= {}^t(1, 2, 0, 0, 1, 0), \\ g_3 &= {}^t(1, 0, 1, 2, 0, 0). \end{aligned}$$

We also define $G^0 \subset G^1$ to be the subgroup generated by $\xi_i = \delta_{i-1}^2 \omega_i \omega_{i+1}$ for $i = 1, 2, 3$. Then G^0 is isomorphic to $(\mathbb{Z}/3)^2$ because $\xi_1 \xi_2 \xi_3 = 1$. We note the relations $g_i = \xi_{i+1} \omega_{i+1}$ for $i = 1, 2, 3$.

Note that by (1), X is singular as a cover of \mathbb{P}^2 , so we blow up the three singular points P_1, P_2, P_3 of Δ , and we view X as a nonsingular $(\mathbb{Z}/3)^2$ -cover of the degree 6 del Pezzo surface Y . Then computations using (2)–(5) show that the Kulikov surface X is a minimal surface of general type with $p_g = 0$ and $K^2 = 6$.

§3. Moduli spaces

3.1. Elliptic curves as abelian covers

This digression on elliptic curves and abelian covers is required for the Inoue construction of Section 3.2.

Let $\varphi: E \rightarrow \mathbb{P}^1$ be a Galois $(\mathbb{Z}/3)^2$ -cover branched in $\Delta = P_0 + P_1 + P_\infty$. Note that E is the (unique) Fermat cubic curve (see Remark 3.1). There are several ways to study φ , but for consistency we prefer to use the techniques of abelian covers. Now $\Phi: H_1(\mathbb{P}^1 \setminus \Delta) \rightarrow (\mathbb{Z}/3)^2$ is unique up to choice of generators and is given by

$$\Phi = \begin{pmatrix} 1 & 2 & 0 \\ 0 & 2 & 1 \end{pmatrix}.$$

Using (3) to study the various eigensheaves and quotients of the cover, we establish the following diagram:

$$(10) \quad \begin{array}{ccc} & E & \\ \pi_1 \swarrow & \downarrow \varphi & \searrow \pi_2 \\ E & & \mathbb{P}^1 \\ \rho_1 \searrow & & \swarrow \rho_2 \\ & \mathbb{P}^1 & \end{array}$$

Here π_1 is the isogeny induced by taking the quotient by the group $\langle \eta \rangle$ of translations by the 3-torsion point η . There are three possibilities for π_2 corresponding to the remaining nontrivial subgroups of $(\mathbb{Z}/3)^2$. We choose the subgroup $\langle \omega \rangle$ which fixes the three points $\varphi^{-1}(P_1)$ on E , and we label these points $0, \eta, 2\eta$. Since we have now fixed the origin of E , we may consider $\omega = e^{2\pi i/3}$ as a rotation of E , under which η is fixed. The map ρ_1 is the $\mathbb{Z}/3$ -cover of \mathbb{P}^1 branched in Δ , and ρ_2 is the $\mathbb{Z}/3$ -cover of \mathbb{P}^1 branched over $P_0 + P_\infty$. Note that in terms of the original basis for $(\mathbb{Z}/3)^2$,

$$(11) \quad \eta = \begin{pmatrix} 1 \\ 2 \end{pmatrix} \quad \text{and} \quad \omega = \begin{pmatrix} 2 \\ 2 \end{pmatrix}.$$

REMARK 3.1. Let $E: (a^3 + b^3 + c^3 = 0) \subset \mathbb{P}^2$ be the Fermat cubic curve. Then one way to write the $(\mathbb{Z}/3)^2$ -action is

$$\eta: \begin{pmatrix} a \\ b \\ c \end{pmatrix} \mapsto \begin{pmatrix} a \\ \omega b \\ \omega^2 c \end{pmatrix}, \quad \omega: \begin{pmatrix} a \\ b \\ c \end{pmatrix} \mapsto \begin{pmatrix} \omega a \\ b \\ c \end{pmatrix}.$$

The map π_2 is the projection to the line $(a = 0) \subset \mathbb{P}^2$ with coordinates (b, c) , so that $\rho_2: (b, c) \mapsto (b^3, c^3)$. Then the three ramification points $(0, 1, -1)$, $(0, 1, -\omega)$, and $(0, 1, -\omega^2)$ of π_2 are $0, \eta$, and 2η , respectively. The image of π_1 is $E: (s^3 + x^2x' + xx'^2 = 0) \subset \mathbb{P}^2$, where $x = b^3$, $x' = c^3$, and $s = abc$. This is the Fermat cubic again, via the relation $s^3 = a^3xx'$.

3.2. The Inoue-type construction

In this section we give an alternative description of the Kulikov surface, which is called the *Inoue-type construction*, after [I]. This sets the Kulikov surface into context alongside the Burniat–Inoue surfaces.

First note that after blowing up P_1, P_2, P_3 , we may think of the Kulikov configuration in Figure 1 as an element of the pencil

$$Y_{1,1,1} \subset \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$$

of degree 6 del Pezzo surfaces, with equation $\lambda x_1 x_2 x_3 = \mu x'_1 x'_2 x'_3$, where x_i, x'_i are the coordinates on the i th factor. Now let $E \xrightarrow{\pi_2} \mathbb{P}^1 \xrightarrow{\rho_2} \mathbb{P}^1$ be a $(\mathbb{Z}/3)^2$ -cover of \mathbb{P}^1 as described in Section 3.1. Taking a direct product of three copies of E , we obtain the following diagram:

$$\begin{array}{ccccc}
 E \times E \times E & \xrightarrow{\pi} & \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1 & \xrightarrow{\rho} & \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1 \\
 \cup & & \cup & & \cup \\
 (12) \quad \widehat{X} & \longrightarrow & Z \cup \cup_i Z_i & \longrightarrow & Y_{1,1,1} \\
 & \searrow \tilde{\psi} & & \nearrow \psi & \\
 & & X & &
 \end{array}$$

where $\pi = \prod \pi_2^i, \rho = \prod \rho_2^i$.

Fix an element Y of our pencil of del Pezzo surfaces. Then $\rho^{-1}(Y)$ splits into three components $Z_i, i = 0, 1, 2$, each of which has stabilizer isomorphic to $(\mathbb{Z}/3)^2$. We fix $Z = Z_0$, with stabilizer $G^0 \cong (\mathbb{Z}/3)^2$. Now define $\widehat{X} = \pi^{-1}(Z)$, so that \widehat{X} is a hypersurface in $E \times E \times E$ of tridegree $(3, 3, 3)$, with stabilizer $G^1 = G^0 \oplus (\mathbb{Z}/3)^3$. The quotient of \widehat{X} by G^1 is Y , and by construction, the branch locus of $\widehat{X} \rightarrow Y$ is exactly the blown-up Kulikov configuration. Thus, \widehat{X} is the maximal $(\mathbb{Z}/3)^5$ -cover of Section 2.2.

LEMMA 3.2. *There is a subgroup $G^2 \cong (\mathbb{Z}/3)^3$ of G^1 such that G^2 acts freely on \widehat{X} , and the quotient X is a Kulikov surface.*

Proof. Since $\widehat{X} \rightarrow Y$ is the maximal $(\mathbb{Z}/3)^5$ -cover of Y , we see that the quotient map $(\rho \circ \pi)|_{\widehat{X}} = \varphi$, where φ is determined by the matrix (6) of Section 2.2. Hence, the group G^1 is generated by the columns δ_i, ω_i of (6) as stated in Definition 2.1. By Definition 2.2, the subgroup G^2 is generated by

$$g_i = \xi_{i+1} \omega_{i+1} = \delta_i^2 \omega_{i+1}^2 \omega_{i+2},$$

for $i = 1, 2, 3$.

For completeness, we reconcile the above construction of G^0 with Definition 2.2. Observe that we subdivided the rows of matrix (6) into pairs, each of which corresponds to the action of G^1 restricted to the respective factor of $E \times E \times E$. For example, by Definition 2.2,

$$\xi_1 = {}^t(2, 1, 1, 2, 0, 0),$$

and comparing this with the definition of $\eta = \begin{pmatrix} 1 \\ 2 \end{pmatrix}$ from (11), we see that

$$\xi_1 = \eta_1^2 \eta_2,$$

where η_i now denotes translation by the 3-torsion point η on the i th factor of the product. We also have

$$\xi_2 = \eta_2^2 \eta_3, \quad \xi_3 = \eta_1 \eta_3^2.$$

This is called the *Inoue-type construction* of Kulikov surfaces (see [BC3] and [I] for the original Inoue surfaces). □

3.3. The moduli space of Kulikov surfaces

Using the Inoue-type construction, it is clear that we obtain a pencil of Kulikov surfaces as the pullback of the pencil of del Pezzo surfaces. In fact, we have the following.

THEOREM 3.3. *The pencil of Kulikov surfaces forms a 1-dimensional irreducible component of the moduli space of surfaces of general type with $p_g = 0$ and $K^2 = 6$.*

Proof. Let X be a Kulikov surface. We calculate $h^1(T_X) = h^1(\psi_* T_X)$ using the splitting of $\psi_* T_X$ into eigensheaves according to [P, Proposition 4.1]:

$$(13) \quad \begin{aligned} (\psi_* T_X)^{\text{inv}} &= T_Y(-\log \Delta), \\ (\psi_* T_X)^{(\chi)} &= T_Y(-\log \Delta_g : g \in S_\chi) \otimes \mathcal{L}_\chi^{-1}, \end{aligned}$$

where $S_\chi = \{g \in G \mid \chi(g) \neq m - 1\}$, m is the order of g , and Δ_g is the sum of the components Δ_i of Δ such that $\Phi(\Delta_i) = g$. For example, when $g = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$ we have

$$\Delta_{\begin{pmatrix} 1 \\ 0 \end{pmatrix}} = \Delta_1 + \Delta_2 + \Delta_3 = 3H - 2 \sum_{i=1}^3 E_i,$$

where E_i is the exceptional curve over P_i . The first equality follows from the definition of Ψ in (7) and by applying (2) to each exceptional curve. The remaining cases are

$$\begin{aligned} \Delta_{(1)} &= \Delta_5 + E_3 = H - E_2 + E_3, \\ \Delta_{(2)} &= \Delta_6 + E_1 = H - E_3 + E_1, \\ \Delta_{(1)}^0 &= \Delta_4 + E_2 = H - E_1 + E_2, \end{aligned}$$

and all other Δ_g are zero.

Given the decomposition (13), it is clear that

$$(14) \quad h^i(T_X) = \sum_{\chi} h^i((\psi_*T_X)^{(\chi)}).$$

We use several methods, which we describe subsequently, to calculate the value of $h^2((\psi_*T_X)^{(\chi)})$ for each χ . For convenience, we present the results in Table 1. Now, since $h^0(T_X) = 0$ for a surface of general type, we have

$$\chi(T_X) = -h^1(T_X) + h^2(T_X) = 2K_X^2 - 10\chi(\mathcal{O}_X) = 2.$$

Using this equality, Table 1, and (14), we get $h^1(T_X) = 1$, which proves the theorem. □

The remainder of this section deals with Table 1. The table contains the data required to write down the eigenspace decomposition of ψ_*T_X

Table 1: Decomposition of ψ_*T_X .

	χ	$A = K_Y + L_{\chi}$	$\{\Delta_g : g \in S_{\chi}\}$	$h^2((\psi_*T_X)^{(\chi)})$
(0)	(0, 0)	$-3H + \sum_i E_i$	Δ	0
(a)	(1, 0)	$-H + E_1$	$\Delta_1, \Delta_2, \Delta_3, \Delta_4, \Delta_5, E_2, E_3$	1
	(1, 2)	$-H + E_2$	$\Delta_1, \Delta_2, \Delta_3, \Delta_5, \Delta_6, E_1, E_3$	
	(1, 1)	$-H + E_3$	$\Delta_1, \Delta_2, \Delta_3, \Delta_4, \Delta_6, E_1, E_2$	
(b)	(0, 1)	$-2H + \sum_i E_i$	$\Delta_1, \dots, \Delta_6, E_1, E_2, E_3$	0
(c)	(0, 2)	$-H + \sum_i E_i$	$\Delta_1, \Delta_2, \Delta_3$	0
(d)	(2, 2)	$-E_1$	$\Delta_5, \Delta_6, E_1, E_3$	0
	(2, 0)	$-E_2$	$\Delta_4, \Delta_6, E_1, E_2$	
	(2, 1)	$-E_3$	$\Delta_4, \Delta_5, E_2, E_3$	

according to (13). We also remark that the divisors Δ_g are always composed of a number of disjoint rational curves. Thus, there is no need to distinguish between logarithmic poles along D_1 and D_2 , and logarithmic poles along $D_1 + D_2$ for any curves D_1, D_2 . The data are grouped into cases $(0), (a), \dots, (d)$ by symmetry considerations, avoiding repetition in our calculations.

First, observe that by Serre duality,

$$(15) \quad h^i((\psi_* T_X)^{(X)}) = h^{2-i}(\Omega_Y(\log \Delta_g : g \in S_X)(A)),$$

where $A = K_Y + L_X$. Thus, the foundation for our calculations of the various $h^1((\psi_* T_X)^{(X)})$ is the standard logarithmic residue sequence twisted by A ,

$$(16) \quad 0 \rightarrow \Omega_Y(A) \rightarrow \Omega_Y(\log \Delta_g : g \in S_X)(A) \rightarrow \bigoplus_{g \in S_X} \mathcal{O}_{\Delta_g}(A) \rightarrow 0,$$

and its associated long exact sequence of cohomology, with connecting homomorphism

$$\delta : \bigoplus_{g \in S_X} H^0(\mathcal{O}_{\Delta_g}(A)) \rightarrow H^1(\Omega_Y(A)).$$

Now if $A = 0$, then the image of δ is $\langle c_1(\Delta_g) : g \in S_X \rangle$. Thus, it is possible to calculate the rank of δ directly, because of the Lefschetz (1, 1) theorem. In turn, this gives us the value of $h^0(\Omega_Y(\log \Delta_g : g \in S_X)(A))$ using the long exact sequence. The crucial part of our proof is to adapt this argument to situations where A is nontrivial.

Note that it is possible to calculate the value of $h^1((\psi_* T_X)^{(X)})$ in each case, using the Riemann–Roch theorem together with the logarithmic residue sequence (16). However, this is not needed for the proof, so we omit this calculation.

3.3.1. The basic method. In this section we compute the last column of Table 1 in cases (0), (b), and (d). We first give the setup. Suppose that $|-A|$ has no fixed part, so there is a morphism $\mathcal{O}_Y(A) \rightarrow \mathcal{O}_Y$ which is nonzero on each irreducible component of the branch divisor Δ . Then we obtain the following commutative diagram:

$$\begin{array}{ccccccc} 0 & \longrightarrow & \Omega_Y(A) & \longrightarrow & \Omega_Y(\log \Delta_g : g \in S_X)(A) & \longrightarrow & \bigoplus_{g \in S_X} \mathcal{O}_{\Delta_g}(A) \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & \Omega_Y & \longrightarrow & \Omega_Y(\log \Delta_g : g \in S_X) & \longrightarrow & \bigoplus_{g \in S_X} \mathcal{O}_{\Delta_g} \longrightarrow 0 \end{array}$$

The connecting homomorphisms for the corresponding long exact sequences fit into the commutative square

$$\begin{array}{ccc}
 \bigoplus_{g \in S_\chi} H^0(\mathcal{O}_{\Delta_g}(A)) & \xrightarrow{\delta} & H^1(\Omega_Y(A)) \\
 \downarrow & \searrow \alpha & \downarrow \beta \\
 \bigoplus_{g \in S_\chi} \mathbb{C}_{\Delta_g} & \xrightarrow{c_1} & H^1(\Omega_Y)
 \end{array}$$

where \mathbb{C}_{Δ_g} denotes $H^0(\mathcal{O}_{\Delta_g})$, c_1 is the Chern class map, δ is the connecting homomorphism whose rank we wish to calculate, and β has kernel $H^0(\Omega_Y|_D)$, where $D \in |-A|$.

As an illustration, consider case (b), so that $\chi = (0, 1)$ and $A = -2H + E_1 + E_2 + E_3$. Clearly, A has no fixed part, and moreover,

$$\bigoplus_{g \in S_{(0,1)}} H^0(\mathcal{O}_{\Delta_g}(A)) = \mathbb{C}_{\Delta_1} \oplus \mathbb{C}_{\Delta_2} \oplus \mathbb{C}_{\Delta_3}.$$

Now since Δ_1, Δ_2 , and Δ_3 are linearly independent in $\text{Pic } Y$, $\text{rank } \alpha = 3$. Thus, $\text{rank } \delta = 3$, so $h^0(\Omega_Y(\log \Delta_g : g \in S_{(0,1)})(A)) = 0$.

Case (0) is also treated using this method, since $\bigoplus_{g \in G} H^0(\mathcal{O}_{\Delta_g}(K_Y)) = 0$, giving $h^0(\Omega_Y(\log \Delta)(K_Y)) = 0$. For case (d), suppose that $\chi = (2, 2)$, so that $A = -E_1$. Then we have a natural inclusion

$$H^0(\Omega_Y(\log \Delta_g : g \in S_{(2,2)})(-E_1)) \subseteq H^0(\Omega_Y(\log \Delta_g : g \in S_{(2,2)})),$$

and we can show that the right-hand side vanishes by using the standard logarithmic residue sequence.

3.3.2. The contraction lemma and fibration method. This section deals with case (a), where the basic method does not work. We contract some branch divisors on Y and then calculate directly. A more general version of the following lemma appears in [BC2, Lemma 2.10] and [BC4, Lemma 4.1]. We quote a simplified version, which is sufficient for our purposes.

LEMMA 3.4. *Let $\pi : Z \rightarrow \mathbb{C}^2$ be the blowup of \mathbb{C}^2 at the origin O , with exceptional curve E . Suppose that L_1, L_2 are lines through O with distinct tangents at O , and let D_i be the strict transform of L_i . Then*

$$\pi_* \Omega_Z(\log D_1, \log D_2, \log E) = \Omega_{\mathbb{C}^2}(\log L_1, \log L_2).$$

In case (a), with $\chi = (1, 0)$, we have $A = -H + E_1$. Note that the basic method does not work here: the rank of α is 3, which is not maximal, so we do not have sufficient information to calculate $\text{rank } \delta$. Instead, we contract Δ_3 and E_3 via $\pi: Y \rightarrow Q = \mathbb{P}^1 \times \mathbb{P}^1$, and we let $f: Q \rightarrow \mathbb{P}^1$ be the projection to the second factor. Then by Lemma 3.4,

$$\pi_*(\Omega_Y(\log \Delta_g: g \in S_{(1,0)})(A)) = \Omega_Q(\log(F_1 + F_2 + F_3), \log(B_1 + B_2))(-F_2),$$

where $F_1 = \pi_*\Delta_2$, $F_2 = \pi_*A$, $F_3 = \pi_*E_2$ are fibers of f , and $B_1 = \pi_*\Delta_1$, $B_2 = \pi_*\Delta_5$ are sections. Now, the standard short exact sequence

$$0 \rightarrow f^*\omega_{\mathbb{P}^1} \rightarrow \Omega_Q \rightarrow \omega_{Q/\mathbb{P}^1} \rightarrow 0$$

specializes to

$$0 \rightarrow f^*\omega_{\mathbb{P}^1}(F) \rightarrow \Omega_Q(\log F, \log B) \rightarrow \omega_{Q/\mathbb{P}^1}(B) \rightarrow 0.$$

Thus, twisting by $-F_2$ and observing that $f^*\omega_{\mathbb{P}^1} = \mathcal{O}_Q(-2F)$ and $\omega_{Q/\mathbb{P}^1} = \mathcal{O}_Q(-2B)$, we get

$$h^0(\Omega_Y(\log \Delta_g: g \in S_{(1,0)})(A)) = h^0(\mathcal{O}_Q) = 1.$$

3.3.3. Final case (c). In case (c), $A = -H + \sum E_i$, so $|-A|$ is empty and it is not possible to use the basic method. We cannot use the contraction lemma either, because when we contract Δ_i , we still have poles along E_i , and further contractions are complicated. Instead, we use the natural inclusion

$$H^0(\Omega_Y(\log(\Delta_1, \Delta_2, \Delta_3)(A))) \subseteq H^0(\Omega_Y(B)),$$

where $B = A + \Delta_1 + \Delta_2 + \Delta_3$. Then the exact sequence

$$0 \rightarrow \sigma^*\Omega_{\mathbb{P}^2} \rightarrow \Omega_Y \rightarrow \bigoplus_{i=1}^3 \mathcal{O}_{E_i}(-2) \rightarrow 0$$

gives $H^0(\Omega_Y(B)) = H^0(\sigma^*\Omega_{\mathbb{P}^2}(B))$, and the pullback of the dual Euler sequence to Y

$$0 \rightarrow \sigma^*\Omega_{\mathbb{P}^2} \rightarrow \mathcal{O}_Y(-H)^3 \rightarrow \mathcal{O}_Y \rightarrow 0$$

gives $H^0(\sigma^*\Omega_{\mathbb{P}^2}(B)) = H^0(\mathcal{O}_Y(-A)^3) = 0$. Hence,

$$h^0(\Omega_Y(\log \Delta_g: g \in S_{(0,2)})(A)) = 0.$$




Figure 2: The complete quadrangle and the cube.

3.4. Degenerate Kulikov surfaces

In this section, we construct three degenerate Kulikov surfaces arising as $(\mathbb{Z}/3)^2$ -covers of special elements of the pencil of del Pezzo surfaces

$$Y_{1,1,1}: (\lambda x_1 x_2 x_3 = \mu x'_1 x'_2 x'_3) \subset \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1.$$

These degenerate surfaces are members of the boundary of the compactified moduli space, in the sense of [KSB].

3.4.1. A hyperelliptic surface. When $\lambda = \mu = 1$, the three lines $\Delta_4, \Delta_5,$ and Δ_6 of the Kulikov configuration meet in a point P_0 . This configuration is called the *complete quadrangle*, and it is illustrated on the left side of Figure 2. The $(\mathbb{Z}/3)^2$ -cover $X \rightarrow Y$ has an elliptic singularity of degree 9 over P_0 . If we blow up Y at P_0 , then the induced $(\mathbb{Z}/3)^2$ -cover \tilde{X} is the resolution.

Note that the pencil of conics in \mathbb{P}^2 passing through P_0, \dots, P_3 gives rise to a pencil of Fermat cubics on \tilde{X} , so \tilde{X} is an elliptic surface. Moreover, the minimal model of \tilde{X} is the $(\mathbb{Z}/3)^2$ -cover of $\mathbb{P}^1 \times \mathbb{P}^1$ obtained by contracting the strict transforms of $\Delta_4, \Delta_5,$ and Δ_6 . The branch locus is transformed to six lines (three in each ruling), and the minimal model of \tilde{X} is the hyperelliptic surface $(E \times E)/(\mathbb{Z}/3)^2$.

One can also see this degeneration in terms of the Inoue-type construction of Section 3.2. The action of G^2 is free on $E \times E \times E$ outside the orbit of the point $(0, 0, 0)$, which has stabilizer $\mathbb{Z}/3$ generated by $g_1 g_2 g_3$. The degenerate Kulikov surface described above is obtained when the hypersurface $\hat{X}_{3,3,3} \subset E \times E \times E$ contains the point $(0, 0, 0)$.

3.4.2. *Two reducible surfaces.* When $\mu = 0$ or $\lambda = 0$, the del Pezzo surface Y breaks into three copies of $\mathbb{P}^1 \times \mathbb{P}^1$. We consider only the case $\mu = 0$, as shown on the right side of Figure 2, because $\lambda = 0$ has a similar treatment by symmetry. The components of Y are called $Y_i: (x_i = 0)$, and Y has normal crossing singularities along the lines $D_k: (x_i = x_j = 0)$ for $i, j, k = \{1, 2, 3\}$, indicated by dashed lines in the figure. The divisors Δ_4, Δ_5 , and Δ_6 break into pairs of lines; for example, Δ_4 consists of the two lines joining E_1 to Δ_1 , intersecting in a single point Q_1 . For clarity, we do not label Δ_4, Δ_5 , and Δ_6 in the figure.

We construct each $(\mathbb{Z}/3)^2$ -cover $\psi_i: X_i \rightarrow Y_i$ separately, before gluing them together to give the degenerate Kulikov surface. For $i = 1$, we see from the figure that ψ_1 is branched over the four lines E_1, Δ_2, Δ_4 , and Δ_5 . Recall from Section 2 that the $(\mathbb{Z}/3)^2$ -cover of Y_1 branched in Δ is governed by the group homomorphism $\Psi_1: H_1(Y_1 \setminus \Delta, \mathbb{Z}) \rightarrow (\mathbb{Z}/3)^2$. Thus, by (7), ψ_1 must also be branched over D_1 and D_2 , with $\Psi_1(D_1) = -\Psi_1(E_1) - \Psi_1(\Delta_5) = \binom{0}{1}$, $\Psi_1(D_2) = -\Psi_1(\Delta_2) - \Psi_1(\Delta_4) = \binom{2}{2}$. Then by equations (1) and (2), X_1 has three singularities of type $(1/3)(1, 1)$ over Q_2 , and three of type $(1/3)(1, 2)$ over Q_1 . If we blow up Q_2 and contract the strict transforms of D_2 and Δ_5 , the induced cover $\tilde{X}_1 \rightarrow \mathbb{P}^2$ is a rational surface with three $(1/3)(1, 2)$ singularities.

The other components of X have similar singularities over the points Q_i . Perhaps the most interesting aspect of this example is that when we glue the components of X back together, we see that over each Q_i , we must attach a $(1/3)(1, 2)$ point to a $(1/3)(1, 1)$ point. This gives rise to orbifold normal crossing singularities, which can be expressed locally as

$$(xy = 0) \subset \frac{1}{3}(1, 2, 1),$$

where the double curve is given by $(x = y = 0)$.

§4. Fundamental groups and homotopy equivalence

In this section, we give an explicit description of the fundamental group of the Kulikov surface. Using this description, we show that the 1-dimensional irreducible component of the moduli space constructed in Section 3 is actually closed under homotopy equivalence. Hence, the Kulikov surfaces form a connected component of the moduli space.

4.1. Fundamental group and homology

Using the Inoue-type construction of Section 3.2, it is possible to lift the action of $G^2 = \langle g_1, g_2, g_3 \rangle$ to the Fermat cubic curves. Indeed, let z_i be a uniformizing parameter for the i th curve in the direct product (12). Recall that in the proof of Lemma 3.2, we showed that $\xi_i = \eta_i^2 \eta_{i+1}$ for $i = 1, 2, 3$. Now by Definition 2.2, we have $g_i = \xi_{i+1} \omega_{i+1}$. Hence, the action of G^2 lifts to the product $E \times E \times E$ as

$$g_1(z) = \begin{pmatrix} z_1 \\ \omega z_2 + 2\eta_2 \\ z_3 + \eta_3 \end{pmatrix}, \quad g_2(z) = \begin{pmatrix} z_1 + \eta_1 \\ z_2 \\ \omega z_3 + 2\eta_3 \end{pmatrix}, \quad g_3(z) = \begin{pmatrix} \omega z_1 + 2\eta_1 \\ z_2 + \eta_2 \\ z_3 \end{pmatrix}.$$

Now write e_i, e'_i for the lattice generators of the i th factor in $E \times E \times E$. In particular, we fix $e_i = 1, e'_i = \omega$ so that the 3-torsion points $\eta_i = (1/3)(2e_i + e'_i)$ and $2\eta_i = (1/3)(e_i + 2e'_i)$ are fixed under rotation by ω . We lift g_i to affine transformations γ_i in $\mathbb{A}(3, \mathbb{C})$, and we define $\Gamma \subset \mathbb{A}(3, \mathbb{C})$ to be the subgroup generated by γ_i, t_i, t'_i for $i = 1, 2, 3$, where t_i, t'_i are affine translations by e_i, e'_i , respectively. Note that since $\gamma_i^3 = t_{3\eta_{i+2}} = t_{i+2}^2 t'_{i+2}$, where i is taken modulo 3, Γ is in fact generated by γ_i and t_i .

THEOREM 4.1. *A Kulikov surface X has $\pi_1(X) = \Gamma$ and $H_1(X, \mathbb{Z}) = (\mathbb{Z}/3)^3$.*

REMARK 4.2. Kulikov [K] proved that $\pi_1(X)$ is infinite and not abelian and that $\text{Tors } X \supset (\mathbb{Z}/3)^3$.

Proof. By the Inoue-type construction of Section 3.2, X has an étale $(\mathbb{Z}/3)^3$ -cover \widehat{X} , which is a smooth hypersurface of tridegree $(3, 3, 3)$ in $E \times E \times E$. Therefore, by Lefschetz’s theorem, $\pi_1(\widehat{X}) = \pi_1(E \times E \times E)$, which is generated by translations t_i and t'_i for $i = 1, 2, 3$.

Now, Γ acts on the universal cover \mathbb{C}^3 of $E \times E \times E$, and it acts freely on the universal cover $\widetilde{X} \subset \mathbb{C}^3$ of \widehat{X} . By construction, $X = \widetilde{X}/\Gamma$, so \widetilde{X} is the universal cover of X and $\pi_1(X) = \Gamma$. Clearly, we have also shown that there is an exact sequence

$$1 \rightarrow \mathbb{Z}^6 \rightarrow \pi_1(X) \rightarrow G^2 \rightarrow 1.$$

The group $H_1(X, \mathbb{Z})$ is the abelianization of Γ , so we calculate its center $[\Gamma, \Gamma]$. Observe that

$$\begin{aligned}
 (17) \quad & \gamma_1 \gamma_2 = t_3 \gamma_2 \gamma_1 = \gamma_2 \gamma_1 (t_3 t'_3)^{-1}, \\
 & \gamma_2 \gamma_3 = t_1 \gamma_3 \gamma_2 = \gamma_3 \gamma_2 (t_1 t'_1)^{-1}, \\
 & \gamma_3 \gamma_1 = t_2 \gamma_1 \gamma_3 = \gamma_1 \gamma_3 (t_2 t'_2)^{-1};
 \end{aligned}$$

hence, all t_i, t'_i are in $[\Gamma, \Gamma]$. Moreover, $\Gamma / \langle t_i, t'_i \rangle$ is abelian, so $[\Gamma, \Gamma] = \langle t_i, t'_i \rangle$ and

$$H_1(X, \mathbb{Z}) = \Gamma^{\text{ab}} \cong (\mathbb{Z}/3)^3,$$

where the generators are the residue classes of γ_i modulo $[\Gamma, \Gamma]$. □

4.2. The moduli space is closed under homotopy equivalence

In this section, we prove that the moduli space of Kulikov surfaces is actually a connected component of the moduli space of surfaces of general type with $K^2 = 6$. In fact, we prove the following stronger result.

THEOREM 4.3. *Any compact surface S which is homotopically equivalent to a Kulikov surface X is itself a Kulikov surface.*

We prove the theorem in several steps, mostly by an explicit analysis of the fundamental group. A similar theorem for Burniat–Inoue surfaces is proved in [BC3]. For the remainder of this section, X is a Kulikov surface and S is a compact surface homotopically equivalent to X .

Now consider the three subgroups of Γ defined by

$$\begin{aligned}
 \Sigma_1 &= \langle t_1, \gamma_1, t_2, \gamma_2, t_3, \gamma_3^3 \rangle, \\
 \Sigma_2 &= \langle t_1, \gamma_1^3, t_2, \gamma_2, t_3, \gamma_3 \rangle, \\
 \Sigma_3 &= \langle t_1, \gamma_1, t_2, \gamma_2^3, t_3, \gamma_3 \rangle.
 \end{aligned}$$

These are of index 3 in Γ , and using the commutation relations (17) as well as

$$(18) \quad \gamma_i t_j = \begin{cases} (t_j \gamma_i) t_j^2 t'_j = t'_j t_j^{-1} (t_j \gamma_i) & \text{if } j = i + 1, \\ t_j \gamma_i & \text{otherwise,} \end{cases}$$

we see that they are normal subgroups. Thus, each Σ_i corresponds to an étale $\mathbb{Z}/3$ -cover $S_i \rightarrow S$.

LEMMA 4.4. *The Albanese variety $\text{Alb}(S_i)$ of each S_i is the Fermat cubic curve. In other words, $\text{Alb}(S_i)$ is the unique elliptic curve which admits complex multiplication by a primitive cube root of unity.*

Proof. We prove the lemma for S_1 , since the other cases are the same by symmetry. Using commutation relations (17) and (18), we have

$$[\Sigma_1, \Sigma_1] \supset \langle t_2^2 t_2', t_2' t_2^{-1}, t_3, t_3' \rangle,$$

and since $\Sigma_1 / \langle t_2^2 t_2', t_2' t_2^{-1}, t_3, t_3' \rangle$ is abelian, the reverse inclusion holds. Hence,

$$H_1(S_1, \mathbb{Z}) = \Sigma_1^{\text{ab}} \cong \langle \bar{t}_1, \bar{\gamma}_2 \rangle \oplus (\mathbb{Z}/3)^2,$$

where $\bar{t}_1, \bar{\gamma}_2$ denote the classes of t_1, γ_2 , respectively, and the torsion summand $(\mathbb{Z}/3)^2$ is generated by the classes of γ_1 and t_2 . This shows that $\text{Alb}(S_1)$ is an elliptic curve.

Now the Γ/Σ_1 -action associated to the cover $S_1 \rightarrow S$ descends to a non-trivial $\mathbb{Z}/3$ -action on $H^0(S_1, \Omega_{S_1})$. Indeed,

$$H^0(S_1, \Omega_{S_1})^{\Gamma/\Sigma_1} = H^0(S, \Omega_S) = H^1(S, \mathcal{O}_S) = 0,$$

by Hodge theory. Hence, $\text{Alb}(S_1)$ admits an automorphism of order 3. This proves that $\text{Alb}(S_1)$ is the Fermat cubic curve. □

Now write $E'_i = \text{Alb}(S_i)$, and let $\Lambda'_i = \langle t_i, \gamma_{i+1} \rangle$ be the lift of the torsion-free part of $H_1(S_i, \mathbb{Z})$ to Γ . Then $\langle t_i, \gamma_{i+1}^3 \rangle = \langle t_i, t'_i \rangle$ is an index 3 normal subgroup of Λ'_i , which we call Λ_i . Viewed as a lattice, Λ_i corresponds to an elliptic curve E_i and a $\mathbb{Z}/3$ -cover $E_i \rightarrow E'_i$. Since E_i and E'_i are isogenous, E_i is also a Fermat cubic curve.

PROPOSITION 4.5. *Let $\widehat{S} \rightarrow S$ be the maximal abelian cover of S . Then*

- (1) *the Albanese variety of \widehat{S} is the product of elliptic curves $E_1 \times E_2 \times E_3$;*
- (2) *the Albanese map $\text{alb}: \widehat{S} \rightarrow E_1 \times E_2 \times E_3$ is a birational morphism onto its image $A_{\widehat{S}}$, and $A_{\widehat{S}}$ is a hypersurface of tridegree $(3, 3, 3)$.*

Proof. Define $\Lambda = \Lambda_1 \oplus \Lambda_2 \oplus \Lambda_3$, and let $\widetilde{\psi}: \widehat{S} \rightarrow S$ be the étale $(\mathbb{Z}/3)^3$ -cover of S with fundamental group $\Lambda \triangleleft \Gamma$. Note that $H_1(S, \mathbb{Z}) = (\mathbb{Z}/3)^3$, so $\widehat{S} \rightarrow S$ is maximal, and thus, $\widetilde{\psi}$ factors through each S_i . Now composing these factorizations with $\text{alb}: S_i \rightarrow E'_i$ and taking the direct product, we obtain a morphism

$$f: \widehat{S} \rightarrow E'_1 \times E'_2 \times E'_3.$$

Moreover, f factors through $E_1 \times E_2 \times E_3$ via the $\mathbb{Z}/3$ -covers $E_i \rightarrow E'_i$. Since $\pi_1(E_1 \times E_2 \times E_3) = \pi_1(\widehat{S})$, this factorization of f is the Albanese map of \widehat{S} . This proves part (1) of the proposition.

To prove part (2), we compare $H^*(\widehat{S}, \mathbb{Z})$ with $H^*(\widehat{X}, \mathbb{Z})$, where \widehat{X} is the maximal $(\mathbb{Z}/3)^3$ -cover of a Kulikov surface X . First let us factor the Albanese map of \widehat{S} through its image $A_{\widehat{S}}$ as follows:

$$(19) \quad \widehat{S} \xrightarrow{a_{\widehat{S}}} A_{\widehat{S}} \xrightarrow{i_{\widehat{S}}} \text{Alb}(\widehat{S}) = E_1 \times E_2 \times E_3.$$

Then recall from Section 3.2 that \widehat{X} is a hypersurface of tridegree $(3, 3, 3)$ in $E \times E \times E$. Now following the procedure described for part (1), we may also factor the Albanese map of \widehat{X} through its image $A_{\widehat{X}}$:

$$(20) \quad \widehat{X} \xrightarrow{a_{\widehat{X}}} A_{\widehat{X}} \xrightarrow{i_{\widehat{X}}} \text{Alb}(\widehat{X}) = E \times E \times E.$$

This time $a_{\widehat{X}}$ is actually an isomorphism.

Now the factorizations (19) and (20) are both constructed using only properties of the fundamental group, and we have a homotopy equivalence between S and X which induces $\pi_1(S) = \pi_1(X) = \Gamma$. Thus, we have compatible identifications $H^4(\widehat{S}, \mathbb{Z}) = H^4(\widehat{X}, \mathbb{Z})$ and $\text{Alb}(\widehat{S}) = \text{Alb}(\widehat{X})$, which we arrange in the following commutative diagram:

$$\begin{array}{ccccc}
 & & i_{\widehat{S}}^* & \rightarrow & H^4(A_{\widehat{S}}, \mathbb{Z}) & \xrightarrow{a_{\widehat{S}}^*} & H^4(\widehat{S}, \mathbb{Z}) \\
 & & \nearrow & & & & \parallel \\
 H^4(T, \mathbb{Z}) & & & & & & \\
 & & i_{\widehat{X}}^* & \rightarrow & H^4(A_{\widehat{X}}, \mathbb{Z}) & \xrightarrow{a_{\widehat{X}}^*} & H^4(\widehat{X}, \mathbb{Z})
 \end{array}$$

Here we have defined $T = \text{Alb}(\widehat{X}) \cong \text{Alb}(\widehat{S})$.

Now $a_{\widehat{X}}^*$ is an isomorphism, and we also note that $i_{\widehat{X}}^*$ cannot be trivial, because $A_{\widehat{X}}$ is a hypersurface of tridegree $(3, 3, 3)$ in T . This implies that $A_{\widehat{S}}$ is codimension 1 in T , because $H^4(A_{\widehat{S}}, \mathbb{Z}) \neq 0$ by commutativity. Let $[A_{\widehat{S}}]$ and $[A_{\widehat{X}}]$ be the fundamental classes of $A_{\widehat{S}}$ and $A_{\widehat{X}}$ in $H^2(T, \mathbb{Z})$, respectively. Then for any $\lambda \in H^4(T, \mathbb{Z})$, we have

$$(\deg a_{\widehat{S}})[A_{\widehat{S}}] \cdot \lambda = \text{alb}^*(\lambda) = a_{\widehat{X}}^*(i_{\widehat{X}}^*(\lambda)) = [A_{\widehat{X}}] \cdot \lambda.$$

By Poincaré duality, there exists $\lambda' \in H^4(T, \mathbb{Z})$ such that $[A_{\widehat{X}}] \cdot \lambda' = 1$, which implies that $\deg a_{\widehat{S}} = 1$. Thus, \widehat{S} and $A_{\widehat{S}}$ are birational, and $a_{\widehat{S}}^*$ is an isomorphism. Therefore, $[A_{\widehat{S}}] \cdot \lambda = [A_{\widehat{X}}] \cdot \lambda$ for all $\lambda \in H^4(T, \mathbb{Z})$ so that $[A_{\widehat{S}}] = [A_{\widehat{X}}]$,

which implies that $A_{\widehat{S}}$ is a hypersurface of tridegree $(3, 3, 3)$ in T . This proves part (2) of the proposition. \square

To complete the proof of the theorem, we prove that the quotient of $A_{\widehat{S}}$ by Γ/Λ is in the same moduli space as S . This follows from the following.

LEMMA 4.6. *The quotient of $A_{\widehat{S}}$ by Γ/Λ is a surface with $p_g = 0$, $K^2 = 6$, and, at worst, rational double points.*

Proof. We compare the invariants of $A_{\widehat{S}}$ and \widehat{S} , and we show that $A_{\widehat{S}}$ has at worst rational double points. Since $\omega_{A_{\widehat{S}}} = \mathcal{O}_{A_{\widehat{S}}}(3, 3, 3)$, we have $K_{A_{\widehat{S}}}^2 = 6 \cdot 3^3$, while $K_{\widehat{S}}^2 = 3^3 \cdot 6$ because $\widehat{S} \rightarrow S$ is an étale $(\mathbb{Z}/3)^3$ -cover. Note that $p_g(\widehat{S}) = 29$ because $\chi(\widehat{S}) = 3^3\chi(S) = 3^3$ and $q(\widehat{S}) = 3$. The short exact sequence

$$0 \rightarrow \mathcal{O}_T \rightarrow \mathcal{O}_T(A_{\widehat{S}}) \rightarrow \omega_{A_{\widehat{S}}} \rightarrow 0$$

gives rise to

$$\begin{aligned} 0 \rightarrow H^0(T, \mathcal{O}_T) &\rightarrow H^0(T, \mathcal{O}_T(A_{\widehat{S}})) \rightarrow H^0(A_{\widehat{S}}, \omega_{A_{\widehat{S}}}) \\ &\rightarrow H^1(T, \mathcal{O}_T) \rightarrow H^1(T, \mathcal{O}_T(A_{\widehat{S}})). \end{aligned}$$

Since $\mathcal{O}_T(A_{\widehat{S}})$ is ample, we see that $|\omega_{A_{\widehat{S}}}|$ is base point free and that $H^1(T, \mathcal{O}_T(A_{\widehat{S}})) = 0$. We also have $h^0(T, \mathcal{O}_T) = 1$, $h^1(T, \mathcal{O}_T) = 3$, and $h^0(T, \mathcal{O}_T(A_{\widehat{S}})) = 3^3 = 27$ by Künneth’s formula. Hence, $p_g(A_{\widehat{S}}) = 29 = p_g(\widehat{S})$, so $\omega_{\widehat{S}} = \text{alb}^* \omega_{A_{\widehat{S}}}$, and $A_{\widehat{S}}$ has at worst rational double points as singularities. Since the induced action of Γ/Λ on $A_{\widehat{S}}$ is free, the quotient has only rational double points, so $p_g = 0$ and $K^2 = 6$. \square

Thus, we have recovered the Inoue-type construction of our surface S using only the fact that S is homotopy equivalent to a Kulikov surface. This proves the theorem.

§5. Degree of the bicanonical map

Let $k_2: X \rightarrow \mathbb{P}^{K_X^2}$ be the bicanonical map of a Kulikov surface X . Then k_2 is a morphism by Reider’s theorem in [Re], and the image is a surface by [X]. In this section we prove that the degree of k_2 is 1.

Let r be the degree of k_2 , and let s be the degree of its image in $\mathbb{P}^{K_X^2}$. Then clearly $4K_X^2 = rs$. Moreover, $r \leq 4$ because $s \geq K_X^2 - 1 \geq 1$ and $r \neq 3, 4$ by [MP2, Theorem 1.1] and [MP1, Theorem 1.1], respectively, so the only possible values for r are 1 or 2. Now to calculate the value of r , we use the following proposition, which is a natural extension of [P, Proposition 4.1].

PROPOSITION 5.1. *Let $\psi: X \rightarrow Y$ be a finite abelian cover with group G and branch divisor Δ , where X and Y are nonsingular varieties of dimension n . Then the direct image of the bicanonical sheaf $\psi_*\omega_X^2$ splits into eigensheaves*

$$(21) \quad \begin{aligned} (\psi_*\omega_X^2)^{\text{inv}} &= \omega_Y^2(\Delta), \\ (\psi_*\omega_X^2)^{(g)} &= \omega_Y^2\left(\sum_{g \in S_\chi} \Delta_g\right) \otimes \mathcal{L}_{\chi^{-1}}, \end{aligned}$$

where $S_\chi = \{g \in G \mid \frac{m}{d}\chi(g) \neq m - 1\}$, m is the order of g , d is the order of χ , and Δ_g is the sum of the components Δ_i of Δ such that $\Phi(\Delta_i) = g$.

Proof. By [P, Lemma 4.1], locally free sheaves on Y which coincide on the complement of a subset of codimension at least 2 subset are in fact equal on Y . Thus, it suffices to prove the statement in a neighborhood of every point which lies on a single irreducible component of Δ . Therefore, we assume that Δ is irreducible and that $\Delta = \Delta_g$ for some $g \in G$.

Let $W = X/\langle g \rangle$. Then ψ factors through W as $\psi = \rho \circ \pi$, where $\pi: X \rightarrow W$ is a cyclic cover branched over $\rho^{-1}(\Delta)$ and $\rho: W \rightarrow Y$ is unramified. Let $\mathcal{M} = \mathcal{O}_W(M)$ be the line bundle on W corresponding to the cyclic cover π , so $\mathcal{M}^m = \mathcal{O}_W(\rho^*\Delta)$ and the formula

$$\pi_*\mathcal{O}_X = \bigoplus_{\mu=0}^{m-1} \mathcal{M}^{-\mu}$$

decomposes \mathcal{O}_X into eigensheaves.

Choose local coordinates b, w_2, \dots, w_n for W , so that b is a local equation for $\rho^*\Delta$, and let z be a local generator of \mathcal{M}^{-1} as an \mathcal{O}_W -module. Then X is defined by the equation $z^m = b$, so that $mz^{m-1}dz = db$, and we have the following local basis for $\pi_*\omega_X^2$ as an \mathcal{O}_W -module:

$$z^{\mu-2m+2}(db \wedge dw_2 \wedge \dots \wedge dw_n)^2, \quad \mu = 0, \dots, m - 1.$$

As in Section 2, we identify $\langle g \rangle^*$ with \mathbb{Z}/m so that the dual character to g is 1. Then g acts on z by $g \cdot z = \varepsilon z$, where $\varepsilon = \exp(2\pi i/m)$, and by considering the action of $\langle g \rangle$ on our basis, $\pi_*\omega_X^2$ splits into eigensheaves:

$$(22) \quad \begin{aligned} (\pi_*\omega_X^2)^{(0)} &= \omega_W^2(\rho^*\Delta), \\ (\pi_*\omega_X^2)^{(1)} &= \omega_W^2 \otimes \mathcal{M}^{m-1}, \\ (\pi_*\omega_X^2)^{(\mu)} &= \omega_W^2(\rho^*\Delta) \otimes \mathcal{M}^{m-\mu} \quad \text{for } 2 \leq \mu \leq m - 1. \end{aligned}$$

Now since ρ is unramified, we have

$$\rho_*\mathcal{M}^i = \bigoplus_{\frac{m}{d}\chi(g)=i} \mathcal{L}_\chi,$$

and clearly, $\psi_*\omega_X^2 = \rho_*\pi_*\omega_X^2$. Combining this with (22), we obtain the required decomposition of $\psi_*\omega_X^2$. \square

In Table 2 we list the eigensheaves of $\psi_*\omega_X^2$ as calculated using Proposition 5.1. For example, when $\chi = (2, 2)$ so that $\chi^{-1} = (1, 1)$, we have $S_{(1,1)} = \left\{ \binom{1}{0}, \binom{0}{1}, \binom{2}{1} \right\}$ and

$$\Delta_{\binom{1}{0}} + \Delta_{\binom{0}{1}} + \Delta_{\binom{2}{1}} = 5H - 2E_1 - E_2 - 3E_3.$$

Now by (3),

$$L_{(1,1)} = \frac{1}{3}(\Delta_{\binom{1}{0}} + 2\Delta_{\binom{1}{1}} + \Delta_{\binom{0}{1}}) = 2H - E_1 - E_2,$$

so by using (21), we have

$$(\psi_*\omega_X^2)^{(2,2)} = H - E_1 - E_3$$

because $K_Y = -3H + \sum E_i$.

Using Table 2, we can calculate the degree of the bicanonical map of a Kulikov surface.

Table 2: Decomposition of $\psi_*\omega_X^2$.

χ	$\sum_{S_{\chi^{-1}}} \Delta_g$	$L_{\chi^{-1}}$	$(\psi_*\omega_X^2)^{(\chi)}$
(0, 0)	Δ	0	0
(1, 0)	$\Delta_{\binom{0}{1}} + \Delta_{\binom{2}{1}}$	$3H - E_1 - 2E_2 - E_3$	$-H + E_1 + E_2$
(1, 2)	$\Delta_{\binom{0}{1}} + \Delta_{\binom{1}{1}}$	$3H - E_1 - E_2 - 2E_3$	$-H + E_2 + E_3$
(1, 1)	$\Delta_{\binom{1}{1}} + \Delta_{\binom{2}{1}}$	$3H - 2E_1 - E_2 - E_3$	$-H + E_1 + E_3$
(0, 1)	$\Delta_{\binom{1}{0}}$	$2H$	$-H$
(0, 2)	$\Delta_{\binom{1}{0}} + \Delta_{\binom{0}{1}} + \Delta_{\binom{1}{1}} + \Delta_{\binom{2}{1}}$	H	H
(2, 2)	$\Delta_{\binom{1}{0}} + \Delta_{\binom{0}{1}} + \Delta_{\binom{2}{1}}$	$2H - E_1 - E_2$	$H - E_1 - E_3$
(2, 0)	$\Delta_{\binom{1}{0}} + \Delta_{\binom{0}{1}} + \Delta_{\binom{1}{1}}$	$2H - E_2 - E_3$	$H - E_1 - E_2$
(2, 1)	$\Delta_{\binom{1}{0}} + \Delta_{\binom{1}{1}} + \Delta_{\binom{2}{1}}$	$2H - E_1 - E_3$	$H - E_2 - E_3$

PROPOSITION 5.2. *The bicanonical morphism k_2 of a Kulikov surface X is birational.*

Proof. It suffices to show that k_2 separates points in X . It is clear from Table 2 that the only summands of $\psi_*\omega_X^2$ with global sections are the eigen-sheaves with characters $(0, 2)$, $(2, 2)$, $(2, 0)$, and $(2, 1)$ together with the invariant eigensheaf. Note that these characters generate $(\mathbb{Z}/3)^{2*}$.

Now choose a basis of $H^0(X, 2K_X)$ comprising eigenfunctions for the decomposition, so that k_2 is defined via this basis. Then for a generic point x in X and any g in $(\mathbb{Z}/3)^2$, if x and gx have the same image under k_2 , we have $(0, 0)(g) = (0, 2)(g) = (2, 2)(g) = (2, 0)(g) = (2, 1)(g)$, which implies that $g = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$. Therefore, k_2 separates points in each fiber of ψ .

Finally, we show that k_2 separates fibers of ψ . Note that $(\psi_*\omega_X^2)^{(0,2)} = \mathcal{O}_Y(H)$, and therefore k_2 separates points in $Y \setminus (E_1 \cup E_2 \cup E_3)$. It follows that k_2 separates fibers of ψ , so k_2 is birational. \square

§6. Bloch conjecture

In this section, we use methods of [IM] to show that the Bloch conjecture is verified for Kulikov surfaces. Let $A_0^0(S)$ denote the group of rational equivalence classes of 0-cycles of degree 0 on a surface S . The conjecture is that for a surface S with $p_g = 0$, $A_0^0(S)$ is canonically isomorphic to $\text{Alb}(S)$. We note that the conjecture is proved for surfaces with Kodaira dimension $\kappa < 2$ in [BKL]. Since the Kulikov surface X is a surface of general type, we must show that $A_0^0(X) = 0$.

We briefly outline the approach. Suppose that G is a finite group of automorphisms of a surface S . Then every element g in G induces an endomorphism $g_*: A_0^0(S) \rightarrow A_0^0(S)$, and this extends linearly to a homomorphism

$$\Gamma: \mathbb{C}G \rightarrow \text{End}(A_0^0(S)).$$

Now for a subgroup H of G , we define

$$z(H) = \sum_{h \in H} h.$$

Then by [IM] we have $A_0^0(S/H) = 0$ if and only if $\Gamma(z(H)) = 0$. This very elegant result was further refined in [Ba], as follows.

LEMMA 6.1. *Let $H, H_i, i = 1, \dots, n$, be subgroups of G , and let I be the ideal of $\mathbb{C}G$ generated by $z(H_i)$. Suppose that $A_0^0(S/H_i) = 0$ for all i . Then if $z(H)$ is in I , we have $A_0^0(S/H) = 0$.*

Now, this lemma is used together with the Inoue construction in [IM] to prove that Burniat–Inoue surfaces verify the Bloch conjecture. We adapt their method to Kulikov surfaces as follows.

THEOREM 6.2. *Kulikov surfaces verify the Bloch conjecture.*

The proof for Kulikov surfaces is slightly more involved than in [IM] because the groups used are larger, so we present the full calculation. First, we establish some notation to streamline the algebraic manipulations which are used in the proof. Recall from Section 3.2 that a Kulikov surface X is a G^2 -quotient of $\widehat{X}_{3,3,3} \subset E \times E \times E$, where $G^2 = \langle g_1, g_2, g_3 \rangle \cong (\mathbb{Z}/3)^3$. Now G^2 is contained in the larger group $G^1 \cong (\mathbb{Z}/3)^5$ of automorphisms of \widehat{X} . Following Lemma 3.2, we choose generators $\xi_1, \xi_2, \xi_3, \omega_1, \omega_2, \omega_3$ for G^1 subject to the relation $\xi_1 \xi_2 \xi_3 = 1$. By Lemma 3.2 and Definition 2.2, $\xi_i = \eta_i^2 \eta_{i+1}$ are composite translations generating G^0 , ω_i are rotations on the i th factor E , and $g_i = \xi_{i+1} \omega_{i+1}$.

LEMMA 6.3. *Let $H \subset G^1$ be one of the following collection of subgroups:*

$$\langle \omega_i, \xi_n^l \omega_j, \xi_n^m \omega_k \rangle \quad \text{for } 0 \leq l, m \leq 2,$$

where $\{i, j, k\} = \{1, 2, 3\}$ and where $n = i$ or $i - 1$. Additionally, we may choose $H = \langle \xi_3 \omega_1, \xi_2 \omega_2, \xi_2 \omega_3 \rangle$. Then the surface \widehat{X}/H is rational.

REMARK 6.4. This is not an exhaustive list of subgroups H which give rise to a rational quotient, but it is sufficient to prove the theorem. The additional subgroup $\langle \xi_3 \omega_1, \xi_2 \omega_2, \xi_2 \omega_3 \rangle$ is necessary for the proof, and this is precisely what makes the calculation harder than the elementary one in [IM].

Proof of Lemma 6.3. Recall the definition of \widehat{X} as a G^1 -cover of the plane branched in the Kulikov line configuration from Section 2.2. In particular, by Definition 2.1, the columns $\delta_1, \delta_2, \delta_3, \omega_1, \omega_2, \omega_3$ of matrix (6) generate G^1 . Then by Definition 2.2, we have

$$\xi_i = \delta_{i-1}^2 \omega_i \omega_{i+1},$$

and since $\xi_1 \xi_2 \xi_3 = 1$, we also have

$$\delta_1 \delta_2 \delta_3 = (\omega_1 \omega_2 \omega_3)^2.$$

Now suppose that H is a subgroup of G^1 . The quotient \widehat{X}/H is a G^1/H -cover of \mathbb{P}^2 branched in the Kulikov line configuration of Figure 1. Thus, we

may use the techniques of abelian covers to prove that \widehat{X}/H is rational. This involves a series of repetitive calculations, and various cases are related to one another by symmetry. As an illustration, we calculate a typical quotient, as follows.

Let $H = \langle \omega_1, \xi_3^2 \omega_2, \xi_3 \omega_3 \rangle$; then $G^1/H \cong (\mathbb{Z}/3)^2$ is generated by the classes $\overline{\delta_1}$ and $\overline{\omega_3}$. We list the class of each generator of G^1 under the quotient:

$$\overline{\delta_2} \equiv \overline{\omega_3^2}, \quad \overline{\delta_3} \equiv \overline{\delta_1^2 \omega_3}, \quad \overline{\omega_1} \equiv 1, \quad \overline{\omega_2} \equiv \overline{\omega_3^2}.$$

Viewing \widehat{X}/H as a $(\mathbb{Z}/3)^2$ -cover of \mathbb{P}^2 branched in the Kulikov line configuration, we must blow up P_2 and P_3 to remove singular points on the cover. Moreover, the point of intersection of Δ_5 and Δ_6 fails the nonsingularity condition of (1). Thus, we blow up this point, introducing another -1 -curve which we call E . Having done this, we note that by (2), the induced cover is unramified over E_2 and E . Finally, we contract the strict transforms of Δ_1 and Δ_5 . We are left with a nonsingular $(\mathbb{Z}/3)^2$ -cover of $\mathbb{P}^1 \times \mathbb{P}^1$ branched in four distinct lines, two in each ruling. This is a rational surface. \square

Proof of Theorem 6.2. Let I be the ideal generated by $z(H)$ for all H listed in Lemma 6.3. Then the quotients \widehat{X}/H are rational surfaces, so by [BKL, p. 135], $A_0^0(\widehat{X}/H) = 0$. Thus, by Lemma 6.1, it is sufficient to prove that $z(G^2)$ is an element of I . Now, the verification of the Bloch conjecture for Burniat–Inoue surfaces (see [IM]) is a series of elementary polynomial manipulations. Unfortunately, the corresponding manipulations do not suffice for the Kulikov surface.

Instead, we prove that $z(G^2)$ is in I using the MAGMA computer algebra script (see [BCP]) below:

```

Q:=Rationals();
RR<w1,w2,w3,xi1,xi2,xi3>:=PolynomialRing(Q,6);

function z(a)
  return (1+a[1]+a[1]^2)*(1+a[2]+a[2]^2)*(1+a[3]+a[3]^2);
end function;

ListH:=[[w1,xi1^i*w2,xi1^j*w3]:i in [0..2],j in [0..2]] cat
        [[w1,xi3^i*w2,xi3^j*w3]:i in [0..2],j in [0..2]] cat
        [[w2,xi1^i*w3,xi1^j*w1]:i in [0..2],j in [0..2]] cat
        [[w2,xi2^i*w3,xi2^j*w1]:i in [0..2],j in [0..2]] cat
    
```

```

[[w3,xi2^i*w1,xi2^j*w2]:i in [0..2],j in [0..2]] cat
[[w3,xi3^i*w1,xi3^j*w2]:i in [0..2],j in [0..2]];

Append(~ListH,[xi3*w1,xi2*w2,xi2*w3]); // extra generator

I:=ideal<RR|[xi1^3-1,xi2^3-1,xi3^3-1,
             w1^3-1,w2^3-1,w3^3-1,xi1*xi2*xi3-1]
      cat [z(H):H in ListH]>;

zG2:=z([w1*xi1,w2*xi2,w3*xi3]);

zG2 in I; // Result: true

```

□

Acknowledgments. Both authors thank Ingrid Bauer and Fabrizio Catanese for their help and encouragement with this paper. We also thank the referees for their comments, which improved our manuscript in various ways.

REFERENCES

- [AP] V. Alexeev and R. Pardini, *Explicit compactifications of moduli spaces of Campedelli and Burniat surfaces*, preprint, arXiv:0901.4431 [math.AG]
- [Ba] R. Barlow, *Rational equivalence of zero cycles for some more surfaces with $p_g = 0$* , Invent. Math. **79** (1985), 303–308. MR 0778129. DOI 10.1007/BF01388975.
- [BC1] I. C. Bauer and F. Catanese, *A volume maximizing canonical surface in 3-space*, Comment. Math. Helv. **83** (2008), 387–406. MR 2390050. DOI 10.4171/CMH/129.
- [BC2] ———, *Burniat surfaces, II: Secondary Burniat surfaces form three connected components of the moduli space*, Invent. Math. **180** (2010), 559–588. MR 2609250. DOI 10.1007/s00222-010-0237-z.
- [BC3] ———, “Burniat surfaces, I: Fundamental groups and moduli of primary Burniat surfaces” in *Classification of Algebraic Varieties*, EMS Ser. Congr. Rep., Eur. Math. Soc., Zürich, 2011, 49–76. MR 2779467. DOI 10.4171/007-1/3.
- [BC4] ———, *Burniat surfaces, III: Deformations of automorphisms and extended Burniat surfaces*, preprint, arXiv:1011.3770v1 [math.AG]
- [BKL] S. Bloch, A. Kas, and D. Lieberman, *Zero cycles on surfaces with $p_g = 0$* , Compos. Math. **33** (1976), 135–145. MR 0435073.
- [BCP] W. Bosma, J. Cannon, and C. Playoust, “The Magma algebra system, I: The user language” in *Computational Algebra and Number Theory (London, 1993)*, J. Symbolic Comput. **24** (1997), 235–265. MR 1484478. DOI 10.1006/jsc.1996.0125.
- [Bu] P. Burniat, *Sur les surfaces de genre $P_{12} > 1$* , Ann. Mat. Pura Appl. (4) **71** (1966), 1–24. MR 0206810.
- [C] F. Catanese, *On the moduli spaces of surfaces of general type*, J. Differential Geom. **19** (1984), 483–515. MR 0755236.

- [IM] H. Inose and M. Mizukami, *Rational equivalence of 0-cycles on some surfaces of general type with $p_g = 0$* , Math. Ann. **244** (1979), 205–217. MR 0553252. DOI 10.1007/BF01420343.
- [I] M. Inoue, *Some new surfaces of general type*, Tokyo J. Math. **17** (1994), 295–319. MR 1305801. DOI 10.3836/tjm/1270127954.
- [KSB] J. Kollár and N. I. Shepherd-Barron, *Threefolds and deformations of surface singularities*, Invent. Math. **91** (1988), 299–338. MR 0922803. DOI 10.1007/BF01389370.
- [K] V. S. Kulikov, *Old examples and a new example of surfaces of general type with $p_g = 0$* (in Russian), Izv. Ross. Akad. Nauk Ser. Mat. **68** (2004), no. 5, 123–170; English translation in Izv. Math. **68** (2004), no. 5, 965–1008. MR 2104852. DOI 10.1070/IM2004v068n05ABEH000505.
- [MP1] M. Mendes Lopes and R. Pardini, *A connected component of the moduli space of surfaces with $p_g = 0$* , Topology **40** (2001), 977–991. MR 1860538. DOI 10.1016/S0040-9383(00)00004-5.
- [MP2] ———, *Surfaces of general type with $p_g = 0$, $K^2 = 6$ and non birational bicanonical map*, Math. Ann. **329** (2004), 535–552. MR 2127989. DOI 10.1007/s00208-004-0524-3.
- [P] R. Pardini, *Abelian covers of algebraic varieties*, J. Reine Angew. Math. **417** (1991), 191–213. MR 1103912. DOI 10.1515/crll.1991.417.191.
- [Re] I. Reider, *Vector bundles of rank 2 and linear systems on algebraic surfaces*, Ann. of Math. (2) **127** (1988), 309–316. MR 0932299. DOI 10.2307/2007055.
- [X] G. Xiao, *Degree of the bicanonical map of a surface of general type*, Amer. J. Math. **112** (1990), 713–736. MR 1073006. DOI 10.2307/2374804.

Tsz On Mario Chan
Lehrstuhl Mathematik VIII
Mathematisches Institut der Universität Bayreuth
95447 Bayreuth
Germany
mariochan@graduate.hku.hk

Stephen Coughlan
Department of Mathematics and Statistics
University of Massachusetts
Amherst, Massachusetts 01003
USA
coughlan@math.umass.edu