

doi:10.1017/S1474748021000608 © The Author(s), 2022. Published by Cambridge University Press. This is an Open Access article, distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike licence (<https://creativecommons.org/licenses/by-nc-sa/4.0>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the same Creative Commons licence is used to distribute the reused or adapted article and the original article is properly cited. The written permission of Cambridge University Press must be obtained prior to any commercial use.

THE p -ADIC GROSS–ZAGIER FORMULA ON SHIMURA CURVES, II: NONSPLIT PRIMES

DANIEL DISEGNI 

Department of Mathematics, Ben-Gurion University of the Negev, Be'er Sheva 84105, Israel

(disegni@bgu.ac.il)

(Received 4 October 2020; revised 16 November 2021; accepted 16 November 2021; first published online 15 February 2022)

Abstract The formula of the title relates p -adic heights of Heegner points and derivatives of p -adic L -functions. It was originally proved by Perrin-Riou for p -ordinary elliptic curves over the rationals, under the assumption that p splits in the relevant quadratic extension. We remove this assumption, in the more general setting of Hilbert-modular abelian varieties.

Keywords and phrases: Heegner points; p -adic heights; p -adic L -functions

2020 Mathematics Subject Classification: Primary 11G40
Secondary 11F33

Contents

1	Introduction and statement of the main result	2200
1.1.	The main ideas in a classical context	2200
1.2.	Statement	2203
1.3.	Organization of the paper	2208
2	Comparison with [9]	2208
2.1.	The p -adic L -function	2209
2.2.	Equivalence of statements	2210
3	Structure of the proof	2211
3.1.	Notation and setup	2211
3.2.	Analytic kernel	2214
3.3.	Geometric kernel	2215
3.4.	Kernel identity	2217
3.5.	Derivative of the analytic kernel	2218
3.6.	Decomposition of the geometric kernel and comparison	2220
4	Local heights at p	2222



4.1. Norm relation for the generating series	2222
4.2. Intersection multiplicities on arithmetic surfaces	2225
4.3. Decay of intersection multiplicities	2226
4.4. Decay of local heights	2231
Appendix A Local integrals	2234
A.1. Interpolation factors	2234
A.2. Toric period at p	2235
Appendix B Errata to [9]	2237
References	2239

1. Introduction and statement of the main result

The p -adic Gross–Zagier formula of Perrin-Riou relates p -adic heights of Heegner points and derivatives of p -adic L -functions. In its original form [26], it concerns (modular) elliptic curves over \mathbf{Q} , and it is proved under two main assumptions: first, that the elliptic curve is p -ordinary; second, that p splits in the field E of complex multiplications of the Heegner points. The formula has applications to both the p -adic and the classical Birch and Swinnerton-Dyer conjecture.

The first assumption was removed by Kobayashi [19] (see also [4]). The purpose of this work is to remove the second assumption.

We work in the context of our previous work [9], which will enable us in [13] to deduce, from the formula presented here, the analogous one for higher-weight (Hilbert-) modular motives, as well as a version in the universal ordinary family with some new applications. Nevertheless, the new idea we introduce is essentially orthogonal to previous innovations, including those of [9] (and in fact it can be applied, at least in principle, to the nonordinary case as well). For this reason we start in §1.1 by informally discussing it in the simplest classical case of elliptic curves over \mathbf{Q} . The general form of our results is presented in §1.2.

1.1. The main ideas in a classical context

Classically, Heegner points on the elliptic curve A/\mathbf{Q} are images of CM points (or divisors) on a modular curve X , under a parametrization $f: X \rightarrow A$. More precisely, choosing an imaginary quadratic field E , for each ring class character $\chi: \text{Gal}(\overline{E}/E) \rightarrow \overline{\mathbf{Q}}^\times$ one can construct a point $P(f, \chi) \in A_E(\chi)$, the χ -isotypic part of $A(\overline{E})_{\overline{\mathbf{Q}}}$. The landmark formula of Gross and Zagier [17] relates the height of $P(f, \chi)$ to the derivative $L'(A_E \otimes \chi, 1)$ of the L -function of a twisted base change of A . The analogous formula in p -adic coefficients,¹

$$\langle P(f, \chi), P(f, \chi^{-1}) \rangle \doteq \frac{d}{ds}|_{s=0} L_p(A_E, \chi \cdot \chi_{\text{cyc}}^s), \quad (1.1.1)$$

relates cyclotomic derivatives of p -adic L -functions to p -adic height pairings $\langle \cdot, \cdot \rangle$. We outline its proof for p -ordinary elliptic curves.

¹We denote by \doteq equality up to a less-important nonzero factor.

Review of Perrin-Riou’s proof. Our basic strategy is still Perrin-Riou’s variant of the one of Gross and Zagier; we briefly and informally review it, ignoring, for simplicity of exposition, the role of the character χ . Throughout the following discussion, we include pointers to corresponding statements in the main body of the paper, as guideposts meant to assist the reader’s navigation through the more technical framework used there.

Denoting by φ the ordinary eigenform attached to A , each side of formula (1.1.1) is expressed as the image under a functional ‘ p -adic Petersson product with φ ’, denoted by $\ell_\varphi = \text{formula}$ (3.1.6), of a certain kernel function (a p -adic modular form).

For the left-hand side of formula (1.1.1), the form in question is the generating series (compare formula (3.3.4))

$$Z = \sum_{m \geq 1} \langle P^0, T_m P^0 \rangle_X \mathbf{q}^m = \sum_v Z_v, \tag{1.1.2}$$

where $P^0 \in \text{Div}^0(X)$ is a degree 0 modification of the CM point $P \in X$, $\langle \cdot, \cdot \rangle_X$ is a p -adic height pairing on X compatible with the one on A , and the decomposition into a sum running over all the finite places of \mathbf{Q} (compare equation (3.6.1)) follows from a general decomposition of the global height pairing into a sum of local ones. More precisely, global height pairings are valued in the completed tensor product $H^\times \backslash H_{\mathbf{A}^\infty}^\times \hat{\otimes} L$ of the finite idèles of the Hilbert class field H of E , and of a suitable finite extension L of \mathbf{Q}_p . The series Z_v collects the local pairings at $w \mid v$, each valued in $H_w^\times \hat{\otimes} L$.

The analytic kernel \mathcal{S}' giving the right-hand side of formula (1.1.1) is the derivative of a p -adic family of mixed theta-Eisenstein series (compare definition (3.2.5)). It also enjoys a decomposition

$$\mathcal{S}' = \sum_{v \neq p} \mathcal{S}'_v,$$

where, unlike in equation (1.1.2), the sum runs over the finite places of \mathbf{Q} *different from p* (compare equation (3.5.2)). Once we have established that $Z_v \doteq \mathcal{S}'_v$ for $v \neq p$ by computations similar to those of Gross and Zagier (compare Theorem 3.6.1), it remains to show that the p -adic modular form Z_p is annihilated by ℓ_φ (compare Proposition 3.6.2).

In order to achieve this, one aims at showing that after acting on Z_p by a Hecke operator to replace P^0 by $P^{[\varphi]}$ (a lift of the component of its image in the φ -part of $\text{Jac}(X)$), the resulting form $Z_p^{[\varphi]}$ is p -critical (compare Proposition 3.6.3). That is, its coefficients

$$a_{mp^s} := \left\langle P^{[\varphi]}, T_{mp^s} P^0 \right\rangle_{X,p}$$

decay p -adically no more slowly than a constant multiple of p^s . The p -shift of Fourier coefficients extends the action on modular forms of the operator U_p – which, in contrast, acts by a p -adic unit on the ordinary form φ : this implies that p -critical forms are annihilated by ℓ_φ .

To study the terms a_{mp^s} , one constructs a sequence of points $P_s \in X_H$ whose fields of definitions are the layers H_s of the anticyclotomic p^∞ -extension of E . The relations they satisfy allow us to express

$$a_{mp^s} = \left\langle P^{[\varphi]}, D_{m,s} \right\rangle_{X,p},$$

where $D_{m,s}$ is a degree 0 divisor supported at Hecke-translates of P_s , which are all essentially CM points of conductor p^s defined over H_s (compare Proposition 4.1.4). By a projection formula for $X_{H_s} \rightarrow X_H$, the height a_{mp^s} is then a sum, over primes w of H above p , of the images

$$N_{s,w}(h_{m,s,w})$$

of heights $h_{m,s,w}$ computed on $X_{H_{s,w}}$, under the norm map $N_{s,w}: H_{s,w}^\times \hat{\otimes} L \rightarrow H_w^\times \hat{\otimes} L$. Moreover, it can be shown that the L -denominators of $h_{m,s,w} \in H_{s,w}^\times \hat{\otimes} L$ are uniformly bounded (compare Proposition 4.4.1), so that here we may ignore them and think of $h_{m,s,w} \in H_{s,w}^\times \hat{\otimes} \mathcal{O}_L$.

A simple observation from [9] is that the valuation $w(h_{m,s,w})$ equals

$$m_{X_{H_w}}(P^{[\varphi]}, D_{m,s}), \tag{1.1.3}$$

the intersection multiplicity of the *flat extensions* (§4.2) of those divisors to some regular integral model \mathcal{X} of X_{H_w} . In the split case, it is almost immediate to see that this intersection multiplicity vanishes. This implies that

$$N_{s,w}(h_{m,s}) \in N_{s,w}(\mathcal{O}_{H_{s,w}}^\times) \hat{\otimes} \mathcal{O}_L \subset H_w^\times \hat{\otimes} \mathcal{O}_L. \tag{1.1.4}$$

Since the extension $H_{s,w}/H_w$ is totally ramified of degree p^s , the subset in formula (1.1.4) is $p^s(\mathcal{O}_{H_w}^\times \hat{\otimes} \mathcal{O}_L) \subset p^s(H_w^\times \hat{\otimes} \mathcal{O}_L)$, as desired.

The nonsplit case. In the nonsplit case, the p -adic intersection multiplicity has no reason to vanish. However, the foregoing argument will still go through if we more modestly show that expression (1.1.3) itself decays at least like a multiple of p^s (compare Lemma 4.4.3). The idea to prove this is very simple: we show that if s is large, then for the purposes of computing intersection multiplicities with other divisors \mathcal{D} on \mathcal{X} , the Zariski closure of a CM point of conductor at least p^s can *almost* be approximated by some irreducible component V of the special fiber of \mathcal{X} ; hence the multiplicity will be 0 if \mathcal{D} arises as a flat extension of its generic fiber. The qualifier ‘almost’ means that the foregoing holds *except* if $|\mathcal{D}|$ contains V itself, which will be responsible for a multiplicity error term equal to a constant multiple of p^s .

The approximation result (Proposition 4.3.3) is precisely formulated in an (ultra)metric space of irreducible divisors on the local ring of a regular arithmetic surface, which we introduce following a recent work of García Barroso, González Pérez, and Popescu-Pampu [15]. The proof of the result is also rather simple (albeit not effective), relying on Gross’s theory of quasicanonical liftings [18]. The problem of effectively identifying the approximating divisor V is treated in [11].

Subtleties. This description ignores several difficulties of a relatively more technical nature, most of which we deal with by the representation-theoretic approach of [9] (in turn adapted from Yuan, Zhang, and Zhang [29]). Namely, we allow for arbitrary modular parametrizations f , resulting in an extra parameter ϕ in the kernels Z and \mathcal{S}' . By representation-theoretic results, one is free to some extent to *choose* the parameter ϕ to work with without losing generality. A fine choice (or rather a pair of choices) for its p -adic component is dictated by the goal of interpolation, while imposing suitable conditions on its other components allows us to circumvent many obstacles in the proof.

1.2. Statement

We now describe our result in the general context in which we prove it – which is the same as that of [9] (and [29]), to which we refer for a less terse discussion of the background. (At some points, we find some slightly different formulations or normalizations from those of [9], to be more natural: see §2.2 for the equivalence.)

Abelian varieties parametrized by Shimura curves. Let F be a totally real field and let A/F be a simple abelian variety of GL_2 -type. Assume that $L(A, s)$ is modular (this is known in many cases if A is an elliptic curve). Let \mathbf{B} be a quaternion algebra over the adèles $\mathbf{A} = \mathbf{A}_F$ of F , whose ramification set $\Sigma_{\mathbf{B}}$ has odd cardinality and contains all the infinite places. To \mathbf{B} is attached a tower of Shimura curves $(X_{U/F})_{U \subset \mathbf{B}^{\infty \times}}$, with respective Albanese varieties J_U . It carries a canonical system of divisor classes $\xi_U \in Cl(X_U)_{\mathbf{Q}}$ of degree 1, providing a system $\iota_{\xi, U} \in \text{Hom}_F(X_U, J_U)_{\mathbf{Q}}$ defined by $P \mapsto P - \text{deg}(P)\xi_U$.

The space

$$\pi = \pi_{A, \mathbf{B}} = \varinjlim_U \text{Hom}^0(J_U, A)$$

is either zero or a smooth irreducible representation of \mathbf{B}^{\times} (trivial at the infinite places), with coefficients in the number field $M := \text{End}^0(A)$. We assume we are in the case $\pi = \pi_{A, \mathbf{B}} \neq 0$, which under the modularity assumption and condition (1.2.2) can be arranged by suitably choosing \mathbf{B} . Then for all places $v \nmid \infty$, we have $L_v(A, s) = L_v(s - 1/2, \pi)$ in $M \otimes \mathbf{C}$. We denote by $\omega: F^{\times} \backslash \mathbf{A}^{\times} \rightarrow M^{\times}$ the central character of π . We have a canonical isomorphism $\pi_{A^{\vee}, \mathbf{B}} \cong \pi_{A, \mathbf{B}}^{\vee}$ [29, §1.2.2], and we denote by $(\ , \)_{\pi}: \pi_{A, \mathbf{B}} \otimes \pi_{A^{\vee}, \mathbf{B}} \rightarrow M$ the duality pairing.

Heegner points. Let E/F be a CM quadratic extension with associated quadratic character η , and assume that E admits an \mathbf{A} -embedding $E_{\mathbf{A}} \hookrightarrow \mathbf{B}$, which we fix. Then E^{\times} acts on the right on $X = \varprojlim_U X_U$. The fixed-points subscheme $X^{E^{\times}} \subset X$ is F -isomorphic to $\text{Spec } E^{\text{ab}}$, and we fix a point $P \in X^{E^{\times}}(E^{\text{ab}})$. Let

$$\chi: E^{\times} \backslash E_{\mathbf{A}^{\infty}}^{\times} \cong \text{Gal}(E^{\text{ab}}/E) \rightarrow L(\chi)^{\times}$$

be a character valued in a field extension of $L(\chi) \supset M$, satisfying

$$\omega \cdot \chi|_{\mathbf{A}^{\infty \times}} = \mathbf{1},$$

and define

$$A_E(\chi) := (A(E^{\text{ab}}) \otimes_M L(\chi)_\chi)^{\text{Gal}(E^{\text{ab}}/E)},$$

where $L(\chi)_\chi$ is an $L(\chi)$ -line with Galois action by χ .

Then we have a Heegner point functional

$$f \mapsto P(f, \chi) := \int_{\text{Gal}(E^{\text{ab}}/E)} f(\iota_\xi(P)^\tau) \otimes \chi(\tau) d\tau \in A_E(\chi) \tag{1.2.1}$$

(integration for the Haar measure of volume 1) in the space of invariant linear functionals

$$\mathbf{H}(\pi_{A, \mathbf{B}}, \chi) \otimes_{L(\chi)} A_E(\chi), \quad \mathbf{H}(\pi, \chi) := \text{Hom}_{E_{\mathbf{A}}^\times}(\pi \otimes \chi, L(\chi)),$$

where $E_{\mathbf{A}}^\times$ acts diagonally. There is a product decomposition $\mathbf{H}(\pi, \chi) = \bigotimes_v \mathbf{H}(\pi_v, \chi_v)$, where similarly $\mathbf{H}(\pi_v, \chi_v) := \text{Hom}_{E_v^\times}(\pi_v \otimes \chi_v, L(\chi))$.

A local unit of measure for invariant functionals. By foundational local results of Waldspurger, Tunnell, and Saito, the dimension of $\mathbf{H}(\pi, \chi)$ (for any representation π of \mathbf{B}^\times) is either 0 or 1. If A is modular and the global root number

$$\varepsilon(A_E \otimes \chi) = -1, \tag{1.2.2}$$

then the set of local root numbers determines a unique quaternion algebra \mathbf{B} over \mathbf{A} , satisfying the conditions already required and containing $E_{\mathbf{A}}$, such that $\pi_{A, \mathbf{B}} \neq 0$ and $\dim_{L(\chi)} \mathbf{H}(\pi_{A, \mathbf{B}}, \chi) = 1$.² We place ourselves in this case; then there is a canonical factorizable generator

$$Q_{(\cdot), dt} = \prod_v Q_{(\cdot)_v, dt_v} \in \mathbf{H}(\pi, \chi) \otimes \mathbf{H}(\pi^\vee, \chi^{-1})$$

depending on the choice of a pairing $(\cdot) = \prod_v (\cdot)_v : \pi \otimes \pi^\vee \rightarrow L(\chi)$ and a measure $dt = \prod_v dt_v$ on $E_{\mathbf{A}}^\times / \mathbf{A}^\times$. It is defined locally as follows. Let us use symbols $V_{(A, \chi)}$ and $V_{(A, \chi), v}$, which we informally think of as denoting (up to abelian factors) the virtual motive over F with coefficients in $L(\chi)$

$$V_{(A, \chi)} = \text{Res}_{E/F}(h_1(A_E) \otimes \chi) \ominus \text{ad}(h_1(A)(1))$$

and its local components (the associated local Galois representation or, if v is Archimedean, Hodge structure). Then for each place v of F and any auxiliary $\iota : L(\chi) \hookrightarrow \mathbf{C}$,³ we define

²If $\varepsilon(A_E \otimes \chi) = +1$, there is no such quaternion algebra, and all Heegner points automatically vanish.

³Explicitly, if v is Archimedean, we have $\mathcal{L}(V_{(A, \chi), v}, 0) = 2$ and $Q_{(\cdot)_v, dt_v}(f_{1, v}, f_{2, v}) = 2^{-1} \text{vol}(\mathbf{C}^\times / \mathbf{R}^\times, dt_v)(f_{1, v}, f_{2, v})$.

$$\mathcal{L}(\iota V_{(A,\chi),v},s) := \frac{\zeta_{F,v}(2)L(1/2+s,\iota\pi_{E,v}\otimes\iota\chi_v)}{L(1,\eta_v)L(1,\iota\pi_v,\text{ad})} \cdot \begin{cases} 1 & \text{if } v \text{ is finite,} \\ \pi^{-1} & \text{if } v \mid \infty \end{cases} \in \iota L(\chi), \tag{1.2.3}$$

$$Q_{(\cdot),dt_v}(f_{1,v},f_{2,v},\chi_v) := \iota^{-1}\mathcal{L}(\iota V_{(A,\chi),v},0)^{-1} \int_{E_v^\times/F_v^\times} \chi(t_v)(\pi_v(t_v)f_{1,v},f_{2,v})_v dt_v. \tag{1.2.4}$$

We make the situation more canonical by choosing $dt = \prod_v dt_v$ to satisfy

$$\text{vol}(E^\times \backslash E_{\mathbf{A}}^\times / \mathbf{A}^\times, dt) = 1$$

and by defining, for any $f_3 \in \pi, f_4 \in \pi^\vee$ such that $(f_3, f_4) \neq 0$,

$$Q\left(\begin{matrix} f_1 \otimes f_2 \\ f_3 \otimes f_4 \end{matrix}; \chi\right) := \frac{Q_{(\cdot),dt}(f_1, f_2, \chi)}{(f_3, f_4)}. \tag{1.2.5}$$

p -adic heights. Let us fix a prime \mathfrak{p} of M and denote by p the underlying rational prime. Suppose from now on that for each $v \mid p$, A_{F_v} has \mathfrak{p} -ordinary (potentially good or semistable) reduction. That is, for a sufficiently large finite extension $L \supset M_{\mathfrak{p}}$, the rational \mathfrak{p} -Tate module $W_v := V_p A \otimes_M L$ is a reducible 2-dimensional representation of $\text{Gal}(\overline{F}_v/F_v)$:

$$0 \rightarrow W_v^+ \rightarrow W_v \rightarrow W_v^- \rightarrow 0. \tag{1.2.6}$$

Fix such a coefficient field L , and for each $v \mid p$ let $\alpha_v : F_v^\times \cong \text{Gal}(F_v^{\text{ab}}/F_v) \rightarrow L^\times$ be the character giving the action on the twist $W_v^+(-1)$. The field $L(\chi)$ will from now on be assumed to be an extension of L . Under those conditions, there is a canonical p -adic height pairing

$$\langle \cdot, \cdot \rangle : A_E(\chi) \otimes A_E^\vee(\chi^{-1}) \rightarrow \Gamma_F \hat{\otimes} L(\chi),$$

where $\Gamma_F := \mathbf{A}^\times / \overline{F^\times \hat{\mathcal{O}}_F^{p,\times}}$ (the bar denotes Zariski closure). It is normalized ‘over F ’ as in [9, §4.1].

For $f_1, f_3 \in \pi, f_2, f_4 \in \pi^\vee$, and $P^\vee : \pi^\vee \otimes \chi \rightarrow A_E^\vee(\chi^{-1})$ the Heegner point functional of the dual, our result will measure the ratio $\langle P(f_1, \chi), P^\vee(f_2, \chi^{-1}) \rangle / (f_3, f_4)_\pi$ against the value at the f_i of the ‘unit’ Q . The size will be given by the derivative of the p -adic L -function that we now define.

The p -adic L -function. We continue to assume that A is \mathfrak{p} -ordinary, and review the definition of the p -adic L -function from [9, Theorem A] (in an equivalent form). We start by defining the space on which it lives. Write $\Gamma_F = \varprojlim_n \Gamma_{F,n}$ as the limit of an inverse system of finite groups, and define

$$\mathcal{Y}_F^{1.c.} := \bigcup_n \text{Spec} L[\Gamma_{F,n}] \subset \mathcal{Y}_F := \text{Spec} \mathcal{O}_L[\Gamma_F] \otimes_{\mathcal{O}_L} L. \tag{1.2.7}$$

Then \mathcal{Y}_F is a space of continuous characters on Γ_F , and the 0-dimensional ind-scheme $\mathcal{Y}_F^{1.c.}$ is its subspace of locally constant (finite-order) characters.

For a character $\chi' : \text{Gal}(E^{\text{ab}}/E) \rightarrow L^\times$ together with an embedding $\iota : L' \rightarrow \mathbf{C}$, we shall interpolate the ratio of complete L -functions

$$\mathcal{L}(\iota V_{(A, \chi')}, s) := \prod_v \mathcal{L}(\iota V_{(A, \chi'), v}, s), \quad \mathcal{L}(\iota V_{(A, \chi'), v}, s) = \text{formula (1.2.3)}, \quad \Re(s) \gg 0,$$

where the product runs over all places of F .

We now define the p -interpolation factors for the p -adic L -function. First, recall that the (inverse) Deligne–Langlands gamma factor of a Weil–Deligne representation W' of $\text{Gal}(\overline{F}_v/F_v)$ over a p -adic field L' , with respect to a nontrivial character $\psi_v : F_v \rightarrow \mathbf{C}^\times$ and an embedding $\iota : L' \hookrightarrow \mathbf{C}$, is defined as⁴

$$\gamma(W', \psi_v)^{-1} := \frac{L(W')}{\varepsilon(W', \psi_v)L(W^*(1))}.$$

Let $\psi = \prod_v \psi_v : F \setminus \mathbf{A} \rightarrow \mathbf{C}^\times$ be the standard additive character such that $\psi_\infty(\cdot) = e^{2\pi i \text{Tr}_{F/\mathbf{R}}(\cdot)}$; let $\psi_E = \prod_w \psi_{E, w} = \psi \circ \text{Tr}_{\mathbf{A}_E/\mathbf{A}}$. For a place $v \mid p$ of F , let d_v be a generator of the different ideal of F_v . For a character $\chi' : \text{Gal}(\overline{E}/E) \rightarrow \mathbf{C}^\times$, we define

$$e_v(V_{(A, \chi')}) = |d_v|^{-1/2} \frac{\prod_{w \mid v} \gamma(\iota \text{WD}(W_{v|\mathcal{G}_{E, w}}^+ \otimes \chi'_w), \psi_{E, w})^{-1}}{\gamma(\iota \text{WD}(\text{ad}(W_v)(1))^{++}, \psi_v)^{-1}} \cdot \mathcal{L}(V_{(A, \chi'), v})^{-1}, \quad (1.2.8)$$

where $\text{ad}(W_v)(1)^{++} := \text{Hom}(W_v^-, W_v^+)(1) = \omega_v^{-1} \alpha_v^2 \mid \mid_v^2$ and ιWD is the functor from potentially semistable Galois representations to complex Weil–Deligne representations of [14].

Theorem A. *There is a function*

$$\mathcal{L}_p(V_{(A, \chi)}) \in \mathcal{O}(\mathcal{Y}_F)$$

characterized by the following property. For each complex geometric point $s = \chi_F \in \mathcal{Y}_F^{1, \text{c}}(\mathbf{C})$, with underlying embedding $\iota : L(\chi_F) \hookrightarrow \mathbf{C}$,

$$\mathcal{L}_p(V_{(A, \chi)}, s) = \iota e_p(V_{(A, \chi')}) \cdot \mathcal{L}(\iota V_{(A, \chi')}, 0), \quad \chi' := \chi \cdot \chi_{F|\text{Gal}(\overline{E}/E)},$$

where $e_p(V_{(A, \chi')}) := \prod_{v \mid p} e_v(V_{(A, \chi')})$.

The factor $e_p(V_{(A, \chi')})$ coincides with the one predicted by Coates and Perrin-Riou [7] for $V_{(A, \chi')}$ (their conjecture motivates the denominator terms in equation (1.2.8), which are constants), up to the removal of a trivial zero from their interpolation factor for $\text{ad}(W_v)(1)$.

The p -adic Gross–Zagier formula. We are almost ready to state our main result. Denote by $0 \in \mathcal{Y}_F$ the point corresponding to $\chi_F = \mathbf{1}$, and define

$$\mathcal{L}'_p(V_{(A, \chi)}, 0) := d\mathcal{L}_p(V_{(A, \chi)}, 0) \in T_0\mathcal{Y}_F \cong \Gamma_F \hat{\otimes} L(\chi).$$

⁴The terms L and ε are normalized as in [28].

We say that χ_p is *sufficiently ramified* if it is nontrivial on a certain open subgroup of $\mathcal{O}_{E,p}^\times$ depending only on ω_p (see Assumption 3.4.1 for the precise definition and a comment).

Theorem B. *Suppose that the abelian variety A/F is modular and that for all $v \mid p$, the $\text{Gal}(\overline{F}_v/F_v)$ -representation $V_p A$ is ordinary and potentially crystalline. Let $\chi: \text{Gal}(E^{\text{ab}}/E) \rightarrow L(\chi)^\times$ be a finite-order character satisfying*

$$\varepsilon(A_E \otimes \chi) = -1,$$

and suppose that χ_p sufficiently ramified.

Then for any $f_1, f_3 \in \pi$, $f_2, f_4 \in \pi^\vee$ such that $(f_3, f_4)_\pi \neq 0$, we have

$$\frac{\langle P(f_1, \chi), P^\vee(f_2, \chi^{-1}) \rangle}{(f_3, f_4)_\pi} = e_p(V_{(A, \chi)})^{-1} \cdot \mathcal{L}'_p(V_{(\mathbf{A}, \chi)}, 0) \cdot Q\left(\frac{f_1 \otimes f_2}{f_3 \otimes f_4}; \chi\right)$$

in $\Gamma_F \hat{\otimes} L(\chi)$.

Remark 1.2.1. The technical assumptions that χ_p is sufficiently ramified and that $V_p A$ is potentially crystalline⁵ are removed by p -adic analytic continuation in [13, Theorem B], and replaced by the (necessary) assumption that χ_p is *not exceptional* for A – that is, $e_p(V_{(A, \chi)}) \neq 0$ (which in our case is implied by the potential crystallinity).

Note that for the removal of the first assumption, one only needs the anticyclotomic formula analogous to [9, Theorem C.4], and not the full generality of the multivariable formula in [13, Theorem D].

Remark 1.2.2. Concrete versions of the formula of Theorem B may be obtained by choosing explicit parametrizations f_i and evaluating the term Q . This is a local problem, solved in [5]. In particular, by starting from Theorem B (as generalized to all characters following Remark 1.2.1) and applying the same steps as in the proofs of [10, Theorems 4.3.1, 4.3.3], we obtain the simple p -adic Gross–Zagier formula in anticyclotomic families for elliptic curves A/\mathbf{Q} proposed in [10, Conjecture 4.3.2], and similarly the direct analogue⁶ of Perrin-Riou’s original result in [26].

The theorem has familiar applications extending to the nonsplit case from [9] (when the other ingredients are available); we leave their formulation to the interested reader, and highlight instead an application specific to this case pointed out in [10], as well as a new application to the nonvanishing conjecture for p -adic heights.

A new proof of a result of Greenberg and Stevens. As noted in [10, Remark 5.2.3], the anticyclotomic formula indicated in the previous remark, combined with a result of

⁵An assumption of this sort is equally necessary in the proof of the main theorem of [9] (see Appendix B).

⁶Of course, this is a long detour to get there; readers interested exclusively in the removal of the ‘ p splits’ assumption from Perrin-Riou’s formula, or from its analogue over totally real fields, may prefer to try and insert the new argument of the present paper into Perrin-Riou’s proof, or respectively into [8].

Bertolini and Darmon, gives yet another proof (quite likely the most complicated so far, but amenable to generalizations) of the following famous result of Greenberg and Stevens [16]. If A/\mathbf{Q} is an elliptic curve of split multiplicative reduction at p , with Néron period Ω_A and p -adic L -function $L_p(A, -)$ on $\mathcal{Y}_{\mathbf{Q}}$, then $L_p(A, \mathbf{1}) = 0$ and

$$L'_p(A, \mathbf{1}) = \lambda_p(A) \cdot \frac{L(A, \mathbf{1})}{\Omega_A}, \quad (1.2.9)$$

where $\lambda_p(A)$ is the \mathcal{L} -invariant of Mazur, Tate, and Teitelbaum [21].

We recall a sketch of the argument, referring to [10] for more details. One chooses an imaginary quadratic field E such that p is inert in E and that the twist $A^{(E)}$ satisfies $L(A^{(E)}, \mathbf{1}) \neq 0$. By the anticyclotomic p -adic Gross–Zagier formula, $L'_p(A_E, \mathbf{1})$ is the value at $\chi = \mathbf{1}$ of the height of an anticyclotomic family \mathcal{P} of Heegner points. It is shown in [2, §5.2] that the value $\mathcal{P}(\mathbf{1})$ equals, in an extended Selmer group, the Tate parameter $q_{A,p}$ of $A_{\mathbf{Q}_p}$ multiplied by a square root of $L(A_E, \mathbf{1})/\Omega_{A_E}$. The height of $q_{A,p}$, in the ‘extended’ sense of [21, 25], essentially equals $\lambda_p(A)$. This shows that after harmless multiplication by $L(A^{(E)}, \mathbf{1})/\Omega_{A^{(E)}}$, the two sides of equation (1.2.9) are equal.

Exceptional cases and nonvanishing results. Suppose that A/\mathbf{Q} has multiplicative reduction at a prime p inert in E , and that $L(A_E, \mathbf{1}) \neq 0$. Then *for all but finitely many anticyclotomic characters χ of p -power conductor, a Heegner point in $A_E(\chi)$ is nonzero and the p -adic height pairing on $A_E(\chi)$ is nondegenerate.* This follows from noting, similarly to before, that in the p -adic Gross–Zagier formula in anticyclotomic families for A_E , both sides are nonzero since the height side specializes, at the character $\chi = \mathbf{1}$, to a nonzero multiple of $\lambda_p(A)$, which is in turn nonzero by [1].

A similar argument, applied to the formula in Hida families of [13], will yield the following result: if A/\mathbf{Q} is an elliptic curve with multiplicative reduction and $L(A, \mathbf{1}) \neq 0$, then the Selmer group of the self-dual Hida family \mathbf{f} through A has generic rank 1, and neither the height regulator nor the cyclotomic derivative of the p -adic L -function of \mathbf{f} vanishes. The details will appear in [13].

1.3. Organization of the paper

In §2, we restate our theorems in an equivalent form, a direct generalization of the statements from [9] (up to a correction involving a factor of 2, discussed in Appendix B). In §3 we recall the proof strategy from [9], with suitable modifications and corrections. The new argument to treat p -adic local heights in the nonsplit case is developed in §4.

We conclude with two appendices, one dedicated to some local results and the other containing a list of errata to [9].

2. Comparison with [9]

We compare Theorems A and B with the corresponding results from [9]. We continue with the setup and notation of §1.2.

2.1. The p -adic L -function

We deduce our Theorem A from [9, Theorem A].

Let σ^∞ be the nearly \mathfrak{p} -ordinary, M -rational [9, Definition 1.2.1] representation of $\mathrm{GL}_2(\mathbf{A})$ attached to A as in [9]. In [9, Theorem A], we constructed a p -adic L -function

$$L_{p,\alpha}(\sigma_E),$$

which is a bounded function on a rigid space $\mathcal{Y}'_{/L}{}^{\mathrm{rig}}$ (denoted by \mathcal{Y}' in [9]). In the construction of that theorem (and in all this paper), we use the same additive character $\psi_p = \prod_{v|p} \psi_v$ as in Theorem A (and see the correction in Appendix B for the exact ring of definition of $L_{p,\alpha}(\sigma_E)$).

The space $\mathcal{Y}'^{\mathrm{rig}} = \mathcal{Y}'_{\omega, V^p}{}^{\mathrm{rig}}$ parametrizes certain continuous p -adic characters of $E^\times \backslash E_{\mathbf{A}}^\times$ invariant under an arbitrarily fixed compact open subgroup $V^p \subset E_{\mathbf{A}^{p\infty}}^\times$. Boundedness means precisely that we may (and do) identify $L_{p,\alpha}(\sigma_E)$ with a function on a corresponding scheme

$$\mathcal{Y}' \subset \mathrm{Spec} \mathcal{O}_L \llbracket E^\times \backslash E_{\mathbf{A}^\infty}^\times / V^p \rrbracket \otimes L \tag{2.1.1}$$

that, when also viewed as a space of characters χ' , is the subscheme cut out by the closed condition $\omega \cdot \chi'|_{\widehat{\mathcal{O}}_F^{p,\times}} = \mathbf{1}$. Similarly to \mathcal{Y}_F , the scheme \mathcal{Y}' contains a 0-dimensional subscheme $\mathcal{Y}'^{\mathrm{l.c.}}$ parametrizing the locally constant characters in \mathcal{Y}' . The function $L_{p,\alpha}(\sigma_E)$ is characterized by the following property. Denote by D_K the discriminant of a number field K . Then at all $\chi' \in \mathcal{Y}'$ with underlying embedding $\iota: L \hookrightarrow \mathbf{C}$, we have

$$L_{p,\alpha}(\sigma_E)(\chi') = \prod_{v|p} Z_v^\circ(\chi'_v, \psi_v) \cdot \frac{\pi^{2[F:\mathbf{Q}]|D_F|^{1/2}}}{2\zeta_F(2)} \cdot \mathcal{L}(\iota V_{(A,\chi')}) \tag{2.1.2}$$

for certain local factors Z_v° .

Fix a finite-order character

$$\chi: E^\times \backslash E_{\mathbf{A}^\infty}^\times \cong \mathrm{Gal}(E^{\mathrm{ab}}/E) \rightarrow L(\chi)^\times$$

satisfying $\omega \cdot \chi|_{\mathbf{A}^\infty \times} = \mathbf{1}$, and consider the map

$$\begin{aligned} j_\chi: \mathcal{Y}_F &\rightarrow \mathcal{Y}', \\ \chi_F &\mapsto \chi \cdot \chi_F \circ N_{E_{\mathbf{A}^\infty \times} / \mathbf{A}^\infty \times}. \end{aligned}$$

Proof of Theorem A. Define

$$C(\chi'_p) := \frac{e_p(V_{(A,\chi')})}{\prod_{v|p} Z_v^\circ(\chi'_v, \psi_v)}. \tag{2.1.3}$$

We show in Proposition A.1.2 that this is a constant in $C \in L$, independent of χ'_p .

Define

$$\mathcal{L}_p(V_{(A,\chi)}) := \frac{2\zeta_F(2)}{\pi^{2[F:\mathbf{Q}]|D_F|^{1/2}}} \cdot C \cdot L(1, \sigma_v, \mathrm{ad}) \cdot j_\chi^* L_{p,\alpha}(\sigma_E), \tag{2.1.4}$$

a function in $\mathcal{O}(\mathcal{Y}_F)$. It is clear from the definition and equation (2.1.2) that it satisfies the required interpolation property. \square

2.2. Equivalence of statements

We now restate Theorem B in a form that directly generalizes [9, Theorem B]. It is the form in which we will prove it, for convenience of reference.

We retain the setup of §1.2. Let d_F be the Γ_F -differential defined before [9, Theorem B]. For all $v \nmid \infty$, let dt_v be the measure on E_v^\times / F_v^\times specified in [9, paragraph after (1.1.2)] if $v \nmid \infty$ and the measure giving $\mathbf{C}^\times / \mathbf{R}^\times$ volume 2 if $v \mid \infty$.

Theorem 2.2.1. *Retain the assumptions of Theorem B, and fix a decomposition $(\cdot)_\pi = \prod_v (\cdot)_v$, with $(1,1)_v = 1$ if $v \mid \infty$. Then for all $f_1 \in \pi, f_2 \in \pi^\vee$,*

$$\langle P(f_1, \chi), P^\vee(f_2, \chi^{-1}) \rangle = c_E \cdot \prod_{v \mid p} Z_v^\circ(\chi_v)^{-1} \cdot d_F L_{p, \alpha}(\sigma_{A, E})(\chi) \cdot \prod_{v \nmid \infty} Q_{(\cdot)_v, dt_v}(f_1, f_2, \chi) \tag{2.2.1}$$

in $\Gamma_F \hat{\otimes} L(\chi)$, where

$$c_E := \frac{\zeta_F(2)}{(\pi/2)^{[F:\mathbf{Q}]} |D_E|^{1/2} L(1, \eta)} \in \mathbf{Q}^\times.$$

Lemma 2.2.2. *Theorem 2.2.1 is equivalent to Theorem B. When every prime $v \mid p$ splits in E , it specializes to [9, Theorem B] as corrected in Appendix B.*

Proof. The second assertion is immediate; we prove the first one. First, we note that equation (2.2.1) is equivalent to

$$\frac{\langle P(f_1, \chi), P^\vee(f_2, \chi^{-1}) \rangle}{(f_3, f_4)_\pi} = c_E \cdot \prod_{v \mid p} Z_v^\circ(\chi_v)^{-1} \cdot d_F L_{p, \alpha}(\sigma_{A, E})(\chi) \cdot 2^{-[F:\mathbf{Q}]} \prod_v \frac{Q_{(\cdot)_v, dt_v}(f_1, f_2, \chi)}{(f_3, v, f_4, v)_v} \tag{2.2.2}$$

for any $f_3 \in \pi, f_4 \in \pi^\vee$ with $f_{3, \infty} = f_{4, \infty} = 1$ and $(f_3, f_4)_\pi \neq 0$ (the extra power of 2 comes from the Archimedean places). The left-hand side of equation (2.2.2) is the same as that of the formula of Theorem B, and the product of the terms after the L -derivative in the right-hand side equals

$$\begin{aligned} & 2^{-[F:\mathbf{Q}]} \prod_v \frac{dt_v}{dt} \cdot Q\left(\frac{f_1 \otimes f_2}{f_3 \otimes f_4}; \chi\right) \\ &= 2^{1-[F:\mathbf{Q}]} |D_{E/F}|^{1/2} |D_F|^{1/2} \pi^{-[F:\mathbf{Q}]} L(1, \eta) \cdot Q\left(\frac{f_1 \otimes f_2}{f_3 \otimes f_4}; \chi\right), \end{aligned}$$

because the measure $\prod_v dt_v$ (resp., dt) gives $E^\times \setminus \mathbf{A}_E^\times / \mathbf{A}^\times$ volume $2 |D_{E/F}|^{1/2} |D_F|^{1/2} \pi^{-[F:\mathbf{Q}]} L(1, \eta)$ (resp., 1).

Next, we have $d_F L_{p,\alpha}(\sigma_E)(\chi) = \frac{1}{2} d(j_\chi^* L_{p,\alpha}(\sigma_E))(\mathbf{1})$, and it is clear from comparing the interpolation properties that

$$\prod_{v|p} Z_v^\circ(\chi_v)^{-1} \cdot \frac{1}{2} d(j_\chi^* L_{p,\alpha}(\sigma_E))(\mathbf{1}) = \frac{\pi^{2[F:\mathbf{Q}]} |D_F|^{1/2}}{2\zeta_F(2)} \cdot e_p(V_{(A,\chi)})^{-1} \cdot \mathcal{L}'_p(V_{(A,\chi)}, 0).$$

It follows that the right-hand side of equation (2.2.1) equals $c \cdot e_p(V_{(A,\chi)})^{-1} \cdot \mathcal{L}'_p(V_{(A,\chi)}, 0) \cdot Q\left(\frac{f_1 \otimes f_2}{f_3 \otimes f_4}; \chi\right)$, where

$$c = c_E \cdot \frac{\pi^{2[F:\mathbf{Q}]} |D_F|^{1/2}}{2\zeta_F(2)} \cdot 2^{1-[F:\mathbf{Q}]} |D_{E/F}|^{1/2} |D_F|^{1/2} \pi^{-[F:\mathbf{Q}]} L(1, \eta) = 1. \quad \square$$

3. Structure of the proof

We review the formal structure of the proof in [9], dwelling only on those points where the arguments need to be modified or corrected. For an introductory description with some more details than are given in §1.1, see [9, §1.7]. Readers interested in a detailed understanding of the present section are advised to keep a copy of [9] handy.

3.1. Notation and setup

We very briefly review some notation and definitions from [9], which will be used throughout the paper.

Galois groups. If K is a perfect field, we denote by $\mathcal{G}_K := \text{Gal}(\overline{K}/K)$ its absolute Galois group.

Local fields. For v finite a place of F , we denote by ϖ_v a fixed uniformizer and by $q_{F,v}$ the cardinality of the residue field of F_v . We denote by d_v a generator of the absolute different of F_v , by D_v a generator of the relative discriminant of E_v/F_v (equal to 1 unless v ramifies in E), and by e_v the ramification degree of E_v/F_v . If $w|v$ is a place of E , we denote by $q_v: E_v \rightarrow F_v$ and $q_w: E_w \rightarrow F_v$ the relative norm maps.

We denote by $\psi = \prod_v \psi_v: F \setminus \mathbf{A} \rightarrow \mathbf{C}^\times$ the additive character fixed before Theorem A.

Base change of rings and schemes. If R is a ring, R' is an R -algebra, M is an R -module, and S is an R -scheme, we denote $M_{R'} = M \otimes_R R', S_{R'} = S \times_{\text{Spec} R} \text{Spec} R'$.

Groups, measures, integration. We adopt the same notation and choices of measures as in [9, §1.9], including a regularized integration \int^* . In particular, $T := \text{Res}_{E/F} \mathbf{G}_{m,E}, Z = \mathbf{G}_{m,F}$, and on the adelic points of T/Z we use two measures dt (the same as introduced before Theorem 2.2.1) and $d^\circ t$. *The measure denoted by dt in the introduction will not be used.*

Operators at p . Let $v | p$ be a place of F . We denote by ϖ_v a fixed uniformizer at v . For $r \geq 1$ we let $K_1^1(\varpi_v^r) \subset \mathrm{GL}_2(\mathcal{O}_{F,v})$ be the subgroup of matrices which become upper unipotent upon reduction modulo ϖ^r . We denote by

$$U_{v,*} = K_1^1(\varpi_v^r) \begin{pmatrix} 1 & \\ & \varpi_v^{-1} \end{pmatrix} K_1^1(\varpi_v^r), \quad U_v^* = K_1^1(\varpi_v^r) \begin{pmatrix} \varpi^r & \\ & 1 \end{pmatrix} K_1^1(\varpi_v^r)$$

the usual double coset operators, and define

$$w_{r,v} := \begin{pmatrix} & 1 \\ -\varpi_v^r & \end{pmatrix} \in \mathrm{GL}_2(F_v).$$

We also define $w_r := \prod_{v|p} w_{r,v} \in \mathrm{GL}_2(F_p)$, and if $(\beta_v)_{v|p}$ are characters of F_v^\times , we denote $\beta_p(\varpi) := \prod_{v|p} \beta_v(\varpi_v)$.

Spaces of characters. We denote by $\mathcal{Y}_F, \mathcal{Y}', \mathcal{Y}$, respectively, the schemes over L defined in formulas (1.2.7) and (2.1.1) and the subscheme of \mathcal{Y} cut out by the condition $\chi|_{\mathbf{A}^\times} = \omega^{-1}$. We add to this notation a superscript ‘l.c.’ to denote the ind-subschemes of locally constant characters (which has a model over a finite extension of M in L).

Let $\mathcal{I}_{\mathcal{Y}|\mathcal{Y}'}$ be the ideal sheaf of $\mathcal{Y} \subset \mathcal{Y}'$. If \mathcal{M} is a coherent $\mathcal{O}_{\mathcal{Y}'}$ -module, we denote

$$d_F: \mathcal{M} \otimes_{\mathcal{O}_{\mathcal{Y}'}} \mathcal{I}_{\mathcal{Y}|\mathcal{Y}'} \rightarrow \mathcal{M} \otimes_{\mathcal{O}_{\mathcal{Y}'}} \mathcal{I}_{\mathcal{Y}|\mathcal{Y}'} / \mathcal{I}_{\mathcal{Y}|\mathcal{Y}'}^2 = \mathcal{M}|_{\mathcal{Y}} \hat{\otimes} \Gamma_F$$

the normal derivative (compare the definition before [9, Theorem B]).

Kirillov models. Let $\sigma^\infty = \bigotimes_{v|\infty} \sigma_v$ be the M -rational automorphic representation of $\mathrm{GL}_2(\mathbf{A})$ attached to A , and denote abusively still by σ^∞ its base change to L . For every place v the representations σ_v of \mathbf{B}_v^\times and π_v of $\mathrm{GL}_2(F_v)$ are Jacquet–Langlands correspondents.

For $v | p$, we denote by

$$\mathcal{H}_{\psi_v}: \sigma_v \rightarrow C^\infty(F_v^\times, L)$$

a fixed rational Kirillov model.

Orthogonal spaces. We define $\mathbf{V} := \mathbf{B}$ equipped with the reduced norm q , a quadratic form valued in \mathbf{A} . The image of $E_{\mathbf{A}}$ is a subspace \mathbf{V}_1 of the orthogonal space \mathbf{V} , and we let \mathbf{V}_2 be its orthogonal complement. The restriction $q|_{\mathbf{V}_1}$ is the adelization of the norm of E/F .

Schwartz spaces and Weil representation. If \mathbf{V}' is any one of the spaces already discussed, we denote by $\overline{\mathcal{S}}(\mathbf{V}' \times \mathbf{A}^\times) = \bigotimes'_v \overline{\mathcal{S}}(\mathbf{V}'_v \times F_v^\times)$ the Fock space of Schwartz functions considered in [9]. (This differs from the usual Schwartz space only at infinity.) There is a *Weil representation*

$$r = r_\psi: \mathrm{GL}_2(\mathbf{A}) \times \mathrm{O}(\mathbf{V}, q) \rightarrow \mathrm{End} \overline{\mathcal{S}}(\mathbf{V} \times \mathbf{A}^\times),$$

defined as in [9, §3.1]. The orthogonal group of \mathbf{V} naturally contains the product $T(\mathbf{A}) \times T(\mathbf{A})$ acting by left and right multiplication on \mathbf{V} . The Weil representation also

depends on a choice of additive characters ψ . The restriction $r|_{T(\mathbf{A}) \times T(\mathbf{A})}$ preserves the decomposition $\mathbf{V}_1 \oplus \mathbf{V}_2$, hence it accordingly decomposes as $r_1 \oplus r_2$.

Special data at p . We list the functions at the places $v \mid p$ that we use.

Define

$$W_v(y) := \mathbf{1}_{\mathcal{O}_{F,v}-0}(y)|y|_v \alpha_v(y), \tag{3.1.1}$$

the ordinary vector in the fixed Kirillov model \mathcal{K}_{ψ_v} of $\sigma_{A,v}$. We consider

$$\varphi_v = \varphi_{v,r} := \mathcal{K}_{\psi_v}^{-1}(\alpha_v(\varpi_v)^{-r} w_r W_v) \in \sigma_v. \tag{3.1.2}$$

Now we consider Schwartz functions. We let

$$\mathbf{B}_v \cong M_2(F_v)$$

be the indefinite quaternion algebra over F_v ; this choice is justified a posteriori by Corollary A.2.3. The following choices of functions correct and modify the ones fixed in [9] (compare the errata in Appendix B); note in particular that we will use two different functions on $\mathbf{V}_{2,v}$.

Decompose orthogonally $\mathbf{V}_v = \mathbf{V}_{1,v} \oplus \mathbf{V}_{2,v}$, where $\mathbf{V}_{1,v} = E_v$ under the fixed embedding $E_{\mathbf{A}^\times} \hookrightarrow \mathbf{B}^\times$. We define the following Schwartz functions on, respectively, F_v^\times and its product with $\mathbf{V}_{1,v}$, $\mathbf{V}_{2,v}$, \mathbf{V}_v :

$$\begin{aligned} \phi_{F,r}(u) &:= \delta_{1,U_{F,r}}(u), & \text{where } \delta_{1,U_{F,r}}(u) &:= \frac{\text{vol}(\mathcal{O}_{F,v}^\times)}{\text{vol}(1 + \varpi^r \mathcal{O}_{F,v})} \mathbf{1}_{1 + \varpi^r \mathcal{O}_{F,v}}(u), \\ \phi_{1,r}(x_1, u) &:= \delta_{1,U_{T,r}}(x_1) \delta_{1,U_{F,r}}(u), & \text{where } \delta_{1,U_{T,r}}(x_1) &= \frac{\text{vol}(\mathcal{O}_{E,v})}{\text{vol}(1 + \varpi^r \mathcal{O}_{E,v})} \mathbf{1}_{1 + \varpi^r \mathcal{O}_{E,v}}(x_1), \end{aligned} \tag{3.1.3}$$

and

$$\begin{aligned} \phi_2^\circ(x_2, u) &:= \mathbf{1}_{\mathcal{O}_{\mathbf{V}_{2,v}}}(x_2) \mathbf{1}_{\mathcal{O}_{F,v}^\times}(u), \\ \phi_{2,r}(x_2, u) &:= e_v^{-1} |d|_v \cdot \mathbf{1}_{\mathcal{O}_{\mathbf{V}_{2,v}} \cap q^{-1}(-1 + \varpi^r \mathcal{O}_{F,v})}(x_2) \mathbf{1}_{1 + \varpi^r \mathcal{O}_{F,v}}(u), \\ \phi_r(x, u) &:= \phi_{1,r}(x_1, u) \phi_{2,r}(x_2, u). \end{aligned} \tag{3.1.4}$$

p -adic modular forms and q -expansions. In [9, §2], we defined the notion of Hilbert automorphic forms and twisted Hilbert automorphic forms (the latter depend on an extra variable $u \in \mathbf{A}^\times$). We also defined the associated space of q -expansions, and a less redundant space of *reduced* q -expansions. When the coefficient field is a finite extension L of \mathbf{Q}_p , these spaces are endowed with a topology. We have an (injective) reduced- q -expansion map on modular forms, denoted by

$$\varphi' \mapsto \mathfrak{q}\varphi'.$$

The image of modular forms (resp., cusp forms) of level $K^p K_1^1(p^\infty) \subset \text{GL}_2(\mathbf{A}^{p^\infty})$, parallel weight 2, and central character ω^{-1} is denoted by $\mathbf{M} = \mathbf{M}(K^p, \omega^{-1})$ (resp., \mathbf{S}). The closure

of \mathbf{M} (resp., \mathbf{S}) in the space of q -expansions with coefficients in L is denoted \mathbf{M}' (resp., \mathbf{S}') and its elements are called p -adic modular forms (resp., cusp forms).

If $\mathcal{Y}' = \mathcal{Y}_F, \mathcal{Y}'$, we define the notion of a \mathcal{Y}' -family of modular forms by copying word for word [9, Definition 2.1.3]; the resulting notion coincides with that of bounded families on the analogous rigid spaces considered there.

For a finite set of places S disjoint from those above p , we have also defined a certain quotient space $\overline{\mathbf{S}}'_S$ of cuspidal reduced q -expansions modulo those whose coefficients of index $a \in F^\times \mathbf{A}^{S \infty \times}$ all vanish. According to [9, Lemma 2.1.2], for any S the reduced- q -expansion map induces an injection

$$\mathbf{S} \hookrightarrow \overline{\mathbf{S}}'_S. \tag{3.1.5}$$

p -adic Petersson product and p -critical forms. For $\varphi^p \in \sigma$, we defined in [9, Proposition 2.4.4] a functional

$$\ell_{\varphi^p, \alpha} : \mathbf{M}(K^p, \omega^{-1}, L) \rightarrow L, \tag{3.1.6}$$

whose restriction to classical modular forms equals, up to an adjoint L -value, the limit as $r \rightarrow \infty$ of Petersson products with antiholomorphic forms $\varphi^p \varphi_{p,r} \in \sigma$ with component $\varphi_{p,r} = \prod_{v|p} \varphi_{v,r}$ as in equation (3.1.2).

Set $v | p$. We say that a form or q -expansion over a finite-dimensional \mathbf{Q}_p -vector space L is v -critical if its coefficients a_* (where $* \in \mathbf{A}^{\infty \times}$) satisfy

$$a_{m\varpi_v^s} = O(q_{F,v}^s) \tag{3.1.7}$$

in L , uniformly in $m \in \mathbf{A}^{\infty \times}$. Here for two functions $f, g : \mathbf{N} \rightarrow L$, we write

$f = O(g) \iff$ there is a constant $c > 0$ such that $|f(s)| \leq c|g(s)|$ for all sufficiently large s .

The space of p -critical forms is the sum of the spaces of v -critical forms for $v | p$. Any element in those spaces is annihilated by $\ell_{\varphi^p, \alpha}$.

3.2. Analytic kernel

The analytic kernel is a p -adic family of theta-Eisenstein series, related to the p -adic L -function. We review its main properties.

Proposition 3.2.1. *There exist p -adic families of q -expansions of modular forms \mathcal{E} over \mathcal{Y}_F and \mathcal{I} over \mathcal{Y}' , satisfying the following:*

1. *For any $\chi_F \in \mathcal{Y}_F^{1.c.}(\mathbf{C})$ and any $r = (r_v)_{v|p}$ satisfying $c(\chi_F) | p^r$, we have the identity of q -expansions of twisted modular forms of weight 1:*

$$\mathcal{E}(u, \phi_2^{p\infty}; \chi_F) = |D_F| \frac{L^{(p)}(1, \eta \chi_F)}{L^{(p)}(1, \eta)} \mathfrak{a} E_r(u, \phi_2, \chi_F),$$

where

$$E_r(g, u, \phi_2, \chi_F) := \sum_{\gamma \in P^1(F) \backslash SL_2(F)} \delta_{\chi_F, r}(\gamma g w_r) r(\gamma g) \phi_2(0, u) \tag{3.2.1}$$

is the Eisenstein series defined in [9, §3.2], with respect to $\phi_2 = \phi_2^{p\infty}(\chi_F)\phi_{2,p\infty}^{\circ}$, with $\phi_{2,v}^{\circ}$ as in formula (3.1.4) for $v \mid p$ and $\phi_{2,v}(\chi_v)$ for $v \nmid \infty$ and $\phi_{2,v}^{\circ}$ for $v \mid \infty$ as defined in [9, §3.2].

- For $\phi_1 \in \overline{\mathcal{F}}(\mathbf{V}_1 \times \mathbf{A}^{\times})$ and $\chi' \in \mathcal{Y}^{\wedge, c}$, consider the twisted modular form of weight 1 with parameter $t \in E_{\mathbf{A}}^{\times}$:

$$\theta(g, (t, 1)u, \phi_1) := \sum_{x_1 \in E} r_1(g, (t, 1))\phi_1(x, u). \tag{3.2.2}$$

For any $\chi' \in \mathcal{Y}^{\wedge, c}$, define $\chi_F := \omega^{-1}\chi'_{|\mathbf{A}^{\times}} \in \mathcal{Y}_F^{\wedge, c}$. Then for any $r = (r_v)_{v \mid p}$ satisfying $r_v \geq 1$ and $c(\chi_F) \mid p^r$, we have

$$\mathcal{I}(\phi^{p\infty}; \chi') = \frac{c_{U_p}|D_E|^{1/2}}{|D_F|^{1/2}} \int_{[T]}^* \chi'(t) \sum_{u \in \mu_{U_p}^{\natural} \setminus F^{\times}} \mathfrak{a}\theta((t, 1), u, \phi_1; \chi') \mathcal{E}(q(t)u, \phi_2^{p\infty}; \chi_F) d^{\circ}t, \tag{3.2.3}$$

where for $v \mid p$, we have $\phi_{1,v} = \phi_{1,v,r}$ is as in formula (3.1.3).

- We have

$$L_{\varphi^p, \alpha}(\mathcal{I}(\phi^{p\infty})) = L_{p, \alpha}(\sigma_E) \cdot \prod_{v \mid p} |d|_v^2 |D|_v \prod_{v \nmid p\infty} \mathcal{R}_v^{\natural}(W_v, \phi_v, \chi'_v), \tag{3.2.4}$$

where the local terms \mathcal{R}_v^{\natural} are as in [9, Propositions 3.5.1, 3.6.1].

Proof. Part 1 is [9, Proposition 3.3.2]. Part 2 summarizes [9, §3.4]. Part 3 is [9, (3.7.1)], with the correction of Appendix B. □

Derivative of the analytic kernel. We denote

$$\mathcal{I}'(\phi^{p\infty}; \chi) := d_F \mathcal{I}(\phi^{p\infty}; \chi), \tag{3.2.5}$$

a p -adic modular form with coefficients in $\Gamma_F \hat{\otimes} L(\chi)$.

3.3. Geometric kernel

The geometric kernel function [9, §§5.2, 5.3] is related to the heights of Heegner points. We recall its construction and modularity.

CM divisors. For any $x \in \mathbf{B}^{\infty \times}$, we have a Hecke translation $T_x: X \rightarrow X$, and a Hecke correspondence $Z(x)_U$ on $X_U \times X_U$. Fix any $P \in X^{E^{\times}}(E^{\text{ab}})$, and for $x \in \mathbf{B}^{\infty \times}$, let $[x] := T_x P$ be the Hecke-translate of P by x , and let $[x]_U$ be its image in X_U . If H/E is any finite extension, the points in $X_{U,H}$ corresponding to Galois orbits of points of the form $[x]_U$ are called *CM points* (for the CM field E).

Let $\text{Cl}(X_{U,\overline{F}})_{\mathbf{Q}} \supset \text{Cl}^0(X_{U,\overline{F}})_{\mathbf{Q}}$ be the space of divisor classes with \mathbf{Q} -coefficients and its subspace consisting of classes with degree 0 on every connected component. Denote by $(\)^0: \text{Cl}(X_{U,\overline{F}})_{\mathbf{Q}} \rightarrow \text{Cl}^0(X_{U,\overline{F}})_{\mathbf{Q}}$ the linear section of the inclusion whose kernel is spanned

by the push-forwards to $X_{U, \overline{F}}$ of the classes of the canonical bundles of the connected components of $X_{U', \overline{F}}$, for any sufficiently small U' .

We define the χ -isotypic CM divisors

$$t_\chi := \int_{[T]}^* \chi(t) [t^{-1}]_U d^\circ t \in \text{Div} \left(X_{U, \overline{F}} \right)_{L(\chi)},$$

$$t_\chi^0 := \int_{[T]}^* \chi(t) [t^{-1}]_U^0 d^\circ t \in \text{Div}^0 \left(X_{U, \overline{F}} \right)_{L(\chi)},$$

where the integrations simply reduce to (normalized) finite sums.

Generating series. For $a \in \mathbf{A}^{\infty \times}$, $\phi^\infty \in \overline{\mathcal{S}}(\mathbf{V} \times \mathbf{A}^\times)$, consider the correspondences

$$\tilde{Z}_a(\phi^\infty) := c_{U^p} w_U |a| \sum_{x \in U \setminus \mathbf{B}^{\infty \times} / U} \phi^\infty(x, aq(x)^{-1}) Z(x)_U, \tag{3.3.1}$$

where $w_U = |\{\pm 1\} \cap U|$ and c_{U^p} is defined in [9, (3.4.3)]. By [9, Theorem 5.2.1] (due to Yuan, Zhang, and Zhang), there is an automorphic form

$$\tilde{Z}(\phi^\infty) \in C^\infty(\text{GL}_2(F) \backslash \text{GL}_2(\mathbf{A}), \mathbf{C}) \otimes_{\mathbf{Q}} \text{Pic}(X_U \times X_U)_{\mathbf{Q}}, \tag{3.3.2}$$

whose a th reduced coefficient is the image of $\tilde{Z}_a(\phi^\infty)$ for each $a \in \mathbf{A}^{\infty \times}$.

Let

$$\langle \cdot, \cdot \rangle = \langle \cdot, \cdot \rangle_X : J^\vee(\overline{F}) \times J(\overline{F}) \rightarrow \Gamma_F \hat{\otimes} L \tag{3.3.3}$$

be the p -adic height pairing defined as in⁷ [9, Lemma 5.3.1]. (We abusively omit the subscript X , as we will no longer need to use the pairing on $A_E(\chi) \otimes A_E^\vee(\chi)$.)

We define the *geometric kernel* to be

$$\tilde{Z}(\phi^\infty, \chi) := \sum_{a \in F^\times} \left\langle \tilde{Z}_a(\phi^\infty) [1]_U^0, t_\chi^0 \right\rangle \mathbf{q}^a. \tag{3.3.4}$$

By [I, Proposition 5.3.2 and the formula after its proof], the series $\tilde{Z}(\phi^\infty, \chi)$ is (the q -expansion of) a weight 2 cuspidal Hilbert modular form of central character ω^{-1} , with coefficients in $\Gamma_F \hat{\otimes} L(\chi)$.

Geometric kernel and Shimizu lifts. Let

$$\theta_{l_p} : (\sigma^\infty \otimes \overline{\mathcal{S}}(\mathbf{V}^\infty \times \mathbf{A}^{\infty, \times})) \otimes_M L \rightarrow (\pi \otimes \pi^\vee) \otimes_{M, l_p} L$$

be Shimizu’s theta lifting defined in [9, §5.1]. Let

$$\text{T}_{\text{alg}} : \pi^U \otimes_M \pi^{\vee, U} \rightarrow \text{Hom}(J_U, J_U^\vee) \otimes M$$

be defined by $\text{T}_{\text{alg}}(f_1, f_2) := f_2^\vee \circ f_1$.

⁷There is a typo in *loc. cit.* (also noted in Appendix B): the left-hand side of the last equation in the statement should be $\langle f_1'(P_1), f_2'(P_2) \rangle_{J, *}$.

Proposition 3.3.1. *If $\phi_v = \phi_{r,v}$ is as in formula (3.1.4) for all $v \mid p$, then for any sufficiently large r' , the geometric kernel*

$$\tilde{Z}(\phi^\infty, \chi)$$

is invariant under $\prod_{v \mid p} K_1^1(\varpi_v^{r'})$, and it satisfies

$$\begin{aligned} & \ell_{\varphi^p, \alpha} \left(\tilde{Z}(\phi^\infty, \chi) \right) \\ &= 2|D_F|^{1/2}|D_E|^{1/2}L(1, \eta) \cdot \left\langle T_{\text{alg}, t_p} \left(\theta_{t_p} \left(\varphi, \alpha_p \mid \cdot \mid_p(\varpi)^{-r'} w_{r'}^{-1} \phi \right) \right) P_{\chi}, P_{\chi}^{-1} \right\rangle_X. \end{aligned} \quad (3.3.5)$$

Proof. The invariance under $\prod_{v \mid p} K_1^1(\varpi_v^{r'})$ follows from the invariance of ϕ_r under the action of $(^1 \mathcal{O}_{F, p}^1)$ and the continuity of the Weil representation. The proof of equation (3.3.5) is indicated in [9, proof of Proposition 5.4.3] (with the correction of Appendix B). \square

3.4. Kernel identity

We state our kernel identity and recall how it implies the main theorem.

Assumptions on the data. Consider the following local assumptions on the data at primes above p :

Assumption 3.4.1. Let $U_{F,v}^\circ := 1 + \varpi_v^n \mathcal{O}_{F,v}$ with $n \geq 1$ be such that ω_v is invariant under $U_{F,v}^\circ$. The character χ_p is *sufficiently ramified* in the sense that it is nontrivial on

$$V_p^\circ := \prod_{v \mid p} q_v^{-1} \mid_{\mathcal{O}_{E,v}} (U_{F,v}^\circ) \subset \mathcal{O}_{E,p}^\times.$$

(Recall from §3.1 that $q_v : E_v \rightarrow F_v$ is the norm map.)

Under this assumption, we have $t_\chi = t_\chi^0$; see [I, Proposition 8.1.1.3], where $\xi_U \in \text{Cl}(X_U)_\mathbf{Q}$ denotes the *Hodge* class defining the section $\text{Cl}(X_U)_\mathbf{Q} \rightarrow \text{Cl}^0(X_U)_\mathbf{Q}$. The technical advantage gained, which is the same as in [I] and is implicitly reaped in Theorem 3.6.1, is that one may analyze the height-generating series purely in terms of pairs of CM divisors of degree 0, thus avoiding a study of ξ_U and the recourse to p -adic Arakelov theory made in [8].

Assumption 3.4.2. For each $v \mid p$, the open compact $U_v \subset \mathbf{B}_v^\times$ satisfies the following:

- $U_v = U_{v,r} = 1 + \varpi^r M_2(\mathcal{O}_{F,v})$ for some $r \geq 1$.
- The integer $r \geq n$ is sufficiently large that the characters χ_v and $\alpha_v \circ q_v$ of E_v^\times are invariant under $U_{v,r} \cap \mathcal{O}_{E,v}^\times$.

Convention on citations from [9]. In [9], we denoted by S_{nonsplit} the set of places of F nonsplit in E , and by S_p the set of places of F above p . When referring to results from [9], we henceforth stipulate that one should read any assumption such as ‘set $v \in S_{\text{nonsplit}}$ ’

or ‘let v be a place in F nonsplit in E ’ as ‘set $v \in S_{\text{nonsplit}} - S_p$ ’. Similarly, the set S_1 fixed in [9, §6.1] should be understood to consist only of places not above p .

Theorem 3.4.3 (kernel identity). *Assume the hypotheses of Theorem B, and that $U, \varphi^p, \phi^{p\infty}, \chi, r$ satisfy the assumptions of [9, §6.1] as well as Assumptions 3.4.1 and 3.4.2. Define $\phi_p := \otimes_{v|p} \phi_{v,r}$, with $\phi_{v,r} = \text{formula (3.1.4)}$. Then*

$$\ell_{\varphi^p, \alpha}(\text{d}_F \mathcal{S}(\phi^{p\infty}; \chi)) = 2|D_F|L_{(p)}(1, \eta) \cdot \ell_{\varphi^p, \alpha}(\tilde{Z}(\phi^\infty, \chi)).$$

The elements of the proof will be gathered in §3.6.

Lemma 3.4.4. *Theorem 3.4.3 implies Theorem 2.2.1.*

Proof. As in [9, Proposition 5.4.3] as corrected in Appendix B, we consider the following equivalent (by [9, Lemma 5.3.1]) form of the identity of Theorem 2.2.1:

$$\begin{aligned} & \langle \text{T}_{\text{alg}, \iota_p}(f_1 \otimes f_2) P_\chi, P_{\chi^{-1}} \rangle_J \\ &= \frac{\zeta_F^\infty(2)}{(\pi^2/2)^{[F:\mathbb{Q}]} |D_E|^{1/2} L(1, \eta)} \prod_{v|p} Z_v^\circ(\alpha_v, \chi_v)^{-1} \cdot \text{d}_F L_{p, \alpha}(\sigma_{A, E})(\chi) \cdot Q(f_1, f_2, \chi), \end{aligned} \tag{3.4.1}$$

where $\iota_p: M \hookrightarrow L(\chi)$, and $P_\chi = \int_{[T]} \text{T}_t(P - \xi_P)\chi(t) dt \in J(\overline{F})_{L(\chi)}$. By linearity, equation (3.4.1) extends to an identity that makes sense for any element $\mathbf{f} \in \pi \otimes \pi^\vee$. By the multiplicity 1 result for $E_{\mathbb{A}^\infty}^\times$ -invariant linear functionals on each of π, π^\vee , it suffices to prove equation (3.4.1) for one element $\mathbf{f} \in \pi \otimes \pi^\vee$ such that $Q(\mathbf{f}, \chi) \neq 0$ (compare [29, Lemma 3.23]).

We claim that Theorem 3.4.3 gives equation (3.4.1) for $\mathbf{f} = \theta(\varphi, \phi)$, where:

- φ_∞ is standard antiholomorphic in the sense of [9] and ϕ_∞ is standard in the sense of [9]; and
- for all $v \nmid p$, we have that $\varphi_v = \text{equation (3.1.2)}$ and $\phi_v = \phi_{v,r} = \text{formula (3.1.4)}$ for any sufficiently large r .

The claim follows from equations (3.2.4) and (3.3.5), the local comparison between \mathcal{R}_v^\natural and $Q_v \circ \theta_v$ for $v \nmid p$ of [9, Lemma 5.1.1], and the local calculation at $v|p$ of Proposition A.2.2.

Finally, the existence of φ, ϕ satisfying both the required assumptions and $Q(\theta(\varphi, \phi)) \neq 0$ follows from [9, Lemma 6.1.6] away from p , and the explicit formula of Proposition A.2.2 at p . □

3.5. Derivative of the analytic kernel

We start by studying the incoherent Eisenstein series $\mathcal{E}(\phi_2^{p\infty})$. For $a \in F_v^\times$, denote by

$$W_{a,v}^\circ$$

the normalized local Whittaker function of $E_r(\phi_2^{p\infty}(\chi_F)\phi_{2,p}^\circ(\chi_F) = \text{formula (3.2.1)})$, defined as in [9, Proposition 3.2.1 and the paragraph after its proof].

The following reviews and corrects [9, Proposition 7.1.1]:

Proposition 3.5.1. *For each $v \mid p$, let $\phi_{2,v} = \phi_{2,r,v}$ be as in formula (3.1.4).*

1. *Let v be a place of F and set $a \in F_v^\times$.*

(a) *If a is not represented by $(\mathbf{V}_{2,v}, uq)$, then $W_{a,v}^\circ(g, u, \mathbf{1}) = 0$.*

(b) *(Local Siegel–Weil formula.) If $v \nmid p$ and there exists $x_a \in \mathbf{V}_{2,v}$ such that $uq(x_a) = a$, then*

$$W_{a,v}^\circ\left(\begin{pmatrix} y & \\ & 1 \end{pmatrix}, u, \mathbf{1}\right) = \int_{E_v^1} r\left(\begin{pmatrix} y & \\ & 1 \end{pmatrix}, h\right) \phi_{2,v}(x_a, u) dh.$$

(c) *(Local Siegel–Weil formula at p .) If $v \mid p$, $a \in -1 + \varpi^r \mathcal{O}_{F,v}$, and $u \in 1 + \varpi^r \mathcal{O}_{F,v}$, let $x_a \in \mathbf{V}_{2,v}$ be such that $uq(x_a) = a$. Then*

$$W_{a,v}^\circ\left(\begin{pmatrix} y & \\ & 1 \end{pmatrix}, u, \mathbf{1}\right) = |d|_v \int_{E_v^1} r\left(\begin{pmatrix} y & \\ & 1 \end{pmatrix}, h\right) \phi_{2,v}(x_a, u) dh. \tag{3.5.1}$$

2. *For any $a \in F_v^\times \cap \prod_{v \mid p} (-1 + \varpi^r \mathcal{O}_{F,v})$, $u \in F^\times \cap \prod_{v \mid p} (1 + \varpi^r \mathcal{O}_{F,v})$, there is a place $v \nmid p$ of F such that a is not represented by (\mathbf{V}_2, uq) .*

Proof. Parts 1(a) and (b) are as in [9, Proposition 7.1.1]. Before continuing, observe that under our assumptions, a is always represented by $(\mathbf{V}_{2,v}, uq_v)$ for all $v \mid p$: this is clear if v splits in E , and up to possibly enlarging the integer r , it may be seen by the local constancy of q_v and the explicit identity $q_v(j_v) = -1$, where j_v = formula (4.1.2) if v is nonsplit. Then part 1(c) follows by explicit computation of both sides (starting, for example, as in [29, proof of Proposition 6.8] for the left-hand side). Explicitly, we have

equation (3.5.1) =

$$\begin{cases} e_v^{-1} |d|_v \text{vol}\left(E_v^1 \cap \mathcal{O}_{E,v}^\times, dh\right) |y|^{1/2} \mathbf{1}_{\mathcal{O}_{F,v}}(ay) \mathbf{1}_{\mathcal{O}_{F,v}^\times}(y^{-1}u) & \text{if } v(a) \geq 0 \text{ and } v(u) = 0, \\ 0 & \text{otherwise.} \end{cases}$$

Finally, part 2 follows from the observation of the previous paragraph and [29, Lemma 6.3]. □

Lemma 3.5.2. *Suppose that for all $v \mid p$, we have that $\phi_{1,v} = \phi_{1,r,v}$ is as in formula (3.1.3). Then for any $t \in T(\mathbf{A})$, the a th q -expansion coefficient of the the theta series of formula (3.2.2) vanishes unless*

$$a \in \bigcap_{v \mid p} 1 + \varpi^r \mathcal{O}_{F,v}.$$

Proof. This is straightforward. □

Denote

$$U_{p,*}^r := \left(\prod_{v \mid p} U_{v,*} \right)^r,$$

an operator on modular forms that extends to an operator on all the spaces of p -adic q -expansions defined so far.

Corollary 3.5.3. *Suppose that $\phi^{p^\infty} \in \mathcal{S}(\mathbf{V}_2^{p^\infty} \times \mathbf{A}^{p^\infty, \times})$ satisfies the assumptions of [9, §6.1]. Set $\chi \in \mathcal{Y}_\omega^{1.c.}$. For any sufficiently large r , we have*

$$U_{p,*}^r \mathcal{I}(\phi^{p^\infty}; \chi) = 0.$$

Proof. For the first assertion, we need to show that the a th reduced q -expansion coefficient of \mathcal{I} vanishes for all a satisfying $v(a) \geq r$ for all $v \mid p$. By the defining property (3.2.3) of $\mathcal{I}(\chi')$ and the choice of ϕ_p , the group $T(F_p) \subset T(\mathbf{A})$ acts trivially on the q -expansion coefficients of $\mathcal{I}(\chi')$. The remaining integration on $T(F)Z(\mathbf{A})V^p \backslash T(\mathbf{A})/T(F_p)$ is a finite sum, so the coefficients a are a sum of products of the coefficients of index a_1 of θ and of index a_2 of $\mathcal{E}(\mathbf{1})$, for pairs (a_1, a_2) with $a_1 + a_2 = a$. When $a = 0$, the vanishing follows from the vanishing of the constant term of \mathcal{E} , which is proved as in [29, Proposition 6.7]. For $a \neq 0$, by Lemma 3.5.2 only the pairs (a_1, a_2) with $a_1 \in \bigcap_{v \mid p} 1 + \varpi^r \mathcal{O}_{F,v}$ contribute. If $v(a) \geq r$ for $v \mid p$, this forces $a_2 \in \bigcap_{v \mid p} -1 + \varpi_v^r \mathcal{O}_{F,v}$. Then the coefficient of index a_2 of $\mathcal{E}(\mathbf{1})$ vanishes by Proposition 3.5.1. \square

We can now proceed as in [9, §7.2], except for the insertion of the operator $U_{p,*}^r$. (This will be innocuous for the purposes of Theorem 3.4.3, since the kernel of $U_{p,*}^r$ is contained in the kernel of $\ell_{\varphi^p, \alpha}$.) We obtain, under the assumptions of [9, §6.1], a decomposition of formula (3.2.5)

$$U_{p,*}^r \mathcal{I}'(\phi^{p^\infty}; \chi) = \sum_{v \in S_{\text{non-split}} - S_p} U_{p,*}^r \mathcal{I}'(\phi^{p^\infty}; \chi)(v) \tag{3.5.2}$$

valid in the space \mathbf{S} of p -adic q -expansions with coefficients in $\Gamma_F \hat{\otimes} L(\chi)$ (see [9, §6.1] for the definition of $\mathcal{I}'(\phi^{p^\infty}; \chi)(v)$).

3.6. Decomposition of the geometric kernel and comparison

Suppose that Assumptions 3.4.1 and 3.4.2, as well as the assumptions of [9, §6.1], are satisfied. Then we may decompose [9, §8.2] the generating series (3.3.4) as

$$\tilde{Z}(\phi^\infty, \chi) = \sum_v \tilde{Z}(\phi^\infty, \chi)(v), \tag{3.6.1}$$

according to the decomposition $\langle , \rangle_X = \sum_v \langle , \rangle_{X,v}$ of the height pairing. Here the sum runs over all finite places of F .

The following is the main result of [9] on the local comparison away from p . Let $\overline{\mathbf{S}}' = \overline{\mathbf{S}}'_{S_1}$ be the quotient of the space of p -adic q -expansions recalled before formula (3.1.5).

Theorem 3.6.1 ([9, Theorem 8.3.2]). *Let $\phi^\infty = \phi^{p^\infty} \phi_p$ with $\phi_p = \prod_{v \mid p} \phi_v$ as in formula (3.1.4). Suppose that Assumptions 3.4.1 and 3.4.2, as well as the assumptions of [9, §6.1], are satisfied. Then we have the following identities of reduced q -expansions in $\overline{\mathbf{S}}'$:*

1. If $v \in S_{\text{split}} - S_p$, then

$$\tilde{Z}(\phi^\infty, \chi)(v) = 0.$$

2. If $v \in S_{\text{nonsplit}} - S_1 - S_p$, then

$$U_{p,*}^r \mathcal{S}'(\phi^{p^\infty}; \chi)(v) = 2|D_F|L_{(p)}(1, \eta) U_{p,*}^r \tilde{Z}(\phi^\infty, \chi)(v).$$

3. If $v \in S_1$, then

$$U_{p,*}^r \mathcal{S}'(\phi^{p^\infty}; \chi)(v), \quad U_{p,*}^r \tilde{Z}(\phi^\infty, \chi)(v)$$

are theta series attached to a quaternion algebra over F .

4. The sum

$$U_{p,*}^r \tilde{Z}(\phi^\infty, \chi)(p) := \sum_{v \in S_p} U_{p,*}^r \tilde{Z}(\phi^\infty, \chi)(v)$$

belongs to the isomorphic image $\bar{\mathbf{S}} \subset \bar{\mathbf{S}}'$ of the space of p -adic modular forms \mathbf{S} .

By this theorem and the decompositions of \mathcal{S}' and \tilde{Z} , the proof of the kernel identity of Theorem 3.4.3 (hence of the main theorem) is now reduced to showing the following proposition. (See [9, §8.3, last paragraph] for the details of the deduction.)

Proposition 3.6.2. *Retain the assumptions of Theorem 3.6.1, and further assume that $V_p A$ is potentially crystalline at all $v \mid p$. Then the p -adic modular form*

$$U_{p,*}^r \tilde{Z}(\phi^\infty, \chi)(p) \in \bar{\mathbf{S}}$$

is annihilated by $\ell_{\varphi^p, \alpha}$.

Let S be a finite set of non-Archimedean places of F such that for all $v \notin S$, all the data are unramified, U_v is maximal, and ϕ_v is standard. Let $K = K^p K_p$ be the level of the modular form $\tilde{Z}(\phi^\infty)$, and let

$$T_{\iota_p}(\sigma^\vee) \in \mathcal{H}^S(L) = \mathcal{H}^S(M) \otimes_{M, \iota_p} L$$

be any σ^\vee -idempotent in the Hecke algebra as in [9, Proposition 2.4.4]. By that result, in order to establish Proposition 3.6.2 it suffices to prove that

$$\ell_{\varphi^p, \alpha} \left(U_{p,*}^r T_{\iota_p}(\sigma^\vee) \tilde{Z}(\phi^\infty, \chi)(p) \right) = 0. \tag{3.6.2}$$

As in [9], we will in fact prove the following, which implies equation (3.6.2):

Proposition 3.6.3. *Let $v \mid p$. Under the assumptions of Proposition 3.6.2, for all $v \mid p$, the element*

$$T_{\iota_p}(\sigma^\vee) \tilde{Z}(\phi^\infty, \chi)(v) \in \bar{\mathbf{S}}'$$

is v -critical in the sense of equation (3.1.7).

The proof will occupy the following section.

4. Local heights at p

The goal of this section is to prove Proposition 3.6.3, whose assumptions we retain throughout except for an innocuous modification to the data at the places $v \mid p$. Namely, let $\tilde{\phi}_v$ be the Schwartz function denoted by ϕ_r in formula (3.1.4), and let $\tilde{U}_v \subset \mathbf{B}_v^\times$ be the open compact subgroup denoted by $U_{v,r}$ in Assumption 3.4.2. Then we define

$$U_v := \tilde{U}_v U_{F,v}^\circ, \quad \phi_v := \int_{U_{F,v}^\circ} r(z,1) \tilde{\phi}_v dz.$$

Since $\chi_{v \mid F_v^\times} = \omega_v^{-1}$ is by construction invariant under $U_{F,v}^\circ$, the geometric kernels $\tilde{Z}(\phi^{p^\infty} \tilde{\phi}_p, \chi)$ and $\tilde{Z}(\phi^{p^\infty} \phi_p, \chi)$ are equal. Therefore we may work on the curve X_U .

We fix a place $v \mid p$. If v splits in E , then the desired result is proven in [9, §9] with the correction in Appendix B. Therefore we may and do assume that v is nonsplit. We denote by w the place of E above v .

We refer the reader to §1.1 for a general sketch of our argument. It is developed here as follows. In §4.1, we prove that after acting by a high power of $U_{v,*}$, the coefficients of the generating series are height pairings with CM points of high v -conductor (*norm relation*). After some general background in §4.2, we use the norm relation to prove the decay property of arithmetic intersection multiplicities in §4.3. Finally, in §4.4 we use again the norm relation and some p -adic Hodge theory to prove the decay property of local heights.

4.1. Norm relation for the generating series

The goal of this subsection will be to show that for s large enough, each q -expansion coefficient of $U_{p,*}^s \tilde{Z}(\phi^\infty, \chi)$ is a height pairing of CM divisors, one of which is supported on Galois orbits of CM points of the ‘pseudo-conductor’ s (as defined later).

We start by considering the $U_{v,*}$ -action on the generating series $\tilde{Z}(\phi^\infty) =$ formula (3.3.2). Recall that d_v (§3.1) is a generator of the different ideal of F_v .

Lemma 4.1.1. *If $a \in \mathbf{A}^{\infty \times}$ satisfies $v(a) \geq -v(d_v)$, the a th reduced q -expansion coefficient of $U_{v,*} \tilde{Z}(\phi^\infty)$ equals*

$$\tilde{Z}_{a\varpi_v}(\phi^\infty),$$

where $\tilde{Z}_{a'}(\phi^\infty)$ is defined in formula (3.3.1).

Proof. After computing the Weil action of $U_{v,*}$ on ϕ , this is a simple change of variables. □

We can factor

$$\tilde{Z}_a(\phi^\infty) = \tilde{Z}_{a^v}(\phi^{v\infty}) Z_{a^v}(\phi_v)$$

as the composition of the commuting correspondences

$$\begin{aligned} \tilde{Z}_a^v(\phi^{v\infty}) &:= c_{U^p} \sum_{x^v \in U^v \backslash \mathbf{B}^{v\infty \times} / U^v} \phi^{v\infty}(x^v, aq(x^v)^{-1}) Z(x^v)_U, \\ Z_{a_v}(\phi_v) &:= \sum_{x_v \in U_v \backslash \mathbf{B}_v^\times / U_v} \phi_v(x_v, aq(x_v)^{-1}) Z(x_v)_U. \end{aligned} \tag{4.1.1}$$

From here until after the proof of Lemma 4.1.2, we work in a local situation and drop *v* from the notation. Let $\theta \in \mathcal{O}_E$ be such that $\mathcal{O}_E = \mathcal{O}_F + \theta \mathcal{O}_F$, and write $T = \text{Tr}_{E/F}(\theta)$, $N = N_{E/F}(\theta)$. Fix the embedding $E \rightarrow \mathbf{B} = M_2(F)$ to be

$$t = a + \theta b \mapsto \begin{pmatrix} a + bT & bN \\ -b & a \end{pmatrix}.$$

Set

$$j := \begin{pmatrix} 1 & T \\ & -1 \end{pmatrix}. \tag{4.1.2}$$

Then $j^2 = 1$, $q(j) = -1$, and for all $t \in E$, $jt = t^c j$; and in the orthogonal decomposition $\mathbf{V} = \mathbf{V}_1 \oplus \mathbf{V}_2$, we have

$$\mathcal{O}_{\mathbf{V}_2} = \mathbf{V}_2 \cap M_2(\mathcal{O}_F) = j\mathcal{O}_E.$$

Let $\Xi(\varpi^r) = 1 + \varpi^r \mathcal{O}_E + j(\mathcal{O}_E \cap q^{-1}(1 + \varpi^r \mathcal{O}_F))$; then ϕ is a fixed multiple of the characteristic function of $\Xi(\varpi^r) \times (1 + \varpi^r \mathcal{O}_F) \subset \mathbf{B} \times F^\times$. For $a \in F^\times$, let

$$\Xi(\varpi^r)_a = \{x \in \Xi(\varpi^r) \mid q(x) \in a(1 + \varpi^r \mathcal{O}_F)\}.$$

Then the local component (4.1.1) of the generating series equals

$$Z_a(\phi) = \sum_{x \in U \backslash \Xi(\varpi^r)_a U_F^\circ / U} Z(x)_U,$$

up to a constant that is independent of *a*.

Lemma 4.1.2. *Let $a \in F^\times$ satisfy $v(a) = s \geq r$. The natural map $\Xi(\varpi^r)_a U_F^\circ / U \rightarrow U \backslash \Xi(\varpi^r)_a U_F^\circ / U$ is a bijection. For either quotient set, a complete set of representatives is given by the elements*

$$x(b) := 1 + jb$$

as *b* ranges through a complete set of representatives for

$$q^{-1}(1 - a(1 + \varpi^r \mathcal{O}_F)) / (1 + \varpi^{r+s} \mathcal{O}_E) \subset (\mathcal{O}_E / \varpi^{r+s} \mathcal{O}_E)^\times.$$

Proof. It is equivalent to prove the same statement for the quotients of $\Xi(\varpi^r)_a$ by the group $\tilde{U} = 1 + \varpi^r M_2(\mathcal{O}_F)$. By acting on the right with elements of $\mathcal{O}_E \cap U$, we can bring any element of Ξ to one of the form $x(b)$. Write any $\gamma \in \tilde{U}$ as $\gamma = 1 + \varpi^r u_1 + j\varpi^r u_2$ with $u_1, u_2 \in \mathcal{O}_E$. Then

$$x(b)\gamma = (1 + jb)(1 + \varpi^r u_1 + j\varpi^r u_2) = 1 + \varpi^r(u_1 + b^c u_2) + j(b + \varpi^r(bu_1 + u_2))$$

is another element of the form $x(b')$ if and only if $u_1 = -b^c u_2$. In this case,

$$b' = b + \varpi^r(1 - q(b))u_2 \in b(1 + \varpi^{r+s}\mathcal{O}_E).$$

Thus the class of b modulo ϖ^{r+s} is the only invariant of the quotient Ξ/\tilde{U} . The \tilde{U} -action on the left similarly preserves this invariant. □

We go back to a global setting and notation, restoring the subscripts v and w . Denote by $\text{rec}_{E_w} : E_w^\times \rightarrow \mathcal{G}_{E,w}^{\text{ab}}$ the reciprocity map of class field theory. Recall that for $x \in \mathbf{B}^{\infty \times}$, we have a point $[x]_U \in X_U$; for a subset $\Xi' \subset \mathbf{B}^{\infty \times}$, we similarly denote $[\Xi']_U = \{[x]_U \mid x \in \Xi'\}$.

Lemma 4.1.3. *Fix $a \in \mathcal{O}_{F,v}$ with $v(a) = s \geq r$.*

1. *Set $b \in (\mathcal{O}_{E,w}/\varpi_v^{r+s}\mathcal{O}_{E,w})^\times$, $t \in 1 + \varpi_v^r\mathcal{O}_{E,w}$. Then*

$$\text{rec}_{E_w}(t)[x(b)]_U = [x(bt^c/t)]_U.$$

2. *We have*

$$[\Xi(\varpi_v^r)_a U_{F,v}^\circ]_U = \bigsqcup_{\bar{b}} \text{rec}_{E_w} \left((1 + \varpi_v^r\mathcal{O}_{E,w}) / (1 + \varpi_v^r\mathcal{O}_{F,v}) (1 + \varpi_v^{r+s}\mathcal{O}_{E,w}) \right) [x(\bar{b})]_U,$$

where the Galois action is faithful, and \bar{b} ranges through a set of representatives for

$$q_v^{-1}(1 - a(1 + \varpi_v^r\mathcal{O}_{F,v})) / (1 + \varpi_v^{r+s}\mathcal{O}_{E,w}) \cdot (1 + \varpi_v^{r+v(\theta-\theta^c)}\mathcal{O}_{E,w} \cap q_v^{-1}(1)). \tag{4.1.3}$$

The size of this set is bounded uniformly in a .

Proof. For part 1, we have

$$\text{rec}_{E_w}(t)[x(b)]_U = [t + tjb]_U = [t + jt^c b]_U = [1 + jbt^c/t]_U.$$

Part 2 follows Lemma 4.1.2 and part 1, noting that the group $(1 + \varpi_v^{r+v(\theta-\theta^c)}\mathcal{O}_{E,w}) \cap q_v^{-1}(1)$ is the image of the map $t \mapsto t^c/t$ on $1 + \varpi_v^r\mathcal{O}_{E,w}$. Finally, the map (projection, q_v) gives an injection from

$$q_v^{-1}(1 - a(1 + \varpi_v^r\mathcal{O}_{F,v})) / (1 + \varpi_v^{r+s}\mathcal{O}_{E,w}) \cdot (1 + \varpi_v^{r+v(\theta-\theta^c)}\mathcal{O}_{E,w} \cap q_v^{-1}(1))$$

to

$$\left(\mathcal{O}_{E,w} / \varpi_v^{r+v(\theta-\theta^c)}\mathcal{O}_{E,w} \right)^\times \times (1 - a(1 + \varpi_v^r\mathcal{O}_{F,v})) / q(1 + \varpi_v^{r+s}\mathcal{O}_{E,w}),$$

whose size is bounded uniformly in a (more precisely, the second factor is isomorphic to $\mathcal{O}_{F,v}/\text{Tr}(\mathcal{O}_{E,v})$ via the map $1 - a(1 + \varpi_v^r x) \mapsto x$). □

We denote by \bar{w} an extension of the place w to E^{ab} . For $s \geq 0$, let

$$H_s \subset E^{\text{ab}}$$

be the finite abelian extension of E with norm group $U_F^\circ U_T^v (1 + \varpi_v^{r+v+s}\mathcal{O}_{E,v})$, where $U_T^v = U^v \cap E_{\mathbf{A}^\infty}^\times$. Let $H_\infty = \bigcup_{s \geq 0} H_s$. If r_v is sufficiently large, for all $s \geq 0$ the extension H_s/H_0 is totally ramified at \bar{w} of degree $q_{F,v}^s$, and

$$\text{Gal}(H_s/H_0) \cong \text{Gal}(H_{s,\bar{w}}/H_{0,\bar{w}}) \cong (1 + \varpi_v^{r_v}\mathcal{O}_{E,v}) / (1 + \varpi_v^{r_v}\mathcal{O}_{F,v}) (1 + \varpi_v^{r+v+s}\mathcal{O}_{E,v}).$$

In particular,

$$[H_s : H_0] = [H_{s,\bar{w}} : H_{0,\bar{w}}] = q_{F,v}^s.$$

We will say that a CM point $z \in X_{H_0}$ has *pseudo-conductor* $s \geq 0$ (at \bar{w}) if $H_{0,\bar{w}}(z) = H_{s,\bar{w}}$.

Proposition 4.1.4. *There exists an integer $d > 0$ such that for all $a \in \mathbf{A}^{\infty,x}$ with $v(a) = s \geq r_v$, there exists a degree 0 divisor $D_a \in d^{-1}\text{Div}^0(X_{U,H_s})$, supported on CM points, such that*

$$\tilde{Z}_a(\phi^\infty)[1]_U = \text{Tr}_{H_s/H_0}(D_a)$$

in $\text{Div}^0(X_{U,H_0})$. All prime divisor components of D_a are CM points of pseudo-conductor s .

Proof. This follows from Lemma 4.1.3(2), by taking D_a to be a fixed rational multiple (independent of a) of

$$\tilde{Z}_a^v(\phi^{v\infty}) \sum_{\bar{b} \in \text{formula (4.1.3)}} [x(\bar{b})]_U.$$

The divisor is of degree 0 by [9, Proposition 8.1.1]. Its prime components are not defined over proper subfields $H_{s'} \subset H_s$, because of the faithfulness statement of Lemma 4.1.3(2). □

4.2. Intersection multiplicities on arithmetic surfaces

Before continuing, we gather some definitions and a key result.

Ultrametricity of intersections on surfaces. Let \mathcal{X} be a 2-dimensional regular Noetherian scheme, finite flat over a field κ or a discrete valuation ring \mathcal{O} with residue field κ . We denote by $(\cdot)_{\mathcal{X}}$ the usual \mathbf{Z} -bilinear intersection-multiplicity pairing of divisors intersecting properly on \mathcal{X} ; for effective divisors D_j ($j = 1, 2$) with $\mathcal{O}_{D_j} = \mathcal{O}_{\mathcal{X}}/\mathcal{I}_j$, it is defined by

$$(D_1 \cdot D_2)_{\mathcal{X}} = \text{length}_{\kappa} \mathcal{O}_{\mathcal{X}} / (\mathcal{I}_1 + \mathcal{I}_2).$$

The subscript \mathcal{X} will be omitted when it is clear from context.

We will need the following result of García Barroso, González Pérez, and Popescu-Pampu:

Proposition 4.2.1. *Let R be a Noetherian regular local ring of dimension 2, which is a flat module over a field or a discrete valuation ring. Let Δ be any irreducible curve in $\text{Spec}R$. Then the function*

$$d_{\Delta}(D_1, D_2) := \begin{cases} (D_1 \cdot \Delta)(D_2 \cdot \Delta) / (D_1 \cdot D_2) & \text{if } D_1 \neq D_2, \\ 0 & \text{if } D_1 = D_2 \end{cases}$$

is an ultrametric distance on the space of irreducible curves in $\text{Spec}R$ different from Δ .

Proof. For those rings R that further satisfy the property of containing \mathbf{C} , this is proved in [15]. The proof only relies on (i) the existence of embedded resolutions of divisors in the spectra of such rings and (ii) the negativity of the intersection matrix of the exceptional divisor of a projective birational morphism between spectra of such rings. Both results still hold under our weaker assumptions: see [20, Theorem 9.2.26] for (i) and [20, Theorem 9.1.27, Remark 9.1.28] for (ii). \square

Arithmetic intersection multiplicities. Suppose now that \mathcal{X} is a 2-dimensional regular Noetherian scheme, proper flat over a discrete valuation ring \mathcal{O} with residue field κ . A divisor on \mathcal{X} is called *horizontal* (resp., *vertical*) if each of the irreducible components of the support $|D|$ is flat over \mathcal{O} (resp., contained in the special fiber \mathcal{X}_κ). We extend $(\cdot) := (\cdot)_{\mathcal{X}}$ to a bilinear form (\bullet) on pairs of divisors on \mathcal{X} sharing no common horizontal irreducible component of the support by

$$(\mathcal{X}_\kappa \bullet V) := 0$$

if V is any vertical divisor.

Denote by X the generic fiber of \mathcal{X} . If $D \in \text{Div}^0(X)$ with Zariski closure \overline{D} in \mathcal{X} , a *flat extension* of D is a divisor $\widehat{D} \in \text{Div}(\mathcal{X})_{\mathbf{Q}}$ such that $\widehat{D} - \overline{D}$ is vertical and

$$(\widehat{D} \bullet V) = 0$$

for any vertical divisor V on \mathcal{X} . A flat extension of D exists and is unique up to addition of rational linear combinations of the connected components of \mathcal{X}_κ .

The arithmetic intersection multiplicity on divisors with disjoint supports in $\text{Div}^0(X)$ is then defined by

$$m_X(D_1, D_2) := (\overline{D}_1 \bullet \widehat{D}_2) = (\widehat{D}_1 \bullet \overline{D}_2) \in \mathbf{Q}.$$

4.3. Decay of intersection multiplicities

We continue using the notation introduced in §4.1. Let $m_{\overline{w}} := m_{X_{H_0, \overline{w}}}$. Developing the approximation argument sketched in the introduction, we will show that for any degree-0 divisor D on X_{H_0} , we have

$$m_{\overline{w}}\left(\widetilde{Z}_{a\varpi^s}(\phi^\infty)[1]_U, D\right) = O(q_{F,v}^s)$$

in L , uniformly in a .

Set $U_{0,v} := \text{GL}_2(\mathcal{O}_{F,v}) \subset \mathbf{B}_v^\times$ and $X_0 := X_{U_v U_{0,v}}$, let \mathcal{X}_0 be the canonical model of X_{0,F_v} over $\mathcal{O}_{F,v}$, which is smooth [6], and let \mathcal{X}'_0 be its base change to $\mathcal{O}_{H_0, \overline{w}}$. Let \mathcal{X} be the integral closure of \mathcal{X}'_0 in $X_{H_0, \overline{w}}$, which is a regular model of $X_{H_0, \overline{w}}$ over $\mathcal{O}_{H_0, \overline{w}}$, and

let $p: \mathcal{X} \rightarrow \mathcal{X}'_0$ be the natural map. Thus we have a diagram

$$\begin{array}{ccc}
 X_{H_0, \bar{w}} & \longrightarrow & \mathcal{X} \\
 \downarrow & & \downarrow p \\
 X_{0, H_0, \bar{w}} & \longrightarrow & \mathcal{X}'_0 = \mathcal{X}_{0, \mathcal{O}_{H, \bar{w}}} \\
 \downarrow & & \downarrow \\
 X_{0, E_v} & \longrightarrow & \mathcal{X}_{0, \mathcal{O}_{E, w}}
 \end{array}$$

of curves and regular integral models. (The bottom row will be used in proving Lemma 4.3.2.)

Some intersection multiplicities. As a preliminary, we first compute the intersection multiplicities of Zariski closures of CM points with the special fiber of \mathcal{X} , then bound their intersections with horizontal divisors.

We denote by κ the residue field of H_0, \bar{w} and by k the algebraic closure of κ . For a scheme \mathcal{C} and a point $y \in \mathcal{C}$, we denote $\mathcal{C}_y := \text{Spec } \mathcal{O}_{\mathcal{C}, y}$.

Lemma 4.3.1. *Let $z_s \in X_{H_0}$ be a CM point with pseudo-conductor s , let \bar{z}_s be its closure in \mathcal{X} , and let $y \in \mathcal{X}_\kappa$ be its reduction modulo \bar{w} . Then*

$$(\bar{z}_s \cdot [\mathcal{X}_{y, \kappa}])_{\mathcal{X}_y} = [\kappa(y) : \kappa] q_{F, v}^s.$$

(Recall that, following §3.1, $\mathcal{X}_{y, \kappa}$ denotes the special fiber of \mathcal{X}_y .)

Proof. We will deduce this from Gross’s theory of quasicanonical liftings [18], which we recall. The situation is purely local, and we drop all subscripts v, w , and \bar{w} . For a finite extension $K \supset E$ contained in E^{ab} , let K^{un} be the maximal unramified extension of K contained in E^{ab} (thus the residue field of K^{un} is identified with k).

By [6, §7.4], for any supersingular point $y_0 \in \mathcal{X}_{0, \mathcal{O}_E^{\text{un}}}$ the completed local ring of $\mathcal{X}_{0, \mathcal{O}_E^{\text{un}}}$ at y_0 is isomorphic to $\mathcal{O}_{E^{\text{un}}}[[u]]$ and is the deformation ring of formal modules studied by Gross. The main result of [18] is that for any CM point $z_0 \in X_{0, E^{\text{un}}}$, there exists a unique integer t (the conductor of z_0) such that the following hold. First, the field $E^{\text{un}}(z_0)$ is the abelian extension $E^{(t)}$ of E^{un} with norm group $(\mathcal{O}_F + \varpi_v^t \mathcal{O}_E)^\times / \mathcal{O}_F^\times$, which is totally ramified of some degree d_t . Second, the inclusion of the Zariski closure $\bar{z}_0 \hookrightarrow \mathcal{X}_{0, \mathcal{O}_E^{\text{un}}}$ gives rise to a map of complete local rings

$$\mathcal{O}_{E^{\text{un}}}[[u]] \rightarrow \mathcal{O}_{E^{(t)}} = \mathcal{O}(\bar{z}_0),$$

which sends u to a uniformizer $\varpi^{(t)}$ of $E^{(t)}$. It follows that if μ_t is the minimal polynomial of $\varpi^{(t)}$,

$$(\bar{z}_0 \cdot \mathcal{X}_{0, k})_{\mathcal{X}_{0, \mathcal{O}_E^{\text{un}}, y_0}} = \dim_k \mathcal{O}_{E^{\text{un}}}[[u]] / (\varpi_E, \mu_t(u)) = \dim_k \mathcal{O}_{E^{(t)}} / \varpi_E = d_t. \tag{4.3.1}$$

Consider now the situation of the lemma. By the projection formula,

$$\begin{aligned}
 (\bar{z}_s \cdot [\mathcal{X}_{y,\kappa}])_{\mathcal{X}_y} &= (\bar{z}_s \cdot \mathbf{p}^* \mathcal{X}_{0,\kappa_0})_{\mathcal{X}'_y} \\
 &= (\mathbf{p}_* \bar{z}_s \cdot \mathcal{X}_{0,\kappa})_{\mathcal{X}'_{0,y'}} = [H_0(z_s) : H_0(z_{0,s})] \cdot (\bar{z}_{0,s} \cdot \mathcal{X}_{0,\kappa})_{\mathcal{X}'_{0,y'}},
 \end{aligned}$$

where $y' = \mathbf{p}(y)$ and $z_{0,s} = \mathbf{p}(z_s) \in X_{0,H_0}$ is a CM point. The last intersection multiplicity is $[\kappa(y) : \kappa]$ times the multiplicities of the base-changed divisors to the ring of integers of H_0^{un} , where $z_{0,s}$ remains an irreducible divisor since $H_0(z_{0,s}) \subset H_0(z_s)$ is totally ramified over H_0 . We perform such base change to $\mathcal{O}_{H_0^{\text{un}}}$ without altering the notation. Let z be the image of $z_{0,s} \in X_{H_0^{\text{un}}}$ in $X_{E^{\text{un}}}$, and let t be the conductor of z ; so $E^{(t)} \subset H_0^{\text{un}}(z_{0,s})$. The fiber above $z \cong \text{Spec } E^{(t)}$ in $X_{H_0^{\text{un}}}$ is

$$\text{Spec } E^{(t)} \otimes_{E^{\text{un}}} H_0^{\text{un}} = \text{Spec } H_0^{\text{un}}(z_{0,s})^{\oplus c}$$

for $c = [E^{(t)} : E^{\text{un}}] \cdot [H_0^{\text{un}} : E^{\text{un}}] / [H_0(z_{0,s}) : H_0]$, and $z_{0,s}$ is one of the factors in the right-hand side. By the projection formula applied to $\mathcal{X}_0 \times \text{Spec } \mathcal{O}_{H_0^{\text{un}}} \rightarrow \mathcal{X}_0 \times \text{Spec } \mathcal{O}_{E^{\text{un}}}$ and equation (4.3.1), we have

$$\begin{aligned}
 [H_0(z_s) : H_0(z_{0,s})] \cdot (\bar{z}_{0,s} \cdot \mathcal{X}_{0,\kappa})_{\mathcal{X}'_{0,y'}} &= [\kappa(y) : \kappa] c^{-1} \cdot [H_0(z_s) : H_0(z_{0,s})] \cdot [H_0^{\text{un}} : E^{\text{un}}] \cdot d_t \\
 &= [\kappa(y) : \kappa] \cdot [H_0(z_s) : H_0] \cdot d_t^{-1} \cdot d_t = [\kappa(y) : \kappa] q_F^s,
 \end{aligned}$$

as desired. □

Lemma 4.3.2. *Let Δ be an irreducible horizontal divisor in \mathcal{X} . The intersection multiplicities*

$$(\Delta \cdot \bar{z})$$

are bounded by an absolute constant as z ranges among CM points of sufficiently large pseudo-conductor reducing to y .

Proof. The intersection multiplicity $(\Delta \cdot \bar{z})$ is bounded by the degree of the natural map $q: \mathcal{X} \rightarrow \mathcal{X}_{0,\mathcal{O}_{E,v}}$ near y , times the intersection multiplicities of the push-forward divisors to $\mathcal{X}_{0,\mathcal{O}_{E,v}}$. Similar to the proof of Lemma 4.3.1, we may estimate this intersection in the base change of \mathcal{X}_0 to $\mathcal{O}_{E^{\text{un}}}$. The base change of the divisor $q_* z$ equals a sum of CM points of $\mathcal{X}_{0,\mathcal{O}_{E^{\text{un}}}}$; because the extensions H_s/H_0 are totally ramified, the number of points in this divisor is bounded by an absolute constant, and the conductors of all those CM points go to infinity with the pseudo-conductor s of z . Thus it suffices to show that if Δ_0 is a fixed horizontal divisor on $\mathcal{X}_{0,E^{\text{un}}}$, its intersection multiplicity with CM points of conductor t is bounded as $t \rightarrow \infty$.

Let $z \in \mathcal{X}_{0,\mathcal{O}_{E^{\text{un}}}}$ be a CM point of conductor t , and let $y_0 \in \mathcal{X}_{0,k}$ be the image of the reduction of t . Now write the image of Δ_0 in the completion of $\mathcal{X}_{0,\mathcal{O}_{E^{\text{un}}}}$ at y_0 as

$$\widehat{\Delta}_0 = \text{Spec } \mathcal{O}_{E^{\text{un}}}[[u]]/(f) \subset \text{Spec } \mathcal{O}_{E^{\text{un}}}[[u]],$$

with $f = \sum_{i=1}^d a_i u^i$ an integral nonconstant monic polynomial. Let $\tilde{f} \in k[u]$ be the reduction of f . Then

$$\begin{aligned} (\Delta_0 \cdot \bar{z}) &= \dim_k \mathcal{O}_{E^{\text{un}}}[[u]]/(f, \mu_t) \\ &= \dim_k \mathcal{O}_{E^{(t)}} / \left(f \left(\varpi^{(t)} \right) \right) = \dim_k \mathcal{O}_{E^{(t)}} / \left(\left(\varpi^{(t)} \right)^{\deg(\tilde{f})} \right) = \deg(\tilde{f}) \leq d \end{aligned}$$

if t is sufficiently large, since the normalized valuations of $\varpi^{(t)}$ decrease to 0 as $t \rightarrow \infty$. This completes the proof of the lemma. \square

Approximation by vertical divisors. The following proposition contains the essential new ingredient of this work. We denote by

$$\text{CM}(X_{H_0})_{\geq s} \supset \text{CM}(X_{H_0})_s$$

respectively the set of CM points of X_{H_0} that have pseudo-conductor at least and equal to a given integer s . We denote by \mathcal{V} the set of irreducible components of \mathcal{X}_κ (henceforth: ‘vertical components’), and if $y \in \mathcal{X}_\kappa$ is a closed point, we denote by $\mathcal{V}_y \subset \mathcal{V}$ the set of vertical components of $\mathcal{X}_{y,\kappa}$. We still use a bar to denote Zariski closure.

Proposition 4.3.3. *There exist an integer $s_0 \geq 1$, depending only on X_{H_0} , and a function*

$$(V, \rho) : \text{CM}(X_{H_0})_{\geq s_0} \longrightarrow \mathcal{V} \times \mathbf{Q}^\times$$

satisfying the following property:

For every divisor $D \in \text{Div}(\mathcal{X})_L$, there exists a constant $s_D \geq s_0$ depending only on the support of D , such that if $z \in X_{H_0}$ is a CM point of conductor $s \geq s_D$, then $(\bar{z} \cdot D)$ may be computed as follows. Let $V = V(z)$, $\rho = \rho(z)$, and write

$$D = c \mathcal{X}_\kappa + D'$$

with $c \in L$ and $D' \in \text{Div}(\mathcal{X})_L$ a divisor whose support does not contain V . Then

$$(\bar{z} \cdot D) = c[\kappa(y) : \kappa] q_{F,v}^s + \rho(V \bullet D), \tag{4.3.2}$$

where $y \in \mathcal{X}_\kappa$ denotes the reduction of z modulo \bar{w} .

Remark 4.3.4. The vertical component $V(z)$ will be characterized as the one maximizing the intersection multiplicity with \bar{z} . We refer the reader to [11, §2, Figure 1 in §1] for an equivalent and possibly more vivid geometric description⁸ of the relation of V to z_s : one can define pairwise disjoint open subsets (‘geometric basins’) of the Berkovich analytification of X , labelled by the irreducible components of the special fiber; then z_s belongs to the basin corresponding to V .

Proof. We will omit all subscripts v, w, \bar{w} and use some of the notation introduced in the proof of Lemma 4.3.1.

⁸Note however that the substantial results of [11] hold for the curve X over the field F_v .

Let $y \in \mathcal{X}_\kappa$ be a closed point, and write $[\mathcal{X}_{y,\kappa}] = \sum_{V' \in \mathcal{Y}_y} e_{V'} V'$ as divisors. By Lemma 4.3.1, the weighted sum

$$\sum_{V'} e_{V'} (\bar{z}_s \cdot V') = (\bar{z}_s \cdot [\mathcal{X}_{y,\kappa}]) = [\kappa(y) : \kappa] q_F^s \tag{4.3.3}$$

is independent of the choice of a CM point $z_s \in X_{H_0}$ of pseudo-conductor s reducing to y . As equation (4.3.3) goes to infinity with s (and the coefficients $e_{V'}$ are independent of s), the quantity

$$\max_{V' \in \mathcal{Y}} (\bar{z}_s \cdot V') \tag{4.3.4}$$

(more precisely, the minimum of those maxima as z_s varies in $\text{CM}(X_{H_0})_s$) goes to infinity with s .

Fix now a CM point $z_s \in X_{H_0}$ of pseudo-conductor s , let $y \in \mathcal{X}_\kappa$ be its reduction, and let $V \in \mathcal{Y}_y$ be a vertical component realizing the maximum in expression (4.3.4). Let $D \neq V$ be an irreducible divisor in \mathcal{X}_y . Pick any irreducible horizontal divisor $\Delta \neq D, \bar{z}_s$ in \mathcal{X}_y , and consider the ultrametric distance d_Δ of Proposition 4.2.1 for $R = \mathcal{O}_{\mathcal{X},y}$. (Note that Δ may be drawn from a finite set independent of z_s and D ; in fact, we may fix any set $\underline{\Delta}$ of at least two irreducible horizontal divisors that are not Zariski closures of CM points, and for given D pick any $\Delta \in \underline{\Delta} - \{D\}$.)

By the choice of V and Lemma 4.3.2, if s is sufficiently large (a condition depending on D), we have

$$d_\Delta(\bar{z}_s, V) = \frac{(\bar{z}_s \cdot \Delta)(V \cdot \Delta)}{(\bar{z}_s \cdot V)} < d_\Delta(V, D),$$

so that by Proposition 4.2.1,

$$d_\Delta(\bar{z}_s, D) = d_\Delta(V, D).$$

Unwinding the definitions,

$$(\bar{z}_s \cdot D) = \rho(V \cdot D) \tag{4.3.5}$$

for $\rho := (\bar{z}_s \cdot \Delta)/(V \cdot \Delta)$. Applied to a vertical component $D = V' \neq V$, formula (4.3.5) together with Lemma 4.3.2 shows the uniqueness of the maximizing $V =: V(z_s)$ for large s ; it is clear that $\rho =: \rho(z_s)$ is then uniquely determined as well. Now the intersection formula (4.3.2) follows by linearity from equation (4.3.5) and Lemma 4.3.1. \square

Corollary 4.3.5. *If $D \in \text{Div}^0(X_{H_0})_L$ is any degree-0 divisor, then for all sufficiently large s and all a ,*

$$m_{\bar{w}}(\tilde{Z}_{a\varpi^s}(\phi^\infty)[1]_U, D) = O(q_{F,v}^s)$$

in L , where the implied constant can be fixed independently of a and s .

Proof. Let \widehat{D} be a flat extension of D to a divisor on \mathcal{X} (with coefficients in L), and abbreviate $Z_{a,s} := \tilde{Z}_{a\varpi^s}(\phi^\infty)[1]_U$. Then by Propositions 4.1.4 and 4.3.3,

$$m_{\bar{w}}(Z_{a,s}, D) = (\bar{Z}_{a,s} \cdot \widehat{D}) = Aq_{F,v}^s + \sum_i \lambda_i (V_i \bullet \widehat{D})$$

for some vertical components $V_i \subset \mathcal{X}$ and some A , $\lambda_i \in L$. By the definition of flat extension, $(V_i \bullet \widehat{D}) = 0$ for all i . The constant A is a linear combination of the constants c in equation (4.3.2) (which depend only on \widehat{D}), with coefficients whose denominators are bounded by those of $Z_{a,s}$; by Proposition 4.1.4, the latter are bounded independently of a and s . \square

4.4. Decay of local heights

Recall that we need to prove (Proposition 3.6.3) that

$$T_{\mathfrak{t}_p}(\sigma^\vee)\widetilde{Z}(\phi^\infty, \chi)(v)$$

is a v -critical element of $\overline{\mathbf{S}}'$.

As in [9, §9.2, proof of Proposition 9.2.1], this is reduced to the following. For any $s \geq 0$, denote by $\langle \cdot, \cdot \rangle_{s, \overline{w}}$ the local height pairing on $X_{H_{s, \overline{w}}}$, which is valued in $H_{s, \overline{w}}^\times \widehat{\otimes} L$; and let

$$\langle \cdot, \cdot \rangle_{\overline{w}} = [H_{s, \overline{w}} : F_v]^{-1} \cdot N_{H_{s, \overline{w}}/F_v} \left(\langle \cdot, \cdot \rangle_{s, \overline{w}} \right),$$

which is valued in $F_v^\times \widehat{\otimes} L \subset \Gamma_F \widehat{\otimes} L$ and is compatible with varying s by [9, (4.1.6)]. Then we will show that for all $\overline{w} \mid v$ and all $a \in \mathbf{A}^{S_{1^\infty, \times}}$ with $v(a) = r_v$, we have

$$\begin{aligned} \left\langle \widetilde{Z}_{a\varpi^s}(\phi^\infty)[1]_{U, T_{\mathfrak{t}_p}}(\sigma^\vee)_{U^\dagger} t_\chi \right\rangle_{\overline{w}} &= O(q_{F,v}^s) \quad \text{in } F_v^\times \widehat{\otimes} L(\chi), \\ \left\langle \widetilde{Z}_{a\varpi^s}(\phi^\infty)[1]_{U, T_{\mathfrak{t}_p}}(\sigma^\vee)_{U^\dagger} t_\chi \right\rangle_{0, \overline{w}} &= O(q_{F,v}^s) \quad \text{in } H_{0, \overline{w}}^\times \widehat{\otimes} L(\chi), \end{aligned} \tag{4.4.1}$$

where the second statement implies the first one. Until Lemma 4.4.3, the argument follows the lines of previous works [9, 24, 27].

The norm relation and heights. Denote by N_s the norm from $H_{s, \overline{w}}$ to $H_{0, \overline{w}}$, set $L' := L(\chi)$, and let $\mathfrak{p}' \subset \mathcal{O}_{L'}$ be the maximal ideal. By the norm relation of Proposition 4.1.4, the aforementioned compatibility [9, (4.1.6)], and the integrality result of [9, Proposition 4.3.2],⁹

$$\begin{aligned} \left\langle \widetilde{Z}_{a\varpi^s}(\phi^\infty)[1]_{U, T_{\mathfrak{t}_p}}(\sigma^\vee)_{U^\dagger} t_\chi \right\rangle_{0, \overline{w}} &= \langle \text{Tr}_{H_s/H_0}(D_{a\varpi^s}), T_{\mathfrak{t}_p}(\sigma^\vee)_{U^\dagger} t_\chi \rangle_{0, \overline{w}} \\ &= N_s \left(\langle D_{a\varpi^s}, T_{\mathfrak{t}_p}(\sigma^\vee)_{U^\dagger} t_\chi \rangle_{s, \overline{w}} \right) \\ &\in \mathfrak{p}'^{-(d_{00} + d_0 + d_{1,s} + d_{2,s})} N_s \left(H_{s, \overline{w}}^\times \widehat{\otimes} \mathcal{O}_{L'} \right) \end{aligned} \tag{4.4.2}$$

for some integers $d_{i,(s)} \geq 0$ that we now define and study.

Boundedness of denominators. Set

$$V' := \pi_{A^\vee}^U \otimes_M V_{\mathfrak{p}} A^\vee(\chi^{-1}) \subset V := V_p J_U \otimes_{\mathbf{Q}_p} L'$$

⁹When comparing with the similar argument of [9, §9.2, proof of Proposition 9.2.1], our field H_s should be assimilated to the H'_s there.

considered as \mathcal{G}_{E_w} -modules; let V'' be its direct complement in the decomposition of V in [9, (9.2.4)], and let $0 \rightarrow V'^+ \rightarrow V' \rightarrow V'^- \rightarrow 0$ be the ordinary filtration analogous to formula (1.2.6). If $? \in \{', '' , '+\}$, define $T^? := T_p J_U \otimes_{\mathbf{Z}_p} \mathcal{O}_{L'} \cap V^?$, and let $T'^- = T'/T'^+$. Then the integers $d_{i,(s)}$ are defined as follows:

- d_{00} accounts for the denominators of the divisors, and it can be taken to be independent of s by Proposition 4.1.4.
- d_0 is such that $\mathfrak{p}^{d_0} T \subset T' \oplus T''$.
- $d_{1,s} := \text{length}_{\mathcal{O}_{L'}} H^1(H_{s,\bar{w}}, T''^*(1))_{\text{tors}}$.
- $d_{2,s} := \text{length}_{\mathcal{O}_{L'}} H_f^1(H_{s,\bar{w}}, T')/N_\infty H_f^1(H_{s,\bar{w}}, T')$, where N_∞ denotes the universal norms ([23, §6], [9, §4.3]) with respect to the infinite abelian extension of $H_{s,\bar{w}}$ cut out by the closure in $\mathcal{G}_{H_{s,\bar{w}}}^{\text{ab}} \supset H_{s,\bar{w}}^\times$ of

$$\text{Ker} \left[H_{s,\bar{w}}^\times \xrightarrow{N_{H_{s,\bar{w}}/F_v}} F_v^\times \rightarrow \Gamma_F \hat{\otimes} L \right].$$

Proposition 4.4.1. *Suppose that $V_p A$ is potentially crystalline as a representation of \mathcal{G}_{F_v} ; then the sequences of integers $(d_{1,s})$ and $(d_{2,s})$ are bounded.*

We will use the following vanishing result, in which \bar{L} denotes an algebraic closure of L' :

Lemma 4.4.2. *Set $\Gamma_\infty := \text{Gal}(H_{\infty,\bar{w}}/E_w) \cong E^\times \backslash E_{\mathbf{A}^\infty}^\times / U^v U_{F,v}^\circ$. For all Hodge–Tate characters $\psi: \mathcal{G}_{E_w} \rightarrow \bar{L}^\times$ factoring through Γ_∞ , and for any*

$$V^? \in \{V, V''^*(1), V'^+, *(1), V'^-\},$$

we have

$$H^0(E_w, V^?(\psi)) = 0.$$

Proof. The proof is largely similar to that of [9, Lemma 9.2.4], to which we refer for the background on the p -adic Hodge theory of characters.

We have

$$H^0(E_w, V^?) = \mathbf{D}_{\text{crys}}(V^?(\psi))^{\varphi=1},$$

where φ is the crystalline Frobenius, and it suffices to prove that $\mathbf{D}_{\text{crys}}(V^?(\psi))^{\varphi^d=1} = 0$ for $d = [E_w : E_{w,0}]$, where $E_{w,0}$ is the maximal unramified extension of \mathbf{Q}_p contained in E_w . As V' has been assumed potentially crystalline, it is pure of weight -1 , hence so are all the subquotients of V' and $V'^*(1)$. In particular, φ^d acts with negative weights on $\mathbf{D}_{\text{crys}}(V^?)$ for $V^? = V'^+, *(1), V'^-$; by [22, Theorem 5.3], this last assertion is also true of $V^? = V \cong V^*(1)$ and its subquotients such as $V^? = V''^*(1)$. Therefore, it suffices to show that φ^d acts with weight 0 on $\mathbf{D}_{\text{crys}}(\psi^m)$ for m such that ψ^m is crystalline.

Since ψ is trivial on U_F° , the Hodge–Tate weights $(n_\tau)_{\tau \in \text{Hom}(E_w, \bar{L})}$ satisfy $n_\tau + n_{\tau c} = 0$, where c is the complex conjugation of E_w/F_v . The action of φ^d on $\mathbf{D}_{\text{crys}}(\psi^m)$ is by

$$\psi \circ \text{rec}_{E,w}(\varpi_w)^{-m} \cdot \prod_{\tau \in \text{Hom}(E_w, \bar{L})} \varpi_w^{mn_\tau}, \tag{4.4.3}$$

where $\varpi_w \in E_w$ is any uniformizer. Choose ϖ_w so that $\varpi_w^{e(E_w/F_v)} = \varpi_v$ is a uniformizer in F_v . Then $\varpi_w^c = \pm \varpi_w$, so that the second factor in expression (4.4.3) is ± 1 . On the other hand, the subgroup $F^\times \backslash F_{\mathbf{A}^\infty}^\times / U_{F,v}^\circ (U^v \cap F_{\mathbf{A}^\infty}^\times) \subset \Gamma_\infty$ is finite, hence ϖ_v and ϖ_w have finite order in Γ_∞ . It follows that the first factor in expression (4.4.3) is a root of unity too, hence φ^d acts with weight 0 on $\mathbf{D}_{\text{crys}}(\psi^m)$. \square

Proof of Proposition 4.4.1. By the long exact sequence attached to

$$0 \rightarrow T''^*(1) \rightarrow V''^*(1) \rightarrow T''^*(1) \otimes L' / \mathcal{O}_{L'} \rightarrow 0$$

and the vanishing of $H^0(H_{s,\bar{w}}, V''^*(1))$ (which follows from Lemma 4.4.2), we have

$$\begin{aligned} H^1(H_{s,\bar{w}}, T''^*(1))_{\text{tors}} &\cong H^0(H_{s,\bar{w}}, T''^*(1) \otimes_{\mathcal{O}_{L'}} L' / \mathcal{O}_{L'}) \\ &= H^0(E_w, T''^*(1) \otimes_{\mathcal{O}_L} \mathcal{O}_L[\text{Gal}(H_{s,\bar{w}}/E_w)]) \otimes_{\mathcal{O}_L} L' / \mathcal{O}_{L'}. \end{aligned}$$

By [23, Theorems 6.6, 6.9] (or strictly speaking, a slightly generalized form thereof which still holds true by the arguments in [9, proof of Proposition 4.3.2]) and the vanishing of $H^0(H_{s,\bar{w}}, V'^{+,*}(1) \oplus V'^-)$ (which follows from Lemma 4.4.2), we have

$$\begin{aligned} d_{2,s} &\leq \text{length}_{\mathcal{O}_{L'}} H^0(H_{s,\bar{w}}, T'^{+,*}(1) \otimes_{\mathcal{O}_{L'}} L' / \mathcal{O}_{L'}) + \text{length}_{\mathcal{O}_{L'}} H^0(H_{s,\bar{w}}, T'^- \otimes_{\mathcal{O}_{L'}} L' / \mathcal{O}_{L'}) \\ &= \text{length}_{\mathcal{O}_{L'}} H^0(E_w, (T'^{+,*}(1) \oplus T'^-) \otimes_{\mathcal{O}_L} \mathcal{O}_L[\text{Gal}(H_{s,\bar{w}}/E_w)]) \otimes_{\mathcal{O}_L} L' / \mathcal{O}_{L'}. \end{aligned}$$

Then the boundedness of $d_{1,s}$ and $d_{2,s}$ follows as in [27, proof of Proposition 8.10] from the vanishing of

$$H^0(H_{\infty,\bar{w}}, V^?) \subset \bigoplus_{\psi: \Gamma_\infty \rightarrow \bar{L}^\times \text{ Hodge–Tate}} H^0(E_w, V^?(\psi))$$

for $V^? \in \{V''^*(1), V'^{+,*}(1), V'^-\}$, which is a consequence of Lemma 4.4.2. \square \square

Completion of the proofs. We are ready to reduce our decay statement for local heights to the decay statement for intersection multiplicities proved in §4.3.

Lemma 4.4.3. *For all $s' \leq s$, the restriction of the \bar{w} -adic valuation yields an isomorphism of $\mathcal{O}_{L'}$ -modules*

$$\bar{w}: N_s \left(H_{s,\bar{w}}^\times \hat{\otimes} \mathcal{O}_{L'} \right) / q^{s'} \cdot \left(H_{0,\bar{w}}^\times \hat{\otimes} \mathcal{O}_{L'} \right) \rightarrow \mathcal{O}_{L'} / q^{s'} \mathcal{O}_{L'}.$$

Proof. We drop all subscripts \bar{w} . Recall that the extension H_s/H_0 is totally ramified of degree q^s . Let $\varpi_s \in \mathcal{O}_{H_s}$ be a uniformizer; then $\omega_0 := N_s(\varpi_s)$ is a uniformizer of H_0 . For $* = 0, s$ we have the decompositions

$$H_*^\times \hat{\otimes} \mathcal{O}_{L'} = \mathcal{O}_{H_*}^\times \hat{\otimes} \mathcal{O}_{L'} \oplus \varpi_* \otimes \mathcal{O}_{L'}.$$

The map N_s respects the decompositions and, by local class field theory, has image

$$q^s \cdot (\mathcal{O}_{H_0}^\times \hat{\otimes} \mathcal{O}_{L'}) \oplus \varpi_0 \otimes \mathcal{O}_{L'}.$$

The valuation map annihilates the first summand and sends the second one isomorphically to $\mathcal{O}_{L'}$. The result follows. □

Proof of Proposition 3.6.3. By the comparison of the valuation component of local heights with arithmetic intersections in [9, Proposition 4.3.1], applied to the curve X_{U, H_0} , the image of the left-hand side of equation (4.4.2) under \bar{w} is

$$m \left(\tilde{Z}_{a\varpi^s}(\phi^\infty)_U[1], T_{\iota_p}(\sigma^\vee)_U^t t_\chi \right). \tag{4.4.4}$$

By Corollary 4.3.4 applied to $D = T_{\iota_p}(\sigma^\vee)_U^t t_\chi$, the right-hand side of expression (4.4.4) is $O(q_{F,v}^s)$. By equation (4.4.2), Proposition 4.4.1, and Lemma 4.4.3, we deduce the desired decay statement (4.4.1). □

Summary We have just completed the proof of Proposition 3.6.3. It implies Proposition 3.6.2, which together with Theorem 3.6.1 implies the kernel identity of Theorem 3.4.3. By Lemma 3.4.4, that implies Theorem 2.2.1, which is in turn an equivalent form of Theorem B by Lemma 2.2.2.

Appendix A. Local integrals

Throughout this appendix, v denotes a place of F above p unless specified otherwise. We use some of the notation introduced in §3.1, in particular the Weil representation r (see [9, §3.1] or [29] for the formulas defining it).

A.1. Interpolation factors

We relate the interpolation factors of the p -adic L -function of this paper with those from [9].

Lemma A.1.1. *Let $\xi: E_w^\times \rightarrow \mathbf{C}^\times$ and $\psi: E_w \rightarrow \mathbf{C}^\times$ be characters, with $\psi \neq \mathbf{1}$. Let dt be a Haar measure on E_w^\times . Then*

$$\int_{E_w^\times} \xi(t)\psi(t)dt = \frac{dt}{d_\psi t} \cdot \xi(-1) \cdot \gamma(\xi, \psi)^{-1}.$$

The left-hand side is to be understood in the sense of analytic continuation from characters $\xi|\cdot|^s$ for $\Re(s) \gg 0$.

Proof. We may fix $dt = d_\psi t$. Then the result follows from the functional equation for GL_1 [3, (23.4.4)]:

$$Z(\phi, \xi) = \gamma(\xi, \psi)^{-1} Z(\hat{\phi}, \xi^{-1} | |), \tag{A.1.1}$$

where for a Schwartz function ϕ on E_w ,

$$Z(\phi, \xi) := \int_{E_w^\times} \phi(t)\xi(t)d_\psi t, \quad \hat{\phi}(t) := \int_{E_w} \phi(x)\psi(xt)d_\psi x.$$

Namely, we insert in equation (A.1.1) the function

$$\hat{\phi} := \delta_{-1+\varpi_v^n \mathcal{O}_F} := \text{vol}(1 + \varpi_v^n \mathcal{O}_{F,v}, d_\psi t)^{-1} \mathbf{1}_{-1+\varpi_v^n \mathcal{O}_{F,v}}, \quad n \geq 1,$$

approximating a delta function at $t = -1$. Then by Fourier inversion, $\phi(t) = \hat{\phi}(-t) = \delta_{1+\varpi_v^n \mathcal{O}_F} * \psi(t)$, which if n is sufficiently large (depending on the conductor of ξ) has the same integral against ξ as $\psi(t)$. \square

Proposition A.1.2. *The ratio $C(\chi'_p)$ defined in formula (2.1.3) is a constant $C \in L$ independent of χ'_p .*

Proof. By the definition of $e_p(V_{(A, \chi')})$ and a comparison of [9, Lemma A.1.1] with Lemma A.1.1 applied to $\prod_{w|v} \chi'_w \alpha_v | \cdot |_v \circ q_w$, we have

$$C(\chi'_p) = \prod_{v|p} \gamma(\text{ad}(W_v(1)^{++}), \psi_v)^{-1} \in L.$$

\square

A.2. Toric period at p

We compare the toric period at a p -adic place with the interpolation factor. Denote by $P_v \subset \text{GL}_2(F_v)$ the upper triangular Borel subgroup.

Lemma A.2.1. *The quotient space $K_1^1(\varpi^{r'}) \backslash \text{GL}_2(F_v) / P_v$ admits the set of representatives*

$$n^-(c) := \begin{cases} \begin{pmatrix} 1 & \\ c & 1 \end{pmatrix} & \text{if } c \neq \infty, \\ \begin{pmatrix} 1 & \\ -1 & 1 \end{pmatrix} & \text{if } c = \infty, \end{cases} \quad c \in \mathcal{O}_{F,v} / \varpi^{r'} \mathcal{O}_{F,v} \cup \{\infty\}.$$

Proposition A.2.2. *Let $\chi \in \mathcal{A}^{1.c.}$ be a finite-order character, let r be sufficiently large (that is, satisfying the v -component of Assumption 3.4.2), let W_v be as in formula (3.1.1), and let $\phi_v = \phi_{v,r}$ be as in formula (3.1.4).*

Let $\pi_v = \sigma_v$ and let $(\ , \)_v : \pi_v \times \pi_v^\vee \rightarrow L$ be a duality pairing satisfying the compatibility of [9, (5.1.2)] with the local Shimizu lift.¹⁰ Finally, let $Z_v^\circ(\alpha_v, \chi_v)$ be the interpolation factor of the p -adic L -function of [9, Theorem A].

Then for all sufficiently large $r' > r$,

$$Q_{(\cdot), v, d^\circ t_v} \left(\theta_v \left(W_v, \alpha | \cdot |_v (\varpi_v)^{-r'} w_{r',v}^{-1} \phi_v \right), \chi_v \right) = |d|_v^2 |D|_v \cdot L(1, \eta_v)^{-1} \cdot Z_v^\circ(\alpha_v, \chi_v),$$

where $Q_{(\cdot), v, d^\circ t_v}$ uses the measure $d^\circ t_v = |d|_v^{-1/2} |D|_v^{-1/2} dt$.

¹⁰There, the pairing $(\ , \)_v$ is denoted by \mathcal{F}_v .

Proof. We drop all subscripts v , and assume as usual that ψ is our fixed character of level d^{-1} . Let

$$Q_{(v)}^\sharp(f_1, f_2, \chi_v) = \int_{E^\times/F^\times} \chi(t)(\pi(t)f_1, f_2) dt, \tag{A.2.1}$$

where dt is the usual Haar measure on E^\times/F^\times , giving volume $|d|^{1/2}|D|^{1/2}$ to $\mathcal{O}_E^\times/\mathcal{O}_F^\times$.

By the definitions and [9, Lemma A.1.1] (which expresses Z° as a normalized integral), it suffices to show that

$$Q^\sharp\left(\theta\left(W, \alpha|\cdot|(\varpi)^{-r'} w_{r'}^{-1} \phi\right), \chi\right) = |d|^{3/2}|D| \cdot L(1, \eta)^{-1} \cdot \int_{E^\times} \alpha|\cdot| \circ q(t) \chi(t) \psi_E(t) dt.$$

By [9, Lemma 5.1.1] (which spells out a consequence of the normalization of the local Shimizu lifting) and Lemma A.2.1, we can write

$$Q^\sharp := Q\left(\theta\left(W, \alpha|\cdot|(\varpi)^{-r'} w_{r'}^{-1} \phi_r\right), \chi_v\right) = \sum_{c \in \mathbf{P}^1(\mathcal{O}_F/\varpi^{r'})} Q^{\sharp(c)},$$

where for each c ,

$$Q^{\sharp(c)} := |d|^{-3/2} \cdot \alpha|\cdot|(\varpi)^{-r} \int_{F^\times} W\left(\begin{pmatrix} y & \\ & 1 \end{pmatrix} n^-(c)\right) \int_{T(F)} \chi(t) \int_{P(\varpi^{r'}) \backslash K_1^1(\varpi^{r'})} |y|r \left(n^-(c)kw_r^{-1}\right) \phi\left(yt^{-1}, y^{-1}q(t)\right) dkdt \frac{d^\times y}{|y|}.$$

Here $P(\varpi^{r'}) = P \cap K_1^1(\varpi^{r'})$.

It is easy to see that $Q^{\sharp(\infty)} = 0$ (observe that $\phi_{2,r}(0) = 0$). For $c \neq \infty$, we have

$$n^-(c)w_{r'}^{-1} = w_{r'}^{-1} \begin{pmatrix} 1 & -c\varpi^{-r'} \\ & 1 \end{pmatrix},$$

and when $x = (x_1, x_2)$ with $x_2 = 0$,

$$r\left(n^-(c)w_{r'}^{-1}\right)\phi(x, u) = \int_{\mathbf{V}} \psi_E(ux_1\xi_1)\psi(-ucq(\xi))\phi_r\left(\xi, \varpi^{r'}u\right) d\xi.$$

On the support of the integrand, we have $v(u) = \varpi^{-r'}$ and $v(q(\xi)) \geq r$, by the definition of ϕ_r . If $v(c) < r' - r - v(d)$, the integration in $d\xi_2$ gives 0; hence $Q^{\sharp(c)} = 0$ in that case.

Suppose from now on that $v(c) \geq r' - r - v(d)$. Then $\psi(-ucq(\xi)) = 1$ and

$$r\left(n^-(c)w_{r'}^{-1}\right)\phi(x, u) = |d|^3|D|L(1, \eta)^{-1}|\varpi|^r \psi_{E,r}\left(\varpi^{-r'}x_1\right) \delta_{1, U_{F,r}(\varpi^{r'}u)},$$

where $\psi_{E,r} := \text{vol}(\mathcal{O}_E)^{-1} \cdot \delta_{1, U_{T,r}} * \psi_E$, and we have noted that $\hat{\phi}_{2,r}(0) = e^{-1}|d| \cdot \text{vol}(q^{-1}(-1 + \varpi^r \mathcal{O}_F) \cap \mathcal{O}_{\mathbf{V}_2}) = |\varpi|^r |d|^3 |D|^{1/2} L(1, \eta)^{-1}$.

If r' is sufficiently large, the Whittaker function W is invariant under $n^-(c)$. Then

$$Q^\sharp(c) = |d|^{3/2} |D| L(1, \eta)^{-1} \cdot \text{Ivert} \varpi^{r-r'} \alpha(\varpi)^{-r'} \cdot |d|^{1/2} \zeta_{F,v}(1)^{-1} \cdot \int_{F^\times} W\left(\begin{pmatrix} y & \\ & 1 \end{pmatrix}\right) \int_{T(F)} \chi(t) \psi_{E,r}(\varpi^{-r'} x_1) \delta_{1, U_{F,r}}(\varpi^{r'} y^{-1} q(t)) dt d^\times y,$$

where $|\varpi|^{-r'} |d|^{1/2} \zeta_{F,v}(1)^{-1}$ appears as $\text{vol}(P(\varpi^{r'}) \backslash K_1^1(\varpi^{r'}))$.

Integrating in $d^\times y$ and summing the foregoing over the $q_F^{r+v(d)} = |d|^{-1} |\varpi|^{-r}$ contributing values of c , we find

$$Q^\sharp = |d|^{3/2} |D| \cdot L(1, \eta)^{-1} \cdot \int_{E^\times} \chi(t) \alpha | \circ q(t) \psi_E(t) dt,$$

as desired. □

Corollary A.2.3. *If χ_p is not exceptional – that is, $e_p(V_{(A,\chi)}) \neq 0$ – then the quaternion algebra \mathbf{B} over \mathbf{A} satisfying $\mathbf{H}(\pi_{\mathbf{B}}, \chi) \neq 0$ is indefinite at all primes $v \mid p$.*

Proof. By Propositions A.2.2 and A.1.2, if χ_p is not exceptional, then for all $v \mid p$ the functional $Q_v \in \mathbf{H}(\pi_{M_2(F_v)}, \chi_v) \otimes \mathbf{H}(\pi_{M_2(F_v)}^\vee, \chi_v^{-1})$ is not identically zero. □

Appendix B. Errata to [9]

The salient mistakes are the following: the statement of the main theorem is off by a factor of 2; the proof given needs a further assumption, (no stronger than) that $V_p A$ is potentially crystalline at all $v \mid p$ (however, the theorem still holds true without the assumption; compare Remark 1.2.1); and the Schwartz function $\phi_{2,p}$ given by the local Siegel–Weil formula at p needs to be *different* from the Schwartz function used to construct the Eisenstein family.

References in *italics* are directed to [9], and references in roman letters to the present paper.

- *Theorem A.* It should be $L_{p,\alpha}(\sigma_E) \in \mathcal{O}(\mathcal{Y}')^b$ (with the interpolation property being correct for the choice of additive character ψ_p as in Theorem A). For a correct discussion of the ring of rationality of $L_{p,\alpha}(\sigma_E)$, within the context of a generalized construction, see [12, Corollary 4.5.4].
- *Theorem B.* The constant factor should be c_E and not $c_E/2$ (the latter is, according to (1.1.3), the constant factor of the Gross–Zagier formula in Archimedean coefficients).¹¹ The mistake is introduced in the proof of *Proposition 5.4.3* (see later).
The proof works under the further assumption that $V_p A$ is potentially crystalline at all $v \mid p$ (see the correction to *Proposition 9.2.1*).
- *Theorem C.* Similarly, the constant factor should be $c_E/2$ in part 3, and c_E in part 4.

¹¹The heuristic reason for the difference is that the direct analogue of $s \mapsto L(1/2 + s, \sigma_E, \chi)$ is $\chi_F \mapsto L_p(\sigma_E)(\chi \cdot \chi_F \circ N_{E/F})$, whose derivative at $\chi_F = \mathbf{1}$ is twice our $d_F L_p(\sigma_E)(\chi)$, as the tangent map to $\chi_F \mapsto \chi' = \chi \cdot \chi_F \circ N_{E/F} \mapsto \omega^{-1} \cdot \chi'_{\mathbf{A}^\times}$ is multiplication by 2.

- §2.1. The space of p -adic modular forms is the closure of $M_2(K^p K_1^1(p^\infty)_p)$, not $M_2(K^p K^1(p^\infty)_p)$.
- Proposition 2.4.4.1. The multiplier in equation (2.4.3) should be $\alpha \cdot |(\varpi^{-r}) = \prod_{v|p} \alpha_v \cdot |_{\nu}(\varpi_v^{-r})$, and the statement holds for forms in $M_2(K^p K_1^1(p^r)_p)$. Similarly, the definition of $R_{r,v}^\circ$ in Proposition 3.5.1 should have an extra $|\varpi_v|^{-r}$. The result of Proposition A.2.2, as modified later, holds true for this definition of $R_{r,v}^\circ$ (there, a complementary mistake appears between the third-last and second-last displayed equations in the proof).
- Lemma 3.2.2. A factor $\eta(y)$ is missing in the right-hand side of the formula.
- Proposition 3.2.3.2. The proposition should be corrected as follows: *Let $v \mid p$ and let $\phi_{2,v} = \phi_{2,v}^\circ$ be as in formula (3.1.4) (of the present paper). Then*

$$W_{a,r,v}^\circ(1,u,\chi_F) = \begin{cases} |d_v|^{3/2} |D_v|^{1/2} \chi_{F,v}(-1) & \text{if } v(a) \geq 0 \text{ and } v(u) = 0, \\ 0 & \text{otherwise.} \end{cases}$$

- Equation (3.7.1). The right-hand side should have an extra factor of $\prod_{v|p} |d_v|^2 |D_v|$ owing to the correction to Proposition A.2.2.
- Lemma 5.3.1. The left-hand side of the last equation in the statement should be $\langle f'_1(P_1), f'_2(P_2) \rangle_{J,*}$.
- Proof of Proposition 5.4.3. The second-last displayed equation should have the factor of 2 on the right hand side, not the left hand side:

$$2\ell_{\varphi^p,\alpha}(\tilde{Z}(\phi^\infty,\chi)) = |D_F|^{1/2} |D_E|^{1/2} L(1,\eta) \langle T_{\text{alg},\iota_p}(\theta_{\iota_p}(\varphi,\alpha(\varpi)^{-r} w_r^{-1} \phi)) P_\chi, P_\chi^{-1} \rangle.$$

Then the argument shows that first,

$$\begin{aligned} & \langle T_{\text{alg},\iota_p}(f_1 \otimes f_2) P_\chi, P_\chi^{-1} \rangle_J \\ &= \frac{\zeta_F^\infty(2)}{(\pi^2/2)^{[F:\mathbf{Q}]}} |D_E|^{1/2} L(1,\eta) \prod_{v|p} Z_v^\circ(\alpha_v,\chi_v)^{-1} \cdot d_F L_{p,\alpha}(\sigma_{A,E})(\chi) \cdot Q(f_1, f_2, \chi) \end{aligned}$$

(without an incorrect factor of 2 introduced in the denominator of the right-hand side of (5.4.1) there); and second, that this equation is equivalent to *Theorem B* as corrected.

An extra factor $\prod_{v|p} |d_v|^2 |D_v|$ should be inserted in the right-hand sides of the last and fourth-last displayed equations; compare the corrections to (3.7.1) and Proposition A.3.1.

- Proposition 7.1.1(b). It should be replaced by Proposition 3.5.1(b) and (c).
- §7.2, third paragraph. The coefficient in the second displayed equation should have $|D_E|^{1/2}$, not $|D_{E/F}|^{1/2}$, in the denominator.
- Lemma 9.1.1. This is corrected by Lemma 4.1.1 (this does not significantly affect the rest).
- Lemma 9.1.5. The extension H_∞ is contained in a relative Lubin–Tate extension. This is the only property used.

- Proposition 9.2.1. The assumption that $V_p A$ is potentially crystalline at all $v \mid p$ should be added. The bounded dependence on s of the integer $d_2 = d_{2,s}$ was not addressed; it holds true by the proofs of Proposition 4.4.1 and Lemma 4.4.2, which work verbatim in the split case (under the comparison given in footnote 8 of §4.4). The definition of $d_{1,s}$ contains an extra $\otimes_{\mathcal{O}_L} L/\mathcal{O}_L$.
- Lemma 9.2.4. The statement should be that $H^0\left(\tilde{H}'_{\infty, \bar{w}}, V_p J_U^*(1)\right)$ vanishes, and it is this group that should appear in the left-hand side of the first displayed equation in the proof.
- Lemma A.2.1. The list of representatives is missing the element $n^-(\infty) = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$ (compare Lemma A.2.1).
- Proposition A.2.2. The statement should be

$$R_{r,v}^{\natural}(W_v, \phi_v, \chi_v^{\iota}) = |d_v^2|D|_v \cdot Z_v^{\circ}(\chi_v^{\iota}) := |d_v^2|D|_v \cdot \frac{\zeta_{F,v}(2)L(1, \eta_v)^2}{L\left(1/2, \sigma_{E,v}^{\iota}, \chi_v^{\iota}\right)} \prod_{w|v} Z_w(\chi_w^{\iota}).$$

The factor $|d_v^2|D|_v$ missing from [9] should first appear in the right-hand side of the displayed formula for $r(w_r^{-1})\phi(x, u)$ in the middle of the proof.

- Proposition A.3.1. It should be replaced by Proposition A.2.2.

Acknowledgments. I would like to thank the referees for a sharp reading. This research was supported by ISF grant 1963/20.

Competing Interests. None

References

- [1] K. BARRÉ-SIRIEIX, G. DIAZ, F. GRAMAIN AND G. PHILIBERT, Une preuve de la conjecture de Mahler-Manin, *Invent. Math.* **124**(1-3) (1996), 1–9.
- [2] M. BERTOLINI AND H. DARMON, The p -adic L -functions of modular elliptic curves, in *Mathematics Unlimited—2001 and Beyond*, pp. 109–170 (Springer, Berlin, 2001).
- [3] C. J. BUSHNELL AND G. HENNIART, *The Local Langlands Conjecture for $GL(2)$* , Grundlehren der Mathematischen Wissenschaften, **335** (Springer-Verlag, Berlin, 2006).
- [4] K. BÜYÜKBODUK, R. POLLACK AND S. SASAKI, ‘ p -adic Gross-Zagier formula at critical slope and a conjecture of Perrin-Riou, Preprint, 2021, <https://arxiv.org/pdf/1811.08216.pdf>.
- [5] L. CAI, J. SHU AND Y. TIAN, Explicit Gross-Zagier and Waldspurger formulae, *Algebra Number Theory* **8**(10) (2014), 2523–2572.
- [6] H. CARAYOL, Sur la mauvaise réduction des courbes de Shimura, *Compos. Math.* **59**(2) (1986), 151–230.
- [7] J. COATES, Motivic p -adic L -functions, in *L -Functions and Arithmetic (Durham, 1989)*, London Mathematical Society Lecture Note Series, **153**, pp. 141–172 (Cambridge University Press, Cambridge, UK, 1991).
- [8] D. DISEGNI, p -adic heights of Heegner points on Shimura curves, *Algebra Number Theory* **9**(7) (2015), 1571–1646.
- [9] D. DISEGNI, The p -adic Gross–Zagier formula on Shimura curves, *Compos. Math.* **153**(10) (2017), 1987–2074.

- [10] D. DISEGNI, On the p -adic Birch and Swinnerton-Dyer conjecture for elliptic curves over number fields, *Kyoto J. Math.* **60**(2) (2020), 473–510.
- [11] D. DISEGNI, ‘ p -adic equidistribution of CM points’, Preprint, 2021, <https://disegni-daniel.perso.math.cnrs.fr/equi.pdf>.
- [12] D. DISEGNI, ‘ p -adic L -functions for $\mathrm{GL}_2 \times \mathrm{GU}(1)$ ’, Preprint, 2021, <https://disegni-daniel.perso.math.cnrs.fr/pLF-GL2.pdf>.
- [13] D. DISEGNI, ‘The universal p -adic Gross–Zagier formula’, Preprint, 2021, <https://disegni-daniel.perso.math.cnrs.fr/univ.pdf>.
- [14] J.-M. FONTAINE, Représentations l -adiques potentiellement semi-stables, *Astérisque* **223** (1994), 321–347.
- [15] E. R. GARCÍA BARROSO, P. D. GONZÁLEZ PÉREZ AND P. POPESCU-PAMPU, Ultrametric spaces of branches on arborescent singularities, in *Singularities, Algebraic Geometry, Commutative Algebra, and Related Topics*, pp. 55–106 (Springer, Cham, 2018).
- [16] R. GREENBERG AND G. STEVENS, p -adic L -functions and p -adic periods of modular forms, *Invent. Math.* **111**(2) (1993), 407–447.
- [17] B. H. GROSS AND D. B. ZAGIER, Heegner points and derivatives of L -series, *Invent. Math.* **84**(2) (1986), 225–320.
- [18] B. H. GROSS, On canonical and quasicanonical liftings, *Invent. Math.* **84**(2) (1986), 321–326.
- [19] S. KOBAYASHI, The p -adic Gross–Zagier formula for elliptic curves at supersingular primes, *Invent. Math.* **191**(3) (2013), 527–629.
- [20] Q. LIU, *Algebraic Geometry and Arithmetic Curves*, Oxford Graduate Texts in Mathematics, **6** (Oxford University Press, Oxford, UK, 2002).
- [21] B. MAZUR, J. TATE AND J. TEITELBAUM, On p -adic analogues of the conjectures of Birch and Swinnerton-Dyer, *Invent. Math.* **84**(1) (1986), 1–48.
- [22] A. MOKRANE, La suite spectrale des poids en cohomologie de Hyodo-Kato, *Duke Math. J.* **72**(2) (1993), 301–337.
- [23] J. NEKOVARŔ, On p -adic height pairings, in *Séminaire de Théorie des Nombres, Paris, 1990–91*, Progress in Mathematics, **108**, pp. 127–202 (Birkhäuser Boston, Boston, 1993).
- [24] J. NEKOVARŔ, On the p -adic height of Heegner cycles, *Math. Ann.* **302**(4) (1995), 609–686.
- [25] J. NEKOVARŔ, *Selmer Complexes*, *Astérisque*, **310** (2006), viii+559.
- [26] B. PERRIN-RIOU, Points de Heegner et dérivées de fonctions Lp -adiques, *Invent. Math.* **89**(3) (1987), 455–510.
- [27] A. SHNIDMAN, p -adic heights of generalized Heegner cycles, *Ann. Inst. Fourier (Grenoble)* **66**(3) (2016), 1117–1174.
- [28] J. TATE, Number theoretic background, *Automorphic Forms, Representations and L -Functions (Corvallis, Ore., 1977)*, Proceedings of Symposia in Pure Mathematics, **XXXIII**, pp. 3–26 (American Mathematical Society, Providence, RI, 1979).
- [29] X. YUAN, S.-W. ZHANG AND W. ZHANG, *The Gross–Zagier Formula on Shimura Curves*, Annals of Mathematics Studies, **184**, (Princeton University Press, Princeton, NJ, 2012).