



# Lie Derivatives and Ricci Tensor on Real Hypersurfaces in Complex Two-plane Grassmannians

Imsoon Jeong, Juan de Dios Pérez, Young Jin Suh, and Changhwa Woo

*Abstract.* On a real hypersurface  $M$  in a complex two-plane Grassmannian  $G_2(\mathbb{C}^{m+2})$  we have the Lie derivation  $\mathcal{L}$  and a differential operator of order one associated with the generalized Tanaka–Webster connection  $\tilde{\mathcal{L}}^{(k)}$ . We give a classification of real hypersurfaces  $M$  on  $G_2(\mathbb{C}^{m+2})$  satisfying  $\tilde{\mathcal{L}}_\xi^{(k)} S = \mathcal{L}_\xi S$ , where  $\xi$  is the Reeb vector field on  $M$  and  $S$  the Ricci tensor of  $M$ .

## 1 Introduction

It is one of the most classical and interesting parts in differential geometry to find geometric properties of submanifolds on a symmetric space equipped with a Kähler structure  $J$ , *i.e.*, a Hermitian symmetric space. Among Hermitian symmetric spaces as a higher rank space of complex projective space  $P_n(\mathbb{C})$ , the authors have investigated the complex two-plane Grassmannian  $G_2(\mathbb{C}^{m+2})$ , which consists of the set of all complex two-dimensional linear subspaces in  $\mathbb{C}^{m+2}$ . The space  $G_2(\mathbb{C}^{m+2})$  is diffeomorphic to the homogeneous space  $SU_{m+2}/S(U_2 \cdot U_m)$ , the special unitary group  $SU_{m+2}$  acts transitively on  $\mathbb{C}^{m+2}$ , and  $S(U_2 \cdot U_m)$  means the isotropic subgroup of  $SU_{m+2}$ . Cartan decomposition of the Lie algebra of  $S(U_2 \cdot U_m)$  is expressed by  $\mathfrak{k} = \mathfrak{su}_2 \oplus \mathfrak{su}_m \oplus \mathfrak{u}_1$ . We have a Kähler structure  $J$  from  $\mathfrak{u}_1$ , the one-dimensional center of  $\mathfrak{k}$ . Remarkably, we also have a quaternionic Kähler structure  $\mathfrak{J}$  from  $\mathfrak{su}_2$  satisfying  $JJ_\nu = J_\nu J$  ( $\nu = 1, 2, 3$ ), where  $\{J_\nu\}_{\nu=1,2,3}$  is an orthonormal basis of  $\mathfrak{J}$ . When  $m = 1$ ,  $G_2(\mathbb{C}^3)$  is isometric to the two-dimensional complex projective space  $\mathbb{C}P^2$  with constant holomorphic sectional curvature eight. When  $m = 2$ , we note that the isomorphism  $\text{Spin}(6) \simeq \text{SU}(4)$  yields an isometry between  $G_2(\mathbb{C}^4)$  and the real Grassmann manifold  $G_2^+(\mathbb{R}^6)$  of oriented two-dimensional linear subspaces in  $\mathbb{R}^6$ . In this paper we assume  $m \geq 3$ .

To classify real hypersurfaces with certain geometric conditions, let us give an explanation of the geometry of real hypersurfaces on  $G_2(\mathbb{C}^{m+2})$ . Let us consider a real hypersurface  $M$  in  $G_2(\mathbb{C}^{m+2})$  and let  $N$  denote a local unit normal vector field on  $M$

Received by the editors March 18, 2017; revised July 31, 2017.

Published electronically June 8, 2018.

This work was supported by Grant Proj. No. NRF-2015-RIA2A1A-01002459 and Proj. No. NRF-2018-R1D1A1B-05040381. Author I. J. is supported by NRF Grant funded by the Korean Government Grant Proj. No. 2017RIA2B4005317. Author J. D. P. is partially supported by MINECO-FEDER Grant MTM2013-47828-C2-1-P. Author C. W. is supported by NRF Grant funded by the Korean Government Grant Proj. No. 2017R1C1B1010265.

AMS subject classification: 53C40, 53C15.

Keywords: real hypersurface, complex two-plane Grassmannian, Hopf hypersurface, shape operator, Ricci tensor, Lie derivation.

in  $G_2(\mathbb{C}^{m+2})$ . The *Reeb vector field*  $\xi = -JN \in T_pM$  at  $p \in M$  is induced from the Kähler structure  $J$ . Let  $\mathcal{C}$  be the distribution given by the orthogonal complement of  $[\xi]$  in  $T_pM$  at  $p \in M$ . If  $\xi$  is invariant under the shape operator  $A$ , it is said to be *Hopf*. The 1-dimensional foliation of  $M$  by the integral manifolds of the Reeb vector field  $\xi$  is said to be a *Hopf foliation* of  $M$ . We say that  $M$  is a *Hopf hypersurface* in  $G_2(\mathbb{C}^{m+2})$  if and only if the Hopf foliation of  $M$  is totally geodesic. It is the complex maximal subbundle of  $T_pM = \mathcal{C} \oplus \mathcal{C}^\perp$ . The real hypersurface  $M$  is said to be *Hopf* if  $A\mathcal{C} \subset \mathcal{C}$ , or equivalently, the Reeb vector field  $\xi$  is principal, where  $A$  is the shape operator of the real hypersurface  $M$ . If  $X$  is a tangent vector on  $M$ , we can put

$$JX = \phi X + \eta(X)N \quad \text{and} \quad J_\nu X = \phi_\nu X + \eta_\nu(X)N$$

where  $\phi X$  (resp.  $\phi_\nu X$ ) is the tangential part of  $JX$  (resp.  $J_\nu X$ ) and  $\eta(X) = g(X, \xi)$  (resp.  $\eta_\nu(X) = g(X, \xi_\nu)$ ) is the coefficient of normal part of  $JX$  (resp.  $J_\nu X$ ). In this case, we call  $\phi$  the structure tensor field of  $M$ . Using the Gauss and Weingarten formulas in [6, Section 1 and 2], the Kähler condition  $\bar{\nabla}J = 0$  gives  $\nabla_X \xi = \phi AX$  for any tangent vector field  $X$  on  $M$ , where  $\nabla$  (resp.  $\bar{\nabla}$ ) denotes the covariant derivative on  $M$  (resp.  $G_2(\mathbb{C}^{m+2})$ ). From this, it can be easily checked that  $M$  is Hopf if and only if the Reeb vector field  $\xi$  is Hopf.

In this case, the principal curvature  $\alpha = g(A\xi, \xi)$  is said to be a *Reeb curvature* of  $M$ .

From the quaternionic Kähler structure  $\mathfrak{J}$  of  $G_2(\mathbb{C}^{m+2})$ , there naturally exist *almost contact 3-structure* vector fields  $\{\xi_1, \xi_2, \xi_3\}$  defined by  $\xi_\nu = -J_\nu N$ ,  $\nu = 1, 2, 3$ . Now let us denote by  $\mathcal{Q}^\perp = \text{Span}\{\xi_1, \xi_2, \xi_3\}$  a 3-dimensional distribution in the tangent space  $T_pM$  at  $p \in M$ . In addition,  $\mathcal{Q}$  stands for the orthogonal complement of  $\mathcal{Q}^\perp$  in  $T_pM$ . Then it becomes a quaternionic maximal subbundle of  $T_pM$ . Thus, the tangent space of  $M$  consists of the direct sum of  $\mathcal{Q}$  and  $\mathcal{Q}^\perp$  as follows:  $T_pM = \mathcal{Q} \oplus \mathcal{Q}^\perp$ .

For two distributions  $\mathcal{C}^\perp$  and  $\mathcal{Q}^\perp$  defined above, we can consider two natural invariant geometric properties under the shape operator  $A$  of  $M$ , that is,  $A\mathcal{C}^\perp \subset \mathcal{C}^\perp$  and  $A\mathcal{Q}^\perp \subset \mathcal{Q}^\perp$ . The following theorem is from a paper due to Suh [13, Theorem 1.1].

**Theorem A** *Let  $M$  be a real hypersurface in  $G_2(\mathbb{C}^{m+2})$ ,  $m \geq 3$ . Then both  $[\xi]$  and  $\mathcal{Q}^\perp$  are invariant under the shape operator of  $M$  if and only if*

- (A)  *$M$  is an open part of a tube around a totally geodesic  $G_2(\mathbb{C}^{m+1})$  in  $G_2(\mathbb{C}^{m+2})$ , or*
- (B)  *$m$  is even, say  $m = 2n$ , and  $M$  is an open part of a tube around a totally geodesic  $\mathbb{H}P^n$  in  $G_2(\mathbb{C}^{m+2})$ .*

In the case of (A), we want to say  $M$  is of Type (A). Similarly, in the case of (B), we say  $M$  is of Type (B).

Until now, many geometers have investigated some characterizations of Hopf hypersurfaces in  $G_2(\mathbb{C}^{m+2})$  that satisfy commuting conditions involving geometric quantities like shape operator, structure (or normal) Jacobi operator, Ricci tensor, and so on. For a tangent vector  $X$ ,  $\phi X$  is the tangential part of  $JX$ ; then  $\phi$  is said to be the structure tensor field. Commuting Ricci means that the Ricci tensor  $S$  and the structure tensor field  $\phi$  commute with each other, that is,  $S\phi = \phi S$ . From such a point

of view, Suh [12] has given a characterization of real hypersurfaces of Type (A) with commuting Ricci tensor.

On the other hand, a Jacobi field along geodesics of a given Riemannian manifold  $(\bar{M}, \bar{g})$  is an important tool in the study of differential geometry. It satisfies a well-known differential equation that inspires Jacobi operators. It is defined by  $(\bar{R}_X(Y))(p) = (\bar{R}(Y, X)X)(p)$ , where  $\bar{R}$  denotes the curvature tensor of  $\bar{M}$  and  $X, Y$  denote any vector fields on  $\bar{M}$ . It is known to be a self-adjoint endomorphism on the tangent space  $T_p\bar{M}$ ,  $p \in \bar{M}$ . Clearly, each tangent vector field  $X$  to  $\bar{M}$  provides a Jacobi operator with respect to  $X$ . Thus, the Jacobi operator on a real hypersurface  $M$  of  $G_2(\mathbb{C}^{m+2})$  with respect to  $\xi$  (resp.  $N$ ) is said to be a *structure Jacobi operator* (resp. *normal Jacobi operator*) and will be denoted by  $R_\xi$  (resp.  $\bar{R}_N$ ).

Among many geometric conditions, in this paper we focus on commuting conditions that have a strong relationship with hypersurfaces of tube type when the Reeb vector field  $\xi$  belongs to  $\mathcal{Q}^\perp$ , that is to say, the commuting conditions between (1,1) type tensor fields on real hypersurfaces in complex two-plane Grassmannians  $G_2(\mathbb{C}^{m+2})$  are used to give same results to isometric Reeb flow.

For a commuting problem concerned with structure Jacobi operator  $R_\xi$  and structure tensor  $\phi$  of  $M$  in  $G_2(\mathbb{C}^{m+2})$ , that is,  $R_\xi\phi = \phi R_\xi$ , Suh and Yang [16] gave a characterization of a real hypersurface of Type (A) in  $G_2(\mathbb{C}^{m+2})$ . Also, concerned with a commuting problem for the normal Jacobi operator  $\bar{R}_N$ , Pérez, Jeong, and Suh [9] gave a characterization of a real hypersurface of Type (A) in  $G_2(\mathbb{C}^{m+2})$ .

Related to the Levi-Civita connection  $\nabla$ , Tanno [18] introduced the generalized Tanaka-Webster connection (GTW connection) for contact metric manifolds as a generalization of the Tanaka-Webster connection. It is defined as a canonical affine connection on a non-degenerate, pseudo-Hermitian CR-manifold (see [17,19]). Then the GTW connection coincides with Tanaka-Webster connection if the associated CR-structure is integrable. Cho defined the GTW connection for a real hypersurface in a Kähler manifold in such a way that

$$\widehat{\nabla}_X^{(k)} Y = \nabla_X Y + \widehat{F}_X^{(k)} Y,$$

where  $k(\in \mathbb{R} \setminus \{0\})$  denotes a non-zero constant and  $\widehat{F}_X^{(k)} Y$  is defined by

$$\widehat{F}_X^{(k)} Y = g(\phi AX, Y)\xi - \eta(Y)\phi AX - k\eta(X)\phi Y.$$

The skew-symmetric (1,1) type tensor  $\widehat{F}_X^{(k)}$  is said to be a *Tanaka-Webster* (or *k-th-Cho*) *operator* with respect to  $X$ . In particular, if the real hypersurface satisfies  $A\phi + \phi A = 2k\phi$ , then the GTW connection  $\widehat{\nabla}^{(k)}$  coincides with the Tanaka-Webster connection (see [1,2]).

On the other hand, we have considered real hypersurfaces in  $G_2(\mathbb{C}^{m+2})$  satisfying  $(\widehat{\mathcal{L}}_X^{(k)} T)Y = 0$  for any vector fields  $X$  and  $Y$  on  $M$  in  $G_2(\mathbb{C}^{m+2})$ , where  $\widehat{\mathcal{L}}^{(k)}$  is the differential operator of order one given by

$$\widehat{\mathcal{L}}_X^{(k)} Y = \widehat{\nabla}_X^{(k)} Y - \widehat{\nabla}_Y^{(k)} X$$

for any vector fields  $X$  and  $Y$  on  $M$ , where  $T$  denotes a tensor field of type (1,1).

The torsion of the GTW connection is given by

$$\widehat{\mathcal{T}}^{(k)}(X, Y) = \widehat{F}_X^{(k)}(Y) - \widehat{F}_Y^{(k)}(X).$$

The operator defined by  $\widehat{\mathcal{T}}_X^{(k)}(Y) = \widehat{\mathcal{T}}^{(k)}(X, Y)$  is called the *torsion operator associated with X*.

Let  $S$  be the Ricci tensor of  $M$ . We will consider real hypersurfaces  $M$  in  $G_2(\mathbb{C}^{m+2})$  satisfying

$$(C-1) \quad \widehat{\mathcal{L}}_X^{(k)}S = \mathcal{L}_X S,$$

for any vector field  $X$  on  $M$ . This is equivalent to the fact  $\widehat{\mathcal{T}}_X^{(k)}S = S\widehat{\mathcal{T}}_X^{(k)}$ , for any  $X$  tangent to  $M$ .

On the other hand, Hopf hypersurfaces  $M$  are those whose Reeb vector field  $\xi = -JN$  is Killing or, equivalently, a principal vector field, verifying  $A\xi = \alpha\xi$ , where the smooth function  $\alpha = g(A\xi, \xi)$  is said to be the *Reeb curvature* of the Reeb vector field  $\xi$ . Then we can give a classification for  $M$  in  $G_2(\mathbb{C}^{m+2})$  satisfying (C-1) in the particular case  $X = \xi$  as follows.

**Theorem 1.1** *Let  $M$  be a Hopf hypersurface in complex two-plane Grassmannians  $G_2(\mathbb{C}^{m+2})$ ,  $m \geq 3$ . The Ricci tensor  $S$  on  $M$  satisfies  $\widehat{\mathcal{L}}_\xi^{(k)}S = \mathcal{L}_\xi S$  if and only if  $M$  is locally congruent to an open part of a tube of some radius  $r \in (0, \frac{\pi}{2\sqrt{2}})$  around a totally geodesic  $G_2(\mathbb{C}^{m+1})$  in  $G_2(\mathbb{C}^{m+2})$ .*

In this case, there are two kinds of focal sets in  $G_2(\mathbb{C}^{m+2})$ , and the distance between them is  $\frac{\pi}{2\sqrt{2}}$ . By virtue of this Theorem, we give another non-existence property as follows.

**Corollary 1.2** *There does not exist any Hopf hypersurface  $M$  in  $G_2(\mathbb{C}^{m+2})$ ,  $m \geq 3$ , satisfying the condition  $\widehat{\mathcal{L}}_X^{(k)}S = \mathcal{L}_X S$  for any vector field  $X$  on  $M$ .*

In this paper, we refer to [6, 7, 11, 12, 14, 15] for Riemannian geometric structures of a complex two-plane Grassmannian  $G_2(\mathbb{C}^{m+2})$ ,  $m \geq 3$ .

## 2 Proof of Theorem

Let us introduce the Ricci tensor  $S$ , briefly. The curvature tensor  $R(X, Y)Z$  of  $M$  in  $G_2(\mathbb{C}^{m+2})$  can be derived from the curvature tensor  $\overline{R}(X, Y)Z$  of  $G_2(\mathbb{C}^{m+2})$ . Then by contracting and using the geometric structure  $JJ_\nu = J_\nu J$  ( $\nu = 1, 2, 3$ ), we can see the Ricci tensor  $S$  given by

$$g(SX, Y) = \sum_{i=1}^{4m-1} g(R(e_i, X)Y, e_i),$$

where  $\{e_1, \dots, e_{4m-1}\}$  denotes a basis of the tangent space  $T_p M$  of  $M$ ,  $p \in M$ , in  $G_2(\mathbb{C}^{m+2})$  (see [12]). From the definition of the Ricci tensor  $S$  and fundamental for-

mulas in [12, section 2], we have

$$\begin{aligned}
 (2.1) \quad SX &= \sum_{i=1}^{4m-1} R(X, e_i)e_i \\
 &= (4m + 7)X - 3\eta(X)\xi + hAX - A^2X \\
 &\quad + \sum_{v=1}^3 \{-3\eta_v(X)\xi_v + \eta_v(\xi)\phi_v\phi X - \eta_v(\phi X)\phi_v\xi - \eta(X)\eta_v(\xi)\xi_v\},
 \end{aligned}$$

where  $h$  denotes the trace of  $A$ , that is,  $h = \text{Tr}A$  (see [10, (1.4)]).

Using equation (2.1), we will prove that the Reeb vector field  $\xi$  of  $M$  belongs either to  $\mathcal{Q}$  or  $\mathcal{Q}^\perp$ . Under the condition of being Hopf, we get

$$(2.2) \quad \widehat{F}_\xi^{(k)}X = -k\phi X.$$

For  $X = \xi$  into (C-1), we have

$$(2.3) \quad \widehat{F}_\xi^{(k)}(SY) + \phi ASY - S\widehat{F}_\xi^{(k)}(Y) - S\phi AY = 0$$

for any  $Y$  tangent to  $M$ . Taking the inner product of (2.3) with  $Z$ , where  $Z$  denotes a vector field tangent to  $M$ , we get

$$g(\widehat{F}_\xi^{(k)}(SY), Z) + g(\phi ASY, Z) - g(S\widehat{F}_\xi^{(k)}(Y), Z) - g(S\phi AY, Z) = 0.$$

Bearing in mind that  $\widehat{F}_\xi^{(k)}$  is skew-symmetric and  $S$  is symmetric, we have

$$g(Y, -S\widehat{F}_\xi^{(k)}(Z) - SA\phi Z + \widehat{F}_\xi^{(k)}(SZ) + A\phi SZ) = 0.$$

Thus, we have  $-S\widehat{F}_\xi^{(k)}(Z)SA\phi Z + \widehat{F}_\xi^{(k)}(SZ) + A\phi SZ = 0$ , and, replacing  $Y$  by  $Z$ , we obtain

$$(2.4) \quad -S\widehat{F}_\xi^{(k)}(Y) - SA\phi Y + \widehat{F}_\xi^{(k)}(SY) + A\phi SY = 0.$$

Using (2.2), (2.3), and (2.4) gives us

$$\begin{aligned}
 (2.5) \quad &-k\phi SY + \phi ASY + kS\phi Y - S\phi AY = 0, \\
 &kS\phi Y - SA\phi Y - k\phi SY + A\phi SY = 0,
 \end{aligned}$$

respectively.

By combining these equations, we have

$$(2.6) \quad S(\phi A - A\phi)Y = (\phi A - A\phi)SY$$

for any  $Y$  tangent to  $M$ .

**Lemma 2.1** *Let  $M$  be a Hopf hypersurface in  $G_2(\mathbb{C}^{m+2})$ ,  $m \geq 3$ . If  $M$  satisfies  $\widehat{\mathcal{L}}_\xi^{(k)}S = \mathcal{L}_\xi S$ , then  $\xi$  belongs to either the distribution  $\mathcal{Q}$  or the distribution  $\mathcal{Q}^\perp$ .*

**Proof** To show this fact, we consider that the Reeb vector field  $\xi$  satisfies

$$(2.7) \quad \xi = \eta(X_0)X_0 + \eta(\xi_1)\xi_1$$

for some unit vectors  $X_0 \in \mathcal{Q}$ ,  $\xi_1 \in \mathcal{Q}^\perp$  and  $\eta(X_0)\eta(\xi_1) \neq 0$ .

Putting  $Y = \xi$  in (2.5) and (2.6), by (2.7) and using basic formulas in [5, Section 2], it follows that

$$(2.8) \quad \begin{aligned} \phi AX_0 &= k\phi X_0, \\ A\phi X_0 &= k\phi X_0. \end{aligned}$$

On the other hand, to prove the lemma, we need the following equation:

$$(2.9) \quad \begin{aligned} \alpha A\phi X + \alpha\phi AX - 2A\phi AX + 2\phi X &= 2 \sum_{v=1}^3 \left\{ -\eta_v(X)\phi\xi_v - \eta_v(\phi X)\xi_v \right. \\ &\quad \left. - \eta_v(\xi)\phi_v X + 2\eta(X)\eta_v(\xi)\phi\xi_v + 2\eta_v(\phi X)\eta_v(\xi)\xi \right\} \end{aligned}$$

([5, Lemma A]).

Putting  $X = X_0$  into (2.9), we have  $\alpha k - k^2 = \eta^2(X_0)$ .

Since  $k$  is non-zero constant, differentiating this with respect to  $\xi$ , we have

$$\begin{aligned} \xi\alpha &= -\frac{4}{k}\eta(X_0)\{g(\nabla_\xi X_0, \xi) + g(X_0, \nabla_\xi \xi)\} = -\frac{4}{k}\eta(X_0)g(\nabla_\xi X_0, \xi_1) \\ &= -\frac{4}{k}\eta(X_0)g(X_0, \phi_1 A\xi) = \frac{4}{k}\eta(X_0)\alpha g(X_0, \phi_1 \xi) = 0 \end{aligned}$$

where we have used  $\nabla_X \xi_v = q_{v+2}(X)\xi_{v+1} - q_{v+1}(X)\xi_{v+2} + \phi_v AX$ .

This gives  $\xi\alpha = 0$ .

Due to [4, Equation (2.10)],  $A\xi_1 = \alpha\xi_1$  is derived from  $\xi\alpha = 0$ . Equation (2.8) becomes

$$(\alpha - k)\phi\xi_1 = 0.$$

As  $k$  is nonzero constant and  $\phi X_0$  never vanishes, we have  $\alpha = k$ . Then by the equation  $Y\alpha = (\xi\alpha)\eta(Y) - 4\sum_{v=1}^3 \eta_v(\xi)\eta_v(\phi Y)$  in [5, Lemma A], we easily obtain that  $\xi$  belongs either to  $\mathcal{Q}$  or to  $\mathcal{Q}^\perp$  (see [10]). ■

Then by Lemma 2.1, we can divide our consideration into two cases being that  $\xi$  belongs to either  $\mathcal{Q}^\perp$  or  $\mathcal{Q}$ , respectively. Then first we consider the case  $\xi \in \mathcal{Q}^\perp$ . We can put  $\xi = \xi_1 \in \mathcal{Q}^\perp$  for our convenience sake.

Then [8, lemma 1.2] tells us Hopf hypersurface  $M$  in  $G_2(\mathbb{C}^{m+2})$  and  $\xi \in \mathcal{Q}^\perp$  gives  $AS = SA$ . Thus, (2.6) is changed into

$$\begin{aligned} 0 &= S(\phi A - A\phi)Y - (\phi A - A\phi)SY = S\phi AY - SA\phi Y - \phi ASY + A\phi SY \\ &= S\phi AY - AS\phi Y - \phi SAY + A\phi SY = (S\phi - \phi S)AY - A(S\phi - \phi S)Y \end{aligned}$$

By virtue of Lemma 2.1 and the above equations, we assert the following:

**Lemma 2.2** *Let  $M$  be a Hopf hypersurface in  $G_2(\mathbb{C}^{m+2})$ . If  $M$  satisfies  $A(\phi S - S\phi) = (\phi S - S\phi)A$  and  $\xi \in \mathcal{Q}^\perp$ , then we obtain  $S\phi = \phi S$ .*

**Proof** Since the shape operator  $A$  and the tensor  $\phi S - S\phi$  are both symmetric operators and commute with each other, by using the method due to Horn and Johnson [3], there exists a common basis  $\{E_i\}_{i=1, \dots, 4m-1}$  that gives a simultaneous diagonalization. Since  $A\xi = \alpha\xi$  and  $(\phi S - S\phi)\xi = 0$ ,  $\xi$  is principal for  $A$  and  $\phi S - S\phi$ . We write

$AE_i = \lambda_i E_i$  and  $(\phi S - S\phi)E_i = \beta_i E_i$ , where eigenvalues  $\lambda_i$  and  $\beta_i$  are real valued functions for all  $i \in \{1, 2, \dots, 4m - 1\}$ .

Bearing in mind that  $\xi = \xi_1 \in \mathcal{Q}^\perp$ , (2.1) is simplified:

$$(2.10) \quad SX = (4m + 7)X - 7\eta(X)\xi - 2\eta_2(X)\xi_2 - 2\eta_3(X)\xi_3 + \phi_1\phi X + hAX - A^2X.$$

As  $\xi$  is principal for both  $A$  and  $\phi S - S\phi$ , we get

**Case 1.** We can restrict  $X \in [\xi]^\perp$ . Here replacing  $X$  by  $\phi X$  in (2.10) (resp. applying  $\phi$  to (2.10)), we have

$$(2.11) \quad \begin{aligned} S\phi X &= (4m + 7)\phi X - \phi_1 X + 2\eta_2(X)\xi_3 - 2\eta_3(X)\xi_2 + hA\phi X - A^2\phi X, \\ \phi SX &= (4m + 7)\phi X - \phi_1 X + 2\eta_2(X)\xi_3 - 2\eta_3(X)\xi_2 + h\phi AX - \phi A^2 X. \end{aligned}$$

Combining equations in (2.11), we get

$$(2.12) \quad S\phi X - \phi SX = hA\phi X - A^2\phi X - h\phi AX + \phi A^2 X.$$

Putting  $X = E_i$  into (2.12) and using  $AE_i = \lambda_i E_i$ , we obtain

$$(2.13) \quad (S\phi - \phi S)E_i = hA\phi E_i - A^2\phi E_i - h\lambda_i\phi E_i + \lambda_i^2\phi E_i.$$

Taking the inner product with  $E_i$  into (2.13), we have

$$\beta_i g(E_i, E_i) = h\lambda_i g(\phi E_i, E_i) - \lambda_i^2 g(\phi E_i, E_i) = 0.$$

Since  $g(E_i, E_i) \neq 0$ ,  $\beta_i = 0$  for all  $i \in 1, 2, \dots, 4m - 2$ . This is equivalent to  $(S\phi - \phi S)E_i = 0$  for all  $i \in 1, 2, \dots, 4m - 2$ .

**Case 2.** For  $X \in [\xi]$ . This gives  $(S\phi - \phi S)\xi = 0$ . It follows that  $S\phi X = \phi SX$  for any tangent vector field  $X$  on  $M$ . ■

Summing up Lemmas 2.1, 2.2 and [12, Theorem], we conclude that if  $M$  is a Hopf hypersurface in complex two-plane Grassmannian  $G_2(\mathbb{C}^{m+2})$  satisfying (C-1) for  $X = \xi$  and  $\xi\alpha = 0$ , then  $M$  satisfies the condition of type (A) real hypersurfaces. Hereafter, let us check whether the Ricci tensor of a model space of type (A) satisfies the given condition (C-1) for  $X = \xi$ .

First let us consider  $X = \xi$ ; then (C-1) becomes

$$(2.14) \quad (\widehat{\mathcal{L}}_\xi^{(k)} S)Y = (\mathcal{L}_\xi S)Y,$$

which is equivalent to

$$(2.15) \quad -k\phi SY - \phi ASY + kS\phi Y - S\phi AY = 0.$$

When  $\xi$  is Hopf vector field and  $\xi \in \mathcal{Q}^\perp$ , the Ricci tensor  $S$  commutes with the structure tensor  $\phi$  and by [8, lemma 1.2],  $M_A$  satisfies (2.15).

If the Reeb vector field  $\xi$  belongs to the maximal quaternionic subbundle  $\mathcal{Q}$ , then a Hopf hypersurface  $M$  in  $G_2(\mathbb{C}^{m+2})$  is locally congruent to one of type (B) by virtue of [6, Main Theorem].

For  $M_B$ , (2.14) is also equivalent to (2.15). So we assume  $M_B$  satisfies (2.15). For each eigenspace, we have

$$SX = \begin{cases} (4m + 4 + h\alpha - \alpha^2)\xi & \text{if } X = \xi \in T_\alpha, \\ (4m + 4 + h\beta - \beta^2)\xi_\ell & \text{if } X = \xi_\ell \in T_\beta, \\ (4m + 8)\phi\xi_\ell & \text{if } X = \phi\xi_\ell \in T_\gamma, \\ (4m + 7 + h\lambda - \lambda^2)X & \text{if } X \in T_\lambda, \\ (4m + 7 + h\mu - \mu^2)X & \text{if } X \in T_\mu. \end{cases}$$

From [13], we obtain the following equations:

$$(2.16) \quad \alpha = -2 \tan(2r), \quad \beta = 2 \cot(2r), \quad \gamma = 0, \quad \lambda = \cot(r), \quad \mu = -\tan(r), \\ \lambda + \mu = \beta \quad \text{and} \quad h = \alpha + 3\beta + (4n - 4)(\lambda + \mu) = \alpha + (4n - 1)\beta.$$

Thus, we get

$$(\widehat{\mathcal{L}}_\xi^{(k)}S)Y - (\mathcal{L}_\xi S)Y = \begin{cases} 0 & \text{if } Y = \xi \in T_\alpha, \\ (4 - h\beta + \beta^2)(\beta - k)\xi_\ell & \text{if } Y = \xi_\nu \in T_\beta, \\ (4 - h\beta + \beta^2)\phi\xi_\ell & \text{if } Y = \phi\xi_\nu \in T_\gamma, \\ (-k + \lambda)(\lambda - \mu)(h - \lambda - \mu)\phi Y & \text{if } Y \in T_\lambda, \\ (-k + \mu)(\mu - \lambda)(h - \mu - \lambda)\phi Y & \text{if } Y \in T_\mu. \end{cases}$$

From the fourth equation of above (resp., fifth), since  $\mu \neq \lambda$ , due to (2.16), we have  $k = \mu$  or  $h = \beta$  (resp.,  $k = \lambda$  or  $h = \beta$ ). However, if  $h = \beta$ , the third one cannot happen. So we have  $k = \mu = \lambda$ . This gives a contradiction.

**Remark 2.3** Let  $M$  be a real hypersurface in complex two-plane Grassmannian  $G_2(\mathbb{C}^{m+2})$ ,  $m \geq 3$ ; then  $M_B$  does not satisfy the given condition  $(\widehat{\mathcal{L}}_\xi^{(k)}S)Y = (\mathcal{L}_\xi S)Y$ , for any  $Y$  tangent to  $M$ .

Thus, we have asserted Theorem 1.1 in the introduction.

Secondly, we assume that  $M_A$  satisfies (C-1). Putting  $Y = \xi$  into (C-1), we obtain

$$-\sigma\phi AX + k\sigma\phi X + S\phi AX - kS\phi X = 0,$$

where  $S\xi = \sigma\xi = (4m + h\alpha - \alpha^2)\xi$ .

From [13], we obtain the following equation:

$$SX = \begin{cases} (4m + h\alpha - \alpha^2)\xi & \text{if } X = \xi \in T_\alpha, \\ (4m + 6 + h\beta - \beta^2)\xi_\nu & \text{if } X = \xi_\nu \in T_\beta, \\ (4m + 6 + h\lambda - \lambda^2)X & \text{if } X \in T_\lambda, \\ (4m + 8)X & \text{if } X \in T_\mu. \end{cases}$$



For  $Y = \xi \in T_\alpha$ , we get

$$(2.17) \quad (\widehat{\mathcal{L}}_X^{(k)} S)\xi - (\mathcal{L}_X S)\xi = \begin{cases} 0 & \text{if } X = \xi \in T_\alpha, \\ (k - \beta)(-h\alpha + \alpha^2 + 6 + h\beta - \beta^2)\xi_3 & \text{if } X = \xi_2 \in T_\beta, \\ (k - \beta)(-h\alpha + \alpha^2 + 6 + h\beta - \beta^2)\xi_2 & \text{if } X = \xi_3 \in T_\beta, \\ (k - \lambda)(h\alpha - \alpha^2 - 6 - h\lambda + \lambda^2)\phi X & \text{if } X \in T_\lambda, \\ (h\alpha - \alpha^2 - 8)\phi X & \text{if } X \in T_\mu. \end{cases}$$

From the fifth equation in (2.17), we obtain

$$(2.18) \quad h\alpha - \alpha^2 - 8 = 0,$$

and from the definition of  $h$ , we obtain  $h = \alpha + 2\beta + (2m - 2)(\lambda + \mu)$ .

Summing these up, by [13] we have

$$(2.19) \quad (m - 1)t^2 - (m + 2)t + 4 = 0,$$

where  $t = \tan^2(\sqrt{2}r)$ .

From the second equation of (2.17) and (2.18), we obtain

$$(2.20) \quad (k - \beta)(h\beta - \beta^2 - 2) = 0.$$

If we assume that  $h\beta - \beta^2 - 2 = 0$ , then by summing up (2.19) with (2.20), we have  $m = -1$ , which gives us a contradiction. Thus,  $k = \beta$ , so from the fourth equation of (2.17) and (2.18), we get

$$h\lambda - \lambda^2 - 2 = 0,$$

which becomes

$$(2.21) \quad (2m - 3)t^2 - 4t + 1 = 0.$$

Combining (2.19) and (2.21) implies

$$t = \frac{-7m + 11}{(m - 2)(2m + 1)}.$$

Since  $m \geq 3$  and  $t \geq 0$ , this gives us a contradiction.

By virtue of Remark 2.3, we also get the fact that  $M_B$  does not satisfy the given condition  $(\widehat{\mathcal{L}}_X^{(k)} S)Y = (\mathcal{L}_X S)Y$ . Thus, we assert Corollary 1.2.

**Acknowledgment** The authors would like to express their deep gratitude to Dr. Lee working at RIRCM for giving us nice comments for this paper.

### References

- [1] J. T. Cho, *CR structures on real hypersurfaces of a complex space form*. Publ. Math. Debrecen 54(1999), no. 3-4, 473-487.
- [2] ———, *Levi parallel hypersurfaces in a complex space form*. Tsukuba J. Math. 30(2006), no. 2, 329-343. <http://dx.doi.org/10.21099/tkbjm/1496165066>
- [3] R. A. Horn and C. R. Johnson, *Matrix analysis*. Cambridge University Press, Cambridge, 1985. <http://dx.doi.org/10.1017/CBO9780511810817>
- [4] I. Jeong, C. Machado, J. D. Pérez, and Y. J. Suh, *Real hypersurfaces in complex two-plane Grassmannians with  $\mathcal{D}^\perp$ -parallel structure Jacobi operator*. Internat. J. Math. 22(2011), no. 5, 655-673. <http://dx.doi.org/10.1142/S0129167X11006957>

- [5] H. Lee, Y. S. Choi, and C. Woo, *Hopf hypersurfaces in complex two-plane Grassmannians with Reeb parallel shape operator*. Bull. Malays. Math. Sci. Soc. 38(2015), no. 2, 617–634.  
<http://dx.doi.org/10.1007/s40840-014-0039-3>
- [6] H. Lee and Y. J. Suh, *Real hypersurfaces of type B in complex two-plane Grassmannians related to the Reeb vector*. Bull. Korean Math. Soc. 47(2010), no. 3, 551–561.  
<http://dx.doi.org/10.4134/BKMS.2010.47.3.551>
- [7] T. H. Loo, *Semi-parallel real hypersurfaces in complex two-plane Grassmannians*. Differential Geom. Appl. 34(2014), 87–102. <http://dx.doi.org/10.1016/j.difgeo.2014.03.011>
- [8] E. Pak, Y. J. Suh, and C. Woo, *Restricted Ricci conditions for real hypersurfaces in complex two-plane Grassmannians*. Houston J. Math. 41(2015), no. 3, 767–783.
- [9] J. D. Pérez, I. Jeong, and Y. J. Suh, *Real hypersurfaces in complex two-plane Grassmannians with commuting normal Jacobi operator*. Acta Math. Hungar. 117(2007), 201–217.  
<http://dx.doi.org/10.1007/s10474-007-6091-9>
- [10] J. D. Pérez and Y. J. Suh, *The Ricci tensor of real hypersurfaces in complex two-plane Grassmannians*. J. Korean Math. Soc. 44(2007), 211–235.  
<http://dx.doi.org/10.4134/JKMS.2007.44.1.211>
- [11] J. D. Pérez, Y. J. Suh, and Y. Watanabe, *Generalized Einstein real hypersurfaces in complex two-plane Grassmannians*. J. Geom. Phys. 60(2010), 1806–1818.  
<http://dx.doi.org/10.1016/j.geomphys.2010.06.017>
- [12] Y. J. Suh, *Real hypersurfaces in complex two-plane Grassmannians with commuting Ricci tensor*. J. Geom. Phys. 60(2010), 1792–1805. <http://dx.doi.org/10.1016/j.geomphys.2010.06.007>
- [13] ———, *Real hypersurfaces in complex two-plane Grassmannians with parallel Ricci tensor*. Proc. Roy. Soc. Edinburgh Sect. A. 142(2012), 1309–1324. <http://dx.doi.org/10.1017/S0308210510001472>
- [14] ———, *Real hypersurfaces in the complex quadric with parallel Ricci tensor*. Adv. Math. 281(2015), 886–905. <http://dx.doi.org/10.1016/j.aim.2015.05.012>
- [15] ———, *Real hypersurfaces in the complex quadric with harmonic curvature*. J. Math. Pures Appl. 106(2016), 393–410. <http://dx.doi.org/10.1016/j.matpur.2016.02.015>
- [16] Y. J. Suh and H. Y. Yang, *Real hypersurfaces in complex two-plane Grassmannians with commuting structure Jacobi operator*. Bull. Korean Math. Soc. 45(2008), no. 3, 495–507.  
<http://dx.doi.org/10.4134/BKMS.2008.45.3.495>
- [17] N. Tanaka, *On non-degenerate real hypersurfaces, graded Lie algebras and Cartan connections*. Japan. J. Math. (N.S.) 2(1976), 131–190.
- [18] S. Tanno, *Variational problems on contact Riemannian manifolds*. Trans. Amer. Math. Soc. 314(1989), 349–379. <http://dx.doi.org/10.1090/S0002-9947-1989-1000553-9>
- [19] S. M. Webster, *Pseudo-Hermitian structures on a real hypersurface*. J. Differential Geom. 13(1978), 25–41. <http://dx.doi.org/10.4310/jdg/1214434345>

*Division of Future Capability Education, Ju Si-Gyeong College, Pai Chai University, 155-40 Baejae-ro, Seo-gu, Daejeon, 35345, Republic of Korea*  
 e-mail: [imsoon.jeong@gmail.com](mailto:imsoon.jeong@gmail.com)

*Departamento de Geometria y Topologia, Universidad de Granada, 18071-Granada, Spain*  
 e-mail: [jdperez@ugr.es](mailto:jdperez@ugr.es)

*Department of Mathematics and Research Institute of Real and Complex Manifold, Kyungpook National University, Daegu 41566, Republic of Korea*  
 e-mail: [yjsuh@knu.ac.kr](mailto:yjsuh@knu.ac.kr)

*Department of Mathematics Education, Woosuk University, 565-701 Wanju, Jeonbuk, Republic Of Korea*  
 e-mail: [legalgwch@naver.com](mailto:legalgwch@naver.com)