

patterns between patient tumor RNA-seq and clinical outcomes and map these associations on to higher-resolution data like spatial and single-cell transcriptomics. We apply DEGAS to prostate and pancreatic cancer spatial transcriptomics samples, as well as one normal sample of prostate tissue. We used the TCGA prostate cancer cohort to with the accompanying survival information and publicly accessible prostate cancer ST data from 10X Genomics to predict survival associations in the ST slides derived from the TCGA patients. Based on these survival associations, we identify higher risk subsections of ST slides which can be further studied. RESULTS/ANTICIPATED RESULTS: We were able to validate our method by comparing it to Scissor and were able to show that the number of high-risk regions in prostate cancer slides increased with the stage of disease. Furthermore, we identify transcriptomic signatures enriched for ontology terms associated with growth regulation and apoptosis, inflammation, immune signaling, and autophagy in histologically normal prostate tissues and adjacent normal pancreatic cancer tissues that were identified as high-risk by DEGAS. The regions highlighted by DEGAS could reflect transcriptional precursors to intraepithelial neoplasia—a well-recognized premalignant morphological change in glandular epithelium. DISCUSSION/SIGNIFICANCE: Identifying biomarkers of tissue stress that precede morphologic diagnosis of high-grade pre-malignant lesions by a pathologist may help triage patients at high risk for future development of cancer, or aid in better understanding whether histologically normal pre-malignant tissues at tumor margins contribute to recurrence.

270

Development of a Predictive Nomogram for Circumferential Resection Margin in Rectal Cancer Surgery

Megan Shroder, Molly M Ford, Fei Ye, Zhiguo Zhao, Aimal Khan, Shannon McChesney, M. Benjamin Hopkins, Alexander T. Hawkins
Vanderbilt University Medical Center

OBJECTIVES/GOALS: A negative circumferential resection margin (CRM) after surgical resection of rectal cancer decreases local recurrence and increases overall survival. While MRI is used to predict this risk, there is no predictive model that incorporates clinical factors to predict the risk of CRM positivity. METHODS/STUDY POPULATION: Utilizing the National Cancer Database from 2010-2014, we performed a retrospective study evaluating factors predictive for positive CRM after surgical resection of rectal cancer. The primary outcome was positive CRM (tumor \leq 1 mm from the surgical margin). Our population included patients with clinical stage I-III rectal cancer who underwent total mesorectal excision. For the primary outcome, multivariable logistic models were used to estimate the probability of a positive CRM. Model performance was evaluated by using the area under the receiver operating characteristic curve (AUC). Model calibration was assessed by examining the calibration plot. Bootstrapping method (300-iteration) was used to internally validate and estimate optimism-adjusted measures of discrimination and overall model fit. RESULTS/ANTICIPATED RESULTS: There were 28,790 patients included. 2,245 (7.8%) had positive CRM. Older age, race, larger tumor size, higher tumor grade, mucinous and signet tumor histology, APR, open operative approach, facility location, higher T stage, lymphovascular invasion, lack of neoadjuvant chemotherapy/radiation, and perineural invasion were all significantly associated with positive CRM (p < 0.05). DISCUSSION/SIGNIFICANCE: An objective model that predicts

positive CRM and associated poor clinical outcomes is possible to be used in conjunction with MRI. Positive CRM is associated with specific patient demographics, tumor characteristics, and operative approach. These factors can be used to predict CRM positivity in the preoperative period and plan accordingly.

271

Diagnosis and Detection of Thrombosis in PCOS

Mariel Miller, Caitlin Raymond, Bharathi Kavuri, Christopher Zahner
University of Texas Medical Branch

OBJECTIVES/GOALS: Identify and stratify clinical presentations of thrombotic pathology in PCOS patients. This will be accomplished by 1) evaluating clinical assays for the detection of hypofibrinolysis, and 2) analyzing clinical symptomology of thrombosis in PCOS via Symptom-Disease Pair Analysis of Diagnostic Error (SPADE). METHODS/STUDY POPULATION: Preliminary study populations include n=3 for each of the following groups PCOS with thrombotic complications, PCOS without thrombotic complications, healthy controls, and healthy control samples treated ex vivo with PAI-1. Coagulation assays include ACL coagulation panel, antiphospholipid antibody assays, viscoelastic testing with TEG and Quantra devices, and global fibrinolytic capacity. Coagulation assays will be performed on three samples taken at 4-week intervals. The SPADE techniques will be used to evaluate symptomology of thrombosis in the study period and patient electronic medical history. Molecular testing will be performed for pro-thrombotic polymorphisms (PAI-1 and Apo E) and mutations (Factor V Leiden and Prothrombin). RESULTS/ANTICIPATED RESULTS: We anticipate PCOS and ex vivo PAI-1 samples to show signs of hypofibrinolysis on TEG and Quantra devices outside of reference ranges and with statistical significance. We also anticipate seeing a statistical significance with ACL coagulation panels, however, these results are expected to still be within the reference ranges, as seen in previous studies. We believe the SPADE method will identify clinical presentations of hypofibrinolysis in PCOS patients coinciding with patterns in laboratory tests that were misdiagnosed due to being within reference ranges. We hope to stratify clinical presentations predictive of thrombosis in PCOS patients with standard clinical assays and with increased precision using viscoelastic assays. DISCUSSION/SIGNIFICANCE: The incidence rate of thrombosis is 40x higher in PCOS compared to healthy populations. However, the mechanisms of thrombosis in PCOS are unknown and are undetectable with current clinical assays. We hypothesize that chronic cellular stress in PCOS disrupts the regulation of interconnected immune pathways, causing hypofibrinolysis.

272

Differential expression of two Plasmodium falciparum variant surface antigen families in Malian children with cerebral malaria compared to mild malaria*

Jonathan G. Lawton¹, Albert E. Zhou¹, Drissa Coulibaly², Emily M. Stucke¹, Antoine Dara^{1,2}, Matthew B. Laurens¹, Joana C. Silva¹, Mahamadou A. Thera², Mark A. Travassos¹

¹University of Maryland School of Medicine, Baltimore, MD, USA

²University of Sciences, Techniques and Technologies, Bamako, Mali

OBJECTIVES/GOALS: Recent in vitro evidence suggests that diverse parasite protein families called RIFINs and STEVORs are displayed

on the surface of infected red blood cells and may have a role in severe malaria, but they remain sparsely studied in natural infections. We measured the RNA expression of these antigens in Malian children with severe or mild malaria illness. **METHODS/STUDY POPULATION:** We collected blood samples from Malian children aged six months to five years, including 14 with cerebral malaria, 10 with severe malarial anemia, and demographic-matched controls with mild, uncomplicated malaria. We extracted total RNA from each patient and used a custom capture array to selectively enrich *Plasmodium falciparum* parasite RNA. We then performed Illumina next-generation RNA sequencing and reconstructed parasite transcriptomes using reference-free de novo assembly. We identified RNA encoding RIFINs and STEVORs using an in-house classifier, then measured the diversity and abundance of gene expression for each infection. Expression diversity was defined as the number of unique variants transcribed. Expression abundance was calculated as transcripts per million (TPM). **RESULTS/ANTICIPATED RESULTS:** Cerebral malaria cases, but not severe malarial anemia cases, had higher diversity and abundance of RIFIN expression compared to mild infections. Type A RIFINs predominated over Type B RIFINs, and the same two RIFINs were predominantly expressed in all disease phenotypes. We anticipate that predominantly expressed RIFINs share high sequence homology with variants previously shown to bind blood antigens or immune inhibitory receptors. STEVOR expression was also higher in cerebral malaria compared to mild malaria, but STEVOR transcripts were sparse overall. **DISCUSSION/SIGNIFICANCE:** Elevated RIFIN expression in cerebral malaria over mild malaria supports a role for these antigens in pathogenesis. Severe malarial anemia may progress through a different pathogenic mechanism. Predominantly expressed RIFIN variants may be promising targets for vaccines and therapeutics to protect children against cerebral malaria.

273

Dystonia is associated with macro and microstructural abnormalities of the cerebellum: A nested case-control study.

Xenos Mason, Elizabeth Haddad, Neda Jahanshad
University of Southern California Keck School of Medicine

OBJECTIVES/GOALS: Dystonia is a brain disorder which causes excessive muscle activation, manifesting as abnormal movements. Neuroimaging studies of dystonia have revealed changes in a network involving the cerebellum. We sought out to determine whether subjects with dystonia in the UK Biobank exhibit MRI-based cerebellar pathology. **METHODS/STUDY POPULATION:** This nested case-control study drew from the UK Biobank, a cohort of >500,000 subjects in the United Kingdom, aged 40-69, enrolled 2006-2010. Eligible subjects must have undergone diffusion-weighted brain MRI. Dystonia cases were ascertained using ICD10 codes. We selected controls without neurologic diagnoses, matched (1:3) on age, sex, imaging site, and medical comorbidities. Mean diffusivity, fractional anisotropy, intracellular and isotropic volume fractions, and orientation dispersion were extracted from four white-matter tracts (inferior, middle, superior cerebellar peduncles; superior thalamic radiations) along with cerebellum, basal ganglia, and whole-brain grey and white-matter volumes. Means were compared using two-tailed t-tests and the Benjamini-Hochberg

procedure (FDR = 0.05). **RESULTS/ANTICIPATED RESULTS:** 23 cases of dystonia and 69 control subjects were selected and ascertained. After correcting for multiple-comparisons (40 volumetric and 35 diffusion-related), intracellular-volume-fraction (ICVF) of the middle- and superior-cerebellar peduncles was significantly lower in subjects with dystonia, suggesting reduced axon density. Volumetric analysis showed significantly reduced volumes of the motor cerebellum (lobules VI and VIII). There were no differences in basal ganglia, cortical, or whole-brain volumes. **DISCUSSION/SIGNIFICANCE:** These findings support the hypothesis that abnormalities of cerebellar networks contribute to dystonia. In future we will use similar techniques to assess these tracts in subjects with dystonia who are undergoing Deep Brain Stimulation, with the goal of guiding stimulation targeting, and predicting therapeutic outcomes.

274

Early life antibiotic exposure and genetic risk in neurodevelopmental disorders: effects on neurogenesis, the gut microbiome, and behavior

Courtney R. McDermott¹, Anya Mirmagles¹, Zhan Gao¹, Katherine Kimbark², Christiana Ntim¹, Xuesong Zhang¹, Xiaofeng Zhou¹, James H. Millonig¹, Emanuel DiCicco-Bloom¹
¹Rutgers University, NJ ²Lebanon Valley College, PA

OBJECTIVES/GOALS: Our long-term goal is to understand how both genetic and environmental (GxE) factors contribute to neurodevelopmental disorders (NDDs) so that we may potentially intervene in disease pathogenesis and design therapies to address functional deficiencies. **METHODS/STUDY POPULATION:** Our studies use a novel GxE model to determine how cephalosporin antibiotic exposure alters the gut microbiome, hippocampal neurogenesis, and behavior in the genetically vulnerable 16p11.2 microdeletion (16pDel) mouse. This mouse models one of the most frequently observed genetic risk variants implicated in NDDs, including ~1% of autism diagnoses. Wildtype and 16pDel littermates were exposed to saline or the cephalosporin, cefdinir, from postnatal days 5-9. We quantified changes in gut microbiota composition using 16S rRNA gene sequencing and utilized immunoblotting, immunohistochemistry, and bulk RNA gene sequencing to assess changes in hippocampal neurogenesis. An additional cohort of saline or cefdinir-exposed mice were subjected to a behavioral battery to assess changes in sociability and anxiety. **RESULTS/ANTICIPATED RESULTS:** We leveraged the next-generation microbiome bioinformatics platform, Quantitative Insights Into Microbial Ecology 2 (QIIME2) to analyze 16S rRNA gene sequencing datasets of P13 cecal samples from saline- and cefdinir-exposed mice. We found successful perturbations to the gut microbiome following early life cefdinir exposure. Further, we found a robust 50% reduction in hippocampal cyclin E protein in cefdinir-exposed 16pDel male mice, which was replicated in a second independent experiment. This reduction extended to the S-phase cell entry and general stem cell population, quantified by EdU+ and Ki67+ cell numbers, respectively. Lastly, in our first cohort of mice for behavioral studies, we found reduced sociability and increased anxiety-like behaviors in cefdinir-exposed mice. **DISCUSSION/SIGNIFICANCE:** The findings from this GxE model will provide mechanistic insights into the causes of NDDs; they may inform practice guidelines so as to reduce this environmental exposure; and may suggest interventions like probiotics for those at risk in order to overcome altered gut microbiome composition and restore hippocampal neurogenesis defects.