

OBSERVATIONS OF THE CMB ON SCALES OF 2° TO 15°

The Tenerife experiments

R.D. DAVIES, R.J. DAVIS, A. WILKINSON, R.A. WATSON, S.J. MELHUISE,
S. DICKER AND S. VEERARAGHAVAN

NRAL Jodrell Bank, University of Manchester, UK

R. REBOLO, C.M. GUTIERREZ AND R.J. HOYLAND

IAC, La Laguna, Tenerife, Spain

AND

A.N. LASENBY, S. HANCOCK, A.W. JONES AND G. ROCHA

MRAO, Cambridge University, UK

Beamswitching has been used at 10, 15 and 33 GHz to map the microwave background over the Declination range 30° to 45°, covering more than one steradian of the sky. The beamwidth is 5° and the beam-throw is $\pm 8^\circ$ at each frequency. The three data sets are used to separate Galactic emission from intrinsic CMB emission. For the scan at Dec = 40° the intrinsic fluctuation level is $\Delta T_{\text{rms}} = 48_{-15}^{+21} \mu\text{K}$ on a coherence scale of 4°; the equivalent analysis for a Harrison–Zeldovich model gives a power spectrum normalisation of $Q_{\text{rms}} = 22_{-6}^{+10} \mu\text{K}$. The value of the fluctuation amplitude calculated from the likelihood analysis of the two-dimensional data set is $\Delta T_{\text{rms}} = 54 \pm 13 \mu\text{K}$ at 10 GHz and $39_{-7}^{+8} \mu\text{K}$ at 15 GHz.

A short baseline interferometer has been operating at 33 GHz. The spacing of the two low-spillover horns is 17λ (152 mm). The bandwidth is 3 GHz giving an rms noise of 200 μK in 2 minutes. With a primary beamwidth of $5^\circ \times 2.5$ the interferometer lobe size is $1.7^\circ \times 2.5$. The interferometer is largely unaffected by atmospheric water vapour. Data at Dec = 41° show significant structure at high signal-to-noise. The section from RA = 08 – 19 hr shows CMB structure at an rms level of $\sim 20 \mu\text{K}$; this corresponds to a sky $\Delta T_{\text{rms}} \sim 100 \mu\text{K}$, at an angular scale of 2°, when corrected for dilution; a value significantly larger than found in our beamswitching experiments on a 5° – 15° angular scale. This work is now being extended in the *Very Small Array, Tenerife*, a collaboration between our three groups. This has 14 antennas operating at baselines up to 2 m in the frequency range 26–36 GHz. The sensitivity will be $\Delta T/T \sim 2 \times 10^{-6}$ in several months, in the range $l = 100 - 1000$. Observations will begin in 1999.