

Illustrations. The same material should not normally be presented in tables and figures. Legends should be typed on a separate sheet.

- (a) Diagrams should be drawn in Indian ink, about twice the size of the finished block, on Bristol board, stout tracing paper or plastic film. Photographed diagrams are also accepted. Lettering inside the framework of the diagram should be avoided as far as possible but if unavoidable it should be included on a fly-leaf. Marginal lettering should be inserted lightly in pencil on the original diagram or on a fly-leaf.
- (b) Photographs intended for half-tone reproduction should be on glossy paper. They will be accepted only if found necessary by the Editors.
- (c) Colour plates are unlikely to be accepted unless authors bear the cost.

Tables should be as simple and as few as possible. Each table should be typed on a separate sheet.

Abstract. Every article should have a short abstract (not more than **250** words) complete in itself and understandable without reference to the paper. *The abstract will be printed at the beginning of the paper.*

References. Only papers closely related to the author's work should be referred to; exhaustive lists should be avoided. The arrangement of references should be as in recent papers in *Animal Production*; details are given in the notes for guidance.

Proofs are supplied once and must be returned corrected to the Editors *within 3 days*. Only essential corrections should be made.

Reprints. Twenty-five reprints of each paper will be supplied free to authors on request. Further copies may be purchased if the order is sent at the proof-stage.

Animal Production is published six times a year in two volumes. Annual subscription is £49.00 (or \$112.00 in the USA and Canada) and the price for a single part is £9.00 (or \$21.00) net.

Proceedings of the British Society of Animal Production (New Series) ceased publication in 1975 and the material is now published as a separate section in the June issue of *Animal Production*.

Business matters, including regular subscriptions and sales (current and back numbers of the Journal and the Proceedings), should be addressed to: Longman Group Limited, Subscriptions (Journals) Department, Fourth Avenue, Harlow, Essex, England CM19 5AA.

Senior Editor

J. A. F. Rook, Agricultural Research Council

Editors

P. J. Avery, University of Reading
R. C. Campling, Wye College, University of London
M. F. Fuller, Rowett Research Institute
R. B. Land, Animal Breeding Research Organisation
T. L. J. Lawrence, University of Liverpool
A. M. Raven, Department of Agriculture and Fisheries for Scotland
J. H. B. Roy, National Institute for Research in Dairying
A. J. F. Webster, University of Bristol

Technical Editors

Jill M. Read
P. D. Wilson

CONTENTS

	PAGE
CAMPBELL, R. G. The effects of carbadox on the performance and carcass characteristics of early-weaned pigs given high- and low-protein diets	111
JONES, P. W. and TARRANT, M. E. The effect of various factors on the efficacy of tylosin as a growth promoter in clinically healthy pigs	115
GIBB, M. J. and TREACHER, T. T. The effect of body condition and nutrition during late pregnancy on the performance of grazing ewes during lactation	123
VIPOND, J. E., HUNTER, E. A. and KING, MARGARET E. Effects of cereal and protein supplements to swedes (<i>Brassica napus</i>) on intake and performance of pregnant and lactating ewes kept indoors	131
ARORA, R. C. and PANDEY, R. S. Plasma concentrations of progesterone, oestradiol-17 β and luteinizing hormone in relation to repeat breeding in buffalo (<i>Bubalus bubalis</i>)	139
PETERS, A. R. and RILEY, G. M. Milk progesterone profiles and factors affecting <i>post partum</i> ovarian activity in beef cows	145
SOUTHGATE, J. R., COOK, G. L. and KEMPSTER, A. J. A comparison of the progeny of British Friesian dams and different sire breeds in 16- and 24-month beef production systems. 1. Live-weight gain and efficiency of food utilization	155
KEMPSTER, A. J., COOK, G. L. and SOUTHGATE, J. R. A comparison of the progeny of British Friesian dams and different sire breeds in 16- and 24-month beef production systems. 2. Carcass characteristics, and rate and efficiency of meat gain	167
KAISER, A. G., OSBOURN, D. F., ENGLAND, P. and DHANOA, M. S. The utilization by calves of formaldehyde-treated maize silages and the response to supplementary protein	179
MACDEARMID, A., INNES, G. M., WILLIAMS, P. E. V. and KAY, M. Kale for beef production	191
POOLE, D. A. The effects of milking cows three times daily	197
DUCKER, M. J., YARROW, N. and MORANT, S. V. The effect of change and level of nutrition on the reproductive performance of group-fed dairy heifers	203
BAKER, R. D., LE DU, Y. L. P. and BARKER, J. M. The influence of winter nutrition, grazing system and stocking rate on the performance of spring-calving Hereford \times Friesian cows and their calves. 1. Winter nutrition	213
BAKER, R. D., BARKER, J. M. and LE DU, Y. L. P. The influence of winter nutrition, grazing system and stocking rate on the performance of spring-calving Hereford \times Friesian cows and their calves. 2. Grazing system and stocking rate	225
WOOD, P. D. P. and FRAPPELL, J. P. A note on the relationship between milk yield and reproductive performance in some British Friesian sire progeny groups	239