



Nowhere constant families of maps and resolvability

István Juhász and Jan van Mill

Abstract. If X is a topological space and Y is any set, then we call a family \mathcal{F} of maps from X to Y *nowhere constant* if for every non-empty open set U in X there is $f \in \mathcal{F}$ with $|f[U]| > 1$, i.e., f is not constant on U . We prove the following result that improves several earlier results in the literature.

If X is a topological space for which $C(X)$, the family of all continuous maps of X to \mathbb{R} , is nowhere constant and X has a π -base consisting of connected sets then X is c -resolvable.

1 Introduction

The question about how resolvable are crowded locally connected spaces has been around for some time. Costantini proved in [1] that *regular* such spaces are ω -resolvable. Actually, it is proved there that local connectedness can be weakened to having a π -base consisting of connected sets. We simply call such spaces π -connected.

In [5], it is stated that Yaschenko proved that every crowded locally connected Tykhonov space is c -resolvable, however, as far as we know, no proof of this has been published.

Dehghani and Karavan claim in [2] that every crowded locally connected functionally Hausdorff space is c -resolvable. They get this as a corollary of their more general Theorem 2.6, in the proof of which, however, we found a gap, and we do not know if this gap can be fixed. On page 88 of their paper, lines -7 and -6, they claim that $\bigcup_{\gamma \neq \alpha} A_\gamma = f^{-1}(\bigcup_{\gamma \neq \alpha} D_\gamma)$ is not closed in any nonempty connected open subset V of U . But this can only be concluded in case the restriction of f to V is not constant, and they only assume that some continuous function $g : V \rightarrow \mathbb{R}$ is not constant.

It follows, however, from our results below that their Corollary 2.7 of Theorem 2.6 is correct. In fact, the following property weaker than being crowded and functionally Hausdorff suffices: for every non-empty open set U , there is a continuous map of the whole space to \mathbb{R} that is not constant on U .

We are going to get this from a more general result that will make use of the following concept.

Received by the editors December 12, 2023; revised January 20, 2024; accepted January 21, 2024.

Published online on Cambridge Core February 6, 2024.

The first author was supported by the NKFIH (Grant No. K129211).

AMS subject classification: 54A25, 54C30, 54D05.

Keywords: Nowhere constant family of maps, resolvable space, π -base, connected, locally connected.



Definition 1.1 Let X be a topological space and Y any set. We call a family \mathcal{F} of maps from X to Y *nowhere constant (NWC)* if for every non-empty open set U in X , there is $f \in \mathcal{F}$ with $|f[U]| > 1$, i.e., f is not constant on U .

We note the trivial fact that any space that admits a NWC family of maps is crowded, i.e., has no isolated points.

2 The results

We first present a very general result that connects NWC families of maps and resolvability.

Theorem 2.1 Let \mathcal{F} be a NWC family of maps of the topological space X to the set Y . Moreover, \mathcal{B} is a π -base of X and \mathcal{A} is a disjoint family of subsets of Y such that, putting $\mathcal{B}_f = \{B \in \mathcal{B} : |f[B]| > 1\}$ and $U_f = \bigcup (\mathcal{B} \setminus \mathcal{B}_f)$ for any $f \in \mathcal{F}$, we have

$$\forall f \in \mathcal{F} \forall B \in \mathcal{B}_f \forall A \in \mathcal{A} (A \cap f[B \setminus U_f] \neq \emptyset).$$

Then X is $|\mathcal{A}|$ -resolvable.

Proof We are going to use transfinite recursion to produce a disjoint collection $\{D(A) : A \in \mathcal{A}\}$ of dense subsets of X .

To do that, we first introduce the following piece of notation. For $f \in \mathcal{F}$, $B \in \mathcal{B}_f$, and $A \in \mathcal{A}$, we let

$$S(f, B, A) = f^{-1}[A] \cap (B \setminus U_f).$$

It follows from our assumptions that $S(f, B, A) \neq \emptyset$.

We start our recursive construction by choosing $f_0 \in \mathcal{F}$ such that $\mathcal{B}_{f_0} \neq \emptyset$ and then define

$$D_0(A) = \bigcup \{S(f_0, B, A) : B \in \mathcal{B}_{f_0}\}$$

for all $A \in \mathcal{A}$. Then $\{D_0(A) : A \in \mathcal{A}\}$ is disjoint because $D_0(A) \subset f_0^{-1}[A]$ for each $A \in \mathcal{A}$, moreover $B \cap D_0(A) \neq \emptyset$ whenever $B \in \mathcal{B}_{f_0}$ and $A \in \mathcal{A}$.

Now, assume that $\alpha > 0$ and we have already defined $f_\beta \in \mathcal{F}$ and the family $\{D_\beta(A) : A \in \mathcal{A}\}$ for all $\beta < \alpha$, and consider

$$\mathcal{B}_\alpha = \bigcup \{\mathcal{B}_{f_\beta} : \beta < \alpha\}.$$

If $\mathcal{B}_\alpha = \mathcal{B}$, then our construction stops. Otherwise, we may pick $f_\alpha \in \mathcal{F}$ such that $\mathcal{B}_{f_\alpha} \setminus \mathcal{B}_\alpha \neq \emptyset$. In this case, we define

$$D_\alpha(A) = \bigcup \{D_\beta(A) : \beta < \alpha\} \cup \bigcup \{S(f_\alpha, B, A) : B \in \mathcal{B}_{f_\alpha} \setminus \mathcal{B}_\alpha\}$$

for each $A \in \mathcal{A}$.

Note that we clearly have $\bigcup (\mathcal{B}_{f_\alpha} \setminus \mathcal{B}_\alpha) \subset \bigcap \{U_{f_\beta} : \beta < \alpha\}$, hence it may be verified by straightforward transfinite induction that $\{D_\alpha(A) : A \in \mathcal{A}\}$ is disjoint, moreover, we have $B \cap D_\alpha(A) \neq \emptyset$ whenever $A \in \mathcal{A}$ and $B \in \mathcal{B}_{f_\beta}$ with $\beta \leq \alpha$.

Of course, this construction must stop at some ordinal α , in which case putting $D(A) = \bigcup \{D_\beta(A) : \beta < \alpha\}$ the disjoint family $\{D(A) : A \in \mathcal{A}\}$ consists of dense subsets of X because $B \cap D(A) \neq \emptyset$ for any $B \in \mathcal{B}$ and $A \in \mathcal{A}$ and \mathcal{B} is a π -base of X . ■

Now, we deduce from Theorem 2.1 our promised result concerning π -connected spaces.

Corollary 2.2 *If X is any π -connected space for which $C(X)$, the family of all continuous maps of X to \mathbb{R} , is NWC then X is \mathfrak{c} -resolvable.*

Proof To apply Theorem 2.1, we of course put $Y = \mathbb{R}$ and $\mathcal{F} = C(X)$. For \mathcal{B} , we choose a π -base of X consisting of connected sets, and let \mathcal{A} be any disjoint collection of dense subsets of \mathbb{R} with $|\mathcal{A}| = \mathfrak{c}$.

Note that for any $f \in C(X)$ and $B \in \mathcal{B}_f$, we have $|f[B]| > 1$, hence $f[B]$ is a nondegenerate interval in \mathbb{R} , being a non-singleton connected subset of \mathbb{R} , and thus $A \cap f[B] \neq \emptyset$ holds for all $A \in \mathcal{A}$. Consequently, we shall be done if we prove the following claim.

Claim For any $f \in C(X)$ and $B \in \mathcal{B}_f$, we have

$$f[B] = f[B \setminus U_f].$$

To see this, pick any $t \in f[B]$ and note that $B \cap f^{-1}(t)$ is a non-empty proper closed subset of the connected subspace B . Properness follows from $|f[B]| > 1$. Consequently, the boundary H_t of $B \cap f^{-1}(t)$ in B is non-empty. Also, we have $H_t \subset B \cap f^{-1}(t)$, the latter set being closed in B .

Now, take any $B' \in \mathcal{B} \setminus \mathcal{B}_f$, then $f[B'] = \{t'\}$ for some $t' \in \mathbb{R}$. If $t' \neq t$, then we have $B' \cap f^{-1}(t) = \emptyset$, hence $B' \cap H_t = \emptyset$ as well. If, on the other hand, $t' = t$, then $B' \cap B$ is in the interior of $B \cap f^{-1}(t)$ in B , hence we have $B' \cap H_t = \emptyset$ again. This, however, means that $H_t \cap U_f = \emptyset$, consequently, $t \in f[H_t] \subset f[B \setminus U_f]$, completing the proof of the claim. ■

A closer inspection of this proof reveals that we did not use the full force of continuity for the members f of the NWC family $\mathcal{F} = C(X)$. What we used was that f preserves connectedness for $B \in \mathcal{B}$, i.e., $f[B]$ is connected in \mathbb{R} for $B \in \mathcal{B}$, moreover, that $f^{-1}(t)$ is closed in X for all $t \in \mathbb{R}$.

3 Discussion and questions

Corollary 2.2 is clearly sharp in the sense that the cardinal \mathfrak{c} cannot be replaced with anything bigger because there are very nice crowded and locally connected spaces of cardinality \mathfrak{c} . The natural question arises, however, if such a space X is maximally resolvable, i.e., $\Delta(X)$ -resolvable, where $\Delta(X)$ is the minimum cardinality of a non-empty open set in X .

In a recent arXiv preprint [4], Lipin proved that if $2^{\mathfrak{c}} = 2^{(\mathfrak{c}^+)}$, then there is a locally connected and pseudocompact Tychonov space X such that $\Delta(X) > \mathfrak{c}$ but X is not \mathfrak{c}^+ -resolvable. So, at least consistently, Corollary 2.2 is sharp in that sense as well.

As we wrote in the Introduction, Costatini [1] proved that crowded π -connected regular spaces are ω -revolvable. As usual, in his treatment regular implies T_1 . But his proof actually works for all crowded π -connected *quasi-regular* spaces as well, without the use of any additional separation axiom. Recall that a space is quasi-regular if for every non-empty open U , there is a non-empty open V such that $\bar{V} \subset U$. The only slight modification we need is that in this case, the definition of *crowded* has to be replaced by the following assumption: there is no finite indiscrete open subspace. For T_0 -spaces, this assumption is clearly equivalent with the usual one, i.e., not having any isolated points.

Hewitt [3] gave an example of a regular connected space X of cardinality ω_1 . Let $Y = \lambda_f(X)$ be the superextension of X consisting of all finitely generated maximal linked systems consisting of closed subsets of X . Then Y also has size ω_1 , since the collection of all finite subsets of X has size ω_1 , moreover, Y is Hausdorff, connected and locally connected (see [7, IV.3.4(v) and (viii)]).

We claim that Y is also regular and adopt the terminology of [7]. To prove this, let $m \in Y$ be arbitrary, and let A be a closed subset of Y not containing m . Let $F \subseteq X$ be a finite defining set for m , and let $n = m \restriction F$. There is a finite collection of open subsets \mathcal{U} of X such that $m \in \bigcap_{U \in \mathcal{U}} U^+ \subseteq Y \setminus A$. For each $U \in \mathcal{U}$, let $N_U \in n$ be such that $N_U \subseteq U$. By regularity of X , we may pick an open neighborhood V_U of N_U whose closure is contained in U . Then $\bigcap_{U \in \mathcal{U}} V_U^+$ is a closed neighborhood of m that misses A .

Hence Y is a connected, locally connected regular space which by Costantini's result is ω -resolvable. (In this special case, there is also a direct proof of this.) But Y has cardinality ω_1 , hence it is not \mathfrak{c} -resolvable if the Continuum Hypothesis fails. So, for crowded locally connected regular spaces, a positive answer to the following question is the best we can hope for.

Problem 3.1 Let X be regular, crowded, and locally connected (respectively, π -connected). Is X_{ω_1} -resolvable?

We cannot resist mentioning here that it is still a widely open question whether non-singleton connected regular spaces are resolvable (see e.g., [6] for details and references). This seems to be one of the most central open problems in the area.

We call a space X *nowhere 0-dimensional* if no non-empty open subspace of X is 0-dimensional. Clearly, any crowded locally connected, even π -connected, space is nowhere 0-dimensional.

Assume that X is a nowhere 0-dimensional Tychonov space. We may assume that X is a subspace of \mathbb{R}^I for some set I . For every $i \in I$, let $\pi_i: \mathbb{R}^I \rightarrow \mathbb{R}$ be the i th projection. Then, for every nonempty open subset U of X , there exists $i \in I$ such that $\pi_i(U)$ has nonempty interior. For otherwise U would be a subspace of a product of 0-dimensional spaces, and so would itself be 0-dimensional. Hence the collection of projections $\mathcal{P} = \{\pi_i \restriction X : i \in I\}$ is NWC in a (very) strong sense. This observation lead us to the following problem.

Problem 3.2 Is every nowhere 0-dimensional Tychonov space resolvable (ω -resolvable, \mathfrak{c} -resolvable)?

References

- [1] C. Costantini, *On the resolvability of locally connected spaces*. Proc. Amer. Math. Soc. 133(2005), no. 6, 1861–1864.
- [2] A. Dehghani and M. Karavan, *Every locally connected functionally Hausdorff space is c -resolvable*. Topology Appl. 183(2015), 86–89.
- [3] E. Hewitt, *On two problems of Urysohn*. Ann. Math. 47(1946), 503–509.
- [4] A. Lipin, <https://arxiv.org/pdf/2308.01259.pdf>.
- [5] O. Pavlov, *On resolvability of topological spaces*. Topology Appl. 126(2002), 37–47.
- [6] O. Pavlov, *Problems on (ir)resolvability*. In: E. Pearl (ed.), Open problems in topology II, Elsevier, Amsterdam, 2007, pp. 51–59.
- [7] A. Verbeek, *Superextensions of topological spaces*, MC tract 41, Mathematisch Centrum, Amsterdam, 1972.

HUN-REN Alfréd Rényi Institute of Mathematics, Reáltanoda utca 13-15, 1053 Budapest, Hungary

e-mail: juhasz@renyi.hu

Korteweg-de Vries Institute for Mathematics, University of Amsterdam, Science Park 105-107, P. O. Box 94248, 1098 XG Amsterdam, Netherlands

e-mail: j.vanMill@uva.nl