

## Research Article

\*Authors contributed equally to this work.

**Cite this article:** Cao G *et al.* (2025) A survey and analysis of germplasm resources of food crops in Hainan Province. *Plant Genetic Resources: Characterization and Utilization* 1–9. <https://doi.org/10.1017/S1479262125100063>

Received: 18 November 2024

Revised: 30 April 2025

Accepted: 7 May 2025


### Keywords:

crop breeding; germplasm resource  
conservation; local food crop

**Corresponding author:** Qingjie Tang;

Email: [tangqingjie@hnaas.org.cn](mailto:tangqingjie@hnaas.org.cn)

# A survey and analysis of germplasm resources of food crops in Hainan Province

Guangping Cao<sup>1,\*</sup>, XiaoWei Yan<sup>1,2,\*</sup>, Linan Zhai<sup>1</sup>, Jing Xu<sup>1</sup>, Liqiong Tang<sup>1</sup>, Honglin Zhu<sup>1</sup>, Li Lin<sup>1</sup>, Mingchao Zhao<sup>1,2</sup>, Xiaorong Xiao<sup>1,2</sup>, Yapeng Li<sup>1,2</sup>, Qingyu Wang<sup>1</sup> and Qingjie Tang<sup>1,2</sup> 

<sup>1</sup>Institute of Food Crops, Hainan Academy of Agricultural Sciences/Hainan Key Laboratory of Crop Genetics and Breeding, Haikou, China and <sup>2</sup>Sanya Institute, Hainan Academy of Agricultural Sciences, Sanya, China

## Abstract

Germplasm resources are the foundation for improving crop varieties and a strategic asset for global food security. They also advance plant breeding, agricultural biotechnology and the production of essential agricultural goods. To assess the distribution, diversity and conservation status of food crop germplasm in the Hainan Province, China, we conducted a detailed survey of the Hainan Island. Between 2017 and 2022, we collected 330 food crop germplasm resources, encompassing 16 cereal crops, including rice, maize, sweet potato. The collected germplasm resources exhibited traits of high resistance to both biotic and abiotic stresses, including common diseases and drought stress, as well as superior quality and adaptability to poor soil conditions such as sandy land. However, challenges such as low productivity and hybrid degradation were identified. These resources were primarily found in Haikou City, Baisha County, Danzhou City, Wuzhishan City and Sanya City. Additionally, we collected several ancient local varieties and endangered germplasm resources such as ‘Jiezi rice’ and ‘Wuzhishan maize’. This study serves as a reference for the conservation, development and utilization of local food crop germplasm resources in Hainan Province and lays the foundation for breeding and developing new varieties.

## Introduction

Germplasm resources are the foundation for improving crop varieties and a strategic asset for global food security. They also drive advancements in plant breeding, agricultural biotechnology and the production of essential agricultural goods (Wang 2002; Li *et al.* 2015). Enriching crop germplasm diversity prevents the extinction of high-potential varieties and provides essential genetic resources to support the development of the plant breeding-related industry. Recently, environmental changes, resource overexploitation and shifts in crop production, and land use have caused the rapid disappearance of many local crop varieties (Liu *et al.* 2017). For example, three wild rice species are now classified under China’s second-level endangered plant protection catalog (Fan *et al.* 2023). Many governments worldwide, such as China, designate germplasm resource protection as a top priority, emphasizing the critical role of germplasm surveys as the cornerstone of seed industry revitalization and highlighting the importance of investigating, collecting and protecting crop germplasm resources (Chen *et al.* 2018; Zeng *et al.* 2019; Li and Yang 2022; Lei *et al.* 2023).

China conducted two nationwide surveys and collection campaigns of food crop germplasm resources in 1956 and 1981, respectively. These efforts laid a solid foundation for the conservation and utilization of germplasm resources in China. The third nationwide survey and collection campaign of crop germplasm resources was launched in July 2015 and fully completed between 2021 and 2023. This campaign covered 2323 agricultural counties (cities and districts) across China, aiming to comprehensively assess the national status of crop germplasm resources, rescue and collect endangered and rare germplasm, and provide a robust foundation for agricultural breeding innovation, ecological civilization construction and sustainable agricultural development.

Hainan Province, situated at China’s southernmost tip, lies between 108°37′ to 111°03′ east longitude and 18°10′ to 20°10′ north latitude. Positioned on the northern edge of the tropics, Hainan experiences a tropical monsoon climate, earning it the nickname ‘Natural Greenhouse’. The annual average temperature varies between 22.5 and 25.6°C, with the lowest in central Qiongzong county and the highest in Sanya city in the south. The accumulated growing degree days (temperatures ≥10°C) reach 8200°C. Even in January, the coldest month, temperatures remain between 17°C and 24°C. Additionally, the province receives abundant sunlight, with 1750 to 2650 annual sunshine hours and a light exposure rate of 50–60%.



Hainan also experiences abundant rainfall, with annual precipitation ranging from 1000 to 2600 mm. The rainy season, spanning May to October, contributes 70–90% of the annual precipitation, totalling approximately 1500 mm. The dry season, lasting from November to April, provides only 10–30% of the annual rainfall. The soils in the study area are sandy loam with low organic matter content, thus qualifying them as poor soils (Zhang *et al.* 2005). In line with this, the main food crops include rice, wheat, maize and soybean. In this study, food crops refer to cultivated varieties, traditional landraces and wild relatives. However, due to the favourable light, temperature and water availability, the area also supports a diverse range of other agricultural and biological resources, such as fruit trees and vegetables, characterized by their strong adaptability, drought resistance and the ability to thrive in poor soils (Zheng and Gao 2016; Zheng *et al.* 2016).

Previous surveys of food crop germplasm resources in Hainan Province were conducted in 1951, 1981 and the 1990s. These efforts documented the distribution and characteristics of local germplasm resources, providing a valuable historical baseline (Wang *et al.* 1992). Over the next two decades, these resources underwent significant changes. In 2015, China's Ministry of Agriculture launched the Third National Crop Germplasm Resources Survey and Collection Campaign. As part of this initiative, the Institute of Food Crops (IFC) of the Hainan Academy of Agricultural Sciences (HNAAS), acting as the Hainan survey team, conducted extensive investigations and collections across the province. Between 2017 and 2021, the team focused on assessing the current distribution and cultivation status of genetic resources, prioritizing the collection and conservation of ancient landraces and endangered germplasm (Ministry of Agriculture of the PRC, National Development and Reform Commission, Ministry of Science and Technology of the PRC, 2015). Field investigations and resource collection were carried out in 24 cities, counties and districts, including Haikou, Sanya and Danzhou. These efforts provided valuable insights into the genetic diversity of Hainan's food crops. Rice has historically been the dominant food crop in Hainan Province, consistently accounting for over 30% of the total cultivated area since 1951. Its significance lies not only in its role as a staple food but also in its cultural importance and adaptability to Hainan's diverse ecological conditions. This study emphasizes rice to highlight its central role in Hainan's agricultural landscape and its potential for genetic improvement.

## Materials and methods

### Survey tools

In the process of germplasm resource collection, we used a variety of tools to ensure the accurate documentation and secure handling of specimens. Seed bags, mesh bags and plastic bags were used for the safe storage. A GPS device, such as the GPS Tool Kit mobile app, was employed to record precise geographic coordinates of collection sites, enabling accurate tracking and mapping of native habitats. Notebooks and pencils (or marker pens) were used by researchers to take detailed field notes, such as environmental conditions and species-specific observations. Tags were used to label specimens and maintain proper identification during storage and transport. Scissors and knife were used to facilitate the clean cutting of plant materials, such as branches or leaves, required for analysis. Lastly, high-resolution digital cameras were employed to capture visual records of the overall site, population structure, individual species and surrounding ecological

environment. Together, these tools ensured that the collection process was systematic, thorough and scientifically reliable (Liang *et al.* 2013; Yun *et al.* 2015).

### Survey routes and collection methods

The survey spanned across 18 cities on Hainan Island, covering approximately 9000 km (Fig. 1A). Special emphasis was placed on cities and villages with poor transportation, complex terrain, distinctive local customs and predominantly subsistence lifestyles to ensure representative surveys in regions with unique ecological and cultural characteristics. With the assistance of local agricultural technicians, the survey team visited villages and households, conducting in-depth interviews with experienced farmers to collect information on germplasm resources.


During the survey, a crop germplasm resource collection form was completed, recording detailed information such as the sample number, collection date and location (including GPS coordinates, latitude, longitude and altitude), germplasm (variety) name, category, origin and habitat, characteristics and traits of the germplasm resources, their uses, local sowing and harvesting times, cultivation area and duration, as well as information about the collector and the sample provider. Additionally, historical and cultural information related to the germplasm resources was documented, and photographs of the samples were taken for reference and preservation (Shi *et al.* 2018; Tang *et al.* 2018).

After assessing resource abundance and habitat conditions, a radial sampling method was used within a 1 km radius centred on each sampling site. Once a target sample was identified, relevant information was immediately recorded and seeds were collected. To ensure genetic diversity, at least 50 individual plants were selected from different areas of the field and a panicle was collected from each plant and stored separately. The collected seeds were dried in a ventilated and shaded environment for approximately 7–10 days to reduce moisture content and prevent mould or deterioration. After drying, the seeds were packaged in aluminium foil bags. Each sample was carefully labelled, sealed and the associated information meticulously recorded to ensure sample accuracy and to prevent mixing (Lu *et al.* 2008; Yao *et al.* 2021). From all the collected germplasm resources, a subset of samples for each sexually reproducing species was selected and sent to the cold store of the Hainan Academy of Agricultural Sciences for preservation. The number of samples preserved was determined according to the relevant standards of the Hainan Academy of Agricultural Sciences (Table S1). All resources were subjected to both short-term (4°C for 6 months) and medium-term (0°C for 3 years) cold storage. During storage, seed germination rates were monitored after 6 months and 3 years, respectively, to assess preservation effectiveness and sample viability (Lu *et al.* 2008).

## Results

### The novelty of the collected resources in this survey

A total of 1180 materials were collected during this survey, encompassing 7 families and 237 species (Table 1). None of these materials are recorded in the China National Repository. This suggests that this collection of germplasm resources is both diversity and unique. Of the germplasm collected, the Poaceae family was the most abundant, contributing 207 samples (17.54% of the total) across six genera, including *Oryza*, *Coix*, *Setaria* and *Zea* (Table 2). The Cucurbitaceae family, primarily melons, constituted the least of the germplasm with only 74 resources. The



**Figure 1.** The collected food crop germplasm resources in Hainan. (A) The geographic distribution of the collected food crop germplasm resources and (B) the number of collected food crop germplasm resources. In (A), the numbers on the left-side map correspond to the order of cities on the right, while the numbers in parentheses on the right indicate the number of collected germplasm resources. The map of Hainan Island is a modified version based on the map issued by the Ministry of Natural Resources of China, available at (<http://bzdt.Ch.Mnr.Gov.Cn/browse.Html?picid=%224628b0625501ad13015501ad2bfc0185%22>).

'Other' category encompassed a wide variety of germplasm, including fruit crops such as areca nut (*Areca catechu*), guava (*Psidium guajava*) and passion fruit (*Passiflora edulis*); The vegetable crops included taro (*Colocasia esculenta*), eggplant (*Solanum melongena*) and Chinese yam (*Dioscorea polystachya*). The economic crops were sesame (*Sesamum indicum*), camellia (*Camellia*) and large-leaf tea (*Camellia sinensis* var. *assamica*) are also included.

#### *Investigation and collection of germplasm resources of food crops in Hainan Island*

Of the 1180 germplasm resources, 330 were food crop germplasm resources, including 17 types of food crops, such as local rice, corn and soybean (Fig. 1B). The germplasm resources were found in farmlands and open spaces in front of and behind houses, with

**Table 1.** Summary of collected germplasm resources

Family	Collected varieties	Proportion (%)	Genus
Poaceae	207	17.54	6 genera including <i>Oryza</i> , <i>Coix</i> , <i>Setaria</i> , <i>Zea</i> , etc.
Fabaceae	188	15.93	20 genera including <i>Glycine</i> , <i>Cajanus</i> , <i>Arachis</i> , etc.
Moraceae	96	8.14	3 genera including <i>Artocarpus</i> , <i>Ficus</i> and <i>Morus</i>
Solanaceae	83	7.03	3 genera including <i>Solanum</i> , <i>Capsicum</i> and <i>Nicotiana</i>
Euphorbiaceae	84	7.12	3 genera including <i>Manihot</i> , <i>Ricinus</i> L. and <i>Sauropus</i>
Cucurbitaceae	74	6.27	9 genera including <i>Cucurbita</i> , <i>Cucumis</i> L., <i>Luffa</i> , etc.
Others	448	37.97	4 genera including <i>Areca</i> L., <i>Colocasia</i> schott, <i>Amaranthus</i> L. and <i>Camellia</i> L.

altitudes ranging from 2.73 to 657.12 m across 18 cities in Hainan Island (Fig. 1A).

According to experienced local farmers, many of the collected germplasm resources were characterized by stress resistance traits, including disease resistance, pest resistance and drought tolerance (Table 2). Among these materials, 28, 20, 27, 20 and 17 varieties showed disease resistance, pest resistance, drought tolerance, tolerance to poor soil conditions and high-quality traits, respectively. In addition, 11 varieties demonstrated health-promoting properties and unique applications, with 1 variety notably enriched in selenium and multiple vitamins (Table 3).

### *The changes of Hainan agriculture industrial structure*

Hainan Province conducted crop germplasm resource surveys in 1951 and 1981. A comparison with the latest survey in 2014 revealed significant changes in crop planting structure: the food crop area steadily declined, while the areas for vegetables, fruit trees and economic crops progressively increased (Fig. 2). In 1956, food crops covered 90.16% of the province's total cultivated area. This share dropped to 81.54% by 1981 and further declined to 47.74% in 2014, marking a cumulative reduction of 42.42% over 68 years. Meanwhile, the share of vegetables, fruit trees and economic crops in the cultivated area increased markedly. Fruit trees grew the most (+17.10%), followed by vegetables (+16.10%) and economic crops (+8.69%). From 1956 to 2014, Hainan's crop planting structure underwent notable shifts. In 1956, major crops consisted of food crops, economic crops and vegetables. By 1981, fruit trees replaced vegetables as a primary crop. By 2014, the structure prioritized food crops, vegetables and fruit trees. These changes illustrate Hainan's transition from a single-crop reliance to a diversified agricultural model, emphasizing the growing importance of non-food crops in the crop structure.

## **Discussion**

### *The local rice variety resources in Hainan have a long history of cultivation*

Rice is the primary staple crop in Hainan. Rice cultivation in Hainan has a history of over 2000 years, originating when ancient inhabitants began planting rice for sustenance and other uses. Early rice cultivation in Hainan was concentrated near rivers and coastal areas, where fertile soil and abundant water resources supported rice growth. Historical records show that large-scale rice cultivation on Hainan Island began during the Eastern Han Dynasty (AD 214). During this period, war and political unrest prompted many people from the Central Plains to migrate to Hainan, bringing

advanced agricultural techniques and experience. As the population grew and agricultural technology advanced, rice cultivation in Hainan expanded into mountainous and hilly areas. Resource surveys reveal that local rice varieties are highly adaptable and can be cultivated in most areas of Hainan. Surveys also indicate that the central mountainous region contains the richest diversity of local rice cultivars. This is attributed to both suitable natural conditions and the long history of mountain-based food crop cultivation, which plays a significant role in local agriculture and provides substantial benefits.

Local food crop varieties in Hainan are primarily used for cooking rice, making porridge and preparing rice rot. One collected variety, 'Jiezi rice' from Wenchang City (P469005018), is rich in selenium as well as vitamins A and B. Local enterprises have developed and marketed this rice at 35 yuan per kilogram, a price significantly higher than regular and other rice varieties, contributing to rural revitalization. Uncovering outstanding local crop varieties such as 'Jiezi rice' is also one of the key objectives of this survey.

### *The relationship between the collected germplasm resources and Hainan ethnic traditional culture, agricultural civilization and geographic environment*

Hainan Island, with its unique geographic environment and rich ethnic diversity, has long been a cradle of diverse germplasm resources. These resources are not merely biological assets; they are deeply intertwined with the island's traditional culture, agricultural civilization and natural landscapes, reflecting a harmonious coexistence between humans and nature.

The collected germplasm resources include various crops, such as upland rice, particularly the traditional Shanlan rice, which holds significant cultural and agricultural importance for ethnic groups like the Li people. Over centuries of cultivation, these crops have adapted to the island's tropical climate and diverse terrain. The Li people, renowned for their sophisticated farming methods, have developed complex systems to manage these resources, ensuring their sustainability and adaptability. For instance, Shanlan rice, a drought-tolerant variety primarily grown in mountainous areas, exemplifies the Li people's deep understanding of and adaptation to the local environment. Cultivating these crops is not merely an economic activity but also a cultural practice, deeply embedded in community rituals and festivals. One prominent example is the 'March 3rd' Festival, the most significant traditional folk celebration for the Li and Miao ethnic groups, as well as a romantic occasion for young people. Every year, on the third day of the third lunar month, Li people don festive attire and bring Shanlan rice



**Table 2.** Different characteristics of collected food crop germplasm resources in Hainan

Index	Collecting number	Variety name	Crop type	Primary characters
1	P469001002	Shanlanxiangnuo rice*	Cereal crop	Rice blast disease-resistant, drought-resistant
2	P469001012	Shanlanhong rice	Cereal crop	High quality, drought-resistant
3	P469030022	Heisishanlan rice	Cereal crop	High quality, rice blast disease-resistant and tolerance to barren soil
4	P469030023	Huangkeshanlannuo rice	Cereal crop	High-quality, rice blast disease-resistant and drought-resistant
5	P469030024	Nianmishanlan rice	Cereal crop	High quality, rice blast disease-resistant and tolerance to barren soil
6	P469027001	Shanlannuo rice	Cereal crop	High quality, rice blast disease-resistant, eurytopicity
7	P469027017	Red rice	Cereal crop	High quality, rice blast disease-resistant, eurytopicity
8	P469005016	Baojiu	Cereal crop	Sheath blight-resistant, striped stem borer-resistant, drought-resistant
9	P469005017	Chigui	Sugar crop	Sheath blight-resistant, striped stem borer-resistant, drought-resistant
10	P469026008	Red rice	Cereal crop	High quality, sheath blight-resistant, striped stem borer-resistant
11	P469025002	Glutinous rice	Cereal crop	High quality, sheath blight-resistant, striped stem borer-resistant
12	P469025004	Aromatic rice	Cereal crop	High quality, sheath blight-resistant, striped stem borer-resistant
13	P460902034	Shanlannuo	Cereal crop	High-yield high-quality drought-resistant
14	P460902035	Shanlan rice	Cereal crop	High-yield high quality drought-resistant
15	P460902036	Texiannian	Cereal crop	High-yield high quality drought-resistant
16	P460902037	Ke13	Cereal crop	High-yield high quality drought-resistant
17	2018461055	Shanlannuo	Cereal crop	Sweet
18	2018461059	Flax Shanlannuo	Cereal crop	Tall
19	2018461065	Shanlan aromatic rice	Cereal crop	Husk, feeding the birds, dwarf
20	P460203006	Black glutinous rice	Cereal crop	Purplish-black pericarp and seed coat
21	P460203025	Honghei rice	Cereal crop	Red-black rice
22	P460203026	Red rice	Cereal crop	Red rice
23	P460203028	Hongxiangnuo rice	Cereal crop	Husk, white rice, aromatic
24	P460201017	Zimang Shanlan rice	Cereal crop	Purple long grain
25	P460201019	Black nuoshanlan rice	Cereal crop	Red rice
26	P460201020	Red Shanlan rice	Cereal crop	Red rice
30	P460201021	Flax Shanlan rice	Cereal crop	Husk, white rice
31	P469024009	Purple sweet potato	Root and tuber crop	High-yield, eurytopicity
32	P469023005	Wild sorghum	Cereal crop	High quality, eurytopicity
33	2018461001	Pigeonpea	Legume crop	Powdery mildew-resistant, pod-boring pest-resistant, drought-resistant, tolerance to barren soil
34	2019461002	The seed of Job's tears*	Cereal crop	Rust disease, stem borer-resistant
35	2017461025	Big potato	Root and tuber crop	Anthrachnose-resistant, root rot-resistant, white grub-resistant, drought-resistant, tolerance to barren soil
36	P469026005	Round-grain Shanlan rice	Cereal crop	Rice blast disease-resistant, drought-resistant, tolerance to barren soil
37	P469026006	Red Shanlan rice	Cereal crop	Rice blast disease-resistant, drought-resistant, tolerance to barren soil
38	P469026007	Long-grain Shanlan rice	Cereal crop	Rice blast disease-resistant, drought-resistant, tolerance to barren soil
39	P469026013	Fragrant glutinous rice	Cereal crop	Sheath blight-resistant, striped stem borer-resistant, tolerance to barren soil
40	P469026016	Black rice	Cereal crop	Sheath blight-resistant, striped stem borer-resistant, tolerance to barren soil
41	P469026017	Black glutinous rice	Cereal crop	Sheath blight-resistant, striped stem borer-resistant, tolerance to barren soil

(Continued)

**Table 2.** (Continued.)

Index	Collecting number	Variety name	Crop type	Primary characters
42	P469026018	Paotai rice	Cereal crop	Sheath blight-resistant, striped stem borer-resistant, tolerance to barren soil
43	P469007012	Menjian upland rice	Cereal crop	Drought-resistant, eurytopicity, heat-resistant
44	P469007013	Zidie upland rice	Cereal crop	Drought-resistant, eurytopicity, heat-resistant
45	P469007017	Mennong Shanlan rice	Cereal crop	Drought-resistant, eurytopicity, tolerance to barren soil
46	2017461008	Sweet potato	Root and tuber crop	Leafroller-resistant, drought-resistant, tolerance to barren soil
47	P469023011	Local peanut	Legume crops	High-quality, Drought-resistant, eurytopicity, cold-resistant, waterlogging-resistance, heat-resistant, crown rot-resistant, damping-off-resistant, red spider mite-resistant
48	P469001006	Cassava	Root and tuber crop	Drought-resistant, bacterial wilt-resistant, angular leaf spot-resistant, mite-resistant, thrip-resistant
49	P469030004	Qiongzong ormosia	Legume crops	Rust-resistant, angular leaf spot-resistant, heat-resistant
50	P469001005	Wuzhishan maize*	Cereal crop	Leaf spot-resistant, drought-resistant
51	2018461026	Job's tears*	Cereal crop	High-quality, smut-resistant, stem borer-resistant, tolerance to barren soil
52	P469006018	Wanning pigeonpea	Legume crops	Pod-boring pest-resistant, drought-resistant, tolerance to barren soil, heat-resistant
53	2017461020	Red taro	Root crop	Soft rot-resistant, target spot-resistant, aphid-resistant, insect-resistant, drought-resistant, tolerance to barren soil
54	P469007018	Dongfang black soya bean	Legume crop	High-quality, leaf spot-resistant, root rot-resistant, leaf roller-resistant, stem borer-resistant, drought-resistant, tolerance to barren soil, heat-resistant
55	2018461006	Changjiang sorghum	Cereal crop	Seedling blight-resistant, smut-resistant, grain midge-resistant, aphid-resistant, drought-resistant, tolerance to barren soil
56	2017461002	Red sesame	Oilseed Crop	Drought-resistant, tolerance to barren soil
57	P460107095	Small white kernel	Horticultural Crops	Drought-resistant, tolerance to barren soil
58	2019461021	Millet*	Cereal crop	Drought-resistant

Note: The asterisk (\*) in the table to indicate rare and endangered varieties.

**Table 3.** Landrace with special purpose in Hainan


Index	Collecting number	Variety name	Crop type	Special purpose
1	P469005018	Jiezi rice*	Cereal crop	Selenium, vitamin A, vitamin B
2	2017461001	Podi rice*	Cereal crop	Health, treating beriberi
3	2018461031	Nuogu rice	Cereal crop	Ancestral worship
4	2017461030	Yajiao millet	Cereal crop	Lower blood pressure, strengthening the spleen and stomach
5	2017461010	Baigeng taro	Root and tuber crop group	Clearing heat and promoting diuresis
6	P469030026	The seed of Job's tears	Cereal crop	Eating, medicinal use
7	2017461005	Willow bean	Legume crop group	Treating beriberi
8	P469030010	Qiongzong sesame	Oilseed crop group	Processing raw material
9	2018461053	Huangke Shanlan sticky rice	Cereal crop group	Bird feed, brewing
10	2018461057	Make Shanlan sticky rice	Cereal crop group	Brewing
11	2018461058	Make Shanlan sticky rice	Cereal crop group	Brewing

Note: The asterisk (\*) in the table to indicate rare and endangered varieties.

and Shanlan rice wine to gather from all directions. The celebrations begin with ancestral worship, followed by singing in groups, dancing and playing musical instruments to mark the joyous season. Shanlan rice is not only a staple food for the Li people but also a crucial offering and dish during this festival. Additionally,

Shanlan rice wine, a symbol of harvest and joy, plays a vital role in expressing respect for ancestors and love for life, making it an indispensable part of the festivities.

Beyond its cultural significance, germplasm resources are closely linked to Hainan's agricultural civilization. Traditional



**Figure 2.** Planting areas of different crop types in the three survey years.

farming techniques, passed down through generations, such as intercropping and crop rotation, aim to enhance yields while maintaining soil fertility. These practices serve as a testament to the island's rich agricultural heritage. Thus, the collected resources are not just samples of biodiversity; they are living records of agricultural wisdom accumulated over centuries. Hainan's geographic environment has played a critical role in shaping these germplasm resources. The island's diverse landscapes, ranging from coastal plains to volcanic mountains, provide a wide array of ecological niches for various crops. This diversity, in turn, supports a wealth of traditional cultures, each with its unique agricultural practices and food systems. As a result, the collected germplasm resources reflect the intricate relationship between the land and its inhabitants. The germplasm resources collected in Hainan are integral to the island's cultural and agricultural identity. They represent the profound connections between ethnic communities, their traditional practices and the natural environment. As efforts to conserve and utilize these resources continue, it is essential to recognize and respect this intricate relationship to ensure that Hainan's agricultural heritage is preserved for future generations.

#### Existing challenges and suggestions in Hainan crop germplasm resource conservation

The existing challenges can be categorized into the following three aspects:

1. Decline of local varieties and challenges in resource collection. Local food crops are mainly derived from seeds saved by farmers, who often lack scientific knowledge of seed preservation. Prolonged cultivation results in significant variety degeneration, including reduced resistance, quality and yield, which leads to the gradual elimination and a risk of extinction. Furthermore, macro-level agricultural policies prioritize advantageous industries while abandoning less competitive ones, promoting monoculture and causing some local varieties to develop in a single direction. Valuable local varieties are increasingly excluded from markets and gradually vanish from production. Meanwhile, rapid advancements in agricultural science and technology have led to the development and promotion of new, superior varieties. In pursuit of maximum economic benefits, farmers have replaced older varieties that no longer meet modern agricultural demands, accelerating the loss of old germplasm resources (Zheng and Yang 2014; Liu *et al.* 2017).

2. Confusion in variety identification due to local naming practices. Hainan, as a major introduction site for food crop resources, hosts numerous local varieties. This has led to a chaotic situation

where the same name may refer to different varieties, or the same variety may have multiple names.

This naming confusion results in mixed planting of different varieties, reduced variety purity and the loss of desirable traits, ultimately leading to the decline of local varieties. Additionally, it increases the complexity of germplasm resource collection, identification and preservation (Tang *et al.* 2018). To address this, phenotypic identification of newly collected resources is essential. When funding permits, SSR fingerprinting technology should be employed for accurate identification (Liu *et al.* 2018). Combining phenotypic verification with molecular biology enables the analysis of differentiation, origins and evolutionary patterns of local resources.

3. Limitations in the food crop germplasm resource preservation system in Hainan Province. Currently, Hainan Province has only one germplasm resource repository, the 'National Tropical Crop Medium-Term Bank', which primarily stores tropical crop resources. The Hainan Academy of Agricultural Sciences has independently funded the construction of a crop cold storage facility. However, due to prolonged construction time and insufficient funding, the facility's equipment and infrastructure remain inadequate. Compared with advanced domestic germplasm repositories, there is a significant gap, limiting the preservation capacity of crop germplasm resources.

It is recommended to prioritize the major needs of agricultural innovation and modern seed industry development. Efforts should focus on comprehensive collection, proper preservation, detailed evaluation, active innovation and shared utilization of germplasm resources. Medium- and long-term storage facilities should be constructed, and germplasm repositories (gardens) should be improved. Additionally, wild, high-quality and rare crop germplasm resources in Hainan and surrounding areas should be systematically collected and preserved. Establishing a complete germplasm resource protection system is critical to ensuring long-term preservation and utilization of crop germplasm resources.

Additionally, *in situ* conservation is a critical component of a comprehensive conservation strategy, complementing *ex situ* conservation to ensure the complete protection of germplasm resources. *In situ* conservation involves preserving crop varieties in their natural habitats, such as traditional farming systems or natural ecosystems. This method allows the resources to continue evolving under local environmental conditions, maintaining their adaptability and resilience. For example, in Hainan, traditional crops like Shanlan rice are cultivated by the Li people in mountainous areas. These practices not only preserve the genetic diversity of the crops but also sustain the cultural and agricultural

heritage of the local communities. Additionally, *in situ* conservation supports the dynamic interaction between crops and their environment, which is essential for long-term sustainability. While *ex situ* conservation (e.g. seed banks and germplasm repositories) provides a secure backup for germplasm resources, it cannot fully capture the ecological and cultural context in which these resources thrive. *In situ* conservation complements this by preserving the living relationship between crops, their environment and the traditional knowledge associated with their cultivation. Together, these approaches ensure a more holistic and robust conservation strategy.

4. A comprehensive and systematic characterization of food crop germplasm resources is essential. Currently, germplasm resource characteristics are primarily obtained through discussions with experienced local farmers, relying on their long-term cultivation practices. However, no in-depth investigation of these characteristics has yet been conducted. Future works will focus on detailed trait investigations and comprehensive documentation of the collected food crop germplasm resources. This will involve systematically recording phenotypic characteristics, including growth habits, pest and disease resistance, yield potential and quality traits. Such investigations will facilitate precise identification and evaluation of germplasm, ensuring valuable traits are preserved. Modern molecular biology techniques, including genome sequencing and SSR fingerprinting, will complement phenotypic data, offering deeper insights into the genetic basis of key traits.

Efforts will also focus on establishing a standardized database integrating phenotypic, genotypic and ecological data for all collected resources. This database will underpin future breeding programs, allowing researchers to efficiently select and utilize germplasm with desirable traits to develop improved crop varieties. These efforts will not only strengthen the germplasm resource preservation system but also promote the sustainable development of agriculture and the seed industry in Hainan Island.

## Conclusion

From 2017 to 2021, the Hainan investigation team systematically surveyed and collected germplasm resources of food crops in Hainan Province, yielding significant findings that enhance the conservation and utilization of agricultural biodiversity in the region. The study documented the distribution of food crop germplasm resources across districts and identified key traits, including superior quality and tolerance to drought and poor soil conditions. The collection of 330 food crop germplasm resources, including rare and endangered varieties, has expanded the genetic pool for future breeding programmes and established a strong foundation for developing new crop varieties.

**Supplementary material.** The supplementary material for this article can be found at <https://doi.org/10.1017/S1479262125100063>

**Author contributions.** Q.T. conceived and designed this research. L.Z. and X.Y. undertook material collection and phenotypic identification. G.C. wrote the original draft. G.C., J.X., H.Z., L.T., L.L. and Q.W. contributed to materials collection, preservation and classification. All authors have read and agreed to the published version of the manuscript.

**Funding statement.** This work was supported by the Hainan Province Science and Technology Special Fund (ZDYF2024KJTPY027), the Department of Agriculture Species and Variety Resources Protection Fund Project (111821301354052033), China Key R&D Program Project (2021YFD1200102-01). The Technical Innovation Professional Project of the Provincial Scientific

Research Institutes in Hainan Province (jscx202003), Key R&D projects in Hainan Province (ZDYF2024XDNY165) and Coordinated scientific research projects of HAAS(2024-LZSQN005).

**Competing interests.** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as potential conflicts of interest.

## References

- Chen H-W, Li L, Liu L-J, Liu C-Y and Wan Z-H (2018) Survey and analysis of crop germplasm resources in Qichun County of Hubei Province. *Hubei Agricultural Sciences* 57(24), 4. In Chinese.
- Fan W-Y, Liu Z-R, Yun Y, Tang Q-J, Zhou S-Z, Xiao X-R, Zheng X-M and Yang Q-W (2023) Collection and preliminary identification of germplasm resources resistant to bacterial blight of wild rice from Hainan Province. *Journal of Plant Genetic Resources* 24(1), 117–125. In Chinese.
- Lei W-H, Ye L-R, Tan X-J, Zhou H-Y and Qin L-Q (2023) Analysis and discussion on the Third National Survey and Collection of Crop Germplasm Resources in Shunchang County, Fujian Province. *China Seed Industry* 2, 62–64. In Chinese.
- Li T-T and Yang Z (2022) Analysis and discussion on survey and collection of crop germplasm resources in Jinan City. *China Seed Industry* 6, 42–45. In Chinese.
- Li Y, Li Y-H, Yang Q-W, Zhang J-P, Zhang J-M, Qiu L-J and Wang T-Y (2015) Genomics-based crop germplasm research: Advances and perspectives. *Scientia Agricultura Sinica* 48(17), 3333–3353. In Chinese.
- Liang S-C, Chen C-B, Liang Y-T, Yang Q-W, Qiao W-H, Zeng H-Z, Xu Z-J, Huang J and Zhang Y (2013) Collection and conservation strategies of *Oryza officinalis* wall in Guangxi. *Journal of Plant Genetic Resources* 14(6), 991–995. In Chinese.
- Liu X-H, Deng J, Li X-X, Yang J-G, Yang S-Z, Wang T-H and Yu Y-H (2017) The progress of conservation and innovation of crop germplasm. *The Progress of Conservation and Innovation of Crop Germplasm* 18(5), 916–922. In Chinese.
- Liu Z-H, Lin Z-J, Li H-W, Xu Y-Q, Li G-L, Qiu Y-X, Qiu S-X and Tang H (2018) Analysis of genetic diversity, origin and evolution of sweetpotato (*Ipomoea batatas* (L. Lam.) by SRAP. *Journal of Plant Genetic Resources* 19(3), 468–477. In Chinese.
- Lu X-X, Chen S-P and Liu X (2008) Technical regulation on conservation for crop germplasm resources. *China Agriculture Press*. In Chinese.
- Ministry of Agriculture of the PRC, National Development and Reform Commission, Ministry of Science and Technology of the PRC (2015) National medium and long-term development plan for protection and utilization of crop germplasm resources (2015-2030). *Bulletin of Ministry of Agriculture of the People's Republic of China* 8, 4–9. In Chinese.
- Shi H-X, Chen Z-H, Li Z-Y, Li P, Yang W-C, Shi T, Liu W-H and Wu X-H (2018) Preliminary survey and collection of germplasm resources of *Bromus* L. in Tibet Autonomous. *Journal of Plant Genetic Resources* 19(4), 612–618. In Chinese.
- Tang Q-J, Yan X-W, Yang G-F, Zhong Z-F and Tang L-Q (2018) Collection and survey of Shanlan upland rice resources in Hainan and recommendations for its development. *Hybrid Rice* 33(1), 20–24. In Chinese.
- Wang G-Y, Chen T-Q, Huang D and Cai X (1992) *Collection of Investigation on Crop (Plant) Germplasm Resources in Hainan Island*. Beijing: Agricultural Publishing House, pp. 13–19.
- Wang S (2002) Conservation and utilization of crop germplasm resources in China. *China Seed Industry* 10, 8–11. In Chinese.
- Yao Z-F, Wu R-X, Zhang X-J, Dai Z-Y, Yang Y-L, Huang L-F, Liu J and Fang B-P (2021) Systematic field collection and identification of sweetpotato resources in Guangdong. *Journal of Plant Genetic Resources* 22(6), 1498–1508. In Chinese.
- Yun Y, Tang Q-J, Yan X-W, Meng W-D, Wang X-N and Lin Y-Z (2015) Field survey and conservation of wild rice resources in Hainan. *Journal of Plant Genetic Resources* 16(4), 715–719. In Chinese.
- Zeng Y-H, Xie H-X, Cheng W-D, Jiang Y-F, Zhou J-G and Xie X-D (2019) Systematic field survey and collection of maize germplasm resources in Guangxi. *Journal of Plant Genetic Resources* 20(3), 654–661. In Chinese.



- Zhang Y-G, Zhang G-L, Gong Z-T and Wan-gang DENG** (2005) Soil types, quality characteristics and suitability of crops under the different geological conditions of Hainan Island. *Quaternary Sciences* **25**(3), 389–395. In Chinese.
- Zheng D-S and Gao A-N** (2016) Proposal concerning conservation and sustainable use of agrobiological resources in minority area in Guizhou Province. *Journal of Plant Genetic Resources* **17**(5), 957–959. In Chinese.
- Zheng D-S, Gao A-N, Li L-H and Liu X** (2016) Rare germplasm resources of crop and wild relatives of minority area in Guizhou Province. *Journal of Plant Genetic Resources* **17**(3), 570–576. In Chinese.
- Zheng D-S and Yang Q-W** (2014) Resources of wild relatives of crops in China. *Journal of Plant Genetic Resources* **153**(1), 1–11. In Chinese.