

STEADY AND FLUCTUATING PARTS OF SUN'S INTERNAL MAGNETIC FIELD

M. H. GOKHALE and K. M. HIREMATH  
 Indian Institute of Astrophysics, Bangalore 560 034, India

**ABSTRACT** We have modeled the 'steady' part of sun's internal poloidal magnetic field in the form of a central dipole and a central hexapole with strengths  $(0.6 \pm 0.1) B_{R_0}^3$  and  $(0.16 \pm 0.05) B_{R_0}^5$  embedded in an asymptotically uniform field  $B_0$ . A small deviation from isorotation seems to indicate: (i) a slow build-up of toroidal field near the base of the convection zone and (ii) presence of torsional MHD perturbations in the outer radiative core, with latitudinal structure and time scales which may be similar to those of solar activity.

1. INTRODUCTION

In this paper we determine a likely configuration of an axi-symmetric poloidal magnetic field that can remain in a 'steady' state with the helio-seismologically determined internal rotation of the sun.

According to Cowling's theorem, the resistive and the inductive terms of this steady part of the real field must vanish separately. Thus the steady part of the internal field must be in isorotation (Ferraro 1937) with the steady part of the rotation of the plasma. This requires existence of functional relation between the 'observed' rotation rate  $\Omega(r)$  at a point  $r$  and the flux function  $\phi(r)$  of the 'steady' poloidal field.

2. METHOD

The relation between the rotation velocity  $\Omega(r, \theta)$  and the magnetic flux due to the internal and external sources is assumed to be linear, so that :

$$\Omega(r, \theta) = \Omega_0 + \Omega_1 x^2 \sin^2 \theta + \Omega_2 [(2\mu_1 x^{-1} + 4\mu_3 x^{-3} + \dots) \sin^2 \theta + (-5\mu_3 x^{-3} + \dots) \sin^4 \theta + \dots] \tag{1}$$

where  $x$  is the distance from the center in the unit of the solar radius  $R_0$ ,  $\vartheta$  is the co-latitude,  $\Omega_0, \Omega_1, \Omega_2$  are constants and  $\mu_\ell$ ,  $\ell=1, 3, 5, \dots$  are the strengths of the multipoles, in the standard expansion of the magnetic potential, in units of  $B_0 R_0^{\ell+2}$  where  $B_0$  is the asymptotically uniform field at large distances. Isorotation requires  $\Omega_1 = \Omega_2$ .

### 3. THE DATA USED

We use the rotation data from the internal rotation  $\Omega(r)$ , given by Christensen-Dalsgaard and Schou (1988). We determine the coefficients  $\Omega_0, \Omega_1, \mu_1, \mu_3$ , etc., by obtaining least square fits for successive combinations of terms in equation (1) using the data points in convective envelope (CE), outer radiative core (ORC) and combination of these data ('ORC+CE').

### 4. RESULTS AND CONCLUSION

The least square fit to the 'observed'  $\Omega(r, \vartheta)$  in 'CE' yields:




Fig.1. Structure of the 'steady' part of the poloidal field given by  $\mu_1=0.6$  and  $\mu_3=0.16$  in one quadrant of a meridian plane. The field lines A, ..., J correspond to various flux values from 0.5 to 2.1 units. The continuous lines represent branches of the separatrix :  $\phi = 1.84$  units.

$\Omega_0 = 326 \pm 12$  nHz/unit flux,  $\Omega_1 = \Omega_2 = 68 \pm 11$  nHz/unit flux,  $\mu_1 = 0.6 \pm 0.1$  and  $\mu_3 = 0.16 \pm 0.05$ . The strength  $B_0$  cannot be determined from this fit itself, but can be independently estimated to be in the range of  $10^4 - 1G$ .

The resulting field structure is shown in Fig.1. Field directions will be relative to that of  $B_0$  which is presently unknown. Outer part of the structure will be deformed by several processes. The field structure also contains a separatrix. Revolution of its branch running pole-wards near the center, and along the inner boundary of 'CE' in low latitudes, gives the critical surface  $S_*$ , which rotates with angular velocity  $\Omega_* = 430$  nHz. Inside  $S_*$  the rotation is 'rigid', with  $\Omega(r, \theta) \approx \Omega_*$ . However the fit in 'ORC+CE' indicates that near  $S_*$  it can be approximated by equation (1) with  $\Omega_2 = 75$  nHz/unit flux. The 'non-isorotation' due to the difference  $\sim 9^7$  nHz/unit flux between  $\Omega_1$  and  $\Omega_2$  gives a time scale  $\sim 10^8 - 10^9$  year for 'winding' of the poloidal field into a toroidal field  $B_T \sim 10^6 G$  near  $S_*$ . The fit in 'ORC+CE' also yields a small  $\mu_5$ , implying presence of time dependent torsional MHD perturbations, in ORC, with dominant term  $\ell = 5$ . For these perturbations the non-isorotation near  $S_*$  yields 'periods' in the range  $\sim 1-100$  year depending upon the ratio of perturbations in the toroidal and poloidal fields. It is known that in solar magnetic cycle the dominant term is  $\ell = 5$  (Gokhale, et.al 1992 ; Stenflo, J.O. 1988).

On long time scales, the field may also provide outward transport of angular momentum and magnetic flux.

## REFERENCES

- Christensen-Dalsgaard, J., & Schou, J. 1988, in Seismology of the Sun and Sun-like Stars, ed. V. Domingo and E.J. Rolfe ESA : SP-286, p.149.
- Ferraro, V.C.A. 1937, M.N.R.A.S., 97, 457.
- Gokhale, M. H., Javaraiah, J., Kutty, K. N., & Varghese, B. A. 1992, Solar Phys, 138, 35.
- Stenflo, J. O. 1988, Astrophys. & Space Sci., 144, 321.