NOTES FOR AUTHORS

Papers submitted for publication in the *Bulletin of Entomological Research* should be sent, in duplicate, to the Director, C·A·B International Institute of Entomology, 56 Queen's Gate, London SW7 5JR. They must deal with original research concerning insects, mites or ticks of economic importance in the agricultural, medical or veterinary fields; they must be of more than purely local interest, and taxonomic papers will not be accepted unless the organisms dealt with are of definite importance in the field of applied entomology. Review articles may also be submitted, but will normally be by invitation. Papers must not have been published or accepted for publication elsewhere. They will normally be published in order of their receipt in definitive form.

For general guidance on presentation and style, including the preparation of illustrations, contributors are urged to consult *Writing Scientific Papers in English. An ELSE-Ciba Foundation Guide for Authors* (by Maeve O'Connor & E. Peter Woodford). 116 pp., Amsterdam, Associated Scientific Publishers, 1975, ISBN 90-219-4035-3.

Text

Papers should be typewritten (double spaced, with ample margins and on one side of the paper only) and a current number of the *Bulletin* should be consulted for details of style and lay-out. Titles should be as brief as is consistent with clarity. Abbreviations should conform to *British Standard 1991*, *Part I*, *1967*, and are alike in singular and plural; stops will be used only where ambiguity might occur, *e.g.* 'in.' not 'in' for inch, but 'in²' for square inch. Metric units are preferred; if non-metric units are used, metric equivalents should be given at least once in each article.

Tables

Tables should be reduced to the simplest form, and should not be used where text or illustrations give the same information. They should be submitted on separate sheets at the end of the article and must on no account be of a size or form that will not conveniently fit into the page of the *Bulletin*.

Illustrations

Line drawings and graphs must be in jet-black indian ink, preferably on Bristol-board or tracing cloth. Graphs may be drawn on paper ruled in pale blue or light grey; paper having green, red, yellow or brown lines should not be used. Any stippling or hatching, as well as any numerals or lettering, on the text-figures must be completed, by the author, in a form suitable for reproduction; this work is *not* undertaken by the printer. Text-figures, when reproduced, cannot exceed 12·7 cm in width; if, therefore, the original drawings are larger, the lettering on them must be in proportion, so that it is still legible when reduced; special attention should be paid to the relative scales of lettering and line thicknesses in relation to the figure as a whole. Photographs for reproduction as figures are admissible when they are a real contribution to the text. Captions to figures should be typed on a separate sheet.

Abstract

Each paper must contain a carefully prepared and accurate informative abstract that shall be complete in itself and intelligible without reference to the text or figures. It should not exceed 250 words, and should be included at the beginning immediately after the title, author and author's affiliation.

References

References must be based on the name and year system and conform to the present usages of the *Bulletin*. Titles of journals should be abbreviated as in the 4th edition of *The World List of Scientific Periodicals*. References to unpublished information should normally be included only in the text—(A. B. Jones, pers. comm.) or (A. B. Jones, Control of sandflies by insecticides. Unpublished report. Ross Institute, London, 1950).

Offprints

Free copies of offprints are allowed as follows: One author—20 copies, two or more authors—30 copies. Further copies may be obtained on payment, and the number required should be specified when the paper is submitted.

DECEMBER 1988 Vol. 78(4)

BULLETIN OF ENTOMOLOGICAL RESEARCH

(World List Abbreviation: Bull. ent. Res.)
ISSN 0007-4853

Original articles	PAGE
LE SUEUR, D. & SHARP, B. L. The breeding requirements of three members of the Anopheles gambiae Giles complex (Diptera: Culicidae) in the endemic malaria area of Natal, South Africa	549
Cox, J. M. & Wetton, M. N. Identification of the mealybug <i>Planococcus halli</i> Ezzat & McConnell (Hemiptera: Pseudococcidae) commonly occurring on yams (<i>Dioscorea</i> spp.) in Africa and the West Indies	561
BARCLAY, H. J. Models for combining methods of pest control: food-baited and pheromone-baited traps containing either insecticide or chemosterilant	573
GREEN, C. H. The effect of colour on trap- and screen-orientated responses in Glossina palpalis palpalis (Robineau-Desvoidy)(Diptera: Glossinidae)	591
UMEH, E. D. N. Development, oviposition, host feeding and sex determination in Epidinocarsis lopezi (De Santis) (Hymenoptera: Encyrtidae)	605
BLACKWELL, A. The significance of proleg disruption by chlordimeform to sublethal control of larvae of <i>Pieris brassicae</i> (L.) (Lepidoptera: Pieridae) in the field	613
GILLETT, S. D. Solitarization in the desert locust, Schistocerca gregaria (Forskål) (Orthoptera: Acrididae)	623
RIDSDILL-SMITH, T. J. & MATTHIESSEN, J. N. Bush fly, Musca vetustissima Walker (Diptera: Muscidae), control in relation to seasonal abundance of scarabaeine dung beetles (Coleoptera: Scarabaeidae) in south-western Australia	633
JEPSON, P. C. & HEALY, T. P. The location of floral nectar sources by mosquitoes: an advanced bioassay for volatile plant odours and initial studies with <i>Aedes aegypti</i> (L.) (Diptera: Culicidae)	641
Healy, T. P. & Jepson, P. C. The location of floral nectar sources by mosquitoes: the long-range responses of <i>Anopheles arabiensis</i> Patton (Diptera: Culicidae) to <i>Achillea millefolium</i> flowers and isolated floral odour	651
BARKER, G. M., PRESTIDGE, R. A. & POTTINGER, R. P. Reproductive phenology of Listronotus bonariensis (Kuschel) (Coleoptera: Curculionidae) in northern New Zealand pastures	659
GILBERT, N. & COAKER, T. H. Differential survival of British and Mediterranean strains of <i>Pieris rapae</i> (L.) (Lepidoptera: Pieridae) on different brassica cultivars	669
McDonald, G. & Smith, A. M. Phenological development and seasonal distribution of the Rutherglen bug, Nysius vinitor Bergroth (Hemiptera: Lygaeidae), on various hosts in Victoria, south-eastern Australia	673
Powell, W. & Wright, A. F. The abilities of the aphid parasitoids Aphidius ervi Haliday and A. rhopalosiphi De Stefani Perez (Hymenoptera: Braconidae) to transfer between different known host species and the implications for the use of alternative hosts in pest control strategies	683
AGOUNKÉ, D., AGRICOLA, U. & BOKONON-GANTA, H. A. Rastrococcus invadens Williams (Hemiptera: Pseudococcidae), a serious exotic pest of fruit trees and other plants in West Africa	695
NARASIMHAM, A. U. & CHACKO, M. J. Rastrococcus spp. (Hemiptera: Pseudococcidae) and their natural enemies in India as potential biocontrol agents for R. invadens Williams	703
WILLINK, E. & MOORE, D. Aspects of the biology of Rastrococcus invadens Williams (Hemiptera: Pseudococcidae), a pest of fruit crops in West Africa, and one of its primary parasitoids, Gyranusoidea tebygi Noyes (Hymenoptera: Encyrtidae)	709
HERATH, P. R. J., JAYAWARDENA, K. G. I., HEMINGWAY, J. & HARRIS, J. DDT resistance in <i>Anopheles culicifacies</i> Giles and <i>A. subpictus</i> Grassi (Diptera: Culicidae) from Sri Lanka: a field study on the mechanisms and changes in gene frequency after cessation of DDT spraying	717
© C·A·B International, 1988	

All rights reserved. No part of this publication may be reproduced, in any form or by any means, electronically, mechanically, by photocopying, recording or otherwise, without prior permission of the copyright owner.