

and conscientiousness; and finally between self-transcendence and agreeableness. As a result, due to their relationship with temperamental dimensions of psychobiological model, the FFM domains could be related to brain monoaminergic activities.

Keywords Personality; Personality Inventory; Temperament; Character

Disclosure of interest The authors declare that they have no competing interest.

Further reading

Costa PT, McCrae RR. The NEO Personality Inventory Manual Revised. Odessa, FL: Psychological Assessment Resources, 1990.

Cloninger CR, Svrakic DM, Przybeck TR. A psychobiological model of temperament and character. *Arch Gen Psychiatry* 1993;50, 975–90. De Fruyt F., Van De Wiele L., Van Heeringen C. Cloninger's psychobiological model of temperament and character and the Five-Factor Model of personality. *Pers Individ Dif* 2000;29:441–52.

<http://dx.doi.org/10.1016/j.eurpsy.2015.09.231>

P042

Validity of the results of psychological autopsies in suicide prevention policy

M. Benabbas^{1,*}, O. Benelmouloud²

¹ HMRUC/5^e RM, Constantine, Algeria

² EHS de psychiatrie, faculté de médecine, université de Constantine 3, Constantine, Algeria

* Corresponding author.

E-mail address: benabbasmalik@gmail.com (M. Benabbas)

Psychological autopsy is a clinical tool and a research tool likely to clarify the circumstances of a death in cases of suicide, of homicide or suspicious death (Fig. 1.1). In our analysis of the methodology and its application, the purpose specifically focus on autopsies psychological in cases of suicide. Psychological autopsy focuses on the psychological aspects of occurrence of death. It incorporates the field of Suicidology. Its goal is to understand the circumstances as the state of mind of the victim of his act. This type of method includes a reconstruction lifestyle behaviors and events experienced by the individual. Thus, the psychological autopsy is a retrospective analysis. The aim of the psychological autopsy is to collect as much information about the circumstances of the death and to update the reasons for suicide eventually help reveal risk factors (Hawton et al., 2003) This is an epidemiological study of suicides in Eastern Algeria (15 wilayas) through psychological autopsies (2003 to 2010) or more variables were studied to establish a typical profile of suicide in Algeria. The studied variables were: age, sex, occupation, place of residence, the existence of life events, psychiatric history and possibly TS history, source of information (to whom we have collected information: father, mother, brother, sister. . .) and the proceeds used for suicide. In total we identified 1263 cases of suicide with age 15 and over occurred in populations of the North-eastern Algeria during the period from 2003 to 2010. It concludes with the emergence of some variables that may be risk factors namely Age between 30 and 45 years, male gender, social and financial difficulties especially bad life, presence of a psychiatric diagnosis on axis 1 of DSM IV and finally the lack of access to primary care in urban areas. The evaluation of the feasibility and reliability of psychiatric autopsies in Algeria remains lapsed due to non-standardized measurement tools and the difficulty of their implementation on the ground. Finally, the authors highlight the prevalence per 100,000 population per city and the average prevalence for all of East of Algeria.

Keywords Suicide; Psychological autopsy; Prevalence

Disclosure of interest The authors declare that they have no competing interest.

Further reading

Cavanagh JT, Owens DG, Johnstone EC. Suicide and undetermined death in south east Scotland. A case-control study using the psychological autopsy method. *Psychol Med* 1999;29:1141–9.

Cavanagh JT, Carson AJ, Sharpe M, Lawrie SM. Psychological autopsy studies of suicide: a systematic review. *Psychol Med* 2003;33:395–405.

Chambers WJ, Puig-Antish J, Hirsch M, et al. The assessment of affective disorders in children and adolescents by semistructured interview: test-retest reliability of the schedule for affective disorders and schizophrenia for school-age children, present episode version. *Arch Gen Psychiatry* 1985;42:696–702.

Cheng AT, Chen TH, Chen CC, Jenkins R. Psychosocial and psychiatric risk factors for suicide. Case-control psychological autopsy study. *Br J Psychiatry* 2000;177:360–5.

<http://dx.doi.org/10.1016/j.eurpsy.2015.09.232>

Clinique

P043

Quelle évaluation psychiatrique dans le bilan pré-chirurgie bariatrique ?

F. Boulet

CHU de Nîmes, Nîmes, France

Adresse e-mail : fabrice.boulet@chu-nimes.fr

L'évaluation psychiatrique dans le bilan pré-chirurgie bariatrique ne bénéficie pas encore de consensus. Pourtant, la recherche des contre-indications et la rédaction d'un certificat sont obligatoires dans le parcours de tout patient voulant bénéficier d'une chirurgie bariatrique. Cette session se propose de rapporter une revue de la littérature internationale concernant les différents bilans proposés et les limites d'une telle évaluation afin de pouvoir mieux aborder ce type de consultation.

Mots clés Évaluation psychiatrique ; Chirurgie bariatrique ; Revue littérature

Déclaration de liens d'intérêts Les auteurs déclarent ne pas avoir de liens d'intérêts.

Pour en savoir plus

Schlick A, Wagner SA, Mühlhans B, Horbach T, Müller A, Mitchell JE, et al. Agreement between clinical evaluation and structured clinical interviews (SCID for DSM-IV) in morbidly obese pre-bariatric surgery patients. *Psychother Psychosom Med Psychol* 2010;60(12):469–73. [Epub 2010 Apr 16. *Surg Obes Relat Dis* 2010;6(6):628–34. Epub 2010 Feb 6. Congruence between clinical and research-based psychiatric assessment in bariatric surgical candidates].

Mitchell JE, Steffen KJ, de Zwaan M, Ertelt TW, Marino JM, Mueller A. Source Neuropsychiatric Research Institute, Fargo, North Dakota, USA.

Vergouwen AC, Aajoud S, van Wagenveld BA, van Tets WF, Honig A. Effects of bariatric surgery not affected by psychiatric comorbidity: a systematic review of studies of morbidly obese patients. *Ned Tijdschr Geneesk* 2010;154:A1678.

<http://dx.doi.org/10.1016/j.eurpsy.2015.09.233>

P044

Hikikomori, un concept pertinent pour penser le retrait des jeunes au domicile ?

A. Couillet*, N. Chauliac, P. Depraz, J.-L. Terra
Hôpital le Vinatier, Bron, France

* Auteur correspondant.

Adresse e-mail : audrey.couillet@ch-le-vinatier.fr (A. Couillet)

Le terme japonais hikikomori désigne un adolescent ou un jeune adulte, le plus souvent de sexe masculin, passant la quasi-totalité