



KALPANA DEIN, GILL LIVINGSTON AND CHRISTOPHER BENCH

'Why did I become a psychiatrist?': survey of consultant psychiatrists

AIMS AND METHOD

To find out why consultant psychiatrists chose psychiatry as a career. A questionnaire was developed and posted to 87 consultant psychiatrists in substantive posts within a London psychiatric training scheme.

RESULTS

The survey had a response rate of 83% (72 out of 87). The majority of

consultants ($n=40$) chose psychiatry as a career after leaving medical school. The most important reasons cited were empathy with the patient group (36.1%), the interface of psychiatry with the neurosciences (25%), the better working conditions expected in psychiatry (20.8%) and medical school teaching of the subject (19.4%).

CONCLUSIONS

The study highlights the need for recruitment efforts after medical school. The findings also reflect the lasting influence of medical school exposure to psychiatry. Interventions for improving recruitment in psychiatry are suggested. The under-recruitment of British medical graduates is masked by overseas recruitment into the specialty.

In 2005, 2% of consultant posts in psychiatry in England were unfilled, with a further 15% of posts filled by locums (13th Annual Census of Psychiatric Staffing 2005, Royal College of Psychiatrists, <http://www.rcpsych.ac.uk/training/census.aspx>). In 2003 psychiatry accounted for 29% of all consultant vacancies in all specialties in England (364 out of 1264; Department of Health, 2003). The overall shortfall of doctors has been partially addressed by increasing the number of medical students and medical schools, but this does not address the more specific issue of shortage within the specialties (Maidment *et al*, 2003). Goldacre *et al* (2004) found that only 4.3% of all respondents in the first year after their graduation signified that psychiatry was their first choice of long-term career, and there had been very little change between 1974 and 2000. Psychiatry was the least popular clinical specialty in a London medical school (Rajagopal *et al*, 2004).

Several studies have considered factors that lead to the choice of psychiatry as a specialty. These have been carried out among potential psychiatrists in their pre-medical years (Maidment *et al*, 2003), in medical school (Schumacher, 1964; Wilkinson *et al*, 1983; Maidment *et al*, 2003), during pre-registration house jobs (Creed & Goldberg, 1987), and among senior house officers (SHOs; Clarke-Smith & Tranter, 2002; Maidment *et al*, 2004). They have found that career plans change both during and after medical school. There has been a suggestion that the most specialty switching takes place in psychiatry (Held & Zimet, 1975). In addition there is a high rate of 'attrition' of SHOs in psychiatry, who do not progress to higher psychiatric training or to consultant grade posts (Cox, 2000).

A study among consultant psychiatrists would indicate the reasons why people become psychiatrists, and might suggest rational interventions for increasing recruitment and retention. To date there have been few studies of psychiatrists at this stage of their career (Prins, 1998). Our study of consultant psychiatrists aimed to explore their reasons for choosing the specialty and to

help clarify the motivation of people who became psychiatrists, as opposed to those who intend to become psychiatrists and may change their mind.

Method

We designed a structured questionnaire using a Likert scale incorporating factors that had been suggested to relate to recruitment in psychiatry in a systematic review of current literature (Wilkinson *et al*, 1983; Prins, 1998; Baxter *et al*, 2001; Galeazzi *et al*, 2003; Maidment *et al*, 2003; McParland *et al*, 2003). We interviewed six consultant psychiatrists and conducted a focus group with SHOs in psychiatry to find out whether we had covered the entire range of reasons; this resulted in the addition of further questions. The questionnaire was then piloted among nine consultant psychiatrists in one of the mental health units within a London psychiatric training scheme, resulting in further changes in order to clarify some questions. The questionnaire was modified at the suggestion of the West London Mental Health Ethics Committee to include an open question, and was then sent to 87 consultant psychiatrists in substantive posts in a London psychiatric training scheme. Locum consultants were not included in the study as non-consultant grade doctors can cover these posts.

Analysis

Data were analysed using descriptive statistics and SPSS version 6 for Windows. The three most important reasons for choosing psychiatry were grouped together in order to have an adequate number of psychiatrists to compare results between groups. Associations between demographic variables and the reasons were assessed using the χ^2 test. As this was an exploratory study where associations between variables were considered, the level of statistical significance was set at $P=0.01$ to avoid spurious results.



education & training

Results

Participants

Out of a total of 87 consultants surveyed, 72 (82.7%) returned completed questionnaires; 68% of participants were male ($n=49$); years of experience working as a psychiatrist ranged from 10 to 40 years (median 21) starting from SHO level or equivalent. The most common sub-specialty was general adult psychiatry (41.6%), followed by old age psychiatry (13.2%), child and adolescent psychiatry (9.7%), learning disability psychiatry and forensic psychiatry (6.9% each). There were 64 participants (90%) who worked full-time, 6 (8.38%) worked part-time and 2 who did not respond to the question. All but 1 of those who worked part-time were female; 51 participants (70.8%) described themselves as 'clinicians only', 12 (16.7%) were also medical managers, and 5 (7%) were clinicians and academics. Three of the respondents were psychotherapists, and of these, two were also medical managers; one participant was a clinician, academic and a medical manager.

Stage of choosing psychiatry

The majority of the participants ($n=40$, 55.5%) chose psychiatry as a specialty after leaving medical school; 30.5% ($n=22$) decided during medical school and 8.3% ($n=6$) decided on a career in psychiatry before medical school. Four participants did not respond to this question.

Country of primary medical qualification

There were 44 participants who qualified as doctors in the UK (61.1%) and 38.9% ($n=28$) who qualified overseas. Of the latter, 9 qualified in South Asia, 7 in Western Europe, 6 in Africa, 4 in Australia and New Zealand, and 2 in South America.

The 'most important reasons' for choosing psychiatry as a career

The 'most important' reasons for choosing psychiatry as a career are shown in Table 1; 36.1% ($n=26$) chose psychiatry as a career because of their empathy for those with a mental disorder; 25% ($n=18$) because of the interface of psychiatry with the neurosciences; 20.8% ($n=15$) because of the expectation of better working conditions in psychiatry; and 19.4% ($n=14$) because of medical school teaching.

Participants were asked to indicate other reasons which influenced their choice of psychiatry. Other reasons cited were a holistic approach ($n=3$), an interest in people ($n=10$), psychodynamic and psychological reasons ($n=3$), an interest in the subject ($n=8$), the importance of team work ($n=4$) and alternative treatment modalities.

There were no significant differences in the importance of medical school teaching as a factor in recruitment among those who decided on the specialty before,

Table 1. 'Most important' reasons for choosing psychiatry

'Most important' factors influencing the choice of psychiatry	Participants <i>n</i> (%)
Psychiatric teaching received as a medical student	14 (19.4)
Exposure to psychiatry as a house officer	2 (2.8)
Psychiatrists met during medical training	10 (13.9)
Members of family working in psychiatry	2 (2.8)
Patients met during medical training (who were neither family or friends)	8 (11.1)
Empathy for patients with a mental disorder	26 (36.1)
Personal experience of mental disorder in yourself, family or friends	4 (5.6)
Desire to understand yourself better	5 (6.9)
Better working conditions expected in psychiatry (e.g. easier progress to consultant jobs and the working hours more compatible with a family)	15 (20.8)
Possibility of treating mental disorder through alternative approaches (e.g. psychotherapy, an anthropological understanding)	4 (5.6)
Interface of psychiatry with the social sciences (e.g. philosophy, anthropology or law)	10 (13.9)
Interface of psychiatry with the neurosciences	18 (25)
Disenchantment with other medical specialties	7 (9.7)
Sense of fulfilment expected from seeing patients improve	11 (15.3)

during or after medical school. There were no differences in importance of 'better working conditions' among male and female psychiatrists. There were no differences based on the country of primary qualifications.

Discussion

Most study participants decided on psychiatry after medical school and nearly a quarter had entered psychiatry after working in other specialties. This is similar to findings in Canada (Cameron & Persad, 1984) where 58.1% decided after medical school and 27% came from family practice. This study also confirms the extent to which overseas recruitment continues to mask the long-standing under-recruitment of British medical graduates into psychiatry. Recently, there has been an increase in the numbers of British medical graduates. It is premature to assume that the foundation year changes would lead to an increase in the number of medical graduates interested in psychiatry compared with other specialties.

It is likely that these results can be generalised to the whole of England as the study population was representative of the consultant population in the UK in terms of gender and choice of sub-specialty (Royal College of Psychiatrists, 2005). However, a greater proportion of our study population qualified overseas (Goldacre *et al*, 2004).

Student attitudes to psychiatry improve after psychiatric attachments (McParland *et al*, 2003). These improvements then decay after graduation, but it is uncertain whether they remain higher than pre-attach-



ment levels (Burra *et al*, 1982; Sivakumar *et al*, 1986; Baxter *et al*, 2001). The decay may be a result of the pull from other specialties, negative propaganda from other specialties (Clarke-Smith & Tranter, 2002) and narrow negative exposure to psychiatry after medical school. In the absence of exposure to psychiatry after medical school there is little to counteract these factors. After medical school, the narrow exposure of junior doctors to psychiatry may leave them with the impression that psychiatric patients are 'difficult' and 'incurable'.

Cameron & Persad (1984) found that the rejection of other specialties, discovery of psychosocial problems in other specialties and discovering the effectiveness of psychiatric therapy influenced the choice of psychiatry. Participants in our study indicated that during their postgraduate years, they recognised that psychiatry offered the most holistic approach to patient care.

Reasons for choosing psychiatry

About a third of participants cited empathy for this patient group as an important reason for choosing psychiatry as a career. Although there was no difference on a humanitarian scale between medical students who showed a preference for psychiatry and those choosing other specialties (Eron, 1955), medical students who did not show a preference for psychiatry found people with mental illness anxiety-provoking and difficult; in contrast to psychiatrists' who tend to have more positive attitudes (Tucker & Reinhardt, 1968).

Direct involvement in patient care is an important factor in improving medical students' attitudes to psychiatry during their attachment (McParland *et al*, 2003). Longer attachments in psychiatry or more direct involvement in patient care might help to reduce the anxiety that medical students have of psychiatric patients and enable them to develop greater empathy for this group. Foundation year posts in psychiatry (Department of Health, 2002) may offer potential, but others have been more sceptical (Eagles, 2004). Increasing the numbers of general practitioner trainees might also improve recruitment in psychiatry.

The interface of psychiatry with the neurosciences was an important factor for 18 consultant psychiatrists in their choice of this specialty. This is consistent with previous work that suggested that the divorce of psychiatry from mainstream medicine had led to its relative unpopularity (Thompson & Sims, 1999). Galeazzi *et al* (2003) found that the perception of psychiatric care as evidence based and recognition of research opportunities in mental health were important factors that led to the choice of psychiatry as a career.

Working conditions experienced in psychiatry were an important factor for 15 of the participants. Following graduation, the choice of careers is influenced more by lifestyle than by specialty content (Sierles & Taylor, 1995). In our study around half of those who felt that working conditions were an important factor cited better on-call working patterns or work-home balance as important. Without exposure to psychiatry after medical school,

doctors are unlikely to recognise the better work-home balance that is possible with a career in psychiatry.

Medical school teaching of psychiatry was an important factor in choice of specialty (19.4%). This was equally true of those who had decided on the specialty after medical school as for those who had decided earlier. Liaison psychiatry, student psychotherapy schemes and community psychiatric postings have been shown to improve recruitment into psychiatry (Clarke-Smith & Tranter, 2002; Yakeley *et al*, 2004). Enthusiastic teachers, encouragement from consultants, curability and direct patient involvement have all been cited as factors in medical school teaching which improved the attitudes of medical students (McParland *et al*, 2003).

Limitations

This study was carried out among consultant psychiatrists in one London psychiatric training scheme. A retrospective survey of this kind may lead to systematic recall bias. Direct questioning will not always reveal motivations behind a career choice (Halford, 2003). Furthermore, consultant psychiatrists are unlikely to admit that they made a mistake in their career choice.

Conclusions

This study confirms the importance of recruitment initiatives during medical school and in the early postgraduate years (Goldacre *et al*, 2004). Attractive medical school teaching stresses the link between psychiatry and the basic sciences, and is evidence-based, it also offers greater direct involvement with patients. Recruitment may be particularly important after medical school to attract doctors or prevent them losing motivation.

Declaration of interest

None.

Acknowledgements

We thank Drs Bindhu Abraham, Rafey Farouqi, Simon Dein, Gillian Mezey and Serab Ozdural for their help.

References

- BAXTER, H., SINGH, S. P. & DUGGAN, C. (2001) Attitudes of 'tomorrow's doctors' toward mental illness and psychiatry: changes during the final undergraduate year. *Medical Education*, **35**, 381–383.
- BURRA, P., KALIN, R., LEICHNER, S., *et al* (1982) The ATP-30: a scale for measuring medical students' attitudes to psychiatry. *Medical Education*, **16**, 31–38.
- CAMERON, P. & PERSAD, E. (1984) Recruitment into psychiatry: a study of the timing and process of choosing psychiatry as a career. *Canadian Journal of Psychiatry*, **29**, 676–680.
- CLARKE-SMITH, L. & TRANTER, R. (2002) Recruitment and retention in psychiatry. *British Journal of Psychiatry*, **181**, 163.
- COX, J. (2000) Person-power: reflections on the mental health National Service Framework for adults of working age. *Psychiatric Bulletin*, **24**, 201–202.
- CREED, F. & GOLDBERG, D. (1987) Doctors' interest in psychiatry as a career. *Medical Education*, **21**, 235–243.



education &
training

- DEPARTMENT OF HEALTH (2002) *Unfinished Business: Proposals for Reform of the Senior House Officer Grade*. Department of Health.
- DEPARTMENT OF HEALTH (2003) *Department of Health Vacancies Surveys, March 2003*. <http://www.dh.gov.uk/assetRoot/64/07/93/75/04079375.pdf>
- EAGLES, J. M. (2004) Modernising medical careers. *Psychiatric Bulletin*, **28**, 425.
- ERON, L. D. (1955) Effect of medical education on medical students' attitudes. *Journal of Medical Education*, **30**, 559–566.
- GALEAZZI, G. M., SECCHI, C. & CURCI, P. (2003) Current factors affecting the choice of psychiatry as a career: an Italian study. *Academic Psychiatry*, **27**, 74–81.
- GOLDACRE, J. M., DAVIDSON J. M. & LAMBERT, T. W. (2004) Country of training and ethnic origin of UK doctors: database and survey studies. *BMJ*, **329**, 597.
- HALFORD, J. (2003) Invited commentary on: recruiting and retaining psychiatrists. *Advances in Psychiatric Treatment*, **9**, 411–413.
- HELD, M. L. & ZIMET, C. N. (1975) A retrospective view of medical speciality choice. *Journal of Medical Education*, **50**, 2044–2051.
- McPARLAND, M., NOBLE, L. M., LIVINGSTON, G., et al (2003) The effect of a psychiatric attachment on students' attitude to and intention to pursue psychiatry as a career. *Medical Education*, **37**, 447.
- MAIDMENT, R., LIVINGSTON, G., KATONA, M., et al (2003) Carry on shrinking: career intentions and attitudes to psychiatry of prospective medical students. *Psychiatric Bulletin*, **27**, 30–32.
- MAIDMENT, R., LIVINGSTON, G., KATONA, M., et al (2004) Changes in attitudes to psychiatry and intention to pursue psychiatry as a career in newly qualified doctors: a follow-up of two cohorts of medical students. *Medical Teacher*, **26**, 565–569.
- PRINS, H. (1998) Characteristics of consultant forensic psychiatrists (a modest survey). *Journal of Forensic Psychiatry*, **9**, 139–149.
- RAJAGOPAL, S., REHILL, K. S. & GODFREY E. (2004) Psychiatry as a career choice compared with other specialties: a survey of medical students. *Psychiatric Bulletin*, **28**, 444–446.
- SCHUMACHER, C. F. (1964) Personal characteristics of students choosing different types of medical careers. *Journal of Medical Education*, **39**, 278–288.
- SIERLES, F. S. & TAYLOR, M. A. (1995) Decline of US medical student career choice of psychiatry and what to do about it. *American Journal of Psychiatry*, **152**, 1416–1426.
- SIVAKUMAR, K., WILKINSON, G., TOONE, B. K., et al (1986) Attitudes to psychiatry in doctors at the end of their post-graduate year: two-year follow-up of a cohort of medical students. *Psychological Medicine*, **16**, 457–460.
- THOMPSON, D. J. & SIMS, A. C. P. (1999) Malaise in psychiatric recruitment and its remedy. *Psychiatric Bulletin*, **23**, 227–229.
- TUCKER, G. J. & REINHARDT, R. F. (1968) Psychiatric attitudes of young physicians: implications for teaching. *American Journal of Psychiatry*, **124**, 986–991.
- WILKINSON, D. G., TOONE, B. K. & GREER, S. (1983) Medical students' attitudes to psychiatry at the end of the clinical curriculum. *Psychological Medicine*, **13**, 655–658.
- YAKELEY, J., SHOENBERG, P. & HEADY, A. (2004) Who wants to do psychiatry? The influence of a student psychotherapy scheme – a 10-year retrospective study. *Psychiatric Bulletin*, **28**, 208–212.

***Kalpna Dein** Specialist Registrar in Forensic Psychiatry, North London Forensic Services, Chase Farm Hospital, The Ridgeway, Enfield EN2 8JL, email: kalthomas@yahoo.co.uk, **Gill Livingston** Professor of Psychiatry of Older People, Department of Mental Health Sciences, University College London, **Christopher Bench** Consultant Psychiatrist, West London Mental Health NHS Trust, The Claybrook Centre, London W6 8LN

Psychiatric Bulletin (2007), **31**, 230–232. doi: 10.1192/pb.bp.106.012286

MEGAN MUNRO, MICHAEL WESSON AND MARK THEOPHANOUS

First experience of recruiting to the new specialist training programme

In 2002 *Unfinished Business*, a report and consultation paper by Sir Liam Donaldson, Chief Medical Officer for England, put forward proposals for the reform of the senior house officer (SHO) grade, including the formation of the new early years foundation posts (Donaldson, 2002). In 2004 *Modernising Medical Careers – The Next Steps* (Department of Health, 2004) outlined specialty and general practitioner (GP) training programmes building on the foundation programme. As a result all medical training will be changing to a competency-based model from August 2007. This will encompass run-through training from specialist training years 1 (ST1) to 6 (ST6). Some regions and specialties have been chosen as pilot sites for specialist training year 1 commencing August 2006.

Mersey Deanery pilot

The Mersey Deanery has embraced the changes with enthusiasm and was a pilot site for foundation year 2 (F2) trainees in 2004, which included psychiatric placements. Psychiatry is one of the specialties that has agreed to take part in the ST1 pilot.

The Merseyside regional rotational training scheme has 93 trainees at SHO level spread over a large

geographical area. Organisation of the scheme is carried out by three scheme coordinators, one from each of the largest local trusts, with input from local college tutors and specialties via the Basic Specialist Training and Education Committee (STEC). They are supported administratively by the medical staffing department of the single employing trust. Until now the scheme has been run much along the lines of most training schemes, with competitive interview, contracts being renewed on satisfactory progress and an expectation that trainees will progress quickly through years 1 to 3 and obtain their Membership of the Royal College of Psychiatrists before taking up higher postgraduate training.

For the new specialist training scheme it was considered that a more robust form of selection would be required if the right individuals were to be found for the 6-year specialist training now proposed. A new recruitment and selection process was set up by the scheme organisers with help and advice from Mersey Deanery. It consisted of an application process, a short-listing procedure and a competitive interview.

Application process

Advertisements appeared on the Mersey Deanery website (<http://www.merseydeanery.nhs.uk>) along with