

ERRATA

Genet. Res., Camb. (1987), 50, pp. 17–21

Heterogeneity of human haptoglobin α chains detected by two-dimensional gel electrophoresis

BY HUW A. JOHN AND IAN F. PURDOM

The running title for this article should have been
Human plasma haptoglobin α chain heterogeneity

Genet. Res., Camb. (1987), 50, pp. 63–68

Additive variance and average effect with partial selfing

BY A. J. WRIGHT

(1) The second line of Table 1 should read

Table 1. *Quadratic components and identity measures*

For one locus	Sum over loci	Coefficient
$m \sigma_D^2 = \sum_k \sum_l p_k p_l (d_{kl})^2$	$\sigma_D^2 = \sum_m \sigma_D^2$	$2\delta_{\bar{x}+\bar{y}} = P_{(i=k \text{ and } j=l)} + P_{(i=l \text{ and } j=k)} - 2P_{(i=j=k=l)}$

(2) The second formula on page 66 should be

$$C_{Pt} = \sigma_{A..At} + (s/2)'[\sigma_{D..D0} + H'].$$

(3)

$$\alpha'_i = [F(y_{ii} - y_{..}) + (1-F)(y_{i.} - y_{..})]/(1+F) \\ = \{[F\gamma_{it} + (1-F)\gamma_{0i}](x_{ii} - x_{..}) + 2[F(\theta_{i0t} - \gamma_{i0i}) \\ + (1-F)(\theta_{0it} - \gamma_{0i0})](x_{i.} - x_{..})\}/(1+F),$$

on page 68 should be

$$\alpha'_i = [F(y_{ii} - y_{..}) + (1-F)(y_{i.} - y_{..})]/(1+F) \\ = \{[F\gamma_{it} + (1-F)\gamma_{0i}](x_{ii} - x_{..}) + 2[F(\theta_{it} - \gamma_{it}) \\ + (1-F)(\theta_{0i} - \gamma_{0i})](x_{i.} - x_{..})\}/(1+F),$$