

**Note on Mr Tweedie's Theorem in Geometry.**

By PETER PINKERTON, M.A.

Let  $ABC$ ,  $A'B'C'$  (Fig. 4) be two triangles equiangular in the same sense. Let  $BC$ ,  $B'C'$  meet in  $X$ . Describe circles round  $BXB'$ ,  $CXC'$  to meet again in  $O$ . Then it is easy to see that the triangles  $BOC$ ,  $COA$ ,  $AOB$  are equiangular in the same sense to the triangles  $B'OC'$ ,  $C'OA'$ ,  $A'OB'$  respectively. Hence the triangles  $AOA'$ ,  $BOB'$ ,  $COC'$  are similar ;

$$\therefore \frac{AA'}{AO} = \frac{BB'}{BO} = \frac{CC'}{CO} ;$$

$\therefore a . AA'$ ,  $b . BB'$ ,  $c . CC'$  are proportional to  $a . AO$ ,  $b . BO$ ,  $c . CO$ , where  $a$ ,  $b$ ,  $c$  are the sides of the triangle  $ABC$ .

From  $O$  draw  $OP$ ,  $OQ$ ,  $OR$  perpendicular to  $BC$ ,  $CA$ ,  $AB$  respectively.

$$\text{Then } QR = AO \sin A \propto a . AO,$$

$$RP = BO \sin B \propto b . BO,$$

$$PQ = CO \sin C \propto c . CO ;$$

$\therefore a . AA'$ ,  $b . BB'$ ,  $c . CC'$ , being proportional to  $a . AO$ ,  $b . BO$ ,  $c . CO$ , are proportional to  $QR$ ,  $RP$ ,  $PQ$ .

But  $PQR$  is a triangle, unless  $O$  is on the circumcircle of  $ABC$  when  $PQR$  is the Simson line of  $O$ .

$\therefore QR + RP > PQ$ , with two similar inequalities, except that *one* of the inequalities becomes an equality if  $O$  is on the circumcircle of  $ABC$ .

$\therefore a . AA' + b . BB' > c . CC'$ , with two similar inequalities ; one of the inequalities becoming an equality when  $O$  lies on the circumcircle of  $ABC$ .

Similarly in the case of an equality  $O$  lies also on the circumcircle of  $A'B'C'$ .

For the case of equilateral triangles  $a = b = c$  ;

$\therefore AA' + BB' > CC'$ , with two similar inequalities ; one of the three inequalities becoming an equality when  $O$  lies on the circumcircles of  $ABC$  and  $A'B'C'$ .

It is obvious that the theorem reduces to Ptolemy's Theorem or its converse.