

objects in the pocket (wilcox_test p 0.00854). When comparing the group of patients with more than three objects and the presence of dementia, significant p values were also found (wilcox_test p 0.0363).

Conclusions: The number of objects in the pocket of patients with cognitive impairment could be a valuable tool for predicting dementia.

The presence of 3 or more objects in the pocket of a patient with cognitive impairment could correlate with a low score on the functionality scales (FAQ less than 6, absence of dementia stage).

P42: Neuropsychological Impact of Mandatory Preventive Social Isolation (Lockdown) on Older Adults with Cognitive Decline

Authors: Natalia Sierra Sanjurjo, Santiago O Neill, Julián Bustin, María Roca

Objectives: Various studies described the psychological impact of the mandatory preventive social isolation (lockdown) caused by COVID-19 across different age groups. However, no research had specifically analyzed the effects of lockdown on the cognitive performance of older adults with preexisting cognitive decline. This study aimed to determine the impact of lockdown on the cognitive functions of individuals with cognitive decline.

Methods: Argentina was one of the countries with the longest lockdown periods. Older adults in Argentina experienced extended lockdowns lasting several months, significantly limiting their social interactions and routine activities. Cognitive reevaluations were conducted on 16 patients who had attended a memory clinic with cognitive complaints and were evaluated before the lockdown (group 1). Their progression was compared with 16 patients evaluated and reevaluated over a similar period but without undergoing a lockdown (group 2). To compare the progression of both groups a "change index" (CI) were calculated for each test ((score assessment 1 – score assessment 2)/number of months between assessment 1 and assessment 2))

Results: Patient from group 1 and 2 were paired by age (M group 1 = 77.2, DS group 1 = 6.2, M group2 = 77.3 DS group2 = 6.6, p = .98), years of education (M group 1 = 13.6, DS group 1 = 2.8, M group2 = 13.9 DS group2 = 3.0, p = .81), Beck depression inventory score (M group 1 = 12.2, DS group 1 = 7.2, M group2 = 13.9 DS group2 = 7.4, p = .54) and Addenbrooke's Cognitive Examination (ACE. M group 1 = 76.2, DS group 1 = 14.4, M group2 = 76.0 DS group2 = 20.7, p = .97) score at the first assessment. Group 1 showed a significant and greater increase of false positive than group 2 in Rey Auditory Verbal Learning Test recognition (M CI group 1 = -.31, DS CI group 1 = .38, M CI group 2 = -.01, DS CI group 2 = .44, p < .05). Results indicated that patients who experienced the lockdown exhibited a greater cognitive decline compared to the control group.

Conclusions: The findings suggest that the lockdown accelerated cognitive deterioration in individuals with memory complaints.