



On the perimeter, diameter and circumradius of ordinary hyperbolic reduced polygons

Ádám Sagmeister 

Abstract. A convex body R in the hyperbolic plane is called reduced if any convex body $K \subset R$ has a smaller minimal width than R . We answer a few of Lassak's questions about ordinary reduced polygons regarding its perimeter, diameter, and circumradius, and we also obtain a hyperbolic extension of a result of Fabińska.

1 Introduction

The concept of reducedness was introduced by Heil [8] in 1978 motivated by some volume minimizing problems. A convex body (i.e., a convex compact set of non-empty interior) K is called reduced if an arbitrary convex body strictly contained in K has smaller minimal width than K . Pál [26] proved in 1921 that for fixed minimal width, the regular triangle has minimal area among convex bodies in the Euclidean plane. This result of his is also known as the isominwidth inequality. The same problem in higher dimensions remains unsolved, as there are no reduced simplices in \mathbb{R}^n for $n \geq 3$ (see [23, 24]), therefore there are no really good candidates for the volume minimizing problems—so far the best one in \mathbb{R}^3 is the so-called Heil body, which has a smaller volume than any rotationally symmetric body of the same minimal width. The problem can be naturally generalized to other spaces, a natural approach is to study the problem in spaces of constant curvature. Bezdek and Blekherman [1] proved that, if the minimal width is at most $\frac{\pi}{2}$, the regular triangle minimizes the area in S^2 . However, for spherical bodies of larger minimal width, the minimizers of the isominwidth problem are polars of Reuleaux triangles. Surprisingly enough, there is no solution of the isominwidth problem in the hyperbolic space for arbitrary dimension (see Böröczky, Freyer, Sagmeister [3]).

A reverse isominwidth problem is about finding the maximal volume if the minimal width is fixed. Naturally, this problem does not have a maximizer in general, but for reduced bodies we can ask for the convex body that maximizes the volume. However, in \mathbb{R}^3 , the diameter of a reduced body of a given minimal width can be arbitrarily large, and hence the Euclidean problem is only interesting on the plane. It is conjectured, that the unique planar reduced bodies maximizing the area and

Received by the editors October 14, 2024; accepted January 17, 2025.

AMS Subject Classification: 51M09, 51M10, 52A55.

Keywords: Convex geometry, hyperbolic geometry, perimeter, diameter, circumradius.



of minimal width $w > 0$ in \mathbb{R}^2 are the circular disk of radius $\frac{w}{2}$ and the quarter of the disk of radius w . A big step towards the proof of this conjecture was made by Lassak, who proved that among reduced k -gons, regular ones maximize the area, and as a consequence, all reduced polygons have smaller area than the circle. Following Lassak's footsteps, the same conclusion was derived in S^2 by Liu, Chan, and Su [22]. Interestingly enough, the characterization of hyperbolic reduced polygons is still unclear, but clearly it must be different from the Euclidean and spherical ones; there exist reduced rhombi on the hyperbolic plane, while Euclidean and spherical reduced polygons are all odd-gons (see Lassak [12, 14]). However, the so-called ordinary reduced polygons can be examined the same way (these are odd-gons whose vertices have distance equal to the minimal width of the polygon from the opposite sides such that the projection of the vertices to these sides are in the relative interior of the sides). In fact, it was shown by the author [27] that, among ordinary reduced n -gons of a fixed width, regular n -gons maximize the area. This answers one of Lassak's questions posed in [13]. One of his other questions was about the extremality of the perimeter, which is addressed in Section 4 where we give an explicit perimeter formula. However, the extremality of the perimeter remains open, but based on the given formula in Theorem 4.1, we propose a conjecture that is surprising given the Euclidean and spherical analogs. Lassak also proposed to find the infimum of the circumradii of ordinary reduced polygons of minimal width w . In order to obtain the best bound for the circumradius, we need the following result.

Theorem 1.1 *If $P \subset H^2$ is an ordinary reduced polygon of minimal width w , then its diameter is at most*

$$\text{diam}(P) \leq 2 \operatorname{arcosh} \left(\frac{\cosh w + \sqrt{\cosh^2 w + 8}}{4} \right)$$

with equality if and only if P is the regular triangle.

With this diameter bound, we obtain the following bound for the circumradius.

Theorem 1.2 *Let P be an ordinary reduced polygon in H^2 of minimal width w . Then its circumradius is at most*

$$\operatorname{arsinh} \left(\frac{2}{\sqrt{3}} \sqrt{\left(\frac{\cosh w + \sqrt{\cosh^2 w + 8}}{4} \right)^2 - 1} \right)$$

with equality if and only if P is a regular triangle.

Finally, we also have the following result which is analogous to the results of Fabińska [6] and Musielak [25].

Theorem 1.3 *Every ordinary reduced hyperbolic polygon of minimal width w is contained in a circular disk of radius w centered at one of its boundary points.*

The article is structured as follows. In Section 2, we describe the fundamental concepts and notations about hyperbolic width and reducedness. In Section 3, we introduce ordinary reduced polygons and explain some of their basic properties, including a few of the key ideas that will be used to obtain some of the main results. In Section 4, we give a perimeter formula for ordinary reduced n -gons. In Section 5, we prove Theorem 1.1, while in Section 6 we prove Theorems 1.2 and 1.3.

2 Preliminaries

We use the notation H^2 for the hyperbolic plane, which is equipped with the geodesic metric. The geodesic distance of two points $x, y \in H^2$ will be denoted as $d(x, y)$. In this section, we introduce hyperbolic convexity. Many of the concepts are identical with their Euclidean analogs, but as we will soon see, there are exceptions.

For a subset X of the hyperbolic plane H^2 , we say that X is *convex*, if for any pair of points x and y , the unique geodesic segment $[x, y]$ connecting x and y is a subset of X (where $[x, x] = \{x\}$). A *convex body* is a convex compact set of non-empty interior. It is clear, that similarly to Euclidean convexity, the intersection of an arbitrary family of convex sets in the hyperbolic plane is also convex, so we define the *convex hull* of a set $X \subseteq H^2$ as the intersection of all convex sets in H^2 containing X as a subset, and we will use the notation $\text{conv}(X)$ for the convex hull of X . The convex body obtained as the convex hull of the finite set $X = \{x_1, \dots, x_n\}$ is called a *polygon*, and we use the notation $[x_1, \dots, x_n]$ for $\text{conv}(X)$. A point $x_j \in X$ is a *vertex* of the polygon X if $x_j \notin \text{conv}(X \setminus \{x_j\})$; a *k-gon* in the hyperbolic plane is a polygon of k vertices.

For convex bodies, width is an important concept. On the hyperbolic plane there are many different notions of width (see Santaló [28], Fillmore [7], Leichtweiss [20], Jerónimo-Castro–Jimenez-Lopez [10], G. Horváth [9], Böröczky, Csépai, and Sagmeister [2], Lassak [15]). We will use the width function introduced by Lassak, but we note that it is identical with the extended Leichtweiss width defined by Böröczky, Csépai, and Sagmeister [2] on supporting lines. A hyperbolic line ℓ is called a *supporting line* of the convex body K if $K \cap \ell \neq \emptyset$, and K is contained in one of the closed half-spaces bounded by ℓ . The *width* of the convex body K with respect to the supporting line ℓ is the distance of ℓ and ℓ' , where ℓ' is a (not necessarily unique) most distant supporting line from ℓ . It is known that this width function is continuous, and its maximal value coincides with the diameter of the convex body, which will be denoted as $\text{diam}(K)$. The *minimal width* (i.e., the minimal value of the width function on the set of all supporting lines, also known as the *thickness*) of K is denoted by $w(K)$. This notion of minimal width is a monotonic function, that is for arbitrary convex bodies K, L such that $K \subseteq L$, we have $w(K) \leq w(L)$. Hence, the concept of hyperbolic reducedness makes perfect sense. A convex body K is called *reduced*, if for any convex body $K' \subsetneq K$, $w(K') < w(K)$. Reduced bodies are well-studied (see Heil [8], Lassak–Martini [16–18], Lassak–Musielak [19]), as they are often extremizers of volume minimizing problems, and bodies of constant width are also reduced.

If we consider the Poincaré disk model of the hyperbolic plane H^2 , hyperbolic lines are either diameters of the unit disk, or circular arcs intersecting the unit circle orthogonally. The boundary points of the unit disk are the *ideal points* of the hyperbolic plane, and hence there is a natural bijection between hyperbolic lines

and pairs of ideal points. Besides the identity, there are three types of orientation preserving isometries of the hyperbolic plane depending the number of fixed points. If there is one fixed point, the isometry is called an *elliptic isometry* (or *rotation*). We call an isometry with exactly one fixed ideal point, it is called a *parabolic isometry*. Finally, isometries with exactly two fixed ideal points are called *hyperbolic isometries*, which maps the line corresponding to the two fixed ideal points to itself.

3 Ordinary reduced polygons

Lassak proved that hyperbolic convex odd-gons of thickness w are reduced if all vertices are of distance w from the opposite sides, and the orthogonal projections of these vertices onto the opposite sides are in the relative interior of these sides (see [13]). Such polygons are called *ordinary reduced polygons*, since this property characterizes reducedness both in \mathbb{R}^2 and in S^2 (see Lassak [12, 14]), but not in the hyperbolic plane. In Böröczky, Freyer, Sagmeister [3] it is shown that, for each $w > 0$ there are reduced rhombi, whose diameters are unbounded. The characterization of hyperbolic reduced polygons is therefore unclear, so we focus on ordinary reduced polygons in this article. For the diameter of an ordinary reduced polygon of thickness w , we have the following by Lassak [13].

Theorem 3.1 *Let $P \subset H^2$ be an ordinary reduced polygon of thickness w and diameter d . Then,*


$$w < d < \operatorname{arccosh} \left(\cosh w \sqrt{1 + \frac{\sqrt{2}}{2} \sinh w} \right).$$

As a consequence, for each n we can expect an n -gon of extremal area among ordinary reduced n -gons of thickness w by Blaschke's Selection Theorem. In the remainder of the section, we will discuss the area of hyperbolic reduced n -gons based on the arguments of Lassak [11] and Liu–Chang–Su [22].

From now on, P denotes an ordinary reduced n -gon in H^2 whose vertices are v_1, \dots, v_n in cyclic order with respect to the positive orientation. For each i , let t_i be the orthogonal projection of v_i on the line through $v_{i+\frac{n-1}{2}}$ and $v_{i+\frac{n+1}{2}}$, where the indices are taken mod n . By definition, t_i is in the relative interior of $[v_{i+\frac{n-1}{2}}, v_{i+\frac{n+1}{2}}]$, and hence the geodesic segments $[v_i, t_i]$ and $[v_{i+\frac{n+1}{2}}, t_{i+\frac{n+1}{2}}]$ intersect in a point p_i . Let B_i be the union of the two triangles:

$$B_i = [v_i, p_i, t_{i+\frac{n+1}{2}}] \cup [v_{i+\frac{n+1}{2}}, p_i, t_i];$$

we will call B_i a *butterfly*. We observe that these butterflies cover the polygon (see Sagmeister [27]).



Lemma 3.2 Let $P \subset H^2$ be an ordinary reduced n -gon, and B_i be its i^{th} butterfly. Then,

$$P = \bigcup_{i=1}^n B_i.$$

We introduce a few additional notations for some angles of the butterflies. Let

$$\varphi_i = \angle \left(v_i, p_i, t_{i+\frac{n+1}{2}} \right) = \angle \left(t_i, p_i, v_{i+\frac{n+1}{2}} \right)$$

and

$$\alpha_i = \angle \left(t_i, v_{i+\frac{n+1}{2}}, p_i \right).$$

The following lemma from Sagmeister [27] shows that the two triangles involved in the butterfly B_i are congruent.

Lemma 3.3 The two triangles $\left[v_i, p_i, t_{i+\frac{n+1}{2}} \right]$ and $\left[v_{i+\frac{n+1}{2}}, p_i, t_i \right]$ defining B_i are congruent.

Another observation from Sagmeister [27] provides an upper bound for the sum of the vertical angles of the butterflies.

Lemma 3.4 For an ordinary reduced n -gon P with the notations from above,

$$\sum_{i=1}^n \varphi_i \leq \pi.$$

We note that for regular n -gons we have equality in the previous lemma. Also, in the Euclidean plane, the angle sum is always π .

Let β_i be the angle $\angle \left(v_{i+\frac{n+1}{2}}, v_i, p_i \right) = \angle \left(p_i, v_{i+\frac{n+1}{2}}, v_i \right)$. Also, let γ denote half of the inner angle of a regular triangle of minimal width w . We have the following.

Lemma 3.5 Let P be an ordinary reduced n -gon of minimal width w . With the notations introduced above, we have $\beta_i \leq \gamma \leq \alpha_i$ with equality if and only if P is a regular triangle.

Proof First, we calculate the side length a of the regular triangle in terms of w . By the hyperbolic Pythagorean theorem, we have

$$(1) \quad \cosh a = \cosh \frac{a}{2} \cosh w.$$

The identity

$$\cosh a = 2 \cosh^2 \frac{a}{2} - 1$$

combined with (1) leads to a quadratic equation for $\cosh \frac{a}{2}$, whose positive solution is

$$\cosh \frac{a}{2} = \frac{\cosh w + \sqrt{\cosh^2 w + 8}}{4}.$$

Now we apply the hyperbolic law of sines for half of the regular triangle:

$$(2) \quad \sin \gamma = \frac{\sinh \frac{a}{2}}{\sinh a} = \frac{1}{2 \cosh \frac{a}{2}} = \frac{-\cosh w + \sqrt{\cosh^2 w + 8}}{4}.$$

From the hyperbolic law of cosines applied for the right triangle $[v_i, t_i, v_{i+\frac{n+1}{2}}]$ and also for the half of the regular triangle, we have

$$(3) \quad \cosh w = \frac{\cos 2\gamma}{\sin \gamma} = \frac{\cos(\alpha_i + \beta_i)}{\sin \beta_i}.$$

We know that $\beta_i \leq \alpha_i$ (this is Theorem 2 (iii) in Lassak's paper [13]), so from (3) we imply

$$\frac{\cos 2\alpha_i}{\sin \alpha_i} \leq \frac{\cos 2\gamma}{\sin \gamma} \leq \frac{\cos 2\beta_i}{\sin \beta_i}.$$

Finally, we observe that

$$\frac{\cos 2x}{\sin x} = \frac{1 - 2 \sin^2 x}{\sin x},$$

and that the function $\frac{1-2x^2}{x}$ strictly monotonically decreases. Hence, $\sin \beta_i \leq \sin \gamma \leq \sin \alpha_i$, which in turn concludes the proof as all of these angles are acute. The case of equality is clear. ■

Lassak also proved [13] the following.

Lemma 3.6 *Let D be the diameter of an ordinary reduced n -gon in H^2 . Then, the endpoints of a chord of length D are vertices of the polygon, where if v_i is one of the endpoints of the chord, the other endpoint is either $v_{i+\frac{n-1}{2}}$ or $v_{i+\frac{n+1}{2}}$.*

4 The perimeter of ordinary reduced polygons

With the same method that was used in Sagmeister [27], we can also investigate the extremality of the perimeter. With the notations introduced in the previous sections, let

$$g_w(x) = \frac{1 + \cos x - \sqrt{(1 + \cos x)^2 - 4 \tanh^2 w \cos x}}{2 \tanh w}$$

and

$$p_w(x) = \operatorname{arcosh} \left(\frac{1 - g_w(x) \tanh w}{\sqrt{1 - \tanh^2 w}} \right).$$

Then, we have the following formula for the perimeter.

Theorem 4.1 *Let P be an ordinary reduced n -gon of thickness w described as above. Then,*

$$\operatorname{perim}(P) = 2 \sum_{i=1}^n p_w(\varphi_i).$$

Proof By the definition of ordinary reduced polygons, t_i is in the relative interior of the side $\left[v_{i+\frac{n-1}{2}}, v_{i+\frac{n+1}{2}} \right]$, so

$$\operatorname{perim}(P) = \sum_{i=1}^n d(v_i, v_{i+1}) = \sum_{i=1}^n \left(d\left(v_i, t_{i+\frac{n-1}{2}}\right) + d\left(t_{i+\frac{n-1}{2}}, v_{i+\frac{n+1}{2}}\right) \right).$$

On the other hand, $d\left(v_i, t_{i+\frac{n+1}{2}}\right) = d\left(t_i, v_{i+\frac{n+1}{2}}\right)$ by Lemma 3.3, so

$$\operatorname{perim}(P) = 2 \sum_{i=1}^n d\left(t_i, v_{i+\frac{n+1}{2}}\right).$$

If $b_i = d(p_i, t_i)$ and $c_i = d\left(p_i, v_{i+\frac{n+1}{2}}\right)$, $\tanh c_i$ can be expressed as

$$\tanh c_i = \frac{\tanh w - g_w(\varphi_i)}{1 - g_w(\varphi_i) \tanh w},$$

where we refer to Sagmeister [27]. Hence,

$$c_i = \operatorname{artanh} \left(\frac{\tanh w - g_w(\varphi_i)}{1 - g_w(\varphi_i) \tanh w} \right).$$

Using the identity

$$\cosh(\operatorname{artanh} x) = \frac{1}{\sqrt{1-x^2}}$$

and the hyperbolic Pythagorean theorem, we get

$$\begin{aligned} \cosh d \left(t_i, v_{i+\frac{n+1}{2}} \right) &= \frac{\cosh c_i}{\cosh b_i} = \frac{(1 - g_w(\varphi_i) \tanh w) \sqrt{1 - g_w^2(\varphi_i)}}{\sqrt{(1 - g_w^2(\varphi_i))^2 - (\tanh w - g_w(\varphi_i))^2}} \\ &= \frac{1 - g_w(x) \tanh w}{\sqrt{1 - \tanh^2 w}}, \end{aligned}$$

concluding the proof. ■

Now we are ready to prove the following.

Theorem 4.2 *The function p_w is strictly monotonically increasing and strictly convex on the interval $(0, \pi)$.*

Proof It is convenient to use the notation $r_w(x) = \sqrt{(1 + \cos x)^2 - 4 \tanh^2 w \cos x}$, so

$$(4) \quad r'_w(x) = \frac{-\sin x}{r_w(x)} \cdot (1 + \cos x - 2 \tanh^2 w)$$

and

$$g'_w(x) = \frac{-\sin x}{r_w(x)} \cdot (\tanh w - g_w(x)).$$

We deduce

$$\begin{aligned} p'_w(x) &= \sqrt{\frac{\tanh w}{\tanh w \cdot g_w(x)^2 + \tanh w - 2g_w(x)}} \cdot \frac{\sin x (\tanh w - g_w(x))}{r_w(x)} \\ &= \frac{\sqrt{\tanh w (\tanh w - g_w(x)) (1 + \cos x)}}{r_w(x)} \\ &= \frac{\cos \frac{x}{2} \sqrt{2 \tanh w (\tanh w - g_w(x))}}{r_w(x)} \\ &= \frac{\cos \frac{x}{2} \sqrt{2 \tanh^2 w + r_w(x) - (1 + \cos x)}}{r_w(x)}. \end{aligned}$$

To calculate the second derivative, we use (4) to substitute $r'_w(x)$, and we get to a common denominator. We also write $\sin x = 2 \sin \frac{x}{2}$ and $2 \cos^2 \frac{x}{2} = 1 + \cos x$. Thus, we

obtain

$$p_w''(x) = \frac{\sin \frac{x}{2} \sqrt{2 \tanh^2 w + r_w(x) - (1 + \cos x)}}{r_w^3(x)} \cdot \left(\cos^2 \frac{x}{2} r_w(x) + 2 \cos^2 \frac{x}{2} \cos x - 2 \tanh^2 w \right).$$

It is not too difficult to verify that this is positive; otherwise we can reorganize $\cos^2 \frac{x}{2} r_w(x) + 2 \cos^2 \frac{x}{2} \cos x - 2 \tanh^2 w$ as an inequality that is quadratic in $\tanh^2 w$, and we get that if p_w'' is not positive, $\tanh^2 w$ is either negative, or greater than 1, but both are impossible. Hence, p_w is strictly convex. ■

The usual argument used on the Euclidean and spherical planes to find the reduced polygon with the minimal perimeter (see Lassak [12] and Liu–Chang [21]) uses Jensen's inequality after deriving a similar formula for the perimeter as in Theorem 4.1. However, if we consider the results of Lemma 3.4 and Theorem 4.2, we find that this approach does not work on the hyperbolic plane. Considering the perimeter of random ordinary reduced polygons given by Theorem 4.1, we have the following conjecture.

Conjecture 4.3 *Let $P \subset H^2$ be an ordinary reduced n -gon of minimal width w . Then,*

$$\text{perim}(P) \leq \text{perim}(\tilde{P})$$

with equality if and only if P is regular.

We note that contrary to the Euclidean and spherical planes, the perimeter of the regular n -gons of minimal width w is not necessarily monotone in n . Depending on w , even the regular triangle can have a larger perimeter than the circle of the same minimal width.

In a recent work of Chen, Hou, and Jin [4], a consequence of the minimality of the perimeter of the regular n -gon of reduced n -gons of minimal width less than $\frac{\pi}{2}$ on the sphere is that the diameter and the circumradius of reduced spherical n -gons is minimized by the regular ones. Their argument gives the same result on the Euclidean plane. However, as we have seen, on the hyperbolic plane, we do not have the same result on the perimeter of ordinary reduced polygons. We propose the following question regarding the diameter and circumradius of ordinary reduced n -gons.

Question. Is the diameter and the circumradius of the regular n -gon extremal among ordinary reduced n -gons of the same minimal width on the hyperbolic plane?

5 The diameter of ordinary reduced polygons

In this section, we strengthen Lassak's Theorem 3.1 by using Lemma 3.5 in order to find the infimum of the circumradii of ordinary reduced polygons of a prescribed width. We have the following result.

Theorem 5.1 *Let P be an ordinary reduced polygon in H^2 . Then,*

$$\text{diam}(P) \leq 2 \operatorname{arccosh} \left(\frac{\cosh w + \sqrt{\cosh^2 w + 8}}{4} \right)$$

with equality if and only if P is a regular triangle.

Proof Lemma 3.6 allows us to assume that $\text{diam } P = d(v_i, v_{i+\frac{n+1}{2}})$ for some i , and we consider the right triangle $[v_i, t_i, v_{i+\frac{n+1}{2}}]$. Recall the notations $\alpha_i = \angle(t_i, v_{i+\frac{n+1}{2}}, t_{i+\frac{n+1}{2}})$ and $\beta_i = \angle(v_{i+\frac{n+1}{2}}, v_i, t_i)$, and that $\angle(t_i, v_{i+\frac{n+1}{2}}, v_i) = \alpha_i + \beta_i$ as a consequence of Lemma 3.3. Since $d(v_i, t_i) = w$, we get

$$(5) \quad \sinh d(t_i, v_{i+\frac{n+1}{2}}) = \frac{\sinh w \cdot \sin \alpha_i}{\sin(\alpha_i + \beta_i)} \leq \frac{\sinh w}{2 \cos \beta_i}$$

from the hyperbolic law of sines, and using the inequality $\alpha_i \geq \beta_i$ (cf. Lassak [13]), the monotonicity of the sin function on the interval $(0, \frac{\pi}{2})$ and the identity $\sin 2x = 2 \sin x \cos x$. From Lemma 3.5, we also have $\beta_i \leq \gamma$ where γ denotes half the angle of a regular triangle of minimal width w , so we obtain

$$(6) \quad \sinh d(t_i, v_{i+\frac{n+1}{2}}) \leq \frac{\sinh w}{2 \cos \gamma}.$$

from (5). Now we use (6) and the identity $\cosh^2 x - \sinh^2 x = 1$ to obtain

$$\cosh d(t_i, v_{i+\frac{n+1}{2}}) \leq \sqrt{1 + \frac{\sinh^2 w}{4 \cos^2 \gamma}}.$$

We now use the hyperbolic Pythagorean theorem with the assumption $d(v_i, v_{i+\frac{n+1}{2}}) = \text{diam } P$:

$$(7) \quad \cosh \text{diam } P = \cosh w \cosh d(t_i, v_{i+\frac{n+1}{2}}) \leq \cosh w \sqrt{1 + \frac{\sinh^2 w}{4 \cos^2 \gamma}}.$$

Clearly, from the equality case of Lemma 3.5 we imply that equality holds if and only if P is a regular triangle. On the other hand, if P is a regular triangle centered at p , in the right triangle $[p, t_1, v_3]$, the angles are $\frac{\pi}{3}$, $\frac{\pi}{2}$ and γ , respectively, while the length of leg opposite to the $\frac{\pi}{3}$ angle is $\frac{\text{diam } P}{2}$, so using the well-know identity $\cosh b = \frac{\cos B}{\sin A}$ for hyperbolic right triangles of acute angles A and B and legs a and b respectively, we obtain

$$\cosh \frac{\text{diam } P}{2} = \frac{1}{2 \sin \gamma}.$$

This together with (7) gives

$$\operatorname{arcosh} \left(\cosh w \sqrt{1 + \frac{\sinh^2 w}{4 \cos^2 \gamma}} \right) = 2 \operatorname{arcosh} \frac{1}{2 \sin \gamma}.$$

Finally, (2) concludes the proof. \blacksquare

From this sharp upper bound we can see that Lassak's following conjecture is true.

Corollary 5.2 *Let P be an ordinary reduced polygon in H^2 . Then,*

$$1 < \frac{\operatorname{diam} P}{w} < 2.$$

Proof The first inequality trivially holds as $\operatorname{diam} P$ is the maximal width of P , and equality holds exactly for bodies of constant width, which are h -convex, so no polygon is of constant width (see Böröczky, Csépai, and Sagmeister [2] for further details).

The second inequality is an easy consequence of Theorem 1.1 and the strict monotonicity of \cosh for $w > 0$.

We can also observe, that both of these constant bounds are the optimal ones: on one hand, regular $(2k+1)$ -gons are ordinary reduced polygons that approximate a disk as $k \rightarrow \infty$, while on the other hand we also have

$$\begin{aligned} \lim_{w \rightarrow +\infty} \frac{2 \operatorname{arcosh} \left(\frac{\cosh w + \sqrt{\cosh^2 w + 8}}{4} \right)}{w} &= 2 \cdot \lim_{w \rightarrow +\infty} \frac{\operatorname{arcosh} \left(\frac{\cosh w}{2} \right)}{w} \\ &= 2 \cdot \lim_{w \rightarrow +\infty} \frac{\operatorname{arcosh} \left(\frac{e^w}{4} \right)}{w} = 2 \cdot \lim_{w \rightarrow +\infty} \frac{\ln \left(\frac{e^w}{4} \right)}{w} = 2. \end{aligned}$$

\blacksquare

6 The circumradius of ordinary reduced polygons

Now we answer another question of Lassak proposed in [13] with the following upper bound for the circumradius.

Theorem 6.1 *Let P be an ordinary reduced polygon in H^2 of width w . Then,*

$$R(P) \leq \operatorname{arsinh} \left(\frac{2}{\sqrt{3}} \sqrt{\left(\frac{\cosh w + \sqrt{\cosh^2 w + 8}}{4} \right)^2 - 1} \right)$$

with equality if and only if P is a regular triangle.

Proof The hyperbolic Jung theorem (see Dekster [5]) says that

$$R(P) \leq \operatorname{arsinh} \left(\frac{2}{\sqrt{3}} \sinh \left(\frac{\operatorname{diam}(P)}{2} \right) \right).$$

Theorem 1.1, and the identity

$$\sinh(\operatorname{arcosh}(x)) = \sqrt{x^2 - 1}$$

concludes the proof. The case of the equality is clear from the equality case of Theorem 1.1. ■

As a consequence, we have the following.

Corollary 6.2 *Let P be an ordinary reduced polygon in the hyperbolic plane. Then,*

$$\frac{1}{2} < \frac{R(P)}{w} < 1.$$

Proof The first inequality follows from the monotonicity of the minimal width: the minimal width of the circumcircle is $2R(P)$, but the circumcircle is reduced, so $w < 2R(P)$.

The second inequality is equivalent with

$$\sqrt{\left(\frac{\cosh w + \sqrt{\cosh^2 w + 8}}{4}\right)^2 - 1} < \frac{\sqrt{3}}{2} \sinh w.$$

After taking the square of both sides, and applying the identity $\sinh^2 x = \cosh^2 x - 1$, the inequality we want to verify takes the form

$$\left(\frac{\cosh w + \sqrt{\cosh^2 w + 8}}{4}\right)^2 < \frac{3}{4} \cosh^2 w + \frac{1}{4}.$$

With a few simple steps, we can reorganize this inequality as

$$0 < 6 \cosh^4 w - 7 \cosh^2 w + 1 = (6 \cosh^2 w - 1)(\cosh^2 w - 1),$$

which clearly holds.

Let us observe that these constants provide the best possible bounds for the ratio of the circumradius and the minimal width. The sharpness of the first inequality comes from considering a convergent sequence P_k of regular $(2k+1)$ -gons of minimal width w , whose limit is a disk of width w and radius $2w$. As for the second inequality, we can see that

$$\begin{aligned} \lim_{w \rightarrow +\infty} \frac{\operatorname{arsinh}\left(\frac{2}{\sqrt{3}} \sqrt{\left(\frac{\cosh w + \sqrt{\cosh^2 w + 8}}{4}\right)^2 - 1}\right)}{w} &= \lim_{w \rightarrow +\infty} \frac{\operatorname{arsinh}\left(\frac{2}{\sqrt{3}} \sqrt{\left(\frac{e^w}{4}\right)^2 - 1}\right)}{w} = \\ &= \lim_{w \rightarrow +\infty} \frac{\operatorname{arsinh}\left(\frac{e^w}{2\sqrt{3}}\right)}{w} = \lim_{w \rightarrow +\infty} \frac{\ln\left(\frac{e^w}{\sqrt{3}}\right)}{w} = 1. \end{aligned}$$

■

In the Euclidean plane, Fabińska [6] proved that for an arbitrary reduced polygon of width w there is some boundary point, such that the circular disk of radius w centered at that boundary point covers the disk. Musielak [25] showed that the same

holds for spherical reduced polygons. The remainder of this section is dedicated to the proof of the same statement for ordinary reduced polygons in the hyperbolic plane. In most of the steps we can repeat Musielak's spherical argument, but we also apply a few minor adjustments. First, we will need a few lemmas.

Lemma 6.3 *For a compact set $X \subseteq H^n$ and a point $z \in \text{conv } X$, we have*

$$\bigcap_{x \in X} B(x, r) \subseteq B(z, r)$$

for any positive radius r .

Proof If $z \in X$, the statement trivially holds, so we assume $z \in \text{conv } X \setminus X$. If we consider the Beltrami–Cayley–Klein model, Euclidean and hyperbolic line segments coincide, so it is easy to see that Minkowski's theorem holds, that is, $\text{conv } X = \text{conv } E(X)$ where $E(X)$ denotes the extreme points of $\text{conv } X$. We also have $E(X) \subseteq X$, so clearly it is sufficient to prove

$$\bigcap_{x \in E} B(x, r) \subseteq B(z, r)$$

for some set $E \subseteq E(X)$.

Considering again the linearity preserving properties of the Beltrami–Cayley–Klein model, by Carathéodory's theorem there are $k \leq n + 1$ points e_1, \dots, e_k in $E(X)$ such that $z \in [e_1, \dots, e_k]$. Naturally, we can assume that $\bigcap_{i=1}^k B(e_i, r)$ is not empty, otherwise the statement is trivial.

It is easy to see that for a compact set Y , the function $d(y, \cdot)$ restricted to Y attains its maximum for some point $e \in E(Y)$. Therefore, if we choose some point $y \in \bigcap_{i=1}^k B(e_i, r)$, and we set $Y = \{e_1, \dots, e_k, z\}$, then

$$d(y, z) \leq \max_i d(y, e_i) \leq r,$$

so $y \in B(z, r)$, and that concludes the proof. ■

For a convex body $K \subset H^n$ and a positive number r , let us introduce the notation $C(K, r)$ for the set of centers such that the closed balls of radius r centered at points in this set contain K , i.e.,

$$C(K, r) = \{x \in H^n : K \subseteq B(x, r)\}.$$

Musielak's following spherical characterization [25] still holds in H^n . We will omit the proof, as his argument only uses metric considerations, and hence it remains true in H^n .

Lemma 6.4 *For a convex body $K \subset H^n$ and a positive number r ,*

$$C(K, r) = \bigcap_{e \in E(K)} B(e, r).$$

Similarly, the following lemma of Musielak [25] remains true in the hyperbolic plane as well. His inductive proof can be repeated to the letter, as hyperbolic balls of

radius r are also r -spindle convex (e.g., this is a trivial corollary of Theorem 1.2 in Böröczky, Csépai, and Sagmeister [2]).

Lemma 6.5 *If $K \subset H^n$ is a convex body obtained as the intersection of finitely many circular disks of radius r , then the boundary of K is the union of finitely many shorter circular arcs of radius r such that they all have different centers.*

For a convex body K , we also introduce the notation $E^*(K)$ as

$$E^*(K) = \{e \in E(K) : |\partial C(K, r) \cap \partial B(e, r)| > 1\},$$

where ∂X stands for the boundary of X and $|X|$ for its cardinality. Then, from Lemmas 6.4 and 6.5 we can immediately derive the following.

Corollary 6.6 *Let $P \subset H^2$ be a convex polygon. Then,*

$$C(P, r) = \bigcap_{e \in E^*(P)} B(e, r).$$

Finally, we have all the tools to prove the following.

Theorem 6.7 *Let $P \in H^2$ be an ordinary reduced polygon of minimal width w . Then, there is a boundary point $z \in \partial P$, such that $P \subset B(z, w)$.*

Proof Assuming that P is an n -gon, we use the same notations as in Section 3. We consider the set $C(P, w)$, and we aim to show that the intersection $C(P, w) \cap \partial P$ is not empty. We prove this by contradiction.

Corollary 6.2 and the hyperbolic Jung theorem (cf. Dekster [5]) implies that $C(P, w)$ intersects the interior of P . If $C(P, w)$ also contains some point in the exterior of P , then the proof is complete by the convexity of $C(P, w)$ (see Lemma 6.4). So let us assume that $C(P, w)$ is a subset of the interior of P .

Let e_1, \dots, e_m be the points of $E^*(P)$ in a positive orientation, where we understand the indices modulo m . By the assumption that $C(P, w)$ lies in the interior of P , it is easy to see from the definition of ordinary reduced polygons that $3 \leq m \leq n$. By Lemma 6.5, to each vertex e_i in $E^*(P)$, there is a shorter circular arc \mathcal{C}_i of $B(e_i, r)$ on the boundary. Let q_i be the intersection of \mathcal{C}_i and \mathcal{C}_{i+1} . Lemma 6.5 and Corollary 6.6 implies that the boundary of $C(P, w)$ is the union of short circular arcs of radius r connecting q_i and q_{i+1} for $1 \leq i \leq m$; let us denote these arcs by $\widehat{q_i q_{i+1}}$.

Each point e_i of $E^*(P)$ is a vertex $v_{\sigma(i)}$ for an injective map $\sigma: \{1, \dots, m\} \rightarrow \{1, \dots, n\}$. We set $s_i = t_{\sigma(i)}$ for $i \in \{1, \dots, m\}$. By our assumption, $s_i \in \partial B(e_i, r)$ is not in $C(P, w)$, and hence it is not contained in the arc $\widehat{q_i, q_{i+1}}$.

Clearly, the points e_1, q_2 and s_1 are not collinear, since $B(e_1, w) \supset P$, both q_2 and s_1 are boundary points of the circle $B(e_1, w)$, on the other hand, while s_1 is a boundary point of P , q_2 lies in the interior. Hence, s_1 is either in the same open hemisphere bound by the line through e_1 and q_2 as q_3 , or as q_m (or equivalently as e_2). By symmetry, we can assume without loss of generality that the first case occurs.

We also consider a diametral chord $[x, y]$ of P . We note that $x = v_i$ and $y = v_{i + \frac{n+1}{2}}$ for some $1 \leq i \leq n$ where in this case the indices are understood modulo n (cf.

Lassak [13]). Let x' and y' be the orthogonal projections of x and y to the opposite sides, respectively (i.e., $x' = t_i$ and $y' = t_{i+\frac{n+1}{2}}$). Let m_1 be the midpoint of $[x, y']$ and m_2 be the midpoint of $[x', y]$. Then, if we consider the boundary of P , one of the triples (x, m_1, y') and (x', m_2, y) are on the boundary in this order with the same orientation, we assume without loss of generality that this is the positive orientation.

We can also assume that the triple (e_1, x, e_2) are on the boundary of P in this order with respect to the positive orientation, where we allow the case $x = e_1$. This implies that (s_1, x', s_2) also have the same order in the positive orientation, as all chords $[v_i, t_i]$ half the perimeter (see Lassak [13] or Sagmeister [27]).

By our assumption that $C(P, w)$ is in the interior of P , so x' is not contained in $B(e_k, r)$ for some $2 \leq k \leq m$ (by the assumption on the position of p_3 and s_1 , we deduce $d(e_1, x') \leq d(x, x') = w$). We can also observe that there is such e_k on the opposite half-plane of the line through e_1 and p_2 as x' . Let z be the interior point of the segment $[x', e_k]$ such that $d(x', z) = d(x', x) = w$. In particular, $z \in P$. Considering the two triangles $[z, x', y]$ and $[x, x', y]$, they have two equal sides with different enclosed angle. This implies

$$d(z, y) > d(x, y) = \text{diam } P,$$

which is clearly a contradiction. ■

References

- [1] K. Bezdek and G. Blekherman, *Danzer–Grünbaum’s theorem revisited*. Period. Math. Hungar. 39(2000), nos. 1–3, 7–15.
- [2] K. J. Böröczky, A. Csépai, and Á. Sagmeister, *Hyperbolic width functions and characterizations of bodies of constant width in the hyperbolic space*. J. Geom. 115(2024), 15.
- [3] K. J. Böröczky, A. Freyer and Á. Sagmeister, *Pál’s isominwidth problem in the hyperbolic space*. Preprint (2025), [arXiv:2502.04427](https://arxiv.org/abs/2502.04427)
- [4] Q. Chen, P. Hou, and H. Jin, *Diameter and circumradius of reduced spherical polygons*. Beitr. Algebra. Geom. 64(2023), 1–9.
- [5] B. V. Dekster, *The Jung theorem for spherical and hyperbolic spaces*. Acta Math. Hungar. 67(1995), no. 4, 315–331.
- [6] E. Fabińska, *Covering a reduced polygon by a disk*. Demonstr. Math. 42(2009), no. 4, 851–856.
- [7] J. P. Fillmore, *Barbier’s theorem in the Lobachevsky plane*. Proc. Amer. Math. Soc. 24(1970), 705–709.
- [8] E. Heil, *Kleinste konvexe Körper gegebener Dicke*. fachbereich mathematik der TH darmstadt, 1978. Preprint, (453), 1–2.
- [9] Á. G. Horváth, *Diameter, width and thickness in the hyperbolic plane*. J. Geom. 112(2021), no. 3, Paper No. 47, 29.
- [10] J. Jerónimo-Castro and F. G. Jimenez-Lopez, *A characterization of the hyperbolic disc among constant width bodies*. Bull. Korean Math. Soc. 54(2017), 2053–2063.
- [11] M. Lassak, *Area of reduced polygons*. Publ. Math. 67(2005), 349–354.
- [12] M. Lassak, *Reduced convex bodies in the plane*. Israel J. Math. 70(1990), no. 3, 365–379.
- [13] M. Lassak, *Reduced polygons in the hyperbolic plane*. Arch. Math. 123(2024) 329–337.
- [14] M. Lassak, *Reduced spherical polygons*. Colloq. Math. 138(2015), no. 2, 205–216.
- [15] M. Lassak, *Width of convex bodies in hyperbolic space*. Results Math. 79(2024), no. 3, 1–15.
- [16] M. Lassak and H. Martini, *Reduced bodies in Minkowski space*. Acta Math. Hungar. 106(2005), nos. 1–2, 17–26.
- [17] M. Lassak and H. Martini, *Reduced bodies in Euclidean space—A survey*. Expo. Math. 29(2011), 204–219.
- [18] M. Lassak and H. Martini, *Reduced convex bodies in finite dimensional normed spaces: A survey*. Results Math. 66(2014), 405–426.

- [19] M. Lassak and M. Musielak, *Reduced spherical convex bodies*. Bull. Polish Acad. Sci. Math. 66(2018), 87–97.
- [20] K. Leichtweiss, *Curves of constant width in the non-Euclidean geometry*. Abh. Math. Sem. Univ. Hamburg 75(2005), 257–284.
- [21] C. Liu and Y. Chang, *Several problems on reduced spherical polygons of thickness less than $\frac{\pi}{2}$* . Preprint (2020). [arXiv:2204.06317](https://arxiv.org/abs/2204.06317).
- [22] C. Liu, Y. Chang, and Z. Su, *The area of reduced spherical polygons*. Rocky Mountain J. Math. 52(2022), no. 2, 599–607.
- [23] H. Martini and W. Wenzel, *Tetrahedra are not reduced*. Appl. Math. Lett. 15(2002), no. 7, 881–884.
- [24] H. Martini and K. J. Swanepoel, *Non-planar simplices are not reduced*. Publ. Math. Debrecen 64(2004), nos. 1–2, 101–106.
- [25] M. Musielak, *Covering a reduced spherical polygon by a disk*. Rend. Circ. Mat. Palermo (2) 73(2024), no. 1, 65–70.
- [26] J. Pál, *Ein minimumproblem für ovale*. Math. Ann. 83(1921), 311–319.
- [27] Á. Sagmeister, *On the area of ordinary hyperbolic reduced polygons*. Publ. Math. 107(2025). accepted.
- [28] L. A. Santaló, *Note on convex curves in the hyperbolic plane*. Bull. Am. Math. Soc. 51(1945), 405–412.

Bolyai Institute, University of Szeged, Aradi vértanúk tere 1, H-6720 Szeged, Hungary

e-mail: sagmeister.adam@gmail.com