## INSTRUCTIONS FOR CONTRIBUTORS

## SUBMISSION OF MANUSCRIPTS

Manuscripts should be submitted online via our manuscript submission and tracking site, http://www.editorialmanager.com/psm/. Full instructions for electronic submission are available directly from this site. To facilitate rapid reviewing, communications for peer review will be electronic and authors will need to supply a current e-mail address when registering to use the system.

Papers for publication from Europe (except those on genetic topics, irrespective of country), and all papers on imaging topics, should be submitted to the UK Office.

Papers from the Americas, Asia, Africa, Australasia and the Middle East (except those dealing with imaging topics), and all papers dealing with genetic topics, irrespective of country, should be sent to US Office.

Generally papers should not have text more than 4500 words in length (excluding these sections) and should not have more than a combined total of 5 tables and/or figures. Papers shorter than these limits are encouraged. For papers of unusual importance the editors may waive these requirements. A structured abstract of no more than 250 words should be given at the beginning of the article using the headings: Background; Methods; Results; Conclusions. The name of an author to whom correspondence should be sent must be indicated and a full postal address given in the footnote. Any acknowledgements should be placed at the end of the text (before the References section).

Declaration of Interest: A statement must be provided in the acknowledgements listing all financial support received for the work and, for all authors, any financial involvement (including employment, fees, share ownership) or affiliation with any organization whose financial interests may be affected by material in the manuscript, or which might potentially bias it. This applies to all papers including editorials and letters to the editor.

Contributors should also note the following:

- 1. S.I. units should be used throughout in text, figures and tables.
- 2. Authors should spell out in full any abbreviations used in their manuscripts.
- 3. Foreign quotations and phrases should be followed by a translation.
- 4. If necessary, guidelines for statistical presentation may be found in: **Altman DG, Gore SM, Gardner MJ & Pocock SJ** (1983). Statistical guidelines for contributors to medical journals. *British Medical Journal* 286, 1489–1493.

**REFERENCES** (1) The Harvard (author-date) system should be used in the text and a complete list of References cited given at the end of the article. In a text citation of a work by more than two authors cite the first author's name followed by et al. (but the names of all of the authors should be given in the References section). Where several references are cited together they should be listed in rising date order. (2) The References section should be supplied in alphabetical order (authors' names in **bold**, journal titles in full), following the text. Some examples follow:

Miller PM, Byrne M, Hodges A, Lawrie SM, Johnstone EC (2002). Childhood behaviour, psychotic symptoms and psychosis onset in young people at high risk of schizophrenia: early findings from the Edinburgh high risk study. *Psychological Medicine* 32, 173–179.

Cleckley HJ (1941). The Mask of Sanity, 2nd edn. Mosby: St. Louis, MO.

Brewer WJ, Wood SJ, DeLuca C, Pantelis C (2006). Models of olfaction for exploring neurodevelopment. In *Olfaction and the Brain* (ed. W. J. Brewer, D. Castle and C. Pantelis), pp. 97–121. Cambridge University Press: Cambridge.

(3) Online citations

doi (when published online prior to printed issue)

Lauritsen MB, Pedersen CB, Mortensen CB (2004). The incidence and prevalence of pervasive developmental disorders: a Danish population-based study. *Psychological Medicine*. Published online: 21 October 2004. doi:10.1017/S0033291704002387.

URL

World Bank (2003). Quantitative techniques for health equity analysis – Technical Notes (http://siteresources.worldbank. org/INTPAH/Resources/Publications/Quantitative-Techniques/health.eq tn07.pdf). Accessed 15 February 2006.

[Authors are requested to print-out and keep a copy of any online-only material, in case the URL changes or is no longer maintained.]

FIGURES AND TABLES Only essential figures and tables should be included. Further tables, figures, photographs and appendices, may be included with the online version on the journal website. To ensure that your figures are reproduced to the highest possible standards, Cambridge Journals recommends the following formats and resolutions for supplying electronic figures. Please ensure that your figures are saved at final publication size and are in our recommended file formats. Following these guidelines will result in high quality images being reproduced in both the print and the online versions of the journal. Line artwork: Format: tif or eps, Colour mode: black and white (also known as 1-bit), Resolution: 1200 dpi; Combination artwork (line/tone): Format: tif or eps, Colour mode: grayscale (also known as 8-bit), Resolution: 800 dpi; Black and white halftone artwork: Format: tif, Colour mode: grayscale (also known as 8-bit), Resolution: 300 dpi; Colour halftone artwork: Format: tif, Colour mode: CMYK colour, Resolution: 300 dpi. All photographs, graphs, and diagrams should be referred to as figures and should be numbered consecutively in Arabic numerals. Captions for figures should be typed double-spaced on separate sheets. Tables Tables should be typed above the table.

**PROOFS AND OFFPRINTS** Page proofs will be sent to the author designated to receive correspondence. corrections other than to printer's errors may be charged to the author. The corresponding author of each paper will receive a PDF file of their article and hard copy offprints may be purchased if they are ordered on the form supplied when the proof is returned.

## PSYCHOLOGICAL MEDICINE

## CONTENTS

1573	Problematic alcohol use and hippocampal volume in a female sample: disentangling cause from consequence using a co-twin control study design Wilson S, Malone SM, Hunt RH, Thomas KM & lacono WG	1673
	Sex similarities and differences in risk factors for recurrence of major depression van Loo HM, Aggen SH, Gardner CO & Kendler KS	1685
1592	Stabilising sleep for patients admitted at acute crisis to a psychiatric hospital (OWLS): an assessor-blind pilot randomised controlled trial	
	Sheaves B. Freeman D. Isham L. McInerney J. Nickless A.	
	Yu L-M, Rek S, Bradley J, Reeve S, Attard C, Espie CA, Foster R, Wirz-Justice A, Chadwick E & Barrera A	1694
1608	Minority stressors, rumination, and psychological distress in monozygotic twins discordant for sexual minority status  Timmins L. Rimes KA & Rahman O	1705
	Clinical predictors of conversion to bipolar disorder in	
1616	a prospective longitudinal familial high-risk sample: focus on depressive features	
	Frankland A, Roberts G, Holmes-Preston E, Perich T, Levy F, Lenroot R, Hadzi-Pavlovic D, Breakspear M & Mitchell PB	1713
1624	Are there distinct cognitive and motivational sub-groups of children with ADHD?  Lambek R. Sonuga-Barke F. Tannock R. Sørensen AV	
	Damm D & Thomsen PH	1722
	Proin glutamata in madication froe depressed nationts.	
	• • • • • • • • • • • • • • • • • • • •	
1634	Godlewska BR, Masaki C, Sharpley AL, Cowen PJ & Emir UE	1731
	A dimensional approach to assessing psychiatric risk in adults born very preterm	
	Kroll J, Froudist-Walsh S, Brittain PJ, Tseng C-EJ, Karolis V,	
	Murray RM & Nosarti C	1738
1644	Correspondence	1745
1655		
	1592 1608 1616 1624 1634	female sample: disentangling cause from consequence using a co-twin control study design  Wilson S, Malone SM, Hunt RH, Thomas KM & lacono WG  Sex similarities and differences in risk factors for recurrence of major depression  van Loo HM, Aggen SH, Gardner CO & Kendler KS  Stabilising sleep for patients admitted at acute crisis to a psychiatric hospital (OWLS): an assessor-blind pilot randomised controlled trial  Sheaves B, Freeman D, Isham L, McInemey J, Nickless A, Yu L-M, Rek S, Bradley J, Reeve S, Attard C, Espie CA, Foster R, Wirz-Justice A, Chadwick E & Barrera A  Minority stressors, rumination, and psychological distress in monozygotic twins discordant for sexual minority status  Timmins L, Rimes KA & Rahman Q  Clinical predictors of conversion to bipolar disorder in a prospective longitudinal familial high-risk sample: focus on depressive features  Frankland A, Roberts G, Holmes-Preston E, Perich T, Levy F, Lenroot R, Hadzi-Pavlovic D, Breakspear M & Mitchell PB  Are there distinct cognitive and motivational sub-groups of children with ADHD?  Lambek R, Sonuga-Barke E, Tannock R, Sørensen AV, Damm D & Thomsen PH  Brain glutamate in medication-free depressed patients: a proton MRS study at 7 Tesla  Godlewska BR, Masaki C, Sharpley AL, Cowen PJ & Emir UE  A dimensional approach to assessing psychiatric risk in adults born very preterm  Kroll J, Froudist-Walsh S, Brittain PJ, Tseng C-EJ, Karolis V, Murray RM & Nosarti C  Correspondence



1664



sample and meta-analysis

Ho R, Kua E-H, Power MJ & Feng Li

Directional associations between memory impairment and depressive symptoms: data from a longitudinal

Yu J, Lim H-Y, Abdullah FN d/o M, Chan H-M, Mahendran R,