

BRIEF REPORT

Evacuation of Intensive Care Units During Disaster: Learning From the Hurricane Sandy Experience

Mary A. King, MD, MPH; Molly V. Dorfman, MD, MPH; Sharon Einav, MD; Alex S. Niven, MD; Niranjan Kissoon, MD; and Colin K. Grissom, MD

ABSTRACT

Objective: Data on best practices for evacuating an intensive care unit (ICU) during a disaster are limited. The impact of Hurricane Sandy on New York City area hospitals provided a unique opportunity to learn from the experience of ICU providers about their preparedness, perspective, roles, and activities.

Methods: We conducted a cross-sectional survey of nurses, respiratory therapists, and physicians who played direct roles during the Hurricane Sandy ICU evacuations.

Results: Sixty-eight health care professionals from 4 evacuating hospitals completed surveys (35% ICU nurses, 21% respiratory therapists, 25% physicians-in-training, and 13% attending physicians). Only 21% had participated in an ICU evacuation drill in the past 2 years and 28% had prior training or real-life experience. Processes were inconsistent for patient prioritization, tracking, transport medications, and transport care. Respondents identified communication (43%) as the key barrier to effective evacuation. The equipment considered most helpful included flashlights (24%), transport sleds (21%), and oxygen tanks and respiratory therapy supplies (19%). An evacuation wish list included walkie-talkies/phones (26%), lighting/electricity (18%), flashlights (10%), and portable ventilators and suction (16%).

Conclusions: ICU providers who evacuated critically ill patients during Hurricane Sandy had little prior knowledge of evacuation processes or vertical evacuation experience. The weakest links in the patient evacuation process were communication and the availability of practical tools. Incorporating ICU providers into hospital evacuation planning and training, developing standard evacuation communication processes and tools, and collecting a uniform dataset among all evacuating hospitals could better inform critical care evacuation in the future. (*Disaster Med Public Health Preparedness*. 2016;10:20-27)

Key words: disaster, evacuation, critical care, hurricanes, transportation of patients, emergency preparedness

The published experience of health care providers involved in disasters over the past decade has repeatedly emphasized the importance of planning, preparation, and evacuation procedures in these situations. Critically ill patients are especially vulnerable during disasters, because they are both medically fragile and their care and evacuation require special equipment and expertise. Insufficient coordination of resources and patient monitoring during movement of critically ill patients are often cited as reasons for clinical deterioration during transport.¹⁻⁵ Simulation of patient preparation and transport using actual transport equipment increases the likelihood of successful individual evacuation performance as well as system-wide cooperation.^{6,7}

Little is known about provider preparedness or the performance of effective and safe evacuation of critically ill patients during disasters. Intensive care

unit (ICU) providers rarely consider the challenges of moving a critically ill patient within or outside of the hospital in the event that their ICU is rendered incapable of supporting appropriate ongoing care. Although ICU providers are familiar with horizontal transport (moving ICU patients laterally on the same floor), they are less familiar with vertical transport (moving patients either up or down floors), especially in disaster situations that require transport up or down stairs. Although many patients have common critical care requirements (such as mechanical ventilation or vasopressor infusion) that would be routine for most receiving hospitals, some patients require specialized care that may exceed the capability of transport teams and receiving locations (eg, extracorporeal life support [ECLS], intracranial pressure management, and micro-prematurity neonatal care). Deliberate planning is the only way to appropriately address the complex challenges that ICU providers may face in a disaster.

Hurricane Sandy was the deadliest storm outside of the southern United States since 1972,⁸ claiming more than 250 lives. Although the extent of its devastation could not have been fully predicted, the New York City area hospitals most affected received ample warning through news media of the cyclone's impending landfall on November 5, 2012. Hurricane Sandy therefore provides a unique opportunity to evaluate critical care provider preparedness and experience with this single major evacuation event from multiple hospital perspectives and to inform future critical care disaster evacuation planning. The objectives of this report were to describe (1) ICU provider preparedness efforts in New York City prior to ICU evacuation, (2) provider roles and activities during ICU evacuation following Hurricane Sandy, and (3) ICU provider perspectives on their evacuation preparedness, tools, and systems.

METHODS

Following study approval by the Seattle Children's Hospital Institutional Review Board, phone or e-mail contact was attempted with all medical ICU directors from identified New York City area hospitals affected by Hurricane Sandy. Respondents were asked to voluntarily participate in a research survey. Directors who consented were asked to distribute an anonymous electronic survey tool to their ICU nursing, respiratory therapy, and physician staff that they knew to be involved in the Hurricane Sandy ICU evacuation.

Study Tool

An electronic cross-sectional survey addressing ICU evacuation preparations and practices was developed by the study principal investigator. Content validity was ensured by focusing questions on issues raised in previous articles describing ICU evacuation.^{6,9-15} Consensual validity was achieved through the review of 7 internal medicine, critical care, and disaster response experts and serial adaptation of the questionnaire on the basis of their suggestions. The final structured questionnaire contained 23 items and included both open and closed questions (see Appendix 1 in the **online data supplement**).

Respondents were asked to provide demographic information including their job title or leadership position and their role during the hurricane. They next answered a series of questions using a 5-point anchored Likert scale on pre-event ICU planning and preparation, evacuation coordination and leadership, ICU evacuation processes, and major performance stressors. Vertical evacuation was a particular area of focus because of the prominent experience and challenge with critically ill patients in this area in other published disaster experiences.^{4,9,11,12} Finally, the respondents were asked to describe the key highlights of their disaster response experience in a series of open-ended questions including the following: (1) major successes, barriers, and problems that they encountered; (2) specific tools, supplies, and equipment

that were helpful or missing; and (3) any ethical concerns that they encountered.

Study Population

Via a literature search of the press and through consultation with ICU physicians and private industry, we identified 7 New York area hospitals that had been forced to evacuate ICU patients as a result of infrastructural damage during Hurricane Sandy. In alphabetical order, those hospitals were (1) Bellevue Hospital, (2) Coney Island Hospital, (3) Good Samaritan Hospital, (4) Manhattan VA Medical Center, (5) New York Downtown Hospital, (6) NYU Langone Medical Center, and (7) Staten Island Hospital. The American College of Chest Physicians provided the research team with the contact details for the medical ICU directors of these institutions (telephone numbers and e-mails). All ICU directors were initially approached by e-mail, and 2 attempts to contact each were made by e-mail and phone. ICU directors who agreed to participate were asked to complete a questionnaire and to electronically distribute a similar survey to the nurses, respiratory therapists, and physicians who had played direct roles in the evacuation. The final sample size of the group was based on the principle of theoretical saturation in qualitative research.¹⁶

Survey Method

A uniquely identified electronic version of the survey (SurveyMonkey, Palo Alto, CA) was e-mailed to each responding ICU director, who in turn distributed the survey to the relevant staff within the institution. The survey opened December 3, 2012, and closed March 28, 2013. All surveys were completed electronically and anonymously.

Data Collection and Statistical Analysis

Questionnaire responses were collected and analyzed by using Microsoft Office EXCEL 2007 (Microsoft Corp, Redmond, WA) and STATA 10.1 (STATA Corp, College Station, TX). Open-ended responses were coded for thematic penetrance by 2 independent reviewers. The number of interviewees repeating each comment was documented. Because there are common themes on evacuation between adult medical, pediatric, and neonatal ICUs, we decided to include all responses in our analysis. Descriptive statistics included counts and proportions. Percentages were calculated from the total number of respondents. Responses are presented primarily by aggregate totals to observe entire cohort trends and by individual hospital for ICU evacuation and leadership.

RESULTS

Four medical ICU directors from 4 different hospitals that had evacuated their ICUs because of Hurricane Sandy agreed to participate in the study. Given the extenuating circumstances during December 2012 for many of these hospitals when this survey was distributed, the ICU directors were not asked to

provide the number of staff members who received the questionnaire; hence, the survey denominator is unknown. Seventy-five medical professionals who had participated in the Hurricane Sandy ICU evacuation initially responded and 68 (91%) completed the questionnaire (hospital A = 7, hospital B = 11, hospital C = 24, hospital D = 26).

The demographics of the respondents are provided in Table 1. Most respondents (n = 56, or 63%) worked in medical or mixed ICUs. Although we had targeted adult ICUs, 3% of the respondents stated that they were working primarily in neonatal or pediatric ICUs (n = 2). More than one-third of the respondents were physicians (n = 9 [13%] attending physicians and n = 17 [25%] physicians-in-training). Almost half of the respondents were ICU nurses (n = 24, or 35%) and respiratory therapists (n = 14, or 21%). A third of the respondents (n = 23, or 34%) held leadership roles during the ICU evacuation defined as (1) Incident Command leadership role, (2) ICU nursing manager (RN), or (3) ICU medical director (MD). The 23 respondents with leadership roles included 9 ICU nurses, 3 respiratory therapists, 7 ICU physicians, 1 chief resident, 2 ICU residents, and 1 chief medical officer.

ICU Evacuation Planning and Preparation

Responses on this topic are summarized in Table 2. Most respondents (n = 41, or 61%) stated that their ICU had been at least “somewhat” involved in regional evacuation planning prior to Hurricane Sandy, and 51% of the respondents felt that they were at least “somewhat” adequately trained to evacuate ICU patients. In contrast, less than one-fourth of the respondents reported participating in an ICU evacuation drill in the past 2 years (n = 11 [22%]; Figure 1), and among these, only 5 reported prior hands-on evacuation experience using actual equipment and simulated patients.

Although all 4 hospitals housed their ICU patients above ground level, very few (n = 19, or 28%) had any vertical evacuation training or prior experience. A total of 72% of respondents (n = 48) reported that they had never participated in any type of vertical evacuation training or experience. As presented in Figure 2, despite the admitted lack of practical training or prior experience, only 23% of respondents (n = 15) felt that they had been inadequately trained to perform a vertical ICU evacuation.

Almost two-thirds of the respondents (n = 43, or 64%) were directly involved in patient preparation or actual patient transfer following Hurricane Sandy. Providers with more direct involvement in patient transfer during Hurricane Sandy were more likely to report that they felt adequately trained to perform these roles. Only 1 of the 16 respondents who prepared patients for transfer and 3 of the 17 responders who performed actual patient transfer felt inadequately trained. These individuals also felt better trained for vertical

TABLE 1

Provider Demographics by Provider Type, Unit Type, and Leadership Role During Hurricane Sandy ICU Evacuation, Both by Hospital and for Total Respondents^a

Respondent Demographics	Hospital A	Hospital B	Hospital C	Hospital D	Total, N (%)
Provider					
ICU RN	4	9	4	7	24 (35)
RT	0	0	8	6	14 (21)
ICU attending	2	1	4	1	8 (12)
Associate CMO	0	0	0	1	1 (1)
ICU fellow	0	0	2	0	2 (3)
Chief resident	0	0	0	2	2 (3)
ICU resident	0	0	6	7	13 (19)
Other	1	1	0	2	4 (6)
Total	7	11	24	26	68 (100)
Unit					
Medical ICU	3	0	10	16	29 (43)
Multi-patient type ICU	1	8	9	8	27 (40)
Neonatal/pediatric ICU	2	0	1	0	2 (3)
Surgery/trauma ICU	1	0	3	1	5 (7)
Cardiac ICU	0	0	1	0	1 (1)
ED	0	1	0	0	2 (1)
Other	0	2	0	1	1 (1)
Total	7	11	24	26	68 (100)
Leadership role					
Yes	4	9	10	8	23 (34)
No	3	2	14	18	45 (66)
Total	7	11	24	26	68 (100)
Vertical evacuation role					
Vertical evacuation leadership	0	1	3	4	8 (12)
Preparing patients for vertical evacuation	0	4	6	6	16 (24)
Actually moving patients/equip	3	3	7	6	19 (28)
I did not participate	3	3	7	8	21 (31)
Other	1	0	1	1	3 (4)
Total	7	11	24	25	67 (100)

^aAbbreviations: CMO, chief medical officer; ED, emergency department; ICU, intensive care unit; RT, respiratory therapist. The total number of respondents was 68.

evacuation compared with respondents who were not directly involved in these activities. The 8 responders who provided direct vertical evacuation leadership, however, were more ambiguous regarding their preparation: 3 felt adequately trained, 3 felt inadequately trained, and 2 were neutral.

ICU Evacuation Coordination and Leadership

Most respondents (n = 57, or 83%) felt that they had good situational awareness during the event (Supplemental Table 1 in the **online data supplement**). More than half (57%) mentioned teamwork as one of their primary successes, although the response varied among hospitals (A: 71%, B: 27%, C: 50%, D: 73%). By contrast, less than one-third of the respondents (29%) mentioned leadership as a primary success (A: 14%, B: 0%, C: 38%, D: 38%). Communication was identified as a major barrier to success by almost half (43%) of respondents (A: 43%, B: 27%, C: 54%, D: 38%), and 7% mentioned the lack of communication tools (ie, walkie-talkies/phones) as an additional barrier. Two-thirds of these ICU respondents (n = 42, or 65%) felt that the ICU

TABLE 2

Provider Responses to ICU Evacuation Planning and Preparation Questions, Both by Hospital and for Total Respondents ^a					
ICU Evacuation Planning and Preparation Responses	Hospital A	Hospital B	Hospital C	Hospital D	Total, N (%)
Our ICU was involved with regional hospital evacuation planning					
Yes	2	7	6	18	33 (49)
Somewhat	0	1	3	4	8 (12)
Not involved	1	0	7	2	10 (15)
I don't know	4	3	8	2	17 (25)
Total	7	11	24	26	68 (100)
I drilled ICU evacuation in the past 2 years using:					
Bodies and equipment	0	1	0	2	3 (4)
Manikins and equipment	0	2	0	0	2 (3)
Manikins only	0	0	0	0	0
Paper patients	0	0	0	1	1 (1)
Tabletop	0	0	4	1	5 (7)
I have not drilled	6	8	19	19	52 (78)
Other	1	0	1	2	4 (6)
Total	7	11	24	25	67 (100)
Prior to Sandy, I participated in a vertical evacuation in the form of:					
Vertical evacuation training	1	3	2	2	8 (12)
Vertical evacuation drill	0	0	0	1	1 (1)
Training and drill	0	1	0	1	2 (3)
Prior real vertical evacuation	0	0	2	6	8 (12)
I have not participated prior	6	7	20	15	48 (72)
Total	7	11	24	25	67 (100)
I felt adequately trained to perform evacuation					
Yes, agree completely	1	3	3	9	16 (25)
Yes, agree somewhat	2	3	7	5	17 (26)
Neutral	1	2	8	6	17 (26)
No, disagree somewhat	1	1	3	4	9 (14)
No, disagree completely	2	1	2	1	6 (9)
Total	7	10	23	25	65 (100)

^aAbbreviation: ICU, intensive care unit. The survey included questions on regional planning participation, ICU evacuation drills and prior experience, and evacuation training. The total number of respondents was 68.

director should be the ICU evacuation leader rather than the chief medical officer, the Incident Commander, emergency medical services, federal authorities, or another source.

ICU Evacuation Processes

Details of survey responses are summarized in Table 3. Almost half of the respondents reported that patient triage and prioritization methods were primarily determined during rather than prior to the disaster (n = 32, or 47%). A quarter reported that patient triage and prioritization was performed by using predetermined ICU criteria (n = 16, or 24%), whereas the rest responded that triage was performed by using "other" criteria (n = 8, or 12%) or random selection (n = 2, or 3%). Some respondents (n = 10, or 15%) were not privy to the triage process. Although the survey did not provide direct insight into the effectiveness of these triage practices, none of the respondents identified the patient distribution process as a primary success, and 6 of the respondents (5 from a single hospital) commented that they had to deal with ethical issues during patient distribution. One best practice identified through ICU director comments was the manual

FIGURE 1





FIGURE 2



documentation of ICU resource requirements for each patient on a ruled pad of paper during evacuation. This documentation tool was considered essential by the respondents at that institution and was used iteratively during evacuation to provide real-time situational awareness of both patient numbers and resource requirements to inform distribution and transport decisions.

TABLE 3

Provider Responses to ICU Evacuation Process Questions, Both by Hospital and for Total Respondents^a

ICU Evacuation Process	Hospital A	Hospital B	Hospital C	Hospital D	Total, N (%)
How were patients prioritized for evacuation?					
Predetermined criteria	2	1	6	7	16 (24)
Determined during disaster	3	6	10	13	32 (47)
Random	0	0	1	1	2 (3)
I don't know	0	4	5	1	10 (15)
Other	2	0	2	4	8 (12)
Total	7	11	24	26	68 (100)
To track and document evacuated patients and equipment, we used:					
Evacuation or disaster forms	0	4	1	9	14 (21)
Typical transfer forms	0	1	8	10	19 (28)
No forms	1	0	1	0	2 (3)
Ad hoc tracking	2	1	5	0	8 (12)
I don't know	3	5	8	6	22 (33)
Other	1	0	1	0	2 (3)
Total	7	11	24	25	67 (100)
We sent medications with the patient:					
For transport time	0	2	3	10	15 (22)
For special meds up to 24 h	3	0	5	2	10 (15)
Did not send medications	0	5	3	7	15 (22)
I don't know	2	4	11	3	20 (30)
Other	2	0	2	3	7 (10)
Total	7	11	24	25	67 (100)
I continued care in transport or receiving facility					
Yes	6	4	7	16	33 (51)
No	1	6	16	9	32 (49)
Total	7	10	23	25	65 (100)
I continued care at receiving facility because:					
Happened to be credentialed	0	2	0	1	3 (5)
Specifically credentialed prior	0	1	0	2	3 (5)
Emergency credentialed	4	0	4	6	14 (22)
I did not continue care	2	7	19	10	38 (58)
Other	1	0	0	6	7 (11)
Total	7	10	23	25	65 (100)

^aAbbreviation: ICU, intensive care unit. The survey included questions on patient prioritization, patient tracking, medications, and delivering ongoing patient care in transport or at receiving facility. The total number of respondents was 68.

During transfer, providers more frequently used transfer forms (n = 19, or 28%) common to their daily practice to track patients instead of special evacuation or disaster forms (n = 14, or 21%). Some respondents reported “I don't know” (n = 22, or 33%) or no forms/ad hoc/other (n = 12, or 18%) to this question, but interestingly, only 5% of respondents mentioned transfer records as a major barrier to successful patient transfer.

More than one-third of the respondents (n = 25, or 37%) reported that they sent medications with patients for the immediate 24 hours after transfer. Over half of the respondents (n = 33, or 51%) continued to provide patient care within their skill set either during transport or at the receiving facility, despite the fact that very few were credentialed (n = 3, or 5%) or specifically pre-credentialed (n = 3, or 5%) to perform these duties. Two thirds of the respondents who performed inter-facility patient transfer

TABLE 4

Main Evacuation Successes and Barriers and Tools, Equipment, and Systems Either Helpful or Needed as Identified by Respondents in Free Text and Categorized^a

Main Evacuation Successes		Main Evacuation Barriers	
Teamwork	57%	Communication	43%
Good Leadership	29%	Disaster knowledge/training	10%
Patient Safety	16%	Walkie-talkies/ phones	7%
Communication	9%		
Manpower	6%		
Helpful Tools, Equipment, Systems		Tools, Equipment, Systems Wish List	
Flashlights	24%	Walkie-talkies/phones	26%
Med Sleds/transport devices	21%	Lighting/electricity	18%
Oxygen tanks/RT supplies	19%	Flashlights	10%
Ambulances	9%	Portable ventilators	9%
Portable ventilators	9%	Elevators	7%
Walkie-talkies/phones	7%	Portable suction/Buretrol	7%
Batteries	7%	Disaster knowledge/training	7%
Tracking forms	7%	Electronic record	6%
Lighting/electricity	6%	Communication	6%
Patient distribution process	6%	Water	6%

^aAbbreviation: RT, respiratory therapy. Data are presented as the mean percentage of respondents mentioning each category and weighted by hospital.

required emergency credentialing at the receiving facility (n = 14, or 22%). Only 15% of respondents across all hospitals mentioned patient safety during the transfer process as a primary success.

Evacuation Tools

Respondents were asked to list which tools were most helpful during the evacuation process and which were most sorely missed. The responses to these questions are presented in Table 4. The most helpful ICU-specific equipment was transport sleds (21%) and oxygen tanks or respiratory therapy supplies (19%), and the most helpful general evacuation tools were flashlights (24%). The ICU-specific wish list most commonly included portable ventilators and suction (16%), whereas the general evacuation wish list included a greater variety of tools, eg, walkie-talkies/phones (26%), lighting/electricity (18%), and flashlights (10%).

Provider Performance Stressors

Respondents listed lack of water, food, and toilets; personal issues (eg, concern regarding family members); and limited appropriate means to dispose of the dead as major stressors affecting their performance during the ICU evacuation process.

Key Thematic Issues

Respondents identified a long list of key considerations in the survey comment section. These are presented in Supplemental Table 2 in the **online data supplement**.

DISCUSSION

To our knowledge, this is the first study to systematically gather survey data from an interdisciplinary group of ICU providers from multiple institutions to analyze a regional evacuation response to a major disaster event. Our study had several strengths, including diligent pursuit of surveys within 4 months of the event, input from ICU providers from diverse disciplines who were directly involved in the ICU evacuation process, and the elaboration of themes concordant with reports on similar events. These facts allowed us to analyze our survey with some confidence to identify important associations and lessons that may help to inform future ICU evacuation practices. Although it has been suggested that practical equipment is integral to evacuation successes, ours is also the first study to highlight that flashlights, water, walkie-talkies, and other specific “wish list” items may be just as important as (if not more so) complicated ICU-specific equipment (such as transport ventilators).¹⁷ These small, simple tools may play a more significant role in successful ICU evacuations than previously appreciated and ICUs should be included in basic disaster tool planning.

Despite a number of natural disasters in recent years and an increasing emphasis on disaster preparedness by regulatory agencies like the Joint Commission,¹⁸ our survey responses reflect common areas for improvement that have been raised by reports following similar events. Although most institutions surveyed had participated in regional evacuation planning and had significant advance warning of Hurricane Sandy’s landfall, leadership coordination and communication remained significant problems. Triage and prioritization decisions for ICU patient evacuation were generally still determined locally, and most providers performing these activities were not designated leaders and had little or no practical training. It is significant to note that institutional evacuation planning may not always involve the ICU director, because disaster preparedness efforts are often focused on the emergency department. This may create a management challenge for critically ill patients, who are generally the most vulnerable during an evacuation and frequently require specific subject matter expertise to effectively deliver their complex care. The insufficient and slow dissemination of critical information may contribute to continued challenges in creating efficient, streamlined disaster evacuations.

Given multiple reports of lack of regional coordination of hospital resources during disaster-induced hospital evacuations^{19–24} as well as the numerous responses in this survey about the lack of patient distribution coordination, we recommend that disaster planning agreements include prevent local mapping of regional critical care capacity, specialty resources, and ground and air transport with critical care capabilities. For large or late evacuations like Hurricane Sandy, these resources should be reported by each hospital within the region to a centralized Incident Command System or regional evacuation command center.²⁵ The Incident

Command System can determine how to best utilize the existing resources and will be responsible for coordination using stereotyped evacuation patterns based on volume, severity, and special needs of critically ill patients.^{22,26} Our survey results also suggest that continuity of provider care throughout the patient transfer process contributed to the overall perception of safer care. A priori regional credentialing of appropriate ICU staff as part of the overall disaster plan could avoid the challenges of emergency credentialing that the Hurricane Sandy ICU providers and hospitals faced.

System-wide cooperation should be encouraged and practiced, especially in health care systems that do not use a standardized coordinating system as part of their daily routine. Drills should include ICU providers in order to test command and communication systems and should include preparation and transport of simulated patients using actual transport equipment.^{6,7,27} In our survey, the reported lack of concern with such technically challenging practices as vertical evacuation underlines the importance of systematic training, for which detailed recommendations and core competencies have already been published.^{28,29} These drills would not only increase provider preparedness but also help to identify important tools that were missing in the Hurricane Sandy experience, such as flashlights, transport sleds, walkie-talkies or phones, portable ventilators, and suction.

The benefits of a systematic approach to patient prioritization and resource requirements demonstrated by the practical experience of one of the participating hospitals underlines the benefits of a evacuation preparation checklist and tracking system, which are readily available.^{30–35} Providers who are stressed will typically revert to familiar daily routines and processes. The fact that respondents more frequently reported the use of routine transfer forms over disaster forms represents either suboptimal preparation, with the potential for process improvement, or poor correlation between the content and structure of the forms and real-time documentation requirements. The use of disaster forms should be either reconsidered or studied more critically in simulated and real-time events. Lessons from Haiti and Toronto suggest that electronic tracking may be more effective than paper forms^{36,37}; however, disaster conditions (such as those that existed in Hurricane Sandy) may necessitate a manual or paper version. If a decision is made to continue working with disaster forms, the staff should undergo appropriate training and routinely practice the use of these forms. Detailed planning and realistic drills using likely disaster scenarios increase successful individual performance.^{30,38} Communication and leadership are frequently cited challenges to effective ICU management and evacuation following a disaster. The experience from survey respondents following Hurricane Sandy highlights the importance of a critical care team or unit leader, who can maintain real-time situational awareness of the quantity, severity, resource requirements, special equipment, and transport needs of multiple critically ill patients and communicate this

information effectively to the Incident Command System. The critical care unit leader role in the Hospital Emergency Incident Command System was eliminated under the new Hospital Incident Command System, and revision^{28,29,38} of this change according to the Hurricane Sandy experience warrants serious consideration. Additionally, integration into the framework as defined by the Institute of Medicine's Crisis Standards of Care may be beneficial for alignment with other regional hospitals and planning bodies.^{25,39,40}

Our survey did not delve into specific assessment of post-traumatic stress or dysfunction. We highlighted some of the ethical obstacles identified by providers, but did not address their functional status or stress symptoms at the time of the survey, which would have been a ripe opportunity to assess long-term effects of such evacuations. We also did not allow providers to self-rate their performance in terms of patient care, although responses in support of excellent patient safety outcomes indirectly addressed this. A complete description and analysis of the complex events and ICU evacuation responses following Hurricane Sandy is clearly not possible by using a simple survey tool. Owing to financial limitations and sheer geographic distance, we were forced to conduct our study remotely. A major limitation to our study was the participation of only 4 of 7 area hospitals affected by this storm, and the associated risk of selection and response biases. Although we attempted to highlight a variety of subspecialty ICU types, we primarily captured responses from medical ICUs as a result of the sampling methods. Future studies could aim to provide more responses from neonatal, pediatric, medical, cardiac, and other ICUs.

Responses by certain specialty providers, such as respiratory therapists, may have biased the prioritization of some of the tools on the wish lists, for example. Even among the hospitals that agreed to participate, we encountered significant limits to available data due to respondent concern regarding the appropriate use of this information. Outside providers, such as emergency medical services or purely administrative staff, were not queried but might provide a different perspective on the events. By definition, data collected in a disaster situation are frequently limited and nonuniform, raising concerns regarding the public relations and medical-legal ramifications of submitting this information to public scrutiny. Developing clear guidelines for minimum elements of data collection to consider in these situations and an anonymous reporting system at the national or international level may help to facilitate more comprehensive study and understanding of best practices for ICU evacuation that may aid providers and improve patient outcomes in future disaster events.

CONCLUSIONS

The experiences shared by ICU providers involved in the response to Hurricane Sandy identified themes common with other published literature in this area. Disaster planning and

preparation at the ICU level continues to represent an area for improvement, especially in the area of vertical evacuation. Early initiation of patient evacuation efforts led to greater communication and coordination both within and between hospitals, but still left room for improvement in the areas of communication and leadership, especially in the area of patient distribution. Triage, prioritization, and documentation processes were highly variable, and emergency credentialing was frequently necessary to provide patients with the care that they needed during and after transport to new facilities. Phones, flashlights, and portable ventilators are important, high-yield resources that could be targeted for preparation in the event of future ICU evacuations. Many other areas for improvement above could be addressed by better efforts to incorporate ICU providers into hospital evacuation planning and training efforts. A possible solution to improve the ICU processes observed in this event includes the creation of standardized evacuation communication processes and documentation tools, managed by a critical care unit leader who coordinates care and plays a defined and integrated role in the Incident Command System. The development of a standard, uniform database utilized by all evacuating hospitals during a disaster could also better inform patient and system outcomes to further advance critical care evacuation practices in the future.

About the Authors

Harborview Medical Center and Seattle Children's, Seattle, Washington (Dr King), Seattle Children's, Seattle, Washington (Dr Dorfman), Shaare Zedek Medical Center, Jerusalem, Israel (Dr Einav), Madigan Army Medical Center, Tacoma, Washington (Dr Niven), British Columbia Children's Hospital, Vancouver, Canada (Dr Kissoon); and Intermountain Medical Center, Murray, Utah (Dr Grissom).

Correspondence and reprint requests to Mary A. King, MD, MPH, Department of Pediatrics, Box 359774, Harborview Medical Center, 325 Ninth Avenue, Seattle WA 98104 (e-mail: maryking@u.washington.edu).

Acknowledgments

We would like to especially acknowledge the willingness and effort of the ICU medical directors who helped facilitate the participation of their ICU staff as well as the ICU providers themselves who were willing to share their experiences with the medical community. We also thank Dr Asha Devereaux, Dr Jeffrey Dichter, and Dr Jeffrey Upperman for their guidance and expertise in generation of the survey questions.

Supplementary material

To view supplementary material for this article, please visit <http://dx.doi.org/10.1017/dmp.2015.94>

Published online: August 27, 2015.

REFERENCES

1. Orlando S, Bernard ML, Mathews P. Neonatal nursing care issues following a natural disaster: lessons learned from the Katrina experience. *J Perinat Neonatal Nurs.* 2008;22(2):147-153. doi:10.1097/01.JPN.0000319102.20593.12

2. Koyama A, Fuse A, Hagiwara J, et al. Medical relief activities, medical resourcing, and inpatient evacuation conducted by Nippon Medical School due to the Fukushima Daiichi Nuclear Power Plant accident following the Great East Japan Earthquake. *J Nippon Med Sch.* 2011; 78(6):393-396.
3. Farmer JC, Carlton PK. Providing critical care during a disaster: the interface between disaster response agencies and hospitals. *Crit Care Med.* 2006;34(3 Suppl):S56-S59. doi:10.1097/01.CCM.0000199989.44467.2E
4. Taylor IL. Hurricane Katrina's impact on Tulane's teaching hospitals. *Trans Am Clin Climatol Assoc.* 2007;118:69-78.
5. Tanaka H, Iwai A, Oda J, et al. Overview of evacuation and transport of patients following the 1995 Hanshin-Awaji earthquake. *J Emerg Med.* 1998;16(3):439-444.
6. Gildea JR, Etengoff S. Vertical evacuation simulation of critically ill patients in a hospital. *Prehosp Disaster Med.* 2005;20(4):243-248.
7. Cybulski P. Evacuation of a critical care unit. *Dynamics.* 2003;14(3): 21-23.
8. Blake ES, Kimberlain TB, Berg RJ, et al. Tropical Cyclone Report Hurricane Sandy (AL182012) 22-29 October 2012. National Hurricane Center Data Archive. http://www.nhc.noaa.gov/data/tcr/AL182012_Sandy.pdf. Published February 12, 2013. Accessed July 30, 2015.
9. Sternberg E, Lee GC, Huard D. Counting crises: US hospital evacuations, 1971-1999. *Prehosp Disaster Med.* 2004;19(2):150-157.
10. Daugherty EL, Rubinson L. Preparing your intensive care unit to respond in crisis: considerations for critical care clinicians. *Crit Care Med.* 2011;39(11):2534-2539. doi:10.1097/CCM.0b013e3182326440
11. Schultz CH, Koenig KL, Lewis RJ. Implications of hospital evacuation after the Northridge, California, earthquake. *N Engl J Med.* 2003; 348(14):1349-1355. doi:10.1056/NEJMsa021807
12. Murphy GRF, Foot C. ICU fire evacuation preparedness in London: a cross-sectional study. *Br J Anaesth.* 2011;106(5):695-698. doi:10.1093/bja/aer033
13. Uppal A, Evans L, Chitkara N, et al. In search of the silver lining: the impact of Superstorm Sandy on Bellevue Hospital. *Ann Am Thorac Soc.* 2013;10(2):135-142. doi:10.1513/AnnalsATS.201212-116OT
14. Iserson KV. Vertical hospital evacuations: a new method. *South Med J.* 2013;106:37-42. doi: 10.1097/SMJ.0b013e31827caef4
15. Manion P, Golden JJ. Vertical evacuation drill of an intensive care unit: design, implementation, and evaluation. *Disaster Manag Response.* 2004;2:14-19.
16. Glaser BG, Strauss AL. The constant comparative methods of qualitative analysis. In Glaser BG, Strauss AL. *The Discovery of Grounded Theory: Strategies for Qualitative Research.* New York, NY: Aldine DeGruyter; 1967.
17. Aghababian R, Lewis CP, Gans L, et al. Disasters within hospitals. *Ann Emerg Med.* 1994;23(4):771-777.
18. The Joint Commission. Standing Together: An Emergency Planning Guide for America's Communities. http://www.jointcommission.org/Standing_Together__An_Emergency_Planning_Guide_for_Americas_Communities/. Published May 17, 2006. Accessed July 30, 2015.
19. Iwashyna TJ, Christie JD, Moody J, et al. The structure of critical care transfer networks. *Med Care.* 2009;47(7):787-793. doi:10.1097/MLR.0b013e318197b1f5
20. Lavon O, Hershko D, Barenboim E. Large-scale air-medical transport from a peripheral hospital to level-1 trauma centers after remote mass-casualty incidents in Israel. *Prehosp Disaster Med.* 2009;24(6):549-555.
21. Carlascio DR, McSharry MC, LeJeune CJ, et al. Air medical response to the 1990 Will County, Illinois, tornado. *J Air Med Transp.* 1991;10(10): 7, 9-11, 13-16.
22. Fuzak JK, Elkon BD, Hampers LC, et al. Mass transfer of pediatric tertiary care hospital inpatients to a new location in under 12 hours: lessons learned and implications for disaster preparedness. *J Pediatr.* 2010; 157(1):138-143.e2. doi:10.1016/j.jpeds.2010.01.047
23. Cryer HG, Hiatt JR, Eckstein M, et al. Improved trauma system multicase incident response: comparison of two train crash disasters. *J Trauma.* 2010;68(4):783-789. doi:10.1097/TA.0b013e3181d03b8c
24. Kanter RK. Regional variation in critical care evacuation needs for children after a mass casualty incident. *Disaster Med Public Health Prep.* 2012;6(2):146-149. doi:10.1001/dmp.2012.30
25. Burkle FM, Hsu EB, Loefer M, et al. Definition and functions of health unified command and emergency operations centers for large-scale bioevent disasters within the existing ICS. *Disaster Med Public Health Prep.* 2007;1(2):135-141. doi:10.1097/DMP.0b013e3181583d66
26. Born CT, Briggs SM, Ciraulo DL, et al. Disasters and mass casualties: I. General principles of response and management. *J Am Acad Orthop Surg.* 2007;15(7):388-396.
27. Yurt RW, Lazar EJ, Leahy NE, et al. Burn disaster response planning: an urban region's approach. *J Burn Care Res.* 2008;29(1):158-165. doi:10.1097/BCR.0b013e31815f2b8c
28. Hospital Incident Command System Guidebook. California Emergency Medical Services Authority website. http://www.emsa.ca.gov/disaster_medical_services_division_hospital_incident_command_system_resources. Accessed March 8, 2015.
29. Incident Command System Resources. Federal Emergency Management Agency website. <https://www.fema.gov/incident-command-system-resources>. Accessed June 8, 2015.
30. Hoskins JD, Graham RF, Robinson DR, et al. Mass Casualty Tracking with Air Traffic Control Methodologies. *J Am Coll Surg.* 2009;208(6): 1001-1008.
31. Blake N, Stevenson K. Reunification: keeping families together in crisis. *J Trauma.* 2009;67(2 Suppl):S147-S151. doi: 10.1097/TA.0b013e3181af0c13
32. Buono CJ, Chan TC, Killeen J, et al. Comparison of the effectiveness of wireless electronic tracking devices versus traditional paper systems to track victims in a large scale disaster. *AMIA Annu Symp Proc.* 2007 Oct11:886.
33. DeMers G, Kahn C, Buono C, et al. Secure scalable disaster electronic medical record and tracking system. *Prehosp Emerg Care.* 2013;28(5): 498-501. doi: 10.1017/S1049023X13008686
34. Hamilton J. An Internet-based bar code tracking system: coordination of confusion at mass casualty incidents. *Disaster Manag Response.* 2003;1:25-28.
35. Maltz J, C, Ng T, Li D, et al. The Trauma Patient Tracking System: implementing a wireless monitoring infrastructure for emergency response. *Conf Proc IEEE Eng Med Biol Soc.* 2005;3:2441-2446.
36. Callaway DW, Peabody CR, Hoffman A, et al. Disaster Mobile Health Technology: Lessons from Haiti. *Prehosp Disaster Med.* 2012;27(02):148-152.
37. MacDonald RD, Farr B, Neill M, et al. An emergency medical services transfer authorization center in response to the Toronto severe acute respiratory syndrome outbreak. *Prehospital Emerg Care.* 2004;8(2):223-231.
38. Verni C. A hospital system's response to a hurricane offers lessons, including the need for mandatory interfacility drills. *Health Aff (Millwood).* 2012;31(8):1814-1821.
39. Powell T, Hanfling D, Gostin LO. Emergency preparedness and public health: the lessons of Hurricane Sandy. *JAMA.* 2012;308(24):2569-2570.
40. *Crisis Standards of Care: A Toolkit for Indicators and Triggers.* Institute of Medicine. <http://www.iom.edu/Reports/2013/Crisis-Standards-of-Care-A-Toolkit-for-Indicators-and-Triggers.aspx>. Published July 31, 2013. Accessed June 10, 2015.