

# ON ALGEBRAS OF DOMINANT DIMENSION ONE

BRUNO J. MUELLER

**Summary.**  $QF$ -3 algebras  $R$  are classified according to their second commutator algebras  $R'$  with respect to the minimal faithful module, which satisfy  $\text{dom.dim. } R' \geq 2$ . The class  $C(S)$  of all  $QF$ -3 algebras whose second commutator is  $S$ , contains besides  $S$  only algebras  $R$  with  $\text{dom.dim. } R = 1$ .  $C(S)$  contains a unique (up to isomorphism) minimal algebra which can be represented as a subalgebra  $S_0$  of  $S$  describable in terms of the structure of  $S$ , and  $C(S)$  consists just of the algebras  $S_0 \subset R \subset S$  (up to isomorphism). A criterion for  $S_0 \neq S$  and various examples are given. Finally it is shown that the injective hull of  $S$  (as left-, right- or bimodule) is at the same time the injective hull for every  $R \in C(S)$ . This result sheds some light on the fact that  $\text{dom.dim. } S \geq 2$  while  $\text{dom. dim. } R = 1$  for all  $R \in C(S)$ ,  $R \neq S$ : We prove that no composition-factor of the  $R$ -module  $R'/R$  is isomorphic to an ideal.

**The classes  $C(R)$ .** We consider finite-dimensional algebras  $R$  with unit over a field  $K$  and unitary finitely generated  $R$ -modules.  $QF$ -3 algebras are characterized by the existence of a minimal faithful right-module  $X$  which is (unique up to isomorphism and) a direct summand in every faithful module.  $X$  is projective-injective and the sum of the isomorphism-types of dominant<sup>1)</sup> right-ideals, hence itself a right-ideal generated by an idempotent:  $X_R \cong eR_R$ . The  $K$ -dual  $X^*$  of  $X$  is the minimal faithful left-module:  ${}_R X^* \cong {}_R Rf$ . With every  $QF$ -3 algebra  $R$  one associates the second commutator  $R'$  of the minimal faithful (right-)module  $X$ , which is again a  $QF$ -3 algebra and contains  $R$  as a subalgebra, with the same unit, in a natural way:  $1 \in R \subset R'$ . The second commutator of the minimal faithful left-module  ${}_R X^*$  is isomorphic to  $R'$ , over  $R$ . Minimal faithful  $R'$ -modules are  $R'f = Rf$ ,  $eR' = eR$ . (cf. Thrall [6], Morita [3], Tachikawa [5])

The following *dominant dimension* is introduced for every algebra  $R$ :

---

Received February 21, 1967.

<sup>1)</sup> A *dominant* right-ideal is an ideal  $e_1R$  generated by a primitive idempotent  $e_1$ , which is injective.

$\text{dom.dim. } R \geq n$  if there exists an exact sequence  $0 \rightarrow R \rightarrow X_1 \rightarrow \cdots \rightarrow X_n$  of projective-injective modules  $X_i$ . It was shown in [4] that the three such dimensions obtained by using left-modules, right-modules or bimodules coincide. *QF-3* algebras are characterized by  $\text{dom.dim. } R \geq 1$ . The following are equivalent for any *QF-3* algebra  $R$ :  $R = R'$ ;  $\text{dom.dim. } R \geq 2$ ;  $R$  is the endomorphism-ring of a finitely generated fully faithful module<sup>2)</sup>. Hence the inclusion  $R \subset R'$  embeds every *QF-3* algebra  $R$  into an algebra  $R'$  with  $\text{dom.dim. } R' \geq 2$ , and the embedding is proper if and only if  $\text{dom.dim. } R = 1$ . This observation suggests the following classification:

**DEFINITION.** For any algebra  $R$  with  $\text{dom.dim. } R \geq 2$ , let  $C(R)$  denote the class of all *QF-3* algebras  $S$  such that  $S' \cong R$ .

**THEOREM 1.** An algebra  $R$  belongs to  $C(R)$  if and only if it is isomorphic to a subalgebra  $R_1$  of  $R$  that contains the unit 1 and suitable minimal faithful ideals  $eR$ ,  $Rf$  of  $R$ .

*Proof.* Morita ([3], Theorem 17.3) has shown that any  $R_1$  satisfying those conditions is *QF-3* and that  $eR = eR_1$ ,  $Rf = R_1f$  are its minimal faithful modules. Hence

$$\text{Endo}(eR_{1R_1}) = eR_1e = eRe = \text{Endo}(eR_R)$$

and

$R = R' = \text{Endo}(eRe eR) = \text{Endo}(eR_1e eR_1) = R'_1$ , proving  $R_1 \in C(R)$ . (We remark for later application (proof of theorem 8) that this identification of  $R$  and  $R'_1$  is compatible with the embeddings of  $R_1$  into  $R$  and  $R'_1$ . For  $R_1 \subset R'_1 = \text{Endo}(eR_1e eR_1)$  by  $R_1 \ni r_1 \rightarrow (x \rightarrow xr_1) \in \text{Endo}(eR_1e eR_1)$  and  $R = \text{Endo}(eRe eR)$  by  $R \ni r \rightarrow (x \rightarrow xr) \in \text{Endo}(eRe eR)$ , thus  $R_1 \subset R = \text{Endo}(eRe eR)$  again by  $R_1 \ni r_1 \rightarrow (x \rightarrow xr_1) \in \text{Endo}(eRe eR)$ .) Conversely, another result by Morita (Theorem 17.5) says that any *QF-3* algebra  $S$ , as subalgebra of  $S'$ , contains suitable minimal faithful ideals  $eS'$ ,  $S'f$  of  $S'$ .

**DEFINITION.** For any algebra  $R$  with  $\text{dom.dim. } R \geq 2$ , let  $C_0(R)$  denote the set of all subalgebras  $R_1$  of  $R$  containing the unit 1 and suitable minimal faithful ideals  $eR$ ,  $Rf$  of  $R$ .

<sup>2)</sup> A module  ${}_A X$  is *fully faithful* if it contains every indecomposable injective or projective module as a direct summand ( $X$  is a generator-cogenerator).

COROLLARY 2.  $C(R)$  and  $C_0(R)$  contain the same isomorphism-types of algebras.  $R_1 \in C_0(R)$ ,  $R_1 \subset R_2 \subset R$  implies  $R_2 \in C_0(R)$ .

Because of these facts it is of particular interest to characterize the minimal algebras in  $C_0(R)$ .

DEFINITION. For any QF-3 algebra  $R$ , a pair of idempotents  $e, f$  will be called properly chosen if

- (1)  $eR, Rf$  are minimal faithful modules,
- (2)  $ef = fe$  (this implies that  $ef$  is again an idempotent),
- (3) the number  $k$  in a decomposition  $ef = e_1 + \dots + e_k$  into indecomposable orthogonal idempotents  $e_i$  is minimal (compared to all other pairs  $e', f'$  satisfying (1) and (2). For fixed  $ef$ ,  $k$  is obviously the same for each such decomposition).

The set of primitive idempotents of any algebra  $R$  falls into finitely many isomorphism-classes  $E_1, \dots, E_n$  where two primitive idempotents  $e_i, e_j$  are called isomorphic if they generate isomorphic right- (equivalent left-) ideals. Every decomposition of the unit 1 into primitive orthogonal idempotents can be written as  $1 = \sum_{i=1}^n \sum_{j_i=1}^{n_i} e_{ij_i}$  where  $e_{ij_i} \in E_i$  and the numbers  $n_i$  are the same for any such decomposition. Given two decompositions

$$1 = \sum_{i=1}^n \sum_{j_i=1}^{n_i} e_{ij_i} = \sum_{i=1}^n \sum_{j_i=1}^{n_i} e_{i^*j_i}^*$$

there exists an inner automorphism of  $R$ , generated by an invertible element  $x \in R$ , that maps  $e_{ij_i}$  onto  $e_{i^*j_i}^*$ :  $xe_{ij_i}x^{-1} = e_{i^*j_i}^*$ .

LEMMA 3. A pair of idempotents  $e, f$  is properly chosen if and only if it is of the form  $e = \sum_{i \in I} e_{i1}$ ,  $f = \sum_{i \in J} e_{in_i}$ , where  $1 = \sum_{i=1}^n \sum_{j_i=1}^{n_i} e_{ij_i}$  is a decomposition into primitive orthogonal idempotents and the sets  $I, J \subset \{1, \dots, n\}$  characterize those classes  $E_i$  that generate dominant right-, left-ideals.

Proof. Suppose that  $e, f$  are properly chosen.  $ef = fe$  implies that  $e - ef, f - ef, ef, 1 - e - f + ef$  constitute a decomposition of 1 into orthogonal idempotents which can be refined to a decomposition into primitive orthogonal idempotents  $1 = \sum_{i=1}^n \sum_{j_i=1}^{n_i} e_{ij_i}$ . A suitable adjustment of the second index  $j_i$  gives  $e = \sum_{i \in I} e_{i1}$  and  $f = \sum_{i \in J} e_{ik_i}$ , hence  $ef = \sum_{\substack{i \in I \cap J \\ k_i=1}} e_{i1}$ . The minimality-requirement (3) implies  $k_i \neq 1$  whenever possible, that is for  $n_i > 1$ .

Therefore the minimal  $k$  in (3) is the number of elements  $i \in I \subset J$  with  $n_i = 1$ , and a further adjustment of the second index leads to  $f = \sum_{i \in J} e_{in_i}$ . Conversely any pair  $e, f$  of this type satisfies (1) and (2):  $ef = \sum_{\substack{i \in I \cap J \\ n_i = 1}} e_{i1} = fe$ , and this decomposition has the minimal number of summands, so (3) holds and  $e, f$  are properly chosen.

**THEOREM 4.** *Any two minimal subalgebras in  $C_0(R)$  are isomorphic under an inner automorphism of  $R$ . A subalgebra  $R_0$  of  $R$  is minimal in  $C_0(R)$  if and only if it is of the form  $R_0 = K + eR + Rf + RfeR$  where  $e, f$  are properly chosen idempotents of  $R$ .*

*Proof.* Let  $R_0$  be a minimal algebra in  $C_0(R)$ . By definition of  $C_0(R)$  there exist minimal faithful ideals  $eR, Rf$  of  $R$ , contained in  $R_0$ ; further the unit  $1$  of  $R$  lies in  $R_0$ . Hence  $R_0 \supset K + eR + Rf + RfeR$ , and as this is an algebra in  $C_0(R)$  too,  $R_0 = K + eR + Rf + RfeR$  because of the minimality of  $R_0$ .

We shall show that  $e, f$  can be replaced by a properly chosen pair. Refine  $1 = e + (1 - e)$  and  $1 = f + (1 - f)$  to decompositions into primitive orthogonal idempotents  $1 = e_1 + \dots + e_m = f_1 + \dots + f_m$  of  $R_0$ . We get an inner automorphism of  $R_0$ :  $xf_i x^{-1} = e_i$ ;  $x, x^{-1} \in R_0$ . Set  $f' = x f x^{-1} \in R_0$ , then  $ef' = f'e$ . Observe that

$$Rf \ni rf \rightarrow r f x^{-1} = r x^{-1} f' \in Rf'$$

is a  $R$ -isomorphism, thus  $Rf'$  is a minimal faithful module for  $R$ . Further  $R_0 \supset R_0 f' = R_0 x f x^{-1} = R_0 f x^{-1} = Rf x^{-1} = Rf'$ ; hence  $K + eR + Rf' + Rf'eR \subset R_0$  and consequently  $K + eR + Rf' + Rf'eR = R_0$ .

$e, f'$  may still not satisfy (3). But as before, the orthogonal idempotents  $e - ef', f' - ef', ef', 1 - e - f' - ef'$  can be refined in  $R$  to  $1 = \sum_{i=1}^n \sum_{j=1}^{n_i} e_{ij_i}$  with  $e = \sum_{i \in I} e_{i1}$ ,  $f' = \sum_{i \in J} e_{ik_i}$ . The second index can be adjusted such that  $k_i = 1$  or  $= n_i$ , and  $k_i \neq n_i$  for  $i \in I \subset J$  at most. There exists an inner automorphism of  $R$  interchanging  $e_{i1}$  and  $e_{in_i}$  for  $i \in I \subset J$ ,  $k_i \neq n_i$  and leaving all other  $e_{ij_i}$  fixed:  $e_{in_i} = z e_{i1} z^{-1}$ . Replacing  $f'$  by  $f'' = \sum_{i \in J} e_{in_i} = z f' z^{-1}$  we get  ${}_R R f'' \cong {}_R R f'$  and  $f'' e = e f''$  so that  $e, f''$  are properly chosen. Finally  $Re_{in_i} = R z e_{i1} z^{-1} = R e_{i1} z^{-1} \subset R f' e R$  for  $k_i \neq n_i$ ; hence  $K + eR + Rf'' +$

$Rf''eR \subset R_0$  and therefore  $K + eR + Rf'' + Rf''eR = R_0$ , proving that every minimal algebra in  $C_0(R)$  is of the form stated in the theorem.

From Lemma 3 it is obvious that whenever  $e, f$  and  $e^*, f^*$  are two properly chosen pairs of idempotents of  $R$ , then there exists an inner automorphism of  $R$  mapping  $e$  onto  $e^*$  and  $f$  onto  $f^*$ . That completes the proof of the theorem.

**DEFINITION.** For any QF-3 algebra  $R$  with  $\text{dom.dim. } R \geq 2$  and any particular minimal subalgebra  $R_0$  in  $C_0(R)$ , let  $C(R; R_0)$  denote the set of all algebras  $R_1$  with  $R_0 \subset R_1 \subset R$ .

**COROLLARY 5.**  $C(R; R_0)$  and  $C(R)$  contain the same isomorphism-types of algebras.

*Proof.* Any  $S \in C(R)$  is isomorphic to some  $R_1 \in C_0(R)$  which contains a minimal subalgebra  $R_{10}$ .  $R_{10}$  is isomorphic to  $R_0$  by an inner automorphism of  $R$  which carries  $R_1$  into an algebra  $R_2$  in  $C(R; R_0)$ .

**REMARKS.** We collect a few additional (obvious) facts about  $C(R)$ .

(i) The (up to isomorphism unique) minimal algebra  $R_0$  in  $C(R)$  is characterized by the fact that its vector-space-dimension over  $K$  is minimal among the algebras in  $C(R)$ .

(ii)  $R$  is characterized in  $C(R)$  by having maximal  $K$ -dimension.

(iii) While  $\text{dom.dim. } R \geq 2$ , we have  $\text{dom.dim. } S = 1$  for all  $S \in C(R)$  that are not isomorphic to  $R$ .

(iv) If a QF-3 algebra is a ring-direct sum  $R = R_1 \oplus R_2$ , then so is  $R' = R'_1 \oplus R'_2$ . On the other hand if  $R' = S_1 \oplus S_2$ , then  $R$  need not decompose accordingly.

(v) For any QF-3 algebra  $R$  a minimal algebra  $R_0$  in  $C(R')$  can be constructed directly as  $R_0 = K + eR + Rf + RfeR$  where  $e, f$  is any properly chosen pair of idempotents in  $R$ .

This may not be quite obvious: Since there exists a minimal subalgebra  $R_0 = K + eR' + R'f + R'feR' \subset R$  with suitable properly chosen idempotents  $e, f$  of  $R'$ , we get  $R_0 = K + eR + Rf + RfeR$  and  $e, f \in R$ .  $R'f = Rf$ ,  $eR' = eR$  are minimal faithful ideals for  $R$  as well as for  $R'$ . A decomposition  $ef = e_1 + \dots + e_k$  into primitive orthogonal idempotents in  $R'$  always lies in  $R$ , hence constitutes such a decomposition with respect to  $R$ ; and

vice versa. Suppose  $k$  be not minimal for  $R$ ; then the isomorphism-type of at least one  $e_i$ , say  $e_1$ , appears more than once in a decomposition of  $1$  in  $R$ , and we have  $e_1 R_R \cong e'_1 R_R$ ,  $e_1 e'_1 = 0$ . We get an inner automorphism of  $R$  that interchanges  $e_1$  and  $e'_1$  and leads to a  $R'$ -isomorphism  $e_1 R' \cong e'_1 R'$ , contrary to the assumption that  $e, f$  be properly chosen in  $R'$ . Thus  $e, f$  automatically are properly chosen with respect to  $R$ . — Any other properly chosen pair  $e^*, f^*$  in  $R$  can be mapped onto  $e, f$  by an inner automorphism of  $R$  and leads to an algebra  $K + e^*R + Rf^* + Rf^*e^*R$  isomorphic to  $R_0$ .

We want to derive a criterion for  $R = R_0$ . For properly chosen idempotents  $e, f$  in  $R$  we set  $ef = d$ ,  $e - ef = e'$ ,  $f - ef = f'$ ,  $1 - e - f + ef = \varepsilon$ . Then evaluation of  $(d + e' + f' + \varepsilon)R(d + e' + f' + \varepsilon) = R = R_0 = K + e'R + Rf' + RdR$  yields the necessary and sufficient condition  $f'Re' + f'R\varepsilon + \varepsilon R'e' + \varepsilon R\varepsilon = f'RdRe' + f'RdR\varepsilon + \varepsilon RdRe' + \varepsilon RdR\varepsilon + K\varepsilon$ , which may be split into the four conditions  $f'Re' = f'RdRe'$ ,  $f'R\varepsilon = f'RdR\varepsilon$ ,  $\varepsilon R'e' = \varepsilon RdRe'$ ,  $\varepsilon R\varepsilon = \varepsilon RdR\varepsilon + K\varepsilon$ . By construction of  $d = ef$ , the isomorphism-types of the idempotents in  $d$  are different from those in  $e', f'$  and  $\varepsilon$ ; hence there doesn't exist any epimorphism of  $dR_R$  onto a direct summand of  $e'R$ ,  $f'R$  or  $\varepsilon R$ ; consequently the image of every homomorphism of  $dR$  into these modules lies in  $e'N$ ,  $f'N$ ,  $\varepsilon N$  ( $N$  being the radical of  $R$ ) and we get  $e'Rd = e'Nd$ ,  $f'Rd = f'Nd$ ,  $\varepsilon Rd = \varepsilon Nd$ . Correspondingly  $dRe' = dNe'$ ,  $dRf' = dNf'$ ,  $dR\varepsilon = dN\varepsilon$  hold; and the above four conditions imply  $f'Re' = f'NdNe' = f'N^2e'$ ,  $f'R\varepsilon = f'NdN\varepsilon = f'N^2\varepsilon$ ,  $\varepsilon R'e' = \varepsilon NdNe' = \varepsilon N^2e'$ ,  $\varepsilon R\varepsilon = \varepsilon NdN\varepsilon + K\varepsilon = \varepsilon N^2\varepsilon + K\varepsilon$ . Then  $e'R$ ,  $f'R$  cannot contain isomorphic direct summands since that would lead to a map  $e'R \rightarrow f'R$  the image of which wouldn't even be contained in  $f'N$ , hence to an element in  $f'Re'$ , not in  $f'Ne'$ . Similarly  $\varepsilon R$ ,  $f'R$  and  $e'R$ ,  $\varepsilon R$  cannot have isomorphic direct summands. Finally  $\varepsilon R$  cannot decompose directly, since  $\varepsilon = \varepsilon_1 + \varepsilon_2$  (orthogonal idempotents) and  $\varepsilon R\varepsilon = \varepsilon N^2\varepsilon + K\varepsilon$  yields  $\varepsilon_1 = x + k\varepsilon$ ,  $x \in N^2$ ; hence either  $k = 0$ ,  $\varepsilon_1 \in N^2$ ,  $\varepsilon_1 = 0$  or  $k \neq 0$ ,  $0 = x\varepsilon_2 + k\varepsilon_2$ ,  $\varepsilon_2 \in N^2$ ,  $\varepsilon_2 = 0$ .

Summarizing: We have shown that  $R = R_0$  implies that  $R$  is selfbasic and that  $\varepsilon$  is either primitive or zero. Therefore  $\varepsilon R\varepsilon$  is local and has radical  $\varepsilon N\varepsilon$ ; and the condition  $\varepsilon R\varepsilon = \varepsilon NdN\varepsilon + K\varepsilon$  gives  $\varepsilon NdN\varepsilon = \varepsilon N\varepsilon$  and  $\varepsilon R\varepsilon / \varepsilon N\varepsilon \cong K$  if  $\varepsilon \neq 0$ .

Thus we have proved one direction of the following

**THEOREM 6.** *A QF-3 algebra  $R$  is minimal in  $C(R')$  if and only if*

- (1)  $R$  is selfbasic,
- (2) there exists at most one type of idempotents  $\varepsilon$  such that  $R\varepsilon, \varepsilon R$  both are not dominant,

(3)  $f'Re' = f'NdNe'$ ;  $f'Re = f'NdN\varepsilon$ ,  $\varepsilon Re' = \varepsilon NdNe'$ ,  $\varepsilon N\varepsilon = \varepsilon NdN\varepsilon$ ,  $\varepsilon R\varepsilon / \varepsilon N\varepsilon \cong K$  (if  $\varepsilon$  exists); where  $d(e', f')$  is the sum of those idempotents  $e_i$  of a decomposition into primitive orthogonal idempotents  $1 = e_1 + \dots + e_n$  for which  $Re_i, e_i R$  are both dominant ( $e_i R$  but not  $Re_i$  is dominant;  $Re_i$  but not  $e_i R$  is dominant).

Conversely these conditions (1) to (3) immediately lead back to the former conditions for  $R = R_0$ . This completes the proof.

REMARKS. (i) We are particularly interested in the case  $\text{dom.dim. } R \geq 2$ . Here the conditions of the theorem characterize those  $R$  for which  $C(R)$  is trivial (to say it contains the isomorphism-type of  $R$  only).

(ii) Applied to  $R_0$  itself the theorem describes properties of the minimal algebras in the classes  $C(R)$ .

(iii) The conditions can be simplified in certain cases, e.g.: If  $NdN = 0$  (in particular if  $d = 0$ , which for  $\text{dom.dim. } R \geq 2$ , hence  $R = \text{Endo}({}_A X)$  means that  $A$  doesn't have any dominant ideals; or if  $N^2 = 0$ ) they reduce to  $f'Re' = f'Re = \varepsilon Re' = 0$ ,  $\varepsilon R\varepsilon = K\varepsilon$ . If  $e' = f' = 0$  (for  $\text{dom.dim. } R \geq 2$  this means that  $A$  is Frobenius) they reduce to  $\varepsilon N\varepsilon = \varepsilon N(1 - \varepsilon)N\varepsilon$ ,  $\varepsilon R\varepsilon / \varepsilon N\varepsilon \cong K$ .

EXAMPLES. The following remarks are obtained by specializing results of Harada [1] for semi-primary rings to our case of algebras; but easy direct proofs could be given as well.  $R$  denotes a QF-3 algebra,  $A$  its endomorphism-ring and  $R'$  its second commutator, both with respect to the minimal faithful module.

(i) These three statements are equivalent:  $R'$  is semi-simple;  $A$  is semi-simple; the socle of  $R$  is projective. Then, if  $e_1 R, \dots, e_k R$  represent the different types of dominant ideals, the  $D^{(i)} = e_i R e_i$  are division-rings and we have  $A = \bigoplus_{i=1}^k D^{(i)}$ ,  $R' = \bigoplus_{i=1}^k D_{n_i}^{(i)}$  (ring-direct sum of  $n_i \times n_i$ -matrix-rings over the  $D^{(i)}$ ) where  $n_i = D^{(i)}\text{-dim } e_i R$ .

(ii) Equivalent:  $R'$  is simple;  $A = D$  is a division-ring; there exists only one dominant type  $eR$  and the unique minimal subideal of  $eR$  is projective. Then  $D = eRe$  and  $R' = D_n$  where  $n = D\text{-dim } eR$ . The minimal

subalgebra  $R_0$  in  $C(D_n)$  is  $R_0 = \sum_{k=1}^n D c_{1k} + \sum_{i=2}^n D c_{in} + K(\sum_{j=2}^{n-1} c_{jj})$ ; observe  $R_0 \neq D_n$  for  $n > 1$ .

(iii)  $R'$  is simple for every indecomposable hereditary QF-3 algebra  $R$  (Mochizuki [2]). Actually  $R \in C(D_n)$  is hereditary if and only if (up to isomorphism)  $T_n \subset R \subset D_n$  where  $T_n$  denotes the algebra of (upper) triangular matrices. Any such  $R$  is of the form

$$R = \begin{pmatrix} D_{n_1} & D_{n_1, n_2} & \cdots & D_{n_1, n_k} \\ 0 & D_{n_2} & \cdots & D_{n_2, n_k} \\ \vdots & & \ddots & \vdots \\ \vdots & & & \vdots \\ 0 & 0 & \cdots & D_{n_k} \end{pmatrix}$$

**Injective hulls.** LEMMA 7. *Let  $S$  be a QF-3 algebra,  $M'$  a  $S'$ -(left-) module hence a  $S$ -module, and  $M$  a  $S$ -submodule of  $M'$  such that  $S'M = M'$ . Suppose that all simple  $S'$ -submodules of  $M'$  are isomorphic to ideals. Then the  $S'$ -injective hull  $H'$  of  $M'$  (considered as  $S$ -module) is the  $S$ -injective hull of  $M$ .*

*Proof.* Consider any simple  $S'$ -submodule  $I'$  of  $M'$ .  $I'$  being isomorphic to a  $S'$ -ideal and  $S'$  being QF-3, we get a  $S'$ -monomorphism of  $I'$  into a minimal faithful ideal  $S'f = Sf$  which yields an epimorphism  $eS = eS' \cong S'f^* \rightarrow I'^*$ . Hence  $I'^*e \neq 0$  and  $eI' \neq 0$ . But  $eM' = eS'M = eSM \subset M$ , consequently  $0 \neq eI' \subset I' \cap M$  and  $I' \cap M \neq 0$ . Furthermore the  $S'$ -injective hull  $H'(I')$  of  $I'$  is isomorphic to some  $S'f_1 = Sf_1$ ,  $f = f_1 + \cdots$ ; hence  $H'(I')$  has a unique minimal submodule  $I$  when considered as  $S$ -module. We get  $I \subset I' \cap M \subset M$  and  $S'I = I'$  since  $S'I$  is a  $S'$ -submodule of the simple  $S'$ -module  $I'$ . The  $S$ -injective hull of  $I$ , being isomorphic to  $Sf_1$ , is isomorphic to  $H'(I')$  as  $S$ -module.

Let  $\bigoplus_{k=1}^n I'_k$  be the  $S'$ -socle of  $M'$ . As we have seen, each  $I'_k$  contains a unique simple  $S$ -submodule  $I_k$  and the  $S'$ -injective hull  $H'$  of  $M'$ , being the direct sum of the  $S'$ -injective hulls  $H'(I'_k)$  of the  $I'_k$ , is isomorphic to the  $S$ -injective hull of  $\bigoplus I_k$  as  $S$ -module. Since  $\bigoplus I_k$  is semi-simple and is contained in  $M$ , it is in the socle of  $M$ :  $\text{socle}(M) = \bigoplus I_k \oplus J$ . Thus the  $S$ -injective hull of  $M$  is isomorphic to the direct sum of  $H'$  and the  $S$ -injective hull  $H(J)$  of  $J$ , as  $S$ -module. On the other hand  $M \subset M' \subset H'$  and the fact that  $H'$  is  $S$ -injective imply that the  $S$ -injective hull of  $M$  is contained in  $H'$ ; hence a  $K$ -vector-space-dimension argument yields  $H(J) = 0$  and the assertion of the lemma.

**THEOREM 8.** *Let  $R$  be a QF-3 algebra. Then the  $R'$ -injective hull  $H'$  of  $R'$  is the  $R$ -injective hull of  $R$  when considered as  $R$ -module, where all modules are either left-, right- or bimodules.*

*Proof.* Applying Lemma 7 to  $S = R$ ,  $M = R$ ,  $M' = R'$  we get the result for left-modules. A similar argument holds for right-modules.

Considering bimodules, to say modules over the enveloping algebra  $R^e = R \otimes_K R^0$ , we show that  $(R^e)'$  can be identified with  $(R')^e$  by an isomorphism which carries  $R^e$  as (natural) subalgebra of  $(R^e)'$  into  $R^e$  as subalgebra of  $(R')^e$  determined by  $R$  as (natural) subalgebra of  $R'$ . Observe  $\text{dom.dim.}(R')^e = \text{dom.dim.} R' \geq 2$  (Mueller [4], Lemma 6). We have  $1 \otimes 1^0 \in R^e \subset (R^e)'$ ; and the  $(R')^e$ -left- resp. right-modules  $R'f \otimes (eR')^0$ ,  $eR' \otimes (R'f)^0$  where  $R'f = Rf$ ,  $eR' = eR$  are minimal faithful  $R'$ - and  $R$ -ideals, are projective-injective-faithful. We have  $R'f \otimes (eR')^0 = Rf \otimes (eR)^0$ ,  $eR' \otimes (R'f)^0 = eR \otimes (Rf)^0 \subset R^e$ ; hence Theorem 1 yields  $R^e \in C((R^e)')$ , to say  $(R')^e \cong (R^e)'$ , and this isomorphism carries  $R^e$  as subalgebra of  $(R')^e$  into  $R^e$  as subalgebra of  $(R^e)'$ , as indicated above (cf. the proof of Theorem 1). Now choose  $S = R^e$ ,  $S' = (R^e)' = (R')^e$ ;  $M = R$ ,  $M' = R'$ . We get  $S'M = (R')^e R = R' R R' = R' = M'$  and a simple  $(R')^e$ -submodule of  $R'$  — a simple two-sided  $R'$ -ideal — is isomorphic to a  $(R')^e$ -ideal since the QF-3 algebra  $R'$  can be embedded as  $(R')^e$ -module into a projective module. Thus Lemma 7 yields the desired result in this case too.

Mochizuki [2] observed that for hereditary QF-3 algebras  $R$  (where  $R'$  is semi-simple),  $R'$  itself is the injective hull of  ${}_R R$  and  $R_R$ . We see that this phenomenon is rather exceptional:

**COROLLARY 9.**  *$R'$  is the injective hull of  $R$  as left- and/or right- $R$ -module if and only if  $R'$  is quasi-Frobenius.  $R'$  is the injective hull of  $R$  as  $R$ - $R$ -bimodule if and only if  $R'$  is separable.*

Theorem 8 allows the construction of the following diagram of left-, right- or bimodules:

$$\begin{array}{ccccccc}
 & & 0 & & 0 & & \\
 & & \downarrow & & \downarrow & & \\
 0 & \longrightarrow & R & \longrightarrow & H' & & \\
 & & \downarrow & & \downarrow & & \\
 0 & \longrightarrow & R' & \longrightarrow & H' & \longrightarrow & X'_2 \longrightarrow \cdots \longrightarrow X'_n \\
 & & & & \downarrow & & \\
 & & & & 0 & & 
 \end{array}$$

where all rows and columns are exact, the bottom row contains  $R'$ -homomorphisms while all other maps are  $R$ -homomorphisms; where  $H', X'_2, \dots, X'_n$  are  $R'$ -injective-projective and therefore also  $R$ -injective-projective; where  $2 \leq n = \text{dom.dim. } R' \leq \infty$ ; and where the top row cannot be extended further by  $R$ -injective-projective modules if  $R \neq R'$ . Hence the socle of the  $R$ -module  $H'/R$  must contain a simple module non-isomorphic to an ideal while the socle of  $H'/R'$  as  $R'$ - or  $R$ -module contains only simple modules isomorphic to ideals (cf. [4], proof of Lemma 7). Consequently since  $H'/R' \cong H'/R/R'/R$  as  $R$ -modules, socle  $(R'/R)$  has to contain a simple  $R$ -module non-isomorphic to an ideal. We show the following stronger fact:

**THEOREM 10.** *Let  $R$  be a QF-3 algebra. Then all composition-factors of  $R'/R$  as  $R$ -left-, right- or bimodule, are not isomorphic to ideals.*

*Proof.* We apply Lemma 7 choosing  $S$  either  $= R$  or  $= R^e$  (then identifying  $S' = (R^e)'$  with  $(R')^e$  as before) and  $M' = R', R \subset M \subset R'$  any  $S$ -submodule of  $R'$ . Then the  $S'$ -injective hull  $H'$  of  $R'$  is the  $S$ -injective hull of  $M$  when considered as  $S$ -module, and we get the exact sequence of  $S$ -modules  $0 \rightarrow M \xrightarrow{\alpha} H'$ . Suppose it can be extended to  $0 \rightarrow M \xrightarrow{\alpha} H' \xrightarrow{\beta} X$  where  $X$  is  $S$ -injective-projective. Then we get a diagram

$$\begin{array}{ccccc}
 0 & \longrightarrow & M & \xrightarrow{\alpha} & H' & \xrightarrow{\beta} & X \\
 & & \downarrow & & \downarrow & & \downarrow \\
 & & S' \otimes_S M & \xrightarrow{1_{S'} \otimes \alpha} & S' \otimes_S H' & \xrightarrow{1_{S'} \otimes \beta} & S' \otimes_S X \\
 & & \downarrow \psi & & \downarrow 1 & & \downarrow 1 \\
 & & R' & \xrightarrow{\varphi} & S' \otimes_S H' & \xrightarrow{1_{S'} \otimes \beta} & S' \otimes_S X
 \end{array}$$

where  $\psi$  is the epimorphism  $s' \otimes m \rightarrow s'm$  and  $\varphi$  is the homomorphism  $R' \rightarrow H' \rightarrow S' \otimes_S H'$ . The maps  $H' \rightarrow S' \otimes_S H', X \rightarrow S' \otimes_S X$  are  $S$ -isomorphisms since  ${}_S H', {}_S X$  are injective-projective. All squares are commutative — the one in the lower left corner because of  $s' \otimes h' = 1' \otimes s'h' \in S' \otimes_S H'$  (use the isomorphism between  $S' \otimes_S H'$  and  $H'$ ). The bottom row is a complex ( $= 0$ ) since the middle row obviously is and  $\psi$  is epimorphic. Finally  $\varphi$  is  $S'$ -monomorphic, for a simple  $S'$ -submodule  $I'$  of  $\text{Ker } \varphi$  gives  $I' \cap M \neq 0$  and  $I' \cap M \subset \text{Ker } \alpha$  since  $M \rightarrow S' \otimes_S M \rightarrow R'$  turns out to be the injection  $M \rightarrow R'$ ; but  $\alpha$  is monomorphic. Now diagram-chasing shows that  $M \rightarrow R'$  is epimorphic which is a contradiction whenever  $M \neq R'$ ; hence in this case an extension  $0 \rightarrow M \rightarrow H' \rightarrow X$  cannot exist, meaning that socle  $({}_S H' / M)$  will

contain a simple module  $J$  non-isomorphic to an ideal. Since  $J$  cannot lie in  $H'/R'$ , it has to be contained in the socle of  $R'/M$ .

Now suppose  $R \subset M \subset R'$  and that all factors of  $M/R$  are non-isomorphic to ideals. Then there exists  $M \subset M_1 \subset R'$  such that  $J \cong M_1/M$  and all factors of  $M_1/R$  are non-isomorphic to ideals. Thus the theorem is proved by induction.

#### REFERENCES

- [1] M. Harada, *QF-3 and semi-primary PP-rings I, II*. Osaka Math. J. **2**, 357–368 (1965); **3**, 21–28 (1966).
- [2] H.Y. Mochizuki, *On the double commutator algebra of QF-3 algebras*. Nagoya Math. J. **25**, 221–230 (1965).
- [3] K. Morita, *Duality for modules and its application to the theory of rings with minimum condition*. Science Reports Tokyo Kyoiku Daigaku **6**, 83–142 (1958).
- [4] B.J. Mueller, *The classification of algebras by dominant dimension*. Can. J. Math., to appear.
- [5] H. Tachikawa, *On dominant dimensions of QF-3 algebras*. Trans. Amer. Math. Soc. **112**, 249–266 (1964).
- [6] R.M. Thrall, *Some generalizations of quasi-Frobenius algebras*. Trans. Amer. Math. Soc. **64**, 173–183 (1948).

*McMaster University, Hamilton, Ontario, Canada*