

# Observations of magnetic fields in intracluster medium

Federica Govoni

INAF - Osservatorio Astronomico di Cagliari, Strada 54, Loc. Poggio dei Pini, 09012  
Capoterra (Ca), Italy  
email: fgovoni@oa-cagliari.inaf.it

**Abstract.** The presence of  $\mu\text{G}$ -level magnetic fields associated with the intracluster medium of galaxy clusters is now widely acknowledged. Our knowledge of their properties has greatly improved in the recent years thanks to both new radio observations and the developments of new techniques to interpret data.

**Keywords.** magnetic fields, galaxies: clusters: general, polarization

---

## 1. Radio halos and Faraday rotation measures

Most of what we know about intracluster magnetic fields derives from the study of radio halos and Faraday rotation measures of polarized radio galaxies located inside or behind galaxy clusters (see e.g. the reviews by Carilli & Taylor 2002, Govoni & Feretti 2004, Ferrari *et al.* 2008, Feretti *et al.* 2012).

Sensitive radio observations have revealed diffuse emission from the central regions of some merging galaxy clusters. These radio sources, which extend over Mpc scales and are called radio halos, are diffuse, low-surface-brightness, and steep-spectrum synchrotron sources with no obvious optical counterparts. To date, approximately 50 radio halos are known. They demonstrate the existence of relativistic electrons and magnetic fields spread in the intracluster medium. The assumption that radio halos have the magnetic energy density comparable to the energy density in relativistic electrons, requires a volume-averaged magnetic field  $\sim 0.1\text{--}1 \mu\text{G}$ . Sometimes the intracluster magnetic fields can be ordered on scales of hundreds of kpc, as revealed in A2255 (Govoni & Feretti 2005, Pizzo *et al.* 2011) and MACS J0717.5+3745 (Bonafede *et al.* 2009), where a polarized signal, possibly associated to the radio halo, has been detected. Actually, total intensity and polarization intensity radio halo surface brightness fluctuations are strictly related to the magnetic field power spectrum (Tribble 1991, Murgia *et al.* 2004). Thus, observations of radio halos have been used to study the structure of the cluster wide magnetic fields by comparing observations with mock halos from turbulent magnetic fields (Murgia *et al.* 2004, Govoni *et al.* 2006, Vacca *et al.* 2010, Xu *et al.* 2012).

High resolution, detailed RM images of extended cluster radio galaxies have been obtained (e.g. Eilek & Owen 2002, Taylor *et al.* 2007, Guidetti *et al.* 2008, Guidetti *et al.* 2010, Bonafede *et al.* 2010, Vacca *et al.* 2012). These data are usually consistent with central magnetic field strengths of a few  $\mu\text{G}$ , but stronger fields, with values exceeding  $\simeq 10 \mu\text{G}$ , are derived in the inner regions of relaxed cooling core clusters. By analyzing the RM of radio galaxies located at different projected distance from the cluster center, it is possible to investigate the radial decrease of the magnetic field with the gas density (e.g. Dolag *et al.* 2001). The RM distributions seen across the radio galaxies present patchy structures. The observed RM fluctuations indicate that the intracluster

magnetic field is not regularly ordered but turbulent on scales ranging from tens of kpc to  $\lesssim 100$  pc. Dedicated software tools and semi-analytical approach have been developed to constrain the magnetic field power spectrum parameters (Enßlin & Vogt, Murgia *et al.* 2004, Laing *et al.* 2008, Kuchar & Enßlin 2011). The magnetic field power spectrum can be approximated with a power law with the slope close to the Kolmogorov index in some clusters but shallower index are also observed.

In addition to detailed RM studies focused on single clusters, magnetic fields in galaxy clusters can be investigated statistically. Clarke *et al.* (2001) analyzed the average RM values as a function of source impact parameter for a sample of Abell clusters. They found a clear broadening of the RM distribution toward small projected distances from the cluster center (see also e.g. Johnston-Hollitt *et al.* 2004, Govoni *et al.* 2010), clearly indicating that most of the RM contribution comes from the intracuster medium and proving that magnetic fields are present in all galaxy clusters. Bonafede *et al.* (2011) selected a sample of massive galaxy clusters and used the NRAO VLA Sky Survey (Condon *et al.* 1998) to analyze the fractional polarization of hundreds radio sources lying at different projected distances from the cluster center. They detected a clear trend of the fractional polarization, being smaller for sources close to the cluster center and increasing with increasing distance from the cluster central regions. This trend is interpreted as the result of an higher depolarization, occurring because of the higher magnetic field and gas density at the cluster center, and can be reproduced by a magnetic field model with a central value of few  $\mu\text{G}$ .

## References

- Bonafede, A., Feretti, L., Giovannini, G., *et al.* 2009, *A&A* 503, 707  
 Bonafede, A., Feretti, L., Murgia, M., *et al.* 2010, *A&A* 513, A30  
 Bonafede, A., Govoni, F., Feretti, *et al.* 2011, *A&A* 530, A24  
 Carilli, C. L. & Taylor, G. B. 2002, *ARA&A* 40, 319  
 Clarke, T. E., Kronberg, P. P., & Böhringer, H. 2001, *ApJ* 547, L111  
 Condon, J. J., Cotton, W. D., Greisen, E. W., *et al.* 1998, *AJ* 115, 1693  
 Dolag, K., Schindler, S., Govoni, F., & Feretti, L. 2001, *A&A* 378, 777  
 Eilek, J. A. & Owen, F. N. 2002, *ApJ* 567, 202  
 Enßlin T. A. & Vogt C. 2003, *A&A* 401, 835  
 Feretti, L., Giovannini, G., Govoni, F., & Murgia, M. 2012, *A&A Rev.* 20, 54  
 Ferrari, C., Govoni, F., Schindler, S., *et al.* 2008, *Space Science Reviews* 134, 93  
 Govoni, F. & Feretti, L. 2004, *Int. J. Mod. Phys. D*, Vol. 13, N.8, p. 1549  
 Govoni, F., Murgia, M., Feretti, L., *et al.* 2005, *A&A* 430, L5  
 Govoni, F., Murgia, M., Feretti, *et al.* 2006, *A&A* 460, 425  
 Govoni, F., Dolag, K., Murgia, M., *et al.* 2010, *A&A* 522, A105  
 Guidetti, D., Murgia, M., Govoni, F., *et al.* 2008, *A&A* 483, 699  
 Guidetti, D., Laing, R. A., Murgia, M., *et al.* 2008, *A&A* 514, A50  
 Johnston-Hollitt, M., Hollitt, C. P., & Ekers, R. D. 2004, *The Magnetized Interstellar Medium*,  
*Eds: B. Uyaniker, W. Reich, and R. Wielebinski p.13*  
 Kuchar, P. & Enßlin, T. A. 2011, *A&A* 529, A13  
 Laing, R. A., Bridle, A. H., Parma, P., & Murgia, M. 2008, *MNRAS* 391, 521  
 Murgia M., Govoni F., Feretti L., *et al.* 2004, *A&A* 424, 429  
 Pizzo, R. F., de Bruyn, A. G., Bernardi, G., & Brentjens, M. A. 2011, *A&A* 525, A104  
 Taylor G. B., Fabian, A. C., Gentile, G., *et al.* 2007, *MNRAS* 382, 67  
 Tribble, P. C. 1991, *MNRAS* 253, 147  
 Vacca, V., Murgia, M., Govoni, F., *et al.* 2010, *A&A* 514, 71  
 Vacca, V., Murgia, M., Govoni, F., *et al.* 2012, *A&A* 540, 38  
 Xu, H., Govoni, F., Murgia, M., *et al.* 2012, *ApJ* 759, 40