

appearance as less attractive, body size as bigger and weight as heavier. Only symptoms of disordered eating predicted higher extent of investment in one's appearance and health. And neither IBM, nor disordered eating predicted self-evaluation of one's health and fitness or the extent of investment in fitness.

Keywords: body image; Body Mass Index; disordered eating

EPP0968

Time perspective and psychological hardiness mediate a COVID-19 related life threat

E. Nikolaev*

Social And Clinical Psychology, Ulianov Chuvash State University, Cheboksary, Russian Federation

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.1223

Introduction: Unpredictable risks of COVID-19 morbidity and mortality make people suffer from threats and fears. Are there any psychological personality traits that correlate with a decrement in such feelings?

Objectives: To specify the relationship between psychological characteristics of time perspective, hardiness and COVID-19-related life threat.

Methods: We used Zimbardo Time Perspective Inventory, Maddi Hardiness Scale and a 17-point Attitude towards COVID-19 Questionnaire to question 327 Russian university students on social networks. The survey was carried out in the second half of May 2020 in the period of increasing COVID-19 morbidity and mortality rates. The mean age of the respondents was 21.24 ± 2.84 ; most of them were females (61.16%).

Results: The survey showed that every fifth respondent had a high level of COVID-19-related life threat (21.10 %). At the same time, every tenth of the respondents (10.09%) saw no threat to their life in the situation of a fast spreading dangerous novel coronavirus infection. According to the correlation analysis, decrement in feeling personal threat related to the spread of COVID-19 was directly associated ($p < 0.05$) with low indicators on the scales of Negative Past ($r = 0.16$), Hedonistic Present ($r = 0.13$) and Fatalistic Present ($r = 0.17$). Certain inverse relation was found between the level of COVID-19 related life threat and such indicators as psychological hardiness – commitment ($r = -0.16$), and challenge ($r = -0.23$).

Conclusions: Dispositional orientation to the present and future, as well as psychological characteristics of hardiness may mediate COVID-19 related life threat; therefore, these may be used as a possible basis for preventing stress and mental disorders in population.

Keywords: time perspective; psychological hardiness; life threat; COVID-19

EPP0969

Mental health conditions in people searching for aesthetic treatments: The importance of early detection.

F. Gonçalves Viegas*, I. Figueiredo, F. Ferreira and A. Lourenço
Departamento De Saúde Mental, Hospital Prof Dr Fernando Fonseca, Amadora, Portugal

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.1224

Introduction: It's established that aesthetic dysfunctions can be associated with psychiatric conditions. In present times, considering the exponential growth of minimally invasive and accessible techniques, alongside with ideals of beauty present in everyday life through exposure in social media, the importance of early detection of mental illness and its impact on the respective outcome should be emphasized.

Objectives: To review evidence regarding psychiatric disorders in people searching for aesthetic treatments and their impact on the outcome.

Methods: Literature review using Medline database.

Results: Around 50% of individuals seeking aesthetic procedures fulfill the diagnostic criteria for psychiatric disorders. The prevalence of Body Dysmorphic Disorder (BDD) can vary from 5-15%, with some studies showing a prevalence of more than 50%. Patients with heightened BDD symptoms are less satisfied with the outcomes of aesthetic procedures which could result in exacerbation of said symptoms. With regards to eating disorders, evidence suggests the initial satisfaction following aesthetic procedures, when observed, is usually transitory, not leading to long-term changes in self-perception relating to body image, nor improving prognosis or quality of life. There's also some evidence suggesting that personality disorders may be a predictor of poor satisfaction with the results of aesthetic treatments.

Conclusions: Awareness should be raised in this matter, since psychiatric conditions are more common in patients seeking aesthetic treatments and early identification can lead to a better prognosis by providing patients with the mental health treatment they need; this could also reduce the probability of dissatisfaction and subsequent aggravation of psychiatric symptoms following aesthetic interventions.

Keywords: psychiatric conditions; aesthetic; early detection; prevention

EPP0971

The relationship between self-disorders (SDS) and depressive/anxious symptoms in a clinical sample of adolescents

E. Monducci¹, G. Colafrancesco^{2*}, C. Battaglia²,
E. Masini¹, G. De Vita² and M. Ferrara²

¹Department Of Human Neurosciences, University of Rome Sapienza, Rome, Italy and ²Department Of Human Neurosciences, section Of Child And Adolescent Neuropsychiatry, University of Rome Sapienza, Rome, Italy

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.1225

Introduction: Self-disorders (SDs) have been described as a core feature of schizophrenia-spectrum disorders. Previous studies conducted on heterogeneous clinical adult and adolescents samples demonstrated that SDs aggregate selectively in the schizophrenia spectrum disorders compared to other disorders.

Objectives: To examine the specificity of SDs for schizophrenia spectrum disorders in adolescent inpatient sample.

Methods: Fifty-five adolescent inpatients admitted to the Child Psychiatry Unit at the Sapienza University in Rome were assessed for psychopathology using Kiddie Schedule for Affective Disorders and Schizophrenia (K-SADS-PL), Structured Interview for Prodromal Syndromes (SIPS/SOPS), Examination of Anomalous Self-Experiences (EASE), Multidimensional Anxiety Scale for Children (MASC), Calgary depression scale for schizophrenia (CDSS)