

Methods Systematic review of the literature in English (Pubmed). Keywords: “adult ADHD”; “bipolar disorder”; “substance abuse”; “personality disorders”.

Results Adult ADHD has a prevalence in Europe of 3.4% and it seems to appear more frequently in developed countries. In adults, the clinical presentation differs from children. There often are symptoms of impulsivity, attention deficit, restlessness, and emotional dysregulation and risk behavior, causing difficulties in everyday functioning. The coexistence of substance abuse and the similarity between bipolar disorder and personality disorders with ADHD, translate in a difficult diagnosis, especially for patients with mild presentations.

Conclusions So far, the authors conclude that a detailed clinic history plus the validated scales for adult ADHD are the most reliable tools for diagnosis. Neuroimaging and EEG studies do not provide conclusive data to consider them as diagnostic methods.

Disclosure of interest The authors have not supplied their declaration of competing interest.

<http://dx.doi.org/10.1016/j.eurpsy.2016.01.2356>

EV1372

Leganés Psychiatric Hospital in the early twentieth century (1900–1931): An approach to healthcare activity

R. Candela Ramírez^{1,*}, A. Conseglieri Gámez², P. Vázquez de la Torre Escalera¹

¹ Fuenlabrada Hospital, Psychiatry, Fuenlabrada, Madrid, Spain

² Parla Hospital, Psychiatry, Parla, Madrid, Spain

* Corresponding author.

Introduction Leganés Psychiatric Hospital has been the subject of several studies about its institutional history and clinical activity. The first decades of the twentieth century are the less explored years; however, important events for the development and establishment of the discipline of psychiatry happened in Spain during this period.

Objectives/aims To describe the clinical and therapeutic management of inpatients admitted to Leganés National Asylum between 1900 and 1931.

Material and methods This is a retrospective case series study. We reviewed medical records found in the Historical Archives of Psychiatric Institute, Germany ($n = 1043$) of inpatients admitted between 1900 and 1931. We analyzed clinical care variables, mainly related to diagnosis and treatment, of the patients who were admitted during this period ($n = 1043$) with SPSS v21. We consulted bibliography, such as asylum documents and diverse primary and secondary literature.

Results The diagnosis of mania was very common in the early twentieth century. Lypemania nearly disappeared after 1910 and monomania was not observed in any patient. Delusions decreased after 1905. The first patient diagnosed with schizophrenia was admitted on 19th November 1921. From that moment, the use of this term increased significantly. Only 4.4% of patients admitted underwent treatment.

Conclusions We confirmed the predominance of French nosology in the early twentieth century. From 1920, an increase in German nosography was observed. The application of treatments was rare.

Disclosure of interest The authors have not supplied their declaration of competing interest.

<http://dx.doi.org/10.1016/j.eurpsy.2016.01.2357>

EV1373

Characteristics of inpatients admitted to National Asylum of Leganés in the early twentieth century (1900–1931)

R. Candela Ramírez^{1,*}, P. Vázquez de la Torre Escalera¹, A. Conseglieri Gámez²

¹ Fuenlabrada Hospital, Psychiatry, Fuenlabrada, Madrid, Spain

² Parla Hospital, Psychiatry, Parla, Madrid, Spain

* Corresponding author.

Introduction Leganés Psychiatric Hospital has been the subject of several studies about its institutional history, clinical activity and demography of its institutionalized population. The first decades of the twentieth century are the less explored years; however, important events for the development and establishment of the discipline of psychiatry happened in Spain during this period.

Objectives/aims To describe the sociodemographic and hospitalization characteristics of the patients who were admitted to Leganés National Asylum between 1900 and 1931.

Methods This is a retrospective case series study. We reviewed medical records found in the Historical Archives of Psychiatric Institute Germain ($n = 1043$) of inpatients admitted between 1900 and 1931. We analyzed sociodemographic and hospitalization related variables of medical records with SPSS v21. We consulted bibliography, such as asylum documents and diverse primary and secondary literature.

Results Most inpatients were male, single, with an average age of 38 years, came from home and were admitted as fee-paying boarders. Circa 64% of them remained in the institution until death and the average stay was 7.92 years.

Conclusions Even though the Leganés Asylum was born amidst debate on the asylum model, it did not meet the expectations. Among other reasons, it presented serious architectural deficiencies and was unable to classify inpatients according to the French tradition (agitated, dirty, quiet) or to separate populations, such as minors or criminal inpatients, thus becoming a charity institution asylum instead of a therapeutic mental hospital.

Disclosure of interest The authors have not supplied their declaration of competing interest.

<http://dx.doi.org/10.1016/j.eurpsy.2016.01.2358>

EV1374

Cyclic vomiting syndrome

P. Cano Ruiz*, S. Cañas Fraile, A. Gómez Peinado, M.D. Sánchez García

HNSPS, Psiquiatría, Albacete, Spain

* Corresponding author.

Introduction Cyclic vomiting syndrome (SVC) is a functional gastrointestinal disorder characterized by paroxysmal episodes of vomiting, recurrent and cyclical presentation. Although this disorder was first described in children, recently it has increasingly been recognized in adults.

Objective To know the pathogenesis of the syndrome and the optimum approach.

Method Theoretical review and cyclic vomiting syndrome brief statement of a case. This is a man of 51 years for 7 months has repeated episodes of vomiting often than 1 episode for week, with vomiting every 10–15 minutes. The patient is admitted to the neurology department for endless instability and multidirectional nystagmus and right hemispheric deficiency symptoms, with acute renal failure prerenal rehydration and study. While entering data semiotic Wernicke disease are objectified. With replacement therapy meeting evolves favorably at the time of discharge from hospital asymptomatic.

Discussion CVS, in conclusion, is a rare disease in adults whose diagnosis is one of exclusion. The adult presentation usually presents more durable, less frequent episodes. The pathogenesis