

Token Motor Task scores and trihexyphenidyl dose ($r_{xy} = -0.496$, p -value = 0.002). At both weeks 2 and 8, there was a negative relationship between age and Symbol Coding scores ($r_{xy} = -0.387$, p -value = 0.018; $r_{xy} = -0.35$, p -value = 0.034, respectively). Verbal Fluency scores were lower in patients with high scores on the PANSS excitement component and at week 2 ($r_{xy} = -0.42$, p -value = 0.01), this trend continued at week 8 ($r_{xy} = -0.31$, p -value = 0.063). Tower of London scores were negatively associated at week 8 with cognitive and positive PANSS scores ($r_{xy} = -0.46$, p -value = 0.004; $r_{xy} = -0.336$, p -value = 0.042, respectively). **Conclusions:** Thus, we have demonstrated that cognitive impairment in patients with schizophrenia is associated with various factors, and not only antipsychotic treatment.

Disclosure of Interest: None Declared

O0128

Changes in inflammatory parameters and their impact on clinical symptoms in patients suffering from schizophrenia

B. Nycz and K. Krysta*

Department of Rehabilitation Psychiatry, Medical University of Silesia, Katowice, Poland

*Corresponding author.

doi: 10.1192/j.eurpsy.2023.329

Introduction: One of the factors influencing the symptoms of schizophrenia, which may indicate changes in the cognitive functioning of patients, is the fluctuating level of inflammatory cytokines.

Objectives: The aim of the review was to analyze the available literature on the importance of selected inflammatory factors [interleukin-1 β (IL-1 β), interleukin-6 (IL-6), interleukin-8 (IL-8), interleukin-10 (IL-10), of tumor necrosis α (TNF- α)] in schizophrenia and the assessment of the impact of changes in cytokine levels on the occurrence of schizophrenia symptoms.

Methods: For this purpose, available scientific publications from following databases: PubMed, Scopus, Google Scholar were used to prove that the levels of selected inflammatory parameters changed in people suffering from schizophrenia. Moreover, fluctuations in cytokine concentrations influenced the occurrence of negative symptoms of schizophrenia, including cognitive disorders, as well as psychotic symptoms.

Results: An increase in the concentration of IL-1 β in the cerebrospinal fluid of patients with the first episode of schizophrenia has been described, which may indicate the involvement of the cytokine in the inflammatory process involving the CNS. The increased level of IL-6 is associated with the occurrence of psychotic disorders, it is also noted in stressful conditions. IL-6 is qualified as an indicator of exacerbation of schizophrenia, which normalizes after antipsychotic treatment. In the blood of patients with paranoid schizophrenia, elevated levels of IL-8 and IL-6 were detected compared to healthy individuals, which indicates the development of an inflammatory process in schizophrenia. The relationship between the level

of IL-8 in women in the second trimester of pregnancy and the risk of developing schizophrenia spectrum disorder in children has been proven. Untreated patients with acute psychotic symptoms showed an increase in the level of TNF- α in the blood serum (compared to healthy subjects). An increase in the level of TNF- α in the blood serum of patients with an acute relapse of schizophrenia or the first episode of psychosis was also demonstrated. In conclusion, the relationship of IL-6 and TNF- α with the occurrence of psychotic disorders, the relationship of IL-1 β with the appearance of changes in mood, behavior, including cognitive dysfunction, the relationship of IL-8 with the risk of developing schizophrenia spectrum disorder in children, the relationship of reduced concentrations of IL-10 with the intensification of negative symptoms, including cognitive deficits.

Conclusions: In conclusion, the analysis showed that patients with schizophrenia fluctuate in the concentration of inflammatory cytokines, which affects the occurrence of clinical symptoms.

Disclosure of Interest: None Declared

O0129

Changes in quality of life in treatment-resistant schizophrenia patients undergoing VR-assisted therapy for auditory verbal hallucinations: A content analysis

M. Beaudoin^{1,2,3*}, S. Potvin^{1,2}, K. Phraxayavong² and A. Dumais^{1,2,4}

¹Psychiatry and addictology, University of Montreal; ²Research Center, Institut universitaire en sante mentale de Montreal; ³Faculty of Medicine, McGill University and ⁴Research Center, Institut national de psychiatrie légale Philippe-Pinel, Montreal, Canada

*Corresponding author.

doi: 10.1192/j.eurpsy.2023.330

Introduction: VR-Assisted Therapy (VRT) for auditory verbal hallucinations has been demonstrated to have a significant impact on the symptoms, beliefs, and quality of life of patients with treatment-resistant schizophrenia. However, little is known about how these changes are implemented into their lives and on which aspects these improvements occur.

Objectives: This study aimed to qualitatively explore changes in the quality of life of patients who underwent VRT in the context of an ongoing clinical trial.

Methods: Ten consecutive patients enrolled in an ongoing clinical trial were assessed using semi-guided interviews before as well as 3 months after VRT. These encounters have been recorded and transcribed. Then, the content of the participants' discourse was thoroughly analyzed, leading to the generation of an extensive theme grid. Each utterance was then coded by at least two members of the research team, and each disagreement was then discussed in a group format until a consensus was reached. As the cases were analyzed, the grid was adapted in a back and forth manner. New participants were included until data saturation occurred.

Results: The content analysis allowed the identification of nine main themes representing different aspects of the patients' quality