

This is a “preproof” accepted article for *Animal Nutriomics*.

This version may be subject to change during the production process.

10.1017/anr.2024.27

Resveratrol activates PGC-1 α pathway via PRAKK1 to regulate mitochondrial biogenesis and alleviate inflammatory responses in bovine mammary epithelial cells

ChunLi Hu [†], YanHao An [†], XueHu Ma, Xue Feng, Yun Ma, Yanfen Ma *

College of Animal Science and Technology, Key Laboratory of Ruminant Molecular and Cellular Breeding of Ningxia Hui Autonomous Region, Ningxia University, Yinchuan 750021, China

[†] These authors contributed equally to this work. * Corresponding Author: Yanfen Ma mayf@nxu.edu.cn.

This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.

Abstract

Mastitis in dairy cows is an important factor restricting the healthy development of dairy industry. Natural extracts have become a research hotspot to alleviate and prevent diseases because of their unique properties. The purpose of this study was to investigate the effects of resveratrol (RES) on the mitochondrial biosynthesis, anti-oxidation and anti-inflammatory in bovine mammary epithelial cells (BMECs) and its mechanism involved. Blood samples were collected from six healthy cows and six mastitis affected cows, respectively, and lipopolysaccharide (LPS) was used to treat bovine mammary epithelial cells (BMECs) to construct inflammation models, gene interference is achieved by transfection. The results showed that mRNA expression of peroxisome proliferator-activated receptor gamma coactivator-1 α (PGC-1 α) was down-regulated and mitochondrial biogenesis-related gene expression was disrupted in the blood of mastitis cows and LPS-induced BMECs. RES is the best active substance to activate PGC-1 α . The addition of RES can effectively alleviate the production of BMECs reactive oxygen species (ROS) and mitochondrial damage induced by LPS, and improve the anti-oxidation and anti-inflammatory ability, while the alleviation effect of RES is inhibited after interfering with protein kinase AMP-activated catalytic subunit α 1 (PRKAA1). In summary, our study emphasizes that PRKAA1 is a key gene mediating the activation of PGC-1 α by RES, which regulates mitochondrial biosynthesis, inhibits ROS release, attenuates mitochondrial damage, and improves mitochondrial antioxidant capacity through the activation of PGC-1 α by PRKAA1, thus attenuating the inflammatory response in BMECs.

Keywords: PRKAA1; PGC-1 α ; resveratrol; inflammation; mitochondrial biogenesis

1. Introduction

Clinical mastitis can lead to metabolic changes, decrease productivity and increase the elimination rate in lactating cows [1]. It is the most common and expensive disease in dairy cows and usually occurs after calving [2]. Mitochondria, as the place where ATP is produced, meets the energy de-

mand of cells through oxidative phosphorylation system. Moreover, mitochondria are the energy centers of cells, which are very important to the life of eukaryotes [3]. Mitochondria are the main places where reactive oxygen species (ROS) are produced, which are essential to fight infection; however, excessive and uncontrolled production of ROS can become deleterious to the cell, leading to mitochondrial and tissue damage [4]. Therefore, mitochondria are also central to the proinflammatory response and play a vital role in dealing with pathogenic infection [5]. Mitochondria are highly dynamic organelles that undergo a coordinated cycle of fission and fusion, known as “mitochondrial dynamics.” Imbalanced mitochondrial dynamics are associated with a range of diseases that are broadly characterized by impaired mitochondrial function and increased cell death [6]. Research has found that the Xuanfei Baidu Formula (XBF) exerts anti-inflammatory effects by restoring mitochondrial kinetics and reducing the activation of inflammasomes [7].

Peroxisome proliferator-activated receptor γ co-activation factor -1α (PGC- 1α) is a co-transcriptional regulator that is a major factor in the regulation of mitochondrial biogenesis and function [8]. PGC- 1α is an important node connecting metabolic regulation, redox control and the inflammatory pathway; its maladjustment changes redox homeostasis in cells and aggravates inflammatory reaction, usually accompanied by metabolic disorder [9]. In the process of inflammation low-level PGC- 1α downregulates the expression of mitochondrial antioxidant genes, induces oxidative stress and promotes the activation of nuclear factor κ B (NF- κ B), PGC- 1α regulates the expression of mitochondrial antioxidant genes, including catalase, peroxidase 3 and 5, uncoupling protein 2 (UCP2), and thioredoxin reductase, to prevent oxidative damage and mitochondrial dysfunction [10]. The AMP-activated protein kinase (AMPK)/PGC- 1α pathway plays an important role in regulating mitochondrial biogenesis [11]. The α 1 catalytic subunit of AMP-activated protein kinase (PRKAA1) the catalase that plays a key role in regulating cell energy metabolism via phosphorylation, and PRKAA1 has been found to be related to inflammatory [12; 13]. Zhu et al. [14] found that

PRKAA1 mediated the clearance of damaged mitochondria in cells, which is necessary for cell maturation and homeostasis.

Resveratrol (RES) is a polyphenol found in grapes, mulberry trees, peanuts, rhubarb and other plants, which plays a beneficial role in preventing chronic diseases related to inflammation, and participates in regulating cell processes, such as gluconeogenesis, lipid metabolism, mitochondrial biogenesis, angiogenesis and apoptosis [15; 16]. Research has indicated that RES could inhibit microcirculation disorder by activating the sirtuin1(SIRT1)-forkhead box O1 (FoxO1) axis, thus relieving the acute pancreatitis in rats [17]. Kong et al. [18] found that RES improved mouse steatohepatitis by regulating the mmu-miR-599/pregnane X receptor (PXR) pathway to inhibit the related inflammatory factors. RES can reduce mitochondrial damage caused by some stress sources, trigger mitochondrial biogenesis and improve the mitochondrial-related bioenergy state in mammalian cells [19]. Research has shown that RES inhibits apoptosis and oxidative stress of bovine mammary epithelial cells (BMECs) induced by aflatoxin B1 and participates in the Nrf2 signaling pathway [20]. RES also improves granular cell activity through mitochondrial biogenesis [21]. Previous studies have shown that RES can reduce LPS-induced inflammatory response via PGC-1 α , but the specific molecular regulatory mechanism remains unclear [22; 23]. This study aims to investigate how RES activates PGC-1 α to regulate mitochondrial biosynthesis and alleviate the inflammatory responses in BMECs and to provide basic data for subsequent research on RES relieving dairy cows with mastitis.

2. Materials and Methods

2.1 Animal ethics

The Animal Experiment Committee of Ningxia University approved the experimental procedures in line with China's Regulations for the Administration of Affairs Concerning Experimental Animals (Ningxia University Ethics No. 22-72). These procedures were rigorously followed accord-

ing to the approved guidelines. Ningxia Xin'ao Agricultural Farm authorized the use of these animals in the study.

2.2 Collection of cows and blood samples

The experimental cows came from an intensive dairy farm in Yingchuan (Ningxia, China). The cows were divided into healthy cows (n=6) and mastitis cows (n=6) according to the California mastitis test (CMT) method. Then, 10 mL of tail root venous blood was collected from each cow. The serum was separated after centrifugation (3,000 g, 15 min) and stored at -80°C for subsequent ELISA detection. Information on healthy and mastitis cows is presented in **Supplementary Table 1**.

2.3 Construction of the inflammatory BMECs

The BMECs cell lines used in this study were obtained from laboratory researchers frozen in liquid nitrogen [24]. BMECs were spread into a 6-well plate. They were washed twice with PBS (Hyclone, USA) and replaced with a new medium when the cell density reached 60~70% confluency. Then, 105 µL of LPS was added to each well to achieve a final concentration of 50 ng/µL [25], thereby inducing inflammation. Cells extracts were collected in centrifuge tubes following disruption with TRIzol at the 0, 3, 6, 12 and 24 h after induction, and stored at -80°C for RNA extraction.

2.4 BMECs culture and treatment

The BMECs were spread into a 6-well plate, and the cells were treated when the cell density reached 70–80%. The transfection reagent was Lipofectamine 3000 (Invitrogen, Thermo Fisher Scientific, USA). Interfering fragments refer to small molecules interfere with RNA fragments (siRNA), siRNA and homologous target RNA complement each other, and specific enzymes degrade target RNA, thus inhibiting and down-regulating gene expression. Three interfering fragments and one control fragment (Negative control) of PRKAA1 gene were designed and synthesized, the interfering fragment sequences are shown in **Supplementary Table 2**. The interference fragment was synthesized by Shenggong Bioengineering Co., Ltd. (Shanghai, China). The cell treatment in each experimental group was as follows: LPS group: 105 µL (final concentration was 50 ng/µL) LPS (Sigma,

USA) was added for the group; si-PRKAA1 group: the PRKAA1 interference fragment was transfected according to the instruction of Lipofectamine 3000, and the cells were collected after transfection for 48 h; RES+si-PRKAA1 group: BMECs were treated using 15 $\mu\text{mol/L}$ RES (G-clone, China) for 12 h after the PRKAA1, and the interference fragment was transferred into the BMECs for 36 h; LPS+si-PRKAA1 group: BMECs were treated using 50 $\text{ng}/\mu\text{L}$ LPS for 12 h after the PRKAA1, interference fragment was transferred into the BMECs for 36 h; LPS+RES+si-PRKAA1 group: BMECs were treated with 15 $\mu\text{mol/L}$ RES for 12 h and then treated with 50 $\text{ng}/\mu\text{L}$ LPS for 12 h after the PRKAA1 interference fragment was transferred into the BMECs for 24 h.

2.5 Real-time fluorescence quantitative PCR (qPCR)

RNA was extracted from BMECs, and blood was extracted using the Trizol (Takara, Japan) method. The integrity of the RNA was detected using gel electrophoresis, and the quality and concentration of RNA were detected using a multifunctional full-wavelength enzyme-labeled instrument (SYNERGY|LX, Bio-Rad, USA) [26]. The cDNA was obtained by the reverse transcription of RNA with a reverse transcription reagent (Takara, Japan). The qPCR (CFX96, Bio-Rad, USA) system was as follows: consisted of 7.5 μL of 2 \times SYBR green qPCR mix, 1.5 μL of cDNA, 0.3 μL of each primer (upstream and downstream), and 5.4 μL of deionized water. Glyceraldehyde-3-phosphate dehydrogenase (GAPDH) was selected as the internal reference gene. Primer information is shown in **Supplementary Table 3**; primers were synthesized by General Biology Co., Ltd. (Anhui, China).

2.6 Nucleoplasm separation and ELISA

The nuclear-cytoplasmic separation kit (Invitrogen, Thermo Fisher scientific, USA) was used for nuclear-cytoplasmic separation of BMECs. The mass of nuclear-mass separation was detected via 1% gel electrophoresis, assessed the quality of RNA separation between the nucleus and cytoplasm, and reverse transcription was performed to produce cDNA. Semi-quantitative detection was used to

analyze the expression levels of PGC-1 α , PRKAA1, Tfam, Drp1, Fis, Mfn1, and Mfn2 in both compartments, followed by gel electrophoresis to measure each gene's expression in the nucleus and cytoplasm. Gray analysis was performed with Image J (version v10.6.2) [25] online software. The BMECs medium supernatant (or blood) from each treatment group was collected and tested for the secretion of cytokines interleukin 6 (IL-6) and interleukin 8 (IL-8) according to the ELISA kit (Ruixin, Quanzhou, China) instructions. Detection of MDA content according to MDA kit (Ruixin, Quanzhou, China) instructions.

2.7 Bioinformatics analysis

The protein sequence of PRKAA1 and PGC-1 α was obtained in NCBI, and the potential phosphorylation site of PGC-1 α was predicted using netphos 3.1 online software. Protein phosphorylation analysis software Scansite was used to predict the protein kinase that can phosphorylate PGC-1 α [27]. The hydrophilicity and hydrophobicity of the PRKAA1 protein were analyzed using Protscale [28] online software, the secondary structure of PRKAA1 protein was analyzed using Sopma, and the tertiary spatial configuration of PRKAA1 protein was obtained using SWISSMODEL [29]. See **Supplementary table 4** for the web addresses of the online sites

2.8 Flow cytometry

For flow cytometry, cells were processed according to the apoptosis kit instructions. In brief, adherent cells were washed with PBS and treated with trypsin to digest them for 3 min. Then, cell culture medium was added to terminate the digestion, followed by the centrifugation of the medium at 1 000 \times g for 5 min and discarding of the supernatant. Next, 1 mL LPS was added to resuspend the cells, and the suspension was again centrifuged at 1000 \times g for 5 min. The supernatant was discarded and 195 μ L of AnnexinV-FITC binding solution and 5 μ L AnnexinV-FITC were added and mixed gently. Finally, 10 μ L propidium iodide staining solution was added and mixed, followed by incuba-

tion in the dark for 10–20 min at room temperature. The stained cells were subjected to flow cytometry. Cells were processed according to the apoptosis kit instructions (Beyotime, Shanghai, China) [30].

2.9 Western blot

Protein was extracted according to the instructions of the whole protein extraction kit (KeyGEN BioTECH, Nanjing, China). In total, 200 μ L of the mixtures were added into 6-well plates for protein extraction. Proteins were quantified using the BCA protein detection kit (KeyGEN BioTECH, Nanjing, China). Dilute and adjust the protein sample to the same amount, add the loading buffer (Yamei, Shanghai), and then boil at 100°C for 10 min. 1.00 mm minigel was prepared according to the kit instructions of the enzyme PAGE gel rapid preparation kit (Yamei, Shanghai). The gels were used to separate the proteins via electrophoresis, and the electrophoretic positions of the protein bands were compared with a 10–250-kDa prestained protein marker (Yamei, Shanghai). The separated proteins were transferred to PVDF membranes (Millipore, USA). After transfer, the membranes were incubated with a rapid sealing solution (Yamei, Shanghai) for 15 min, followed by washing with TBST (Servicebio, Wuhan) solution and overnight incubation in TBST solution containing primary antibodies primary antibody information was as follows: GAPDH (Abways, AB0036, 1:1 000), Tfam (Abways, CY7172, 1:1 000), Fis (Abways, CY8730, 1:1 000), PGC-1 α (Abcam, ab54481, 1:1 000), Mfn2 (Abcam, ab56889, 1:1 000), Bax (Abways, CY5059, 1:1 000), Caspase-3 (Beyotime, AC030, 1:1 000). After incubation with primary antibodies, the membrane was washed with TBST thrice for 10 min each time. Subsequently, the membranes were incubated with horseradish peroxidase (HRP)-coupled secondary antibodies (HRP-labeled goat trypsin antibodies, 1 : 20 000 Shenggong, Shanghai) for 2 h. Finally, the membrane washed thrice with TBST for 10 min each time and imaged using a chemiluminescence imaging system (Tanon-5200, Shanghai) after treatment with an enhanced chemiluminescence (Yamei, China) reagent.

2.10 Mitochondrial membrane potential and ROS detection

Mitochondrial damage was assessed using a mitochondrial membrane potential assay kit (Be-yotime, China). Briefly, the cells in the 6-well plate were allowed to grow to 70–80% confluency, and the medium in the 6-well plate was discarded and the cells washed with PBS for 1–2 times. Then, 1 mL of medium and 1 mL of JC-1 staining working solution were added to each well, respectively, and they were fully and uniformly mixed and placed in an incubator at 37°C and 5% CO₂ for 20 min. After the incubation, the mixture was discarded, and 2 mL pre-cooled JC-1 staining buffer was added into each well twice for 3 min each time. Then, 2 mL medium was added to each well to avoid light and photographed using a fluorescence inverted microscope after the staining buffer was discarded. The content of ROS in cells was detected using an ROS kit (Lablead, China). Briefly, the cells in the 6-well plate were allowed to grow to 70–80%, and the medium in the 6-well plate was discarded and washed with PBS 1–2 times. Then, 1 mL diluted DCFH-DA was added into each well to be fully and uniformly mixed and placed in an incubator at 37°C and 5% CO₂ for 20 min. At the end of the incubation, DCFH-DA was discarded, and 2 mL serum-free medium without antibody was added into each well for washing 3 times, for 3 min each time, in the dark. The pictures were taken under a fluorescence inverted microscope.

2.11 Data analysis

For each group, at least three independent experiments were performed. The qPCR results of fluorescence quantification were analyzed using the $2^{-\Delta\Delta C_t}$ method, and the data are expressed as mean \pm standard error (SEM). Each test ensures three independent technical repeats. T-test for significance of experimental data were performed using GraphPad Prism 8 software. $P < 0.05$ (*), and $P < 0.01$ (**) indicated statistical significance. The protein band grayscale was analyzed using the Image J software.

3. Results

3.1 Activation Disorders of PGC-1 α in dairy cows with mastitis

In order to clarify the relationship between PGC-1 α and mastitis, the mRNA expression level of PGC-1 α in blood was detected in healthy and mastitis dairy cows. The protein expression levels of inflammatory factors in the blood of healthy cows and mastitis cows were detected with ELISA (Figure 1A and Figure 1B). qPCR was used to measure the mRNA expression levels of inflammatory factors and mitochondrial biosynthesis-related genes in the blood of both healthy and mastitis-affected cows (Figure 1C). The results showed that compared with healthy cows, the content of interleukin 6 (IL-6; Figure 1A, $P < 0.01$) and interleukin 8 (IL-8; Figure 1B, $P < 0.01$) in the blood of mastitis cows increased significantly, and the mRNA expression level of the PGC-1 α gene was significantly decreased (Figure 1C, $P < 0.01$); Compared with healthy cows, the mRNA expression levels of transcription factor A (Tfam), mitofusin 1 (Mfn1) and mitofusin 2 (Mfn2) in the blood of mastitis cows were downregulated, while the mRNA expression level of dynamin-related protein 1 (Drp1) was up-regulated (Figure 1C, $P < 0.01$). Lipopolysaccharide (LPS) was used to induce BMECs for 0, 3, 6, 12 and 24 h, and the results showed that the mRNA expressions of inflammatory factors (IL-6 and IL-8) in each time period were all higher than 0 h, indicating that the inflammatory model was successful constructed (Figure 1D-E, $P < 0.05$). The mRNA expression of PGC-1 α and mitochondrial biogenesis genes (Tfam, Mfn1, Mfn2, Drp1 and fission, Fis) in inflammatory cells was detected. In contrast to the control group, the mRNA expression levels of PGC-1 α , Tfam, Mfn1, and Mfn2 in inflammatory cells decreased, whereas the mRNA expression levels of Drp1 and Fis mRNA increased. (Figure 1F, $P < 0.01$).

3.2 PRKAA1 is a PGC-1 α -phosphorylated kinase

To further screen proteins related to PGC-1 α phosphorylation, the NetPhos 3.1 server was used to predict the existence of 129 potential phosphorylation sites of the protein, and the results showed that the main phosphorylated amino acids were serine (Ser), threonine (Thr) and tyrosine (Tyr; Supplementary Figure 1A). Meanwhile, the protein phosphorylation analysis software ScanSite was also used

to predict the protein kinase that might phosphorylate PGC-1 α (Supplementary Figure 1B), and the results showed that the protein kinase with the highest score was PRKAA1, which targeted 254 threonine (T254) on PGC-1 α (Supplementary Table 5). In addition, we used SOPMA online software to predict the secondary structure of the PRKAA1 protein. It was found that the PRKAA1 protein mainly consisted of α -helix (32.75%, indicated by the blue vertical line), irregular curl (44.08%, shown by the purple short line), a small number of extended chains (red middle vertical line area, 16.38%) and β -corner (green middle vertical line area, 6.79%; Supplementary Figure 1C). The hydrophobicity of the protein was analyzed using ProtScal, which showed that amino acid hydrophobic residues accounted for a large proportion of the PRKAA1 protein (Supplementary Figure 1D). SWISSMOD was used to predict the tertiary structure of PRKAA1, and the results showed that most of the protein had an irregular curl and α -helix structure, which was consistent with the secondary structure prediction results (Supplementary Figure 1E).

3.3 PRKAA1 activates PGC-1 α to regulate mitochondrial biogenesis

The activation of PRKAA1 on PGC-1 α was further determined, and the mRNA expression of PRKAA1 in the blood of mastitis cows was detected using qPCR. The results showed that the mRNA expression of PRKAA1 in mastitis cows was decreased compared to that in healthy cows (Figure 2A, $P < 0.01$). Subsequently, we designed three interfering fragments specific for the PRKAA1 gene and selected si-PRKAA1-1384 as the best interfering fragment (Figure 2B, $P < 0.01$). In contrast to the control group, the mRNA levels of PGC-1 α , Tfam, Mfn1 and Mfn2 was downregulated, while the mRNA expression of mitochondrial fission genes (Drp1 and Fis) was upregulated after PRKAA1 interfered (Figure 2C, $P < 0.05$), which shows that mitochondrial biogenesis is disordered. The nuclear localization of PGC-1 α and mitochondrial biogenesis-related genes (Tfam, Mfn1, Mfn2, Drp1 and Fis) results showed that they were mainly located in the cytoplasm (Figure 2D-H).

3.4 RES activates PGC-1 α via PRKAA1

To find the optimal active substance to activate PGC-1 α , four compounds were tested simultaneously. The BMECs were treated with RES (non-flavonoid polyphenol organic compounds), tea polyphenols (TP, the main component is catechin compounds), lycopene (LYC, a carotenoid found in plant foods) and tartary buckwheat (TB, the main component is flavonoids) extract for 12 h (the final concentrations were all 15 $\mu\text{mol/mL}$), and the cells were collected, respectively. The mRNA expression of PGC-1 α in the four treatment groups was detected using qPCR. The results found that all four substances significantly increased the mRNA expression of PGC-1 α , and the mRNA levels of PGC-1 α was the highest in BMECs treated with RES (Figure 3A, $P < 0.01$). The above results showed that RES was the best activator of PGC-1 α . In order to further prove that RES activates PGC-1 α through PRKAA1 to regulate mitochondrial biosynthesis, BMECs were treated with RES for 12 h after the PRKAA1 interference fragment was transfected into BMECs for 36 h. The results showed that the mRNA expression of PGC-1 α increased significantly in BMECs treated with RES but decreased in BMECs in which PRKAA1 was targeted (Figure 3B, $P < 0.05$) compared with the control group. This indicates the activation efficiency of RES on PGC-1 α without PRKAA1. At the same time, the mRNA expressions of mitochondrial synthesis genes and mitochondrial fission genes were detected. The results showed that the addition of RES promoted the mRNA expressions of the mitochondrial synthesis genes (Tfam, Figure 3C; Mfn1, Figure 3F; Mfn2, Figure 3G; $P < 0.01$) and anti-apoptosis genes (B cell lymphoma protein-2, Bcl-2; Figure 3J, $P < 0.05$), and RES inhibited the mRNA expressions of the mitochondrial fission genes (Drp1, Figure 3D; Fis, Figure 3E; $P < 0.01$) and pro-apoptosis genes (Caspase-3, Figure 3H; Bax, Figure 3I; $P < 0.01$). However, the mRNA expression of the mitochondrial synthesis gene and anti-apoptosis gene was significantly inhibited, and the mRNA expression of the fission and apoptosis gene was promoted by adding RES after interfering with PRKAA1 (Figure 3B-I, $P < 0.01$). A western blot was used to detect the protein expressions of the mitochondrial synthesis gene (Tfam), fission gene (Fis) and apoptosis gene (Caspase3 and Bcl-2 associated X, Bax), and the protein expression results tended to be consistent with those of the qPCR (Figure 3K). The mitochondrial

membrane potential of cells in three experimental groups was detected, and the results showed that the red fluorescence intensity of the mitochondria in RES+si-PRKAA1 group was weaker than that of in the RES group, indicating that the protective effect of RES on mitochondria was weakened after interfering with PRKAA1 (Figure 4A, Supplementary figure 2A). The results of ROS detection showed that the ROS content in the RES+si-PRKAA1 group was higher than that of in the RES group (Figure 4B, Supplementary figure 2B), while the results of flow cytometry showed that there was no significant difference in apoptosis among the three experimental groups (Figure 4C, Supplementary figure 2C). In summary, the PRKAA1 is a key gene that can mediate RES to activate PGC-1 α , and it can also regulate the mitochondrial biogenesis and the expression of genes related to mitochondrial function.

3.5 RES regulate mitochondrial biosynthesis to reduce mitochondrial damage and alleviates apoptosis in inflammatory BMECs by activating the PRKAA1

We further explored the effect of PRKAA1-mediated RES on the mitochondrial biosynthesis in inflammatory BMECs. The results showed that the mRNA expression of PGC-1 α was significantly downregulated in BMECs induced by LPS and upregulated after adding RES, compared with the control group, indicating that RES significantly increased the mRNA expression of PGC-1 α in BMECs induced by LPS. RES did not increase the mRNA levels of PGC-1 α in BMECs induced by LPS after interfering with si-PRKAA1, which shows that the promotion efficiency of RES was dependent on the PRKAA1 (Figure 5A, $P<0.05$) compared with the control group. Similarly, the mRNA levels of the mitochondrial synthesis genes (Tfam, Mfn1 and Mfn2) was inhibited in BMECs induced by LPS. The mRNA expression could be promoted by RES, but the promotion effect of RES was not obvious after interfering with PRKAA1 (Figure 5B-D, $P<0.05$). Meanwhile, the mRNA expression of mitochondrion fission genes (Drp1 and Fis) was increased significantly in the inflammatory state, and the expression was inhibited after adding RES. However, the inhibition effects of RES were weakened after interfering with PRKAA1 (Figure 5E-F, $P<0.05$) and the mRNA expressions of the apoptosis gene and anti-apoptosis gene were also detected, the findings indicated that the mRNA

levels of the apoptotic genes promoted in BMECs induced by LPS, and the mRNA expression could be inhibited by RES, but the promotion effect of RES was not obvious after interfering with PRKAA1. The mRNA expression of the anti-apoptosis gene showed the opposite result (Figure 5G-I, $P < 0.01$). A western blot was used to detect the protein expressions of a mitochondrial synthesis gene (Tfam), fission gene (Fis) and apoptosis gene (Caspase-3 and Bax), and the results were consistent with those of qPCR (Figure 6A). The results of mitochondrial membrane potential showed that the red fluorescence intensity in the LPS group was significantly reduced compared with the control group, indicating that mitochondria were seriously damaged, while the red fluorescence in the RES+LPS group was stronger than that of in the LPS group, indicating that RES had the protective effect on mitochondria, and the protective effect of RES on mitochondria was weakened after interfering with PRKAA1 (Figure 6B, Supplementary figure 2D). Apoptosis was detected using flow cytometry, and we found that apoptosis in the RES+LPS group was lower than that in the LPS group, but increased after interfering with PRKAA1 (Figure 6C, Supplementary figure 2E). Therefore, the above results show that PRKAA1 could mediate RES to promote mitochondrial biosynthesis in BMECs induced by LPS, and reduced mitochondrial damage and alleviated the apoptosis of BMECs.

3.6 RES improve antioxidant and anti-inflammatory capacity in inflammatory BMECs by activating the PRKAA1

On the basis of the above research, the anti-inflammatory mechanism of PRKAA1-mediated RES was explored. In contrast to the control group, the mRNA levels of the oxidative gene malondialdehyde (MDA, $P < 0.01$) in BMECs induced by LPS was increased significantly, and inhibited after BMECs treated with RES, but was not significantly inhibited after interfering with PRKAA1 (Figure 7A, $P < 0.05$). However, the mRNA levels of antioxidant genes (superoxide dismutase, SOD, Figure 7B, $P < 0.05$; glutathione peroxidase, GSH-PX, Figure 7C, $P < 0.05$) showed the opposite result with MDA. Simultaneously, mRNA levels of inflammatory factors (IL-6, IL-8, and IL-1 β) was detected, and we found that RES co-treatment with LPS notably reduced their expression. However, the inhib-

itory effect of RES on inflammatory factors decreased after interfering with PRKAA1, in contrast to the control group (Figure 7D-F, $P < 0.01$). Similarly, the detection results of ROS showed that RES could inhibit the release of ROS from the BMECs induced by LPS, and the inhibition effects of RES was weakened after interfering with PRKAA1 (Figure 7G, Supplementary figure 2F).

4. Discussion

The mitochondrion is the energy center of cells and the center of proinflammatory response, playing an important role in the response against pathogenic infection [4]. Mitochondrial dynamics play a central role in the process of proinflammatory signal transduction [31]. Research has shown that mitochondrial dysfunction promotes inflammatory responses. Moreover, retrograde signaling caused by dysfunctional mitochondria can alter gene expression, cell morphology and function, and mitochondrial kinetics may also play an important role in stress signaling. Normal mitochondria are highly dynamic organelles whose size, shape, and network are controlled by cell physiology [32]. When mitochondrial integrity is compromised, mitochondrial damage-associated molecular patterns engage pattern recognition receptors, trigger inflammation, and promote pathology in an expanding list of diseases [33].

Mitochondrial biogenesis refers to the process during which existing mitochondria produce new mitochondria. This biogenesis process is regulated by PGC-1 α , which is activated through phosphorylation or deacetylation, to further activate nuclear respiratory factors 1 (Nrf1) and Nrf2, and then activate mitochondrial Tfam [34]. Similarly, the expressions of PGC-1 α and Tfam were downregulated in vivo and in vitro in our experiment, which suggested that PGC-1 α had an activation disorder and mitochondrial biogenesis was disordered in mastitis cows. Tfam is a protein that participates in the transcription maintenance and replication of mitochondrial DNA (mtDNA); therefore, the regulation of Tfam levels and PGC-1 α activation can control the expression of mtDNA-encoded proteins, which, in turn, regulate mitochondrial biogenesis [35]. Bi et al. [36] indicated that the addition of irisin to

mouse hepatocytes promoted the expression of PGC-1 α and Tfam, inhibited mitochondrial excessive division, promoted mitochondrial biogenesis and led to an increase in mitochondrial content to alleviate liver injury. Resistin destroys mitochondrial biogenesis by inhibiting the PGC-1 α / Nrf1/Tfam signaling pathway. In human neuronal SH-SY5Y, Resistin destroys mitochondrial biogenesis by inhibiting the PGC-1 α / Nrf1/Tfam signaling pathway [37]. The above results have shown that Tfam expression is similar to the parameters of mitochondrial biogenesis, which leads to the wide acceptance of Tfam as a marker of mitochondrial biogenesis. mtROS reduced the abundance of mitochondrial Tfam in HK2 cells by suppressing its transcription and promoting Lon-mediated Tfam degradation, thus promoting mitochondrial dysfunction and inflammation [38]. The limitation of this experiment is that the ROS measured is the total ROS in the cells, whereas studies have shown that the ROS in the cells mainly comes from the mitochondrial respiratory chain, NADPH oxidase, xanthine oxidase and inflammation [39; 40]. Similarly, mitochondrial damage is characterized by decreased metabolic activity, increased ROS production, membrane permeability changes and mitochondrial protein release into the cytoplasm, which increases intracellular oxidative stress. At a higher level of ROS, a longer mPTP opening may release ROS explosion, leading to the destruction of mitochondria [41; 42]. Mitosis is necessary for the generation of new mitochondria. Mitochondrial fission is crucial for controlling mitochondrial function, and Drp1 is the key mediator for mitochondrial division [43]. As the receptor of Drp1, Fis recruited Drp1 from the cytoplasm and formed a spiral around the mitochondria, then cut off the mitochondrial inner and outer membrane [44]. Similarly, our nuclear and cytoplasmic localization of Drp1 and Fis also showed that they were mainly located in the cytoplasm. Yu et al. [45] found that LPS promotes the expression of signal transducer and transcriptional activator 2 (Stat2) and Drp1 in mouse macrophages, leading to mitochondrial division, while knocking out or inhibiting Drp1 reduces the increase in mitochondrial mass and pro-inflammatory differentiation. It was suggested that the upregulation of Drp1 promoted mitochondrial fission. Mfn1

and Mfn2, homologues of the fuzzy onion in yeast and drosophila, are key regulators of mitochondrial fusion in mammalian cells [46]. Mfn2 is a mitochondrial outer membrane protein that regulates mitochondrial fusion, while Mfn1 and Mfn2 act in three independent molecular complexes to promote mitochondrial fusion [47; 48]. Our research revealed that mRNA expression levels of Drp1 and Fis were elevated, whereas Mfn1 and Mfn2 expression levels were reduced in the blood of dairy cows suffering from mastitis, indicating that mitochondrial dynamics were in an imbalanced state and mitochondrial biogenesis was dysfunctional under the inflammatory state. Based on our findings, PRKAA1 is a key gene for RES to activate PGC-1 α , and PRKAA1 is involved in inflammation-related pathways, such as AMPK/PGC-1/SIRT1, mTOR, NF- κ B, which, in turn, provides a new idea for us to further explore the anti-inflammatory mechanism of PRKAA1-mediated RES.

RES is a non-flavonoid polyphenol organic substance with antioxidant, anti-inflammatory and anti-cancer properties [49], which can inhibit the mRNA expression of the toll-like receptor (TLR) and pro-inflammatory genes, and its antioxidant activity and ability to inhibit the production of enzymes related to eicosanoid contribute to its anti-inflammatory properties [50]. In addition, treatment with RES could inhibit the expression of tumor necrosis factor α (TNF α), IL-6 and IL-1 β . RES can be used as an activator of sirtuin-1 (SIRT1), which can reduce the phosphorylation and acetylation of NF- κ B and the signal transducer and activator of transcription 1 (STAT1) by activating the SIRT3 signal, as well as weak Mn-induced oxidative stress and inflammatory cytokines [51]. Our study found that RES can activate PGC-1 α via PRKAA1 to relieve inflammatory reaction in BMECs. PRKAA1, also known as AMP-activated protein kinase α 1 (AMPK α 1), is a catalytic subunit of AMPK and is crucial for regulating cellular energy metabolism through phosphorylation [52]. It has been reported that the combined treatment of quercetin and RES reduced rat obesity and related inflammation induced by a high-fat diet through the AMPK α 1/SIRT1 signaling pathway, in addition, rats showed anti-inflammatory properties and anti-insulin resistance when administered alone or in

combination [53]. RES acts as an activator of sirtuin-1 (SIRT1), which, through activation of the SIRT3 pathway, reduces the phosphorylation and acetylation of NF- κ B and STAT1. This process also diminishes Mn-induced oxidative stress and the levels of inflammatory cytokines [18]. RES activated the AMPK/PGC-1 α axis to promote obesity-damaged mitochondrial biogenesis and muscle regeneration, whereas the effect of RES was eliminated by AMPK α 1, which reversed the inhibition of PRKAA1 phosphorylation and SIRT1 expression, further confirming our research results [53]. The pro-mitochondrial fusion and anti-inflammatory functions of RES were significantly decreased after PRKAA1 intervention, suggesting that RES may regulate mitochondrial biogenesis through PRKAA1 activation of PGC-1 α , which may play an anti-inflammatory role. In our study, we found that RES application alone significantly increased the gene expression of PGC-1 α , Mfn1 and Mfn2, but the effect was slight when combined with LPS, for this result we hypothesized that LPS stimulation activates a strong inflammatory cascade, including the NF- κ B pathway, which can interfere with some of the beneficial effects of RES, especially in terms of mitochondrial gene expression. While RES is able to counteract the inflammation induced by LPS to some extent, it may not completely override the inflammatory suppression of mitochondrial biogenesis in this context. Mitochondrial damage is characterized by decreased metabolic activity, increased ROS production, membrane permeability changes and mitochondrial protein release into the cytoplasm, which increases intracellular oxidative stress. At a higher level of ROS, a longer mPTP opening may release ROS explosion, leading to the destruction of mitochondria [54; 55]. The massive release of ROS activates inflammatory corpuscles, which, in turn, promotes inflammation [56]. RES can also eliminate excessive ROS, increase the activity of SOD, enhance the potential and ATP levels of mitochondrial membrane, reduce the copy number of mitochondrial DNA and reduce the level of MDA. The increase of free radicals leads to the excessive production of MDA in cells, and the MDA level is usually regarded as a sign of oxidative stress and antioxidant status [57]. Meanwhile, the research results showed that

RES could promote mitochondrial biogenesis and function by activating the PGC-1 α signaling pathway [58], which further confirms our experimental results. In this study, ROS content in BMECS was significantly increased and mitochondrial membrane potential was decreased in the inflammatory state. After addition of RES, ROS content was significantly reduced and mitochondrial damage was alleviated, but after PRKAA1 intervention, ROS content and mitochondrial membrane potential were not significantly different from those in the inflammatory group, indicating that PRKAA1 is an important target mediating RES anti-inflammation.

5. Conclusion

Taken together, PRKAA1 is the key gene that mediates the activation of PGC-1 α by RES, which regulates the mitochondrial biogenesis of mammary epithelial cells in dairy cows, inhibits ROS production, relieves mitochondrial damage, improves the antioxidant capacity of BMECs, and then alleviates the inflammatory reaction of BMECs.

Declaration of interest statement

The authors declare they have no competing financial interest and no conflicts of interest.

Ethics approval and consent to participate : The experimental procedures were approved by the Animal Experiment Committee of Ningxia University according to the Regulations for the Administration of Affairs Concerning Experimental Animals in China (Ningxia University Ethics No. 22-72). The experimental procedures were strictly implemented according to the approved guidelines and regulations.

Conflicts of interest: The authors declare they have no competing financial interest and no conflict of interest.

Acknowledgements: National Natural Science Foundation of China (No. 32060765, Beijing, China), Ningxia Natural Science Foundation of China (No. 2022AAC02006, Yinchuan, China), Ningxia Hui Autonomous Region Top-notch Youth Talents Project (No. 2022), Modern Agro-industry Technology

Research System (CARS-36), Ningxia Hui Autonomous Region Key R&D Projects (No. 2021BEF01001) and Ningxia University Graduate Student Innovation Program (CXXM202348).9

Author contributions: Data analysis, writing—original draft, Chunli Hu and Yanhao An.; data curation, software, Xuehu Ma.; review, editing, Xue Feng.; funding acquisition, project administration, review and editing, Yun Ma and Yanfen Ma. All authors read and approved the final version of the manuscript.

References:

- [1] M.A. Ballou, Growth and Development Symposium: Inflammation: Role in the etiology and pathophysiology of clinical mastitis in dairy cows. *J Anim Sci* 90 (2012) 1466-78.
- [2] F. Zandkarimi, J. Vanegas, X. Fern, C.S. Maier, and G. Bobe, Metabotypes with elevated protein and lipid catabolism and inflammation precede clinical mastitis in prepartal transition dairy cows. *J Dairy Sci* 101 (2018) 5531-5548.
- [3] S.J. Annesley, and P.R. Fisher, Mitochondria in Health and Disease. *Cells* 8 (2019).
- [4] P. Andrieux, C. Chevillard, E. Cunha-Neto, and J. Nunes, Mitochondria as a Cellular Hub in Infection and Inflammation. *Int J Mol Sci* 22 (2021).
- [5] S. Missiroli, S. Paternani, N. Carocchia, G. Pedriali, M. Perrone, M. Previati, M.R. Wieckowski, and C. Giorgi, Mitochondria-associated membranes (MAMs) and inflammation. *Cell Death Dis* 9 (2018) 329.
- [6] B.N. Whitley, E.A. Engelhart, and S. Hoppins, Mitochondrial dynamics and their potential as a therapeutic target. *Mitochondrion* 49 (2019) 269-283.
- [7] Z. Li, H. Pan, J. Yang, D. Chen, Y. Wang, H. Zhang, and Y. Cheng, Xuanfei Baidu formula alleviates impaired mitochondrial dynamics and activated NLRP3 inflammasome by repressing NF-kappaB and MAPK pathways

in LPS-induced ALI and inflammation models. *Phytomedicine* 108 (2023) 154545.

- [8] S. Rius-Pérez, I. Torres-Cuevas, I. Millán, Á.L. Ortega, and S. Pérez, PGC-1 α , Inflammation, and Oxidative Stress: An Integrative View in Metabolism, 2020, pp. 1-20.
- [9] P.S. Eisele, and C. Handschin, Functional crosstalk of PGC-1 coactivators and inflammation in skeletal muscle pathophysiology. *Semin Immunopathol* 36 (2014) 27-53.
- [10] N.R. Das, B. Vaidya, P. Khare, M. Bishnoi, and S.S. Sharma, Combination of Peroxisome Proliferator-activated Receptor Gamma (PPAR γ) Agonist and PPAR Gamma Co-Activator 1alpha (PGC-1alpha) Activator Ameliorates Cognitive Deficits, Oxidative Stress, and Inflammation in Rodent Model of Parkinson's Disease. *Curr Neurovasc Res* 18 (2021) 497-507.
- [11] L. Li, L. Xiao, Y. Hou, Q. He, J. Zhu, Y. Li, J. Wu, J. Zhao, S. Yu, and Y. Zhao, Sestrin2 Silencing Exacerbates Cerebral Ischemia/Reperfusion Injury by Decreasing Mitochondrial Biogenesis through the AMPK/PGC-1alpha Pathway in Rats. *Sci Rep* 6 (2016) 30272.
- [12] S. Krishan, D.R. Richardson, and S. Sahni, Gene of the month. AMP kinase (PRKAA1). *J Clin Pathol* 67 (2014) 758-63.
- [13] Q. Yang, Q. Ma, J. Xu, Z. Liu, X. Mao, Y. Zhou, Y. Cai, Q. Da, M. Hong, N.L. Weintraub, D.J. Fulton, D.C.E. Belin, and Y. Huo, Endothelial AMPK α 1/PRKAA1 exacerbates inflammation in HFD-fed mice. *Br J Pharmacol* 179 (2022) 1661-1678.
- [14] H. Zhu, M. Foretz, Z. Xie, M. Zhang, Z. Zhu, J. Xing, J. Leclerc, M. Gaudry, B. Viollet, and M.H. Zou, PRKAA1/AMPK α 1 is required for autophagy-dependent mitochondrial clearance during erythrocyte maturation. *Autophagy* 10 (2014) 1522-34.
- [15] L. Malaguarnera, Influence of Resveratrol on the Immune Response. *Nutrients* 11 (2019).

- [16] T. Meng, D. Xiao, A. Muhammed, J. Deng, L. Chen, and J. He, Anti-Inflammatory Action and Mechanisms of Resveratrol. *Molecules* 26 (2021).
- [17] Y. Rong, J. Ren, W. Song, R. Xiang, Y. Ge, W. Lu, and T. Fu, Resveratrol Suppresses Severe Acute Pancreatitis-Induced Microcirculation Disturbance through Targeting SIRT1-FOXO1 Axis. *Oxid Med Cell Longev* 2021 (2021) 8891544.
- [18] L. Cong, M.Y. Lei, Z.Q. Liu, Z.F. Liu, Z. Ma, K. Liu, J. Li, Y. Deng, W. Liu, and B. Xu, Resveratrol attenuates manganese-induced oxidative stress and neuroinflammation through SIRT1 signaling in mice. *Food Chem Toxicol* 153 (2021) 112283.
- [19] F.R. Jardim, F.T. de Rossi, M.X. Nascimento, S.B.R. Da, P.A. Borges, I.C. Prescilio, and M.R. de Oliveira, Resveratrol and Brain Mitochondria: a Review. *Mol Neurobiol* 55 (2018) 2085-2101.
- [20] Y. Zhou, Y. Jin, H. Yu, A. Shan, J. Shen, C. Zhou, Y. Zhao, H. Fang, X. Wang, J. Wang, Y. Fu, R. Wang, R. Li, and J. Zhang, Resveratrol inhibits aflatoxin B1-induced oxidative stress and apoptosis in bovine mammary epithelial cells and is involved the Nrf2 signaling pathway. *Toxicon* 164 (2019) 10-15.
- [21] K.J. Grive, and M.T. Sauerbrun-Cutler, Resveratrol improves granulosa cell activity through mitochondrial biogenesis. *Fertil Steril* 115 (2021) 909-910.
- [22] Q. Huang, D.H. Liu, C.F. Chen, Y. Han, Z.Q. Huang, J.W. Zhang, and X.M. Zeng, Pgc-1alpha Promotes Phosphorylation, Inflammation, and Apoptosis in H9c2 Cells During the Early Stage of Lipopolysaccharide Induction. *Inflammation* 44 (2021) 1771-1781.
- [23] W. Xiao-Li, L. Ting, L. Ji-Hong, M. Shu-Ying, and X. Xian-Zhong, The Effects of Resveratrol on Inflammation and Oxidative Stress in a Rat Model of Chronic Obstructive Pulmonary Disease. *Molecules* 22 (2017).
- [24] J. Yang, Q. Hu, J. Wang, Q. Ren, X. Wang, Z. Luoreng, D. Wei, and Y. Ma, RNA-Seq Reveals the Role of

miR-29c in Regulating Inflammation and Oxidative Stress of Bovine Mammary Epithelial Cells. *Front Vet Sci* 9 (2022) 865415.

[25] Y. Strandberg, C. Gray, T. Vuocolo, L. Donaldson, M. Broadway, and R. Tellam, Lipopolysaccharide and lipoteichoic acid induce different innate immune responses in bovine mammary epithelial cells. *Cytokine* 31 (2005) 72-86.

[26] L. He, X. Feng, C. Hu, S. Liu, H. Sheng, B. Cai, and Y. Ma, HOXA9 gene inhibits proliferation and differentiation and promotes apoptosis of bovine preadipocytes. *BMC Genomics* 25 (2024) 358.

[27] Y.F. Ma, Z.H. Wu, M. Gao, and J.J. Loo, Nuclear factor erythroid 2-related factor 2-antioxidant activation through the action of ataxia telangiectasia-mutated serine/threonine kinase is essential to counteract oxidative stress in bovine mammary epithelial cells. *Journal of Dairy Science* 101 (2018) 5317-5328.

[28] M.R. Wilkins, E. Gasteiger, A. Bairoch, J.C. Sanchez, K.L. Williams, R.D. Appel, and D.F. Hochstrasser, Protein identification and analysis tools in the ExPASy server. *Methods Mol Biol* 112 (1999) 531-52.

[29] A. Waterhouse, M. Bertoni, S. Bienert, G. Studer, G. Tauriello, R. Gumienny, F.T. Heer, T. de Beer, C. Rempfer, L. Bordoli, R. Lepore, and T. Schwede, SWISS-MODEL: homology modelling of protein structures and complexes. *Nucleic Acids Res* 46 (2018) W296-W303.

[30] C. Hu, M. Yang, X. Feng, S. Wang, Y. Ma, and Y. Ma, miR-10167-3p targets TCF7L1 to inhibit bovine adipocyte differentiation and promote bovine adipocyte proliferation. *Genomics* 116 (2024) 110903.

[31] S. Marchi, E. Guilbaud, S. Tait, T. Yamazaki, and L. Galluzzi, Mitochondrial control of inflammation. *Nat Rev Immunol* 23 (2023) 159-173.

[32] S. Srinivasan, M. Guha, A. Kashina, and N.G. Avadhani, Mitochondrial dysfunction and mitochondrial dynamics-The cancer connection. *Biochim Biophys Acta Bioenerg* 1858 (2017) 602-614.

- [33] A.P. West, Mitochondrial dysfunction as a trigger of innate immune responses and inflammation. *Toxicology* 391 (2017) 54-63.
- [34] P.A. Li, X. Hou, and S. Hao, Mitochondrial biogenesis in neurodegeneration. *J Neurosci Res* 95 (2017) 2025-2029.
- [35] A. Picca, and A.M. Lezza, Regulation of mitochondrial biogenesis through TFAM-mitochondrial DNA interactions: Useful insights from aging and calorie restriction studies. *Mitochondrion* 25 (2015) 67-75.
- [36] J. Bi, J. Zhang, Y. Ren, Du Z, Q. Li, Y. Wang, S. Wei, L. Yang, J. Zhang, C. Liu, Y. Lv, and R. Wu, Irisin alleviates liver ischemia-reperfusion injury by inhibiting excessive mitochondrial fission, promoting mitochondrial biogenesis and decreasing oxidative stress. *Redox Biol* 20 (2019) 296-306.
- [37] Z. Chen, S. Tao, X. Li, and Q. Yao, Resistin destroys mitochondrial biogenesis by inhibiting the PGC-1alpha/NRF1/TFAM signaling pathway. *Biochem Biophys Res Commun* 504 (2018) 13-18.
- [38] M. Zhao, Y. Wang, L. Li, S. Liu, C. Wang, Y. Yuan, G. Yang, Y. Chen, J. Cheng, Y. Lu, and J. Liu, Mitochondrial ROS promote mitochondrial dysfunction and inflammation in ischemic acute kidney injury by disrupting TFAM-mediated mtDNA maintenance. *Theranostics* 11 (2021) 1845-1863.
- [39] A. Sarniak, J. Lipinska, K. Tytman, and S. Lipinska, Endogenous mechanisms of reactive oxygen species (ROS) generation. *Postepy Hig Med Dosw (Online)* 70 (2016) 1150-1165.
- [40] Q.M. Chen, Nrf2 for protection against oxidant generation and mitochondrial damage in cardiac injury. *Free Radic Biol Med* 179 (2022) 133-143.
- [41] D.B. Zorov, M. Juhaszova, and S.J. Sollott, Mitochondrial reactive oxygen species (ROS) and ROS-induced ROS release. *Physiol Rev* 94 (2014) 909-50.
- [42] R.E. Koch, C.C. Josefson, and G.E. Hill, Mitochondrial function, ornamentation, and immunocompetence.

Biol Rev Camb Philos Soc 92 (2017) 1459-1474.

- [43] H. Yang, C. Sibilla, R. Liu, J. Yun, B.A. Hay, C. Blackstone, D.C. Chan, R.J. Harvey, and M. Guo, Clueless/CLUH regulates mitochondrial fission by promoting recruitment of Drp1 to mitochondria. *Nat Commun* 13 (2022) 1582.
- [44] K. Schmitt, A. Grimm, R. Dallmann, B. Oettinghaus, L.M. Restelli, M. Witzig, N. Ishihara, K. Mihara, J.A. Ripperger, U. Albrecht, S. Frank, S.A. Brown, and A. Eckert, Circadian Control of DRP1 Activity Regulates Mitochondrial Dynamics and Bioenergetics. *Cell Metab* 27 (2018) 657-666.e5.
- [45] W. Yu, X. Wang, J. Zhao, R. Liu, J. Liu, Z. Wang, J. Peng, H. Wu, X. Zhang, Z. Long, D. Kong, W. Li, and C. Hai, Stat2-Drp1 mediated mitochondrial mass increase is necessary for pro-inflammatory differentiation of macrophages. *Redox Biol* 37 (2020) 101761.
- [46] S. Zhao, N. Heng, H. Wang, H. Wang, H. Zhang, J. Gong, Z. Hu, and H. Zhu, Mitofusins: from mitochondria to fertility. *Cell Mol Life Sci* 79 (2022) 370.
- [47] M. Harland, S. Torres, J. Liu, and X. Wang, Neuronal Mitochondria Modulation of LPS-Induced Neuroinflammation. *J Neurosci* 40 (2020) 1756-1765.
- [48] H. Chen, S.A. Detmer, A.J. Ewald, E.E. Griffin, S.E. Fraser, and D.C. Chan, Mitofusins Mfn1 and Mfn2 coordinately regulate mitochondrial fusion and are essential for embryonic development. *J Cell Biol* 160 (2003) 189-200.
- [49] G. Reposi, U.N. Das, and A.R. Eynard, Molecular Basis of the Beneficial Actions of Resveratrol. *Arch Med Res* 51 (2020) 105-114.
- [50] H. Capiralla, V. Vingtdoux, H. Zhao, R. Sankowski, Y. Al-Abed, P. Davies, and P. Marambaud, Resveratrol mitigates lipopolysaccharide- and Abeta-mediated microglial inflammation by inhibiting the

TLR4/NF-kappaB/STAT signaling cascade. *J Neurochem* 120 (2012) 461-72.

- [51] L. Cong, M.Y. Lei, Z.Q. Liu, Z.F. Liu, Z. Ma, K. Liu, J. Li, Y. Deng, W. Liu, and B. Xu, Resveratrol attenuates manganese-induced oxidative stress and neuroinflammation through SIRT1 signaling in mice. *Food Chem Toxicol* 153 (2021) 112283.
- [52] Y. Zhang, X. Zhou, L. Cheng, X. Wang, Q. Zhang, Y. Zhang, and S. Sun, PRKAA1 Promotes Proliferation and Inhibits Apoptosis of Gastric Cancer Cells Through Activating JNK1 and Akt Pathways. *Oncol Res* 28 (2020) 213-223.
- [53] L. Zhao, F. Cen, F. Tian, M.J. Li, Q. Zhang, H.Y. Shen, X.C. Shen, M.M. Zhou, and Du J, Combination treatment with quercetin and resveratrol attenuates high fat diet-induced obesity and associated inflammation in rats via the AMPKalpha1/SIRT1 signaling pathway. *Exp Ther Med* 14 (2017) 5942-5948.
- [54] D.B. Zorov, M. Juhaszova, and S.J. Sollott, Mitochondrial reactive oxygen species (ROS) and ROS-induced ROS release. *Physiol Rev* 94 (2014) 909-50.
- [55] R.E. Koch, C.C. Josefson, and G.E. Hill, Mitochondrial function, ornamentation, and immunocompetence. *Biol Rev Camb Philos Soc* 92 (2017) 1459-1474.
- [56] L. Diebold, and N.S. Chandel, Mitochondrial ROS regulation of proliferating cells. *Free Radic Biol Med* 100 (2016) 86-93.
- [57] A.N. Torun, S. Kulaksizoglu, M. Kulaksizoglu, B.O. Pamuk, E. Isbilen, and N.B. Tutuncu, Serum total anti-oxidant status and lipid peroxidation marker malondialdehyde levels in overt and subclinical hypothyroidism. *Clin Endocrinol (Oxf)* 70 (2009) 469-74.
- [58] J. Zhou, Z. Yang, R. Shen, W. Zhong, H. Zheng, Z. Chen, J. Tang, and J. Zhu, Resveratrol Improves Mitochondrial Biogenesis Function and Activates PGC-1 α Pathway in a Preclinical Model of Early Brain Injury

Following Subarachnoid Hemorrhage. *Frontiers in Molecular Biosciences* 8 (2021).




Figure 1. Activation disorders of PGC-1α in dairy cows with mastitis. A–B. The content of the IL-6 and IL-8 in the blood of dairy cows with mastitis. C. The mRNA expression of PGC-1α, Tfam, Drp1, Fis, Mfn1 and Mfn2 in the blood of mastitis cows. D–E. The mRNA expression of IL-6 and IL-8 in BMECs induced by LPS for 0, 3, 6, 12 and 24 h. F. The mRNA expression of PGC-1α, Tfam, Drp1, Fis, Mfn1 and Mfn2 in BMECs induced by LPS. * $P < 0.05$, ** $P < 0.01$.




Figure 2. PRKAA1 activates PGC-1 α to regulate mitochondrial biogenesis. A. The mRNA expression of PRKAA1 in blood of healthy cows and mastitis cows. B. Screening of interfering fragments of PRKAA1 gene. C. The mRNA levels of PGC-1 α , Tfam, Drp1, Fis, Mfn1, and Mfn2 in BMECs treated with the PRKAA1 interference fragment were measured by qPCR. D. Detection of RNA isolated from nuclear and cytoplasm of BMECs using gel electrophoresis. E-F. Semi-quantitative detection of PRKAA1, PGC-1 α , Tfam, Drp1, Fis, Mfn1 and Mfn2 localization. G-H. Image J was used to quantify the gel electrophoresis patterns of each gene. * $P < 0.05$, ** $P < 0.01$.




Figure 3. RES activates PGC-1 α via PRKAA1. A. BMECs were treated with RES, TP, LYC and TB for 12 h, and the final concentration was 15 μ mol/L. B–J. The mRNA levels of PGC-1 α , Tfam, Drp1, Fis, Mfn1, Mfn2, Caspase-3, Bax and Bcl-2 in control group, RES group and RES+si-PRKAA1 group. K. Western blot. * $P < 0.05$, ** $P < 0.01$.




Figure 4. RES activates PGC-1 α via PRKAA1 relieve mitochondrial damage in inflammatory BMECs, reduce ROS release, and inhibit cell apoptosis. A. Mitochondrial membrane potential detection with JC-1. B. ROS detection with ROS detection kit. C. Apoptosis was detected using flow cytometry. * $P < 0.05$, ** $P < 0.01$. Scale bar = 200 μm .




Figure 5. PRKAA1 mediates RES to regulate mitochondrial biosynthesis in BMECs induced by LPS, reduce mitochondrial damage and alleviate BMECs apoptosis. BMECs were treated with RES for 12 h and then treated with LPS for 12 h after the PRKAA1 interference fragment was transfected into BMECs for 24 h. A-I. The mRNA expression of PGC-1 α , Tfam, Mfn1, Mfn2, Drp1, Fis, Caspase-3, Bax and Bcl-2 in the control, LPS, RES+LPS and RES+LPS+si-PRKAA1 groups, respectively. * $P < 0.05$, ** $P < 0.01$.




Figure 6. A. Western blot. B. Detection of mitochondrial membrane potential using JC-1. C. Apoptosis was detected using flow cytometry. Scale bar = 200 μ m.




Figure 7. PRKAA1 mediates RES to reduce ROS production, improve antioxidant capacity and then alleviate inflammatory reaction in BMECs. A. MDA content. B-C. The mRNA expression of GSH-PX and SOD. D-F. The mRNA expression of IL-6, IL-8 and IL-1 β . G. ROS production in BMECs detected by ROS detection kit. Scale bar = 200 μ m. * $P < 0.05$, ** $P < 0.01$.