

EPP589

Detecting electrophysiological alterations in psychiatric disorders through event-related microstates: a systematic review

F. F. Marzocchi^{1*}, A. Perrottelli¹, G. Di Lorenzo², C. D'Amelio³, E. Caporusso¹, G. M. Giordano¹, L. Giuliani¹, A. Melillo¹, P. Pezzella¹, P. Bucci¹, A. Mucci¹ and S. Galderisi¹

¹University of Campania "Luigi Vanvitelli", Naples; ²University of Rome "Tor Vergata", Rome and ³University of L'Aquila, L'Aquila, Italy

*Corresponding author.

doi: 10.1192/j.eurpsy.2025.815

Introduction: Measurements of event-related potentials (ERP), recorded through electrophysiology (EEG) during sensory and cognitive processing tasks, have been widely employed to investigate the pathophysiological basis of mental health disorders. In the last two decades, the analysis of EEG microstates (MS) has been increasingly applied to ERP data. This methodology allows the detection of spatio-temporal changes in neuronal activity potential at the scalp level, providing new insights into neurobiological alterations detectable in psychiatric disorders.

Objectives: The current systematic review aims at providing an extensive and detailed description of the studies that characterized alterations in ERP-microstates in psychiatric and neurodevelopmental disorders.

Methods: A systematic review of English articles using PubMed, Scopus and Web of Science databases was undertaken in April 2024. The review included case-control studies that employed ERP-microstates analysis to compare MS features between subjects (age > 10 years) diagnosed with a mental health disorder and healthy controls.

Results: Out of the 756 records screened, 15 studies were included in the final qualitative synthesis. The studies included patients with schizophrenia (n=7), alcohol use disorder (n=2), borderline personality disorder (n=1), panic disorder (n=1), autism (n=1), major depressive disorder (n=1), post-traumatic stress disorder (n=1) and attention-deficit/hyperactivity disorder (n=1).

The studies investigated alterations of MS characteristics through different types of tasks. Cartool and RAGU were the main software used for MS analysis. Only rarely studies used similar tasks, showing comparable microstate maps. Fourteen of 15 studies showed a significant difference ($p < 0.05$) in MS characteristics between patients and healthy controls. Main differences regarded parameters such as duration, area under the curve and the order of occurrence of MS.

Conclusions: The present literature review aims to highlight the effectiveness in using microstate analysis to identify spatio-temporal alterations in brain electrical activity in subjects with psychiatric disorders, showing the possible implications of the use of this technique in clinical practice and its advantages, as compared to ERP peak analysis.

Disclosure of Interest: None Declared

Psychopathology

EPP590

Procrastination as a transdiagnostic construct in psychiatric disorders: a conceptual scope review

M. S. M. Barbosa^{1,2}, E. L. Guidugli², G. S. M. Barbosa², M. E. G. Melati³, N. B. Pagliari³, P. B. E. da Silva⁴, P. H. P. Dutra⁴, V. R. Xavier⁴, E. H. Grevet⁵ and Y. A. Ferrão^{2,6*}

¹Psychiatry, USP-Ribeirão Preto, Ribeirão Preto; ²Psychiatry, UFCSPA, Porto Alegre; ³Medicine, UNISINOS, São Leopoldo; ⁴Medicine, UFCSPA; ⁵Psychiatry, UFRGS and ⁶Psychiatry, HMIPV, Porto Alegre, Brazil

*Corresponding author.

doi: 10.1192/j.eurpsy.2025.816

Introduction: Procrastination is commonly conceptualized as an irrational, conscious, and sometimes intentional tendency to delay the start or to complete tasks or decisions, despite consequences. The behavior may be present in several psychiatric disorders, especially OCD, ADHD and anxiety disorders. However, there is no consensus definition of procrastination, nor established pathophysiology or motivation.

Objectives: To analyze the descriptive psychopathology as well as the presence or absence of associated symptoms, motivations, and consequences of procrastination. In sequence, to conceptualize the construct of "procrastination" based on the literature published and define the motivations and consequences of the behavior for the individuals.

Methods: The scoping review complies with the recommendations of the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) and included articles published in English, Spanish and Portuguese, between 1973-2023 and indexed in PubMed/MEDLINE, LILACS, Scielo, EMBASE, Scopus, Web of Science, and PsycINFO database. Research conducted in animals, poster publications, conferences in scientific conclaves, and letters to the author were excluded.

Results: We found 387 studies with different designs (295 [76,2%] were cross-sectional, 12 [3,1%] systematic reviews and 5 [1,3%] were meta-analysis) and published in English (86,6%), Spanish (9,3%) and Portuguese (3,9%). Only 5 studies (1.3%) used populations with ADHD and 1 study (0.02%) used a population with OCD, 7 studies (1.8%) used populations with other disorders (Depression, Anxiety) and 374 studies (96.4%) involved students (mainly university students), workers, or the general population. Of the population the majority was female (63,98%), single, young and has some occupation. In 323 articles, at least one scale was used to assess procrastination, with a total of 40 different ones. Furthermore, 337 studies define procrastination and 305 use the definition of other authors.

Conclusions: Procrastination is a complex event, so far without a defined concept. More investigation is necessary for a more comprehensive understanding of this phenomenon.

Disclosure of Interest: None Declared