

## STRUCTURES, COMPOSITIONS, AND X-RAY DIFFRACTION IDENTIFICATION OF DIOCTAHEDRAL CHLORITES

S. W. BAILEY AND JUDITH S. LISTER

Department of Geology & Geophysics, University of Wisconsin  
Madison, Wisconsin 53706

**Abstract**—Al-rich di,triocahedral chlorite exists as the species cookeite and sudoite. Di,diocahedral chlorite exists as the species donbassite. Cookeite has essential Li in its structure, sudoite has essential Mg, and donbassite has only small amounts of either element. To date, sudoite has been reported to have only II*b* structural units and donbassite to have only Ia structural units. Cookeite is based primarily on Ia structural units, but II*b* units are present in specimens from two localities. Most Al-rich chlorite species have regular-stacking “*r*” or “*s*” 2-layer stacking sequences, but 1-layer Ia-2 and Ia-6 polytypes also are known. The structural units (Ia or II*b*) and the specific stacking sequences can be explained by a combination of local charge balance and minimization of cation-cation repulsion involving the interlayer and tetrahedral cations. X-ray powder diffraction data are adequate to differentiate Al-rich chlorite from triocahedral chlorite and to identify the type of structural unit present, but single crystal study is necessary to identify the 2-layer and 1-layer sequences with certainty.

**Key Words**—Chemical composition, Chlorite, Cookeite, Crystal structure, Donbassite, Sudoite, X-ray powder diffraction.

### INTRODUCTION

Al-rich chlorite is being reported in the literature with increasing frequency. Three well-crystallized species have been defined to date—cookeite, sudoite, and donbassite. These species can be differentiated from triocahedral chlorites by their compositions, smaller  $d(060)$  values (1.49 to 1.51 Å compared with 1.52 to 1.56 Å for triocahedral species), slightly smaller  $d(001)$  values, and a distinctly more intense 003 reflection near 4.70 Å. The three species are more difficult to distinguish from one another by X-ray diffraction data alone. Cookeite and sudoite are di,triocahedral chlorites, i.e., the 2:1 layer is dioctahedral, but the interlayer is triocahedral. Donbassite is di,diocahedral with two dioctahedral sheets. No examples are known as yet having a triocahedral 2:1 layer and a dioctahedral interlayer. The present paper summarizes the compositions and several polytypic forms known to occur for these Al-rich species and presents indexed X-ray powder diffraction data to facilitate their identification.

### COOKEITE

Cookeite is a di,triocahedral chlorite commonly found as an alteration product of Li-bearing minerals in pegmatites and hydrothermal veins. The ideal composition is  $\text{Al}_2(\text{Si}_3\text{Al})\text{O}_{10}(\text{OH})_2 \cdot (\text{Al}_2\text{Li})(\text{OH})_6$ . Its essential Li content (~3–4 wt. %  $\text{Li}_2\text{O}$ ) distinguishes it from sudoite and donbassite, which contain only minor amounts of Li. Černý (1970), in a study of the most reliable analyses of cookeite, found tetrahedral Si to be nearly constant at 3.0 Si per 4.0 positions, but with tetrahedral Al of some specimens substituted by small amounts of B or Be. Li ranges between 0.8 and 1.4 atoms per formula unit and is concentrated mainly in

the interlayer sheet. Fluorine sometimes substitutes for OH in small amounts. Small amounts of Ca + Na + K are often reported in chemical analyses of dioctahedral chlorites as cations residing between the 2:1 layers and the interlayer sheet, but they are more likely to be impurities according to Peacor *et al.* (1988). Formula units of 13 cookeite specimens, modified from Černý (1970), are presented in Table 1. The total octahedral occupancy ranges between 4.8 and 5.3 atoms per 6.0 sites in these formulas, but if the Ca + Na + K totals are excluded as impurities, both the total occupancy and the individual atom values require small revisions.

By far the great majority of cookeite specimens are based on Ia structural units in the terminology of Bailey and Brown (1962). The symbol I indicates that the interlayer octahedral sheet has the same orientation or slant as the octahedral sheet within the 2:1 layer, whereas the symbol II indicates opposite orientations. The symbol *a* indicates that the interlayer sheet is positioned by long hydrogen bonds so that the interlayer cations project normal to the layers onto tetrahedral cations and the centers of 6-fold rings of the 2:1 layers above and below. The symbol *b* indicates lack of such superposition. A following integer (1 to 6) then indicates the precise location of the 6-fold rings of the upper 2:1 layer relative to the symmetry plane of the layer plus interlayer below for regular-stacking polytypes. Several polytypic modifications are known. Vrublevskaja *et al.* (1975) described an unusual occurrence of cookeite in metamorphosed bauxite from Djalair in Middle Asia, where it formed as a result of interaction of Li-rich solutions with pyrophyllite-2*M*. The structure is unique also in that it is a regular-stacking 1-layer triclinic struc-

Table 1. Structural formulae of cookeite specimens.

Sample	Locality	Composition
1	Hebron, Maine	$(\text{Na}_{0.03}\text{K}_{0.015})(\text{Al}_{3.795}\text{Fe}^{3+}_{0.03}\text{Li}_{1.44})_{5.265}(\text{Si}_{3.045}\text{Al}_{0.955})\text{O}_{10}(\text{OH})_{7.94}\text{F}_{0.06}$
2	Ward's Natural Science Establishment, unknown locality	$(\text{Al}_{3.825}\text{Li}_{1.365})_{5.19}(\text{Si}_{3.165}\text{Al}_{0.835})\text{O}_{10}(\text{OH})_8$
3	Buckfield, Maine	$(\text{Na} + \text{K})_{0.075}(\text{Al}_{3.825}\text{Fe}^{3+}_{0.045}\text{Li}_{1.265})_{5.135}(\text{Si}_{3.06}\text{Al}_{0.94})\text{O}_{10}(\text{OH})_{7.985}\text{F}_{0.015}$
4	Northwest U.S.S.R.	$(\text{Na}_{0.12}\text{K}_{0.075})(\text{Al}_{3.63}\text{Fe}^{3+}_{0.115}\text{Fe}^{2+}_{0.14}\text{Mg}_{0.05}\text{Li}_{1.22})_{5.155}(\text{Si}_{2.98}\text{Al}_{1.02})\text{O}_{10}(\text{OH})_8$
5	Dobrá Voda, Czechoslovakia	$(\text{Ca}_{0.055}\text{Na}_{0.01})(\text{Al}_{3.80}\text{Fe}^{3+}_{0.015}\text{Mg}_{0.14}\text{Li}_{1.13})_{5.085}(\text{Si}_{3.04}\text{Al}_{0.96})\text{O}_{10}(\text{OH})_8$
6	Muiane, Mozambique	$(\text{Ca}_{0.01}\text{Na}_{0.01})(\text{Al}_{4.025}\text{Li}_{1.12})_{5.145}(\text{Si}_{2.98}\text{Al}_{0.74}\text{Be}_{0.22}\text{B}_{0.06})\text{O}_{10}(\text{OH})_{7.955}\text{F}_{0.045}$
7	Kalbinski Range, U.S.S.R.	$(\text{Ca}_{0.04}\text{Na}_{0.01}\text{K}_{0.01})(\text{Al}_{3.89}\text{Fe}^{3+}_{0.05}\text{Mg}_{0.025}\text{Li}_{1.11})_{5.075}(\text{Si}_{2.95}\text{Al}_{1.05})\text{O}_{10}(\text{OH})_8$
8	Lipovka, Urals, U.S.S.R.	$(\text{Ca}_{0.025}\text{Na}_{0.19}\text{K}_{0.015})(\text{Al}_{3.75}\text{Fe}^{3+}_{0.005}\text{Fe}^{2+}_{0.015}\text{Mg}_{0.175}\text{Li}_{0.97})_{4.915}(\text{Si}_{3.135}\text{Al}_{0.865})\text{O}_{10}(\text{OH})_8$
9	Manono, Katanga	$(\text{Ca}_{0.025}\text{Na}_{0.02})(\text{Al}_{3.94}\text{Fe}^{3+}_{0.005}\text{Mg}_{0.025}\text{Li}_{0.945})_{4.915}(\text{Si}_{3.09}\text{Al}_{0.905}\text{B}_{0.005})\text{O}_{10}(\text{OH})_8$
10	North Little Rock, Arkansas	$(\text{Ca}_{0.01})(\text{Al}_{4.02}\text{Fe}^{3+}_{0.005}\text{Mg}_{0.005}\text{Li}_{0.86})_{4.89}(\text{Si}_{3.035}\text{Al}_{0.965})\text{O}_{10}(\text{OH})_8$
11	Ogofau, Carmarthenshire, England	$(\text{Na}_{0.01})(\text{Al}_{4.00}\text{Fe}^{3+}_{0.02}\text{Fe}^{2+}_{0.065}\text{Li}_{0.855})_{4.94}(\text{Si}_{3.035}\text{Al}_{0.965})\text{O}_{10}(\text{OH})_8$
12	Radkovic, Czechoslovakia	$(\text{Ca}_{0.055})(\text{Al}_{3.90}\text{Fe}^{3+}_{0.02}\text{Fe}^{2+}_{0.01}\text{Mg}_{0.11}\text{Li}_{0.815})_{4.855}(\text{Si}_{3.085}\text{Al}_{0.66}\text{B}_{0.255})\text{O}_{10}(\text{OH})_8$
13	Wait-a-bit Creek, British Columbia	$(\text{Ca}_{0.16}\text{Na}_{0.12}\text{K}_{0.01})(\text{Al}_{3.88}\text{Mg}_{0.11}\text{Li}_{0.77})_{4.76}(\text{Si}_{2.93}\text{Al}_{1.07})\text{O}_{10}(\text{OH})_8$

Modified from Černý (1970). See that author for references to samples.

ture of the *Ia*-6 type. The space group is  $C\bar{1}$ , and the crystallographic angle  $\alpha$  is distorted to  $90^\circ 33'$ . Most cookeite specimens are less well crystallized and are based on monoclinic 2-layer structures. Lister (1966) pointed out that most of the cookeite specimens observed in her survey of the globular rosettes in Li-bearing pegmatites appeared to be imperfect 2-layer structures similar to the monoclinic "s" structure derived by Mathieson and Walker (1954) in a study of vermiculite. The symbol of Lister and Bailey (1967) for this 2-layer structure is  $\bar{X}_1$ -*Ia*-4: $\bar{X}_1$ -*Ia*-6. This symbol indicates that the *a*/3 shift within each 2:1 layer is directed along  $-X_1$  and successive 2:1 layers are positioned so that the centers of the 6-fold rings are located alternately at points 4 and 6 on opposite sides of the symmetry plane below (related to one another by a *b*/3 shift). The 2-layer nature is only apparent by observation of weak diffuse reflections of index  $k \neq 3n$  on single crystal patterns. These reflections do not show up well on Debye-Scherrer patterns (Table 2a).

Lister (1966) also identified a better crystallized 2-layer cookeite from Wait-a-bit Creek, British Columbia, and at several localities in the Magnet Cove district and around North Little Rock, Arkansas. The latter includes cookeite from the Jeffrey quarry, which is part of the Source Clays Repository of The Clay Minerals Society. The structure can be correlated with the monoclinic "r" and "q" structures of Mathieson and Walker (1954), which have polytypic symbols "r" =  $\bar{X}_1$ -*Ia*-4: $\bar{X}_3$ -*Ia*-4 and "q" =  $\bar{X}_1$ -*Ia*-6: $\bar{X}_3$ -*Ia*-2. Here the *a*/3 shift is directed along  $-X_1$  of the first 2:1 layer and along  $-X_3$  of the second layer. Six-fold rings of successive 2:1 layers are positioned at points 4 and 4 for the "r" structure and at points 6 and 2 for the "q" structure. The two structures are enantiomorphous in space group *Cc*. Bailey (1975) summarized results of an incomplete structural refinement of one of the North Little Rock crystals. In this structure Li was found to

be ordered in one interlayer site, where it apparently achieves the best local charge balance. Interlayer  $\text{Li}^+$  lies on a line directly between a tetrahedral cation in the 2:1 layer below and a tetrahedral cation in the 2:1 layer above. The more highly charged interlayer  $\text{Al}^{3+}$  cations, however, only have a tetrahedral cation on one side and the center of a 6-fold ring on the other side. Although different T-O bond lengths were determined during structural refinement, the poor quality of the film data does not permit a firm conclusion as to ordering of tetrahedral Si and Al, nor has it been possible to obtain a better crystal to date. The powder pattern in Table 2b shows clearly  $k \neq 3n$  reflections that identify the 2-layer nature of the specimen, but the more intense  $k = 3n$  lines are very similar to those of the more common but less well crystallized cookeite specimens.

Černý *et al.* (1971) described a cookeite from a pegmatite at Dobrá Voda, Czechoslovakia, as having *IIB* structural units. The present writers have confirmed a second occurrence of cookeite-*IIB* as pink rosettes on elbaite from a pegmatite at Norway, Maine (University of Wisconsin Museum #6001/1). The powder pattern (Table 2c) is distinctly different from those of cookeite specimens based on *Ia* units. Single crystals were not available, but a comparison of observed and calculated intensities of  $k \neq 3n$  reflections suggests 2-layer "r" or "q" structures.

Frank-Kamenetskii (1960) and Černý (1970) mentioned manandonite from Madagascar as a possible di,triocahedral Al-Li chlorite having substantial tetrahedral boron ( $\text{B}^{\text{IV}} \geq \text{Al}^{\text{IV}}$ ). Guggenheim *et al.* (1983) confirmed the high  $\text{B}^{\text{IV}}$  content of this material by ion probe analysis and stated that its X-ray powder pattern was that of a *Ia* type chlorite. The type manandonite defined by Lacroix (1912, 1922), however, has since been shown to be a trioctahedral 1:1 layer silicate of approximate composition  $(\text{Al}_2\text{Li})(\text{Si}_{1.0}\text{Al}_{0.5}\text{B}_{0.5})\text{O}_5(\text{OH})_4$

(A.-M. Fransolet, University Liège, Liège, Belgium, personal communication, 1988; Ranorosa, 1986). This 1:1 type material is intermixed with a B-rich Al-Li "cookeite", which Strunz (1957) and later workers showed to be a chlorite and mistakenly called manandonite. This chloritic material will probably qualify as a new species when the details of its composition and structure are better known.

### SUDOITE

Sudoite is also a di,triocahedral chlorite, and has been identified in hydrothermal alteration zones, sediments, soils, fissure veins, and low-grade metamorphic rocks from many localities. The ideal composition is  $\text{Al}_2(\text{Si}_3\text{Al})\text{O}_{10}(\text{OH})_2 \cdot (\text{Mg}_2\text{Al})(\text{OH})_6$ . Tetrahedral Al is known to range from about 0.4 to 1.1 atoms per formula unit, octahedral Al from 2.5 to 3.4, and Mg from 1.2 to 2.5. Smaller amounts of Fe, Mn, and Li are also present. Total octahedral occupancy ranges from 4.5 to 5.1 atoms per 6.0 sites. Sudoite contains only a minor amount of Li and less Al and considerably more Mg than cookeite. Representative structural formulae for sudoite are listed in Table 3.

Stanton (1984) reported two groups of iron-rich "sudoites" at the Geco mine, Manitouwadge, Ontario, with  $\text{Fe}^{2+} > \text{Mg}$  for one group. The materials are impure, however, and the average formulae of the two groups differ from those in Table 3 in several respects. Tetrahedral Al is low (0.1 to 0.3 atoms as recalculated by the present authors on the basis of 28 positive charges), the  $\text{Ca} + \text{Na} + \text{K}$  totals are moderately high (0.27 to 0.29 atoms), total octahedral occupancy is low (4.3 to 4.4 atoms per 6.0 sites, excluding  $\text{Ca} + \text{Na} + \text{K}$ ), and the total octahedral charge is negative (rather than positive) relative to an ideal complement of anions. Purer material is needed to verify the nature of this mineral.

Eggleton and Bailey (1967) and Shirozu and Higashi (1976) attempted partial refinements of the structures of different fine-grained sudoite specimens on the basis of X-ray powder diffraction data. Their results confirmed that these specimens are based on *I1b* structural units in which the 2:1 layer is dioctahedral and the interlayer is triocahedral. Drits and Lazarenko (1967) and Lin and Bailey (1985) used single crystals to show that specimens from the Ural Mountains and Belgium, respectively, have 2-layer structures of the "s" type. The powder pattern in Table 4 is indexed best by incorporation of a slight distortion to triclinic geometry. Di,dioctahedral and di,triocahedral chlorites have  $d(001)$  values that are systematically smaller than those for triocahedral specimens of similar tetrahedral compositions, thus invalidating the usage of existing  $d(001)$  graphs and equations for estimating tetrahedral Si,Al contents of such specimens. Lin and Bailey (1985) concluded from 1-dimensional electron density projections of sudoite that the smaller  $d(001)$  value is due

primarily to a thinner dioctahedral sheet within each 2:1 layer.

### DONBASSITE

Donbassite is defined as having two dioctahedral sheets and an ideal composition near  $\text{Al}_2(\text{Si}_3\text{Al})\text{O}_{10}(\text{OH})_2 \cdot \text{Al}_{2.33}(\text{OH})_6$ . For a variable amount ( $x$ ) of tetrahedral substitution the formula can be written as  $\text{Al}_2(\text{Si}_{4-x}\text{Al}_x)\text{O}_{10}(\text{OH})_2 \cdot \text{Al}_{2+\frac{x}{3}}(\text{OH})_6$ . It has been reported from soils, sediments, hydrothermal alteration zones, and fissure veins. Substitution of Al for Si in the tetrahedral sheets of the best crystallized specimens ranges from 0.6 to 1.3 atoms per 4.0 positions. The octahedral cation total is always greater than 4.0 (excluding  $\text{Ca} + \text{Na} + \text{K}$ ), ranging from about 4.2 to 4.5 atoms and indicating that the positive charge on the interlayer sheet arises primarily from the presence of  $\text{Al} + \text{Mg} + \text{Fe}$  cations in excess of 2.0 in that sheet, i.e., partly filling the "vacant" octahedron. Small amounts of octahedral Li are commonly present (Rozinova and Dubik, 1983). Representative structural formulae are listed in Table 5.

Whereas sudoite has only been identified to date as having *I1b* structural units, donbassite has only been identified as having *Ia* structural units. This statement takes into account the evolution of the definition of these names over time, i.e., some sudoite specimens as presently defined have been called donbassite in the past and vice-versa. Drits and Lazarenko (1967) identified a donbassite specimen from the Donbass region, U.S.S.R., as having a 2-layer structure like those of the equivalent "r" and "q" structures and consisting of *Ia* structural units. Aleksandrova *et al.* (1972) refined the crystal structure of a low-Li donbassite from Novaya Zemlya, U.S.S.R. They reported a regular-stacking 1-layer structure of space group *C2*, which is a distorted version of the *Ia-2* polytype of ideal symmetry *C2/m*. The vacant octahedral site in the 2:1 layer lies on the symmetry plane of the layer, but the partly vacant octahedral site in the interlayer sheet lies off the projection of the mirror plane (thereby eliminating the mirror for the crystal as a whole). Adjacent layers adopt the mica configuration. Different mean T-O bond lengths observed in this refinement using film-collected data suggest ordering of tetrahedral Si and Al, but a more accurate refinement using counter-collected data is desired for proof of this point.

No accurate powder data for purified donbassite specimens are available for comparison with those of sudoite and cookeite. Observed  $d(060)$  values are between 1.49 and 1.50 Å for donbassite and between 1.49 and 1.51 Å for sudoite and cookeite. Table 6 lists a calculated XRD pattern for donbassite based on the atomic coordinates reported by Aleksandrova *et al.* (1972) for the *Ia-2* structure. The strong  $k = 3n$  reflections are similar in positions and intensities to those

Table 2. X-ray powder diffraction patterns of cookeite specimens.

<i>hkl</i>	<i>l</i>	<i>d</i> (obs.) (Å)	<i>d</i> (calc.) (Å)	<i>hkl</i>	<i>l</i>	<i>d</i> (obs.) (Å)	<i>d</i> (calc.) (Å)						
<b>(a) Cookeite-1a(s)</b>													
002	75	14.10	14.056	13,12	7	1.680	1.681						
004	60	7.05	7.028	13,14	20	1.636	1.634						
006	100	4.68	4.685	20,12	10	1.627	1.629						
020	}	}	}	20,16	5	1.545	1.542						
110				30	4.44	13,14	10	1.538	1.537				
021				}	}	}	33 $\bar{2}$	}	}	}			
113	}	}	}				060,33 $\bar{1}$				50	1.490	1.489
112							5				4.12	336	}
008				60	3.513	3.514	064,332	8	1.458	1.457			
00,10	20	2.812	2.811	338	8	1.421	1.422						
200,13 $\bar{2}$	30	2.560	2.561	066	}	}	}						
204	}	}	}	334						1.419			
132				50				2.505	2.510	20,18			1.415
134				}	}	}	13,16	8	1.409	1.410			
202	3	2.47	2.469				13,18;33,10	6	1.375	1.373 Å			
206	}	}	}				068	}	}	}			
134				1	2.37	2.377	20,16;336						1.371
136				}	}	}	33,12						1.370
204	90	2.317	2.319				06,10	5	1.314	1.319			
208	}	}	}				338	}	}	}			
136				5	2.202	2.206	20,20				25	1.300	1.314
138				}	}	}	13,18						1.303
206	2	2.140	2.143							1.299			
20,10	}	}	}										
138				5	2.020	2.024							
00,14				10	2.010	2.017							
13,10	25	1.962	1.962										
208	12	1.953	1.956										
00,16	5	1.760	1.759										
20,14	3	1.689	1.687										
<b>(b) Cookeite-1a(r)</b>													
002	90	14.06	14.077	138	}		}						
004	35	7.03	7.039	20,10				20	2.109	2.021			
006	60	4.70	4.692	208	}	B	}						
11 $\bar{1}$	}	}	}	13,10				45	1.960	1.964			
110				20				4.47	4.459	20,10	1	1.790	1.958
112				}	}	}	13,12			1.787			
111	10	4.35	4.366				00,16	5	1.760	1.760			
113	}	}	}				13,12	}	B	}			
112				10	4.17	4.184	20,14				10	1.686	1.685
023				}	}	}	20,12				35	1.638	1.681
113	5	3.87	3.866				13,14	}	B	}			
024	}	}	}				13,14				20	1.542	1.637
115				1	3.66	3.674	20,16				15	1.538	1.632
008				60	3.518	3.519	33 $\bar{2}$	}	}	}			
116	3	3.410	3.404	060	75	1.487	1.489						
116	3	3.060	3.073	33 $\bar{1}$	}	}	}						
00,10	45	2.815	2.815	332						1.457			
200	}	}	}	064				15	1.456	1.456			
132				60	2.560	2.558	336			1.455			
132				}	}	}	334	}	}	}			
204	80	2.505	2.503				338				15	1.419	1.420
202	}	}	}				066				}	}	}
134				3	2.465	2.469	13,16	10	1.412	1.418			
134				}	}	}	20,18	}	}	}			
206	2	2.370	2.372				20,16				7	1.374	1.414
204	}	}	}				13,18				7		1.411
136				100	2.316	2.314	336	}	B	}			
02,11				1	2.220	2.220	068				7	1.371	1.377
136	}	}	}	33,10			1.373						
208				5	1.200	2.202				1.370			
206				}	}	}				1.369			
138	3	2.140	2.144										
02,12	1	2.070	2.077										

Specimen from Waisanen-Tamminen pit, Greenwood, Maine, has an imperfect "s" structure. Cell dimensions: *a* = 5.163, *b* = 8.935, *c* = 28.34 Å,  $\beta$  = 97.2°. Pattern of crystal mounted in a Gandolfi camera, 114.6-mm diameter, graphite-monochromatized FeK $\alpha$  radiation. Indexing by comparison with single crystal data, intensities estimated visually.

Specimen from North Little Rock, Arkansas, has a 2-layer "r" structure with 1a layers. Cell dimensions: *a* = 5.158, *b* = 8.927, *c* = 28.351 Å,  $\beta$  = 96.8°. Conditions as for cookeite-1a(s).

Table 2. Continued.

<i>hkl</i>	I	d(obs.) (Å)	d(calc.) (Å)	<i>hkl</i>	I	d(obs.) (Å)	d(calc.) (Å)	
(c) Cookeite-IIb								
002	80	14.10	14.100	204,136	20	2.319	2.319	
004	60	7.05	7.050	208,136	20	2.203	2.203	
006	80	4.70	4.700	206,13,8	1	2.140	2.143	
111	65	4.45	4.463	20,10;138	15	2.021	2.022	
110				208;13,10	75	1.962	1.962	
112				208;12;13,10	12	1.848	1.847	
111	5	4.35	4.321	20,10;13,12	40	1.792	1.791	
112				00,16	5	1.762	1.762	
024	5	B	3.775	20,14;13,12	15	1.686	1.686	
115				3.684	20,12;13,14	3	1.638	1.636
008	40	3.52	3.525	20,16;13,14	55	1.542	1.542	
026	1	3.22	3.239	060,332	90	1.490	1.490	
118	1	2.92	2.915	064,332,336	15	1.458	1.458	
00,10	30	2.820	2.820	066,334,338	20	1.420	1.420	
119	5	2.705	2.700	068;336;33,10	40	B	1.375	
118	1	2.62	2.630				1.372	
200,132	25	B	2.560	2.561	06,10;338;33,12	8	1.319	1.317
204,132	15		2.505	2.507				
202,134	100		2.468	2.468				
206,134	80		2.373	2.373				

Specimen from Norway, Maine. Cell dimensions:  $a = 5.161$ ,  $b = 8.938$ ,  $c = 28.41$  Å,  $\beta = 97^\circ$ . Conditions as for cookeite-1a(s).

Table 3. Structural formulae of sudoite specimens.

Specimen	Locality	Composition	Reference
1	Tracy mine, Michigan	$(Al_{2.7}Mg_{2.3})_{5.0}(Si_{3.3}Al_{0.7})O_{10}(OH)_8$	Bailey and Tyler (1960)
2	Colorado Plateau	$(R_{5.1})(Si_{3.4}Al_{0.6})O_{10}(OH)_8$	Schultz (1963)
3	Honko deposit, Kamikita mine, Japan	$(Ca_{0.11})(Al_{3.017}Fe^{3+}_{0.345}Fe^{2+}_{0.003}Mn_{0.004}Mg_{1.175})_{4.544}(Si_{3.261}Al_{0.739})O_{10}(OH)_8$	Hayashi and Oinuma (1964); Sudo and Sato (1966)
4	Furutobe mine, Japan	$(Ca_{0.060}Na_{0.17}K_{0.159})(Al_{3.042}Fe^{3+}_{0.037}Mg_{1.594})_{4.673}(Si_{3.242}Al_{0.758})O_{10}(OH)_8$	Tsukahara (1964)
5	Kaiserbach Valley, Germany	$(Ca_{0.09})(Al_{2.70}Fe^{3+}_{0.09}Fe^{2+}_{0.04}Mg_{1.97})_{4.80}(Si_{3.43}Al_{0.57})O_{10}(OH)_8$	Frenzel and Schembra (1965)
6	Berizovsk, Urals, U.S.S.R.	$(Ca_{0.03})(Al_{2.9}Fe_{0.22}Mg_{1.96})_{5.08}(Si_{2.86}Al_{1.14})O_{10}(OH)_8$	Drits and Lazarenko (1967)
7	Kazan Mineral Museum, unknown location	$(Ca_{0.14})(Al_{3.04}Fe^{2+}_{0.18}Mg_{1.61})_{4.83}(Si_{3.02}Al_{0.98})O_{10}(OH)_8$	Vlasov and Drits (1967)
8	Niida mine, Japan	$(Na_{0.03}K_{0.03})(Al_{3.03}Fe^{3+}_{0.06}Mg_{1.78})_{4.87}(Si_{3.01}Al_{0.99})O_{10}(OH)_8$	Kimbara and Nagata (1974)
9	Uchinotai deposit, Kosaka mine, Japan	$(Ca_{0.06}Na_{0.03}K_{0.05})(Al_{3.13}Fe^{3+}_{0.01}Fe^{2+}_{0.01}Mg_{1.60})_{4.75}(Si_{3.19}Al_{0.81})O_{10}(OH)_8$	Tsuzuki and Honda (1977)
10	Okunosawa deposit, Kamikita mine, Japan	$(Ca_{0.01}Na_{0.11}K_{0.01})(Al_{3.19}Fe_{0.03}Mg_{1.37})_{4.59}(Si_{3.08}Al_{0.92})O_{10}(OH)_8$	Honda (1975)
11	Hanaoka mine, Japan	$(Al_{3.2}Mg_{1.6})_{4.8}(Si_{3.2}Al_{0.8})O_{10}(OH)_8$	Shirozu and Higashi (1976)
12	Rydney Altai, U.S.S.R.	$(Ca_{0.17}Na_{0.02}K_{0.06})(Al_{2.45}Fe^{3+}_{0.03}Fe^{2+}_{0.01}Mg_{2.54})_{5.03}(Si_{3.06}Al_{0.94})O_{10}(OH)_8$	Alysheva <i>et al.</i> (1977)
13	Venn-Stavelot Massif, Ardennes, Belgium	(1) $(Al_{2.84}Fe^{3+}_{0.19}Fe^{2+}_{0.02}Mn_{0.02}Mg_{1.91})_{4.98}(Si_{3.01}Al_{0.99})O_{10}(OH)_8$ (2) $(Al_{2.86}Fe^{3+}_{0.17}Mn_{0.02}Mg_{1.94})_{4.99}(Si_{2.99}Al_{1.01})O_{10}(OH)_8$	Fransolet and Bourguignon (1978)
14	Harz Mountains, German Democratic Republic	(1) $(Al_{2.864}Fe^{3+}_{0.136}Fe^{2+}_{0.136}Mn_{0.032}Mg_{1.886})_{5.054}(Si_{3.028}Al_{0.972})O_{10}(OH)_8$ (2) $(Al_{2.871}Fe^{3+}_{0.129}Fe^{2+}_{0.136}Mn_{0.035}Mg_{1.875})_{5.046}(Si_{3.037}Al_{0.963})O_{10}(OH)_8$	Kramm (1980)



Table 4. Powder pattern of sudoite-IIb, Ottre, Belgium.

<i>hkl</i>	I	d(obs.) (Å)	d(calc.) (Å)	<i>hkl</i>	I	d(obs.) (Å)	d(calc.) (Å)
002	40	14.2	14.16	13,10	5	1.870	B
004	80	7.1	7.08	2,0,12			
006	100	4.72	4.721	1,3,10			
020	4	4.50	4.547	1,3,12	12	1.820	}
021			4.496	20,10			
021			4.483	1,3,12			
111	10	4.35	4.387	0,0,16	1	1.770	}
111			4.384	312			
022			4.340	311			
022	10	4.01	4.318	0,4,11	10	1.711	}
114			4.024	1,3,12			
114			4.004	153			
113	5	3.87	3.915	314	5	1.705	}
113			3.905	153			
025			3.561	245			
008	40	3.535	3.540	154	5	1.673	}
025			3.531	246			
115			3.361	2,0,12			
115	1	3.36	3.350	13,14	1	1.650	}
026			3.289	245			
026			3.260	157			
116	2	3.10	3.102	1,3,14	15	1.560	}
116			3.092	2,0,16			
0,0,10			2.932	1,3,14			
202	25	2.835	2.932	332	25	1.549	}
130			2.619	060			
132			2.618	332			
132	10	2.615	2.610	332	50	1.515	}
204			2.550	332			
132			2.549	336			
132	20	2.545	2.544	332	20	1.481	}
202			2.503	064			
134			2.502	338			
134	90	2.502	2.502	338	10	1.422	}
206			2.412	066			
134			2.412	334			
134	35	2.408	2.402	338	10	1.422	}
136			2.364	334			
0,0,12			2.360	066			
1,1,11	15	2.358	2.357	0,0,20	7	1.419	}
204			2.348	068			
136			2.347	3,3,10			
1,1,11	10	2.345	2.346	336	20	1.398	}
136			2.270	068			
208			2.269	1,5,13			
206	20	2.237	2.269		15	1.384	}
138			2.165				
138			2.164				
138	5	1.165	2.051		1	2.050	}
2,0,10			2.050				
0,0,14			2.023				
13,10	10	2.020	2.023		10	2.020	}
208			1.996				
1,3,10			1.980				
1,3,10	18	1.978	1.979		18	1.978	}
			1.979				

Specimen from Ottre, Belgium.  $a = 5.247$ ,  $b = 9.094$ ,  $c = 28.557$  Å,  $\alpha = 90.5^\circ$ ,  $\beta = 97.3^\circ$ , and  $\gamma = 89.9^\circ$ . Structure is of the *s*-type with *I*b layers. Conditions as listed in Table 2. B = broad.

of cookeite-Ia. The strong 021 reflection at 4.269 Å is diagnostic of the Ia-2 polytype.

EVALUATION OF STRUCTURES

The dominance of the Ia structure in cookeite and donbassite and of the IIb structure in sudoite and trioctahedral chlorites is linked directly to the magnitude and location of the positive electrostatic charge in the interlayer sheet. Bailey and Brown (1962) and Shirozu

and Bailey (1965) showed that the IIb structure is by far the most stable structural form of trioctahedral chlorite as a result of the complete absence of any cation-cation repulsion between sheets of the layered assemblage. The Ia and IIa structures, on the contrary, have exact vertical superposition of interlayer and adjacent tetrahedral cations with large resultant cation-cation repulsion values. Nevertheless, the Ia structure would have a very high structural stability rating if the

Table 5. Structural formulae for donbassite specimens.

Specimen	Locality	Composition	Reference
1	Donbass region, U.S.S.R.	(1) $(\text{Ca}_{0.17}\text{Na}_{0.19})(\text{Al}_{3.92}\text{Fe}^{3+}_{0.05}\text{Mg}_{0.21}\text{Li}_{0.05})_{4.23}(\text{Si}_{3.09}\text{Al}_{0.91})\text{O}_{10}(\text{OH})_8$ (2) $(\text{Ca}_{0.19}\text{Na}_{0.22})(\text{Al}_{3.88}\text{Fe}^{3+}_{0.08}\text{Mg}_{0.23})_{4.18}(\text{Si}_{3.06}\text{Al}_{0.94})\text{O}_{10}(\text{OH})_8$ (3) $(\text{Al}_{4.16}\text{Fe}^{3+}_{0.07}\text{Mg}_{0.11})_{4.34}(\text{Si}_{3.10}\text{Al}_{0.90})\text{O}_{10}(\text{OH})_8$	Lazarenko (1940)
2	Alberni soil, British Columbia	$(\text{Ca}_{0.02}\text{K}_{0.23}\text{X}_{0.24})(\text{Al}_{3.22}\text{Fe}^{3+}_{0.38}\text{Ti}_{0.09}\text{Mg}_{0.76})_{4.45}(\text{Si}_{2.80}\text{Al}_{1.20})\text{O}_{10}(\text{OH})_8$	Brydon <i>et al.</i> (1961)
3	Saint-Paul-de-Fenouillet, East Pyrenees	$(\text{Al}_{4.3}\text{Fe}^{3+}_{0.03}\text{Fe}^{2+}_{0.02})_{4.35}(\text{Si}_{2.70}\text{Al}_{1.30})\text{O}_{10}(\text{OH})_8$	Caillère <i>et al.</i> (1962)
4	Kesselberg, Federal Republic of Germany	(1) $(\text{Al}_{4.17}\text{Mg}_{0.10})_{4.27}(\text{Si}_{3.30}\text{Al}_{0.70})\text{O}_{10}(\text{OH})_8$ (2) $(\text{Al}_{4.27})(\text{Si}_{3.2}\text{Al}_{0.8})\text{O}_{10}(\text{OH})_8$	Müller (1961, 1963)
5	Nagol Tarasovki, Donbass, U.S.S.R.	$(\text{Ca}_{0.2})(\text{Al}_{4.0}\text{Mg}_{0.24})_{4.24}(\text{Si}_{3.12}\text{Al}_{0.88})\text{O}_{10}(\text{OH})_8$	Drits and Lazarenko (1967)
6	Namivu, Alto Ligonho, Mozambique	$(\text{Ca}_{0.04})(\text{Al}_{3.96}\text{Fe}^{3+}_{0.02}\text{Fe}^{2+}_{0.016}\text{Mn}_{0.008}\text{Mg}_{0.04}\text{Li}_{0.27})_{4.314}(\text{Si}_{3.45}\text{Al}_{0.55})\text{O}_{10}(\text{OH})_8$	Gomes (1967)
7	Novaya Zemlya, U.S.S.R.	$(\text{Al}_{4.10}\text{Fe}^{3+}_{0.04}\text{Fe}^{2+}_{0.01}\text{Mg}_{0.08}\text{Li}_{0.26})_{4.49}(\text{Si}_{3.14}\text{Al}_{0.86})\text{O}_{10}(\text{OH})_8$	Aleksandrova <i>et al.</i> (1972)
8	Donbass area, U.S.S.R.	$(\text{Al}_{3.96}\text{Fe}^{2+}_{0.08}\text{Mg}_{0.14}\text{Li}_{0.20})_{4.38}(\text{Si}_{3.36}\text{Al}_{0.64})\text{O}_{10}(\text{OH})_8$	Rozinova and Dubik (1983)
9	Szabo Bluff, Scott Glacier, Antarctica	$(\text{Al}_{4.32}\text{Fe}^{3+}_{0.07})_{4.39}(\text{Si}_{2.83}\text{Al}_{1.17})\text{O}_{10}(\text{OH})_8$	Ahn and Buseck (1988)

cation-cation repulsion factor could be minimized. This is exactly what happens in vermiculite, which contains only a few exchangeable interlayer cations and for which the structural type has been shown to be *Ia* without exception. This favorable rating of the *Ia* structure carries over to dioctahedral chlorite species in which one of the three interlayer cation sites is partly vacant (donbassite) or has a low-charge  $\text{Li}^+$  cation present (cookeite).

In the regular-stacking 1-layer chlorite polytypes involving *Ia* and *IIa* layers, two different dispositions of the 2:1 layers exist on opposing sides of the interlayer sheet. The first, termed type A here, is found only in the monoclinic structures *Ia-2* and *IIa-1*, in which successive 2:1 layers adopt the mica configuration with their 6-fold rings opposed. One interlayer cation site thus lies between the centers of these opposed fold rings, and the other two interlayer sites lie on vertical straight lines between tetrahedral cations of the two sets of rings. One form of donbassite adopts the *Ia-2* structure. The structural refinement by Aleksandrova *et al.* (1972) places the partly vacant interlayer site M between one set of superimposed tetrahedral cations and an  $\text{Al}^{3+}$  cation between the other set of two tetrahedra. The second  $\text{Al}^{3+}$  interlayer cation lies between the centers of the 6-fold rings in vertical projection. This distribution of interlayer cations gives the minimum amount of cation-cation repulsion possible for the *Ia-2* structure. The authors stated that repulsion is minimized further by tetrahedral cation ordering and local charge balance such that the source of positive interlayer charge is located between the sources of the tetrahedral negative charges, i.e., between Al-rich tetrahedra above and below. In this arrangement, the

source of positive interlayer charge is the partly vacant M octahedron. Thus, that cation-cation repulsion would be minimized even further by the opposite pattern of tetrahedral cation ordering in which the partly filled and low-charge M site would be between  $\text{Si}^{4+}$ -rich tetrahedra and the high-charge interlayer  $\text{Al}^{3+}$  would lie between Al-rich tetrahedra. Verification of the ordering pattern is desirable.

The other type A structure having a mica configuration of layers is *IIa-1*, which is not known in true chlorites, but is present in franklinfurnaceite, a hybrid chlorite-mica in which the chlorite is of the tri-dioctahedral type. In this mineral there is excellent local charge balance, with interlayer  $\text{R}^{3+}$  cations positioned midway between tetrahedral  $\text{R}^{2+}$ , interlayer  $\text{R}^{2+}$  between tetrahedral  $\text{R}^{4+}$ , and interlayer vacancies between the extra Ca that lie in the 6-fold ring openings (Peacor *et al.*, 1988).

A second disposition of 2:1 layers, termed here type B, exists in the 1-layer triclinic polytypes *Ia-4* and *IIa-3*. Here, the unique interlayer site lies on a center of symmetry between tetrahedral sites above and below, and the other two interlayer sites are asymmetric, with a tetrahedral cation on one side and the center of a 6-fold ring on the other side. Cookeite in metamorphosed bauxite from Djalair, Middle Asia, adopts the *Ia-6 = Ia-4* 1-layer structure (Vrublevskaja *et al.*, 1975). Although the cation ordering scheme is not known for that specimen, logically the low-charge  $\text{Li}^+$  should exist on the center of symmetry between tetrahedral sites above and below and the two high-charge  $\text{Al}^{3+}$  cations should be in the asymmetric locations with a tetrahedral cation on only one side. Cookeite and donbassite also adopt either or both of the 2-layer *Ia* "s" and

Table 6. Calculated powder pattern of Ia-2 donbassite, Novaya Zemlya, U.S.S.R.

<i>hkl</i>	I	<i>d</i> (calc.) (Å)	<i>hkl</i>	I	<i>d</i> (calc.) (Å)
001	60	14.127	20 $\bar{7}$	21	1.702
002	65	7.063	31 $\bar{1}$		1.693
003	100	4.709	15 $\bar{1}$		1.688
020	54	4.478	136		1.683
11 $\bar{1}$	35	4.395	151	11	1.670
021	49	4.269	15 $\bar{2}$		1.661
111	14	4.106	241		1.657
11 $\bar{2}$	93	3.983	137		1.645
022	16	3.782	24 $\bar{3}$	42	1.635
112	25	3.578	206		1.627
004	60	3.532			
11 $\bar{3}$	11	3.444	20 $\bar{8}$	16	1.557
113	28	3.058	137		1.539
114	4	2.941	33 $\bar{1}$	48	1.493
005	28	2.825	060		1.493
024	5	2.773	062	6	1.460
114	4	2.619	331		1.457
13 $\bar{1}$	47	2.569	063	21	1.423
200		2.563	332		1.418
20 $\bar{2}$	52	2.522	227	7	1.414
131		2.508	138		1.413
13 $\bar{2}$	4	2.479	00,10	31	1.413
201		2.463	33 $\bar{5}$		1.382
20 $\bar{3}$	7	2.392	064	1.375	
025		2.390	20,10	1.315	
132	97	2.371	139	1.302	
13 $\bar{3}$		2.331			
202	10	2.310			
204		2.223			
133	4	2.200			
134		2.155			
042	20	2.134			
203		2.132			
20 $\bar{5}$	49	2.042			
134		2.018			
007		2.018			
13 $\bar{5}$		1.974			
204		1.952			

Specimen from Novaya Zemlya, U.S.S.R., is a Ia-2 polytype. Pattern calculated from atomic coordinates of Aleksandrova *et al.* (1972). Cell dimensions:  $a = 5.174$ ,  $b = 8.956$ ,  $c = 14.26$  Å,  $\beta = 97.83^\circ$ . Space group =  $C2$ .

"*r*" structures (terminology of Mathieson and Walker, 1954) that have the same interlayer-layer type B configurations. The "*r*" structure has been reported for donbassite by Drits and Lazarenko (1967), although the cation-ordering pattern was not investigated. Both structures are known for cookeite. Our refinement of the cookeite "*r*" structure places the low-charge interlayer  $\text{Li}^+$  between tetrahedral cations on both sides and the two high-charge interlayer  $\text{Al}^{3+}$  cations in the asymmetric locations with a tetrahedral cation on only one side. The "*s*" structure has not been refined, but a similar local charge balance arrangement can be anticipated. The Ia "*s*" structure is also the most stable form of vermiculite (Shirozu and Bailey, 1966), in which the exchangeable cation lies vertically between tetrahedral cations above and below, with vacancies in the two asymmetric locations. At one time, the tetrahedral cation ordering was thought to place Al-rich tetrahedra

above and below the exchangeable cation to give optimum local charge balance (Shirozu and Bailey, 1966), but later work (de la Calle *et al.*, 1976) ruled out the existence of long-range tetrahedral cation ordering in vermiculite. Some evidence exists for short-range ordering in local domains (Thompson, 1984).

In contrast to the preceding species, sudoite has a trioctahedral interlayer sheet of composition  $(\text{Mg}_2\text{Al})$ , in which all of the cations have medium to high charges. And, just as in trioctahedral chlorites, the IIb structure is favored to avoid cation-cation repulsion present in an *a* position of the interlayer sheet. The presence of a vacancy in the dioctahedral 2:1 layer has little or no effect on the position of the interlayer.

It is not prudent to assume that all sudoite is of the IIb structural type and that all donbassite is of the Ia structural type, because study of more samples may show structural variations in each, just as it has in cookeite. Most cookeite and most donbassite specimens, however, are based on Ia structural units and most sudoite specimens are based on IIb units. These structural units can be identified by X-ray powder diffraction patterns. The Ia structural unit in a 2-layer dioctahedral chlorite is characterized by a strong  $d(204)$  XRD line near 2.32 Å. The IIb structural unit for a 2-layer dioctahedral chlorite is characterized by a strong  $d(202)$  line near 2.47–2.50 Å and a strong  $d(20\bar{6})$  line near 2.37–2.41 Å. Most dioctahedral chlorite specimens appear to be based on regular-stacking 2-layer structures of the "*s*" or "*r*" and "*q*" types, but regular 1-layer forms are known as well. Single crystal study is usually necessary to identify the specific stacking arrangement present.

Environment of occurrence also can be helpful in identification. The presence of essential Li in cookeite usually restricts its occurrence to Li-rich environments, rather than in sediments and soils. Essential Mg in sudoite is not very restrictive and leads to its occurrence in many different environments. An analysis for MgO should be definitive. Donbassite lacks both substantial Mg and Li, and although this restricts its occurrence to very Al-rich environments, it does not necessarily rule out occurrences in soils, sediments, hydrothermal veins, metamorphosed aluminous sediments, etc. Donbassite is less abundant in nature than sudoite and cookeite, presumably because of its unique composition. X-ray diffraction study should be supplemented by chemical analysis for suspected occurrences of all Al-rich chlorites.

#### ACKNOWLEDGMENTS

This research was supported in part by NSF grant EAR-8614868 and in part by Petroleum Research Fund grant 17966-AC2-C, administered by the American Chemical Society.



## REFERENCES

- Ahn, J. H. and Buseck, P. R. (1988) Alteration of andalusite to sheet silicates in a pegmatite: *Amer. Mineral.* **73**, 559–567.
- Aleksandrova, V. A., Drits, V. A., and Sokolova, G. V. (1972) Structural features of dioctahedral one-packet chlorite: *Soviet Physics Crystallogr.* **17**, 456–461 (Engl. transl.)
- Alysheva, E. I., Rusinova, O. V., and Chekvaidze, V. B. (1977) On sudoite from polymetal deposits of Rydnyy Altai: *Dokl. Akad. Nauk S.S.S.R.* **236**, 722–724 (in Russian).
- Bailey, S. W. (1975) Cation ordering and pseudosymmetry in layer silicates: *Amer. Mineral.* **60**, 175–187.
- Bailey, S. W. and Brown, B. E. (1962) Chlorite polytypism: I. Regular and semi-random one-layer structures: *Amer. Mineral.* **47**, 819–850.
- Bailey, S. W. and Tyler, S. A. (1960) Clay minerals associated with the Lake Superior iron ores: *Econ. Geol.* **55**, 150–175.
- Brydon, J. E., Clark, J. S., and Osborne, V. (1961) Dioctahedral chlorite: *Canadian Mineral.* **6**, 595–609.
- Caillère, S., Hénin, S., and Pobeguin, T. (1962) Présence d'un nouveau type de chlorite dans les "bauxites" de Saint-Paul-de-Fenouillet (Pyrénées-Orientales): *C.R. Acad. Sci. Paris* **254**, 1657–1658.
- de la Calle, C., Dubernat, J., Suquet, H., Pézerat, H., Gaultier, J., and Mamy, J. (1976) Crystal structure of two-layer Mg-vermiculites and Na,Ca-vermiculites: in *Proc. Internat. Clay Conf., Mexico City, 1975*, S. W. Bailey, ed., Applied Publishing, Wilmette, Illinois, 201–210.
- Černý, P. (1970) Compositional variations in cookeite: *Canadian Mineral.* **10**, 636–547.
- Černý, P., Povondra, P., and Stanek, J. (1971) Two cookeites from Czechoslovakia: A boron-rich variety and a IIb polytype: *Lithos* **4**, 7–15.
- Drits, V. A. and Lazarenko, E. K. (1967) Structural and mineralogical characteristics of donbassites: *Mineralog. Sb., L'vov* **21**, 40–48 (in Russian).
- Eggleton, R. A. and Bailey, S. W. (1967) Structural aspects of dioctahedral chlorite: *Amer. Mineral.* **52**, 673–689.
- Frank-Kamenetskii, V. A. (1960) A crystallochemical classification of simple and interstratified clay minerals: *Clay Minerals Bull.* **4**, 161–172.
- Fransolet, A.-M. and Bourguignon, P. (1978) Di/trioctahedral chlorite in quartz veins from the Ardenne, Belgium: *Canadian Mineral.* **16**, 365–373.
- Frenzel, G. and Schembra, F. W. (1965) Ein dioktaedrischer Chlorit vom Kaiserbachtal (Südfalz): *N. Jahrb. Mineral., Monats.* 108–114.
- Gomes, C. S. F. (1967) Alteration of spodumene and lepidolite with formation of dioctahedral chlorite plus dioctahedral chlorite-dioctahedral montmorillonite interstratifications: *Publ. Museu Lab. Mineral. Geol. Univ. Coimbra, Memórias Notícias* **64**, 32–57.
- Guggenheim, S., Schulze, W. A., Harris, G. A., and Lin, J.-C. (1983) Noncentric layer silicates: An optical second harmonic generation, chemical, and X-ray study: *Clays & Clay Minerals* **31**, 251–260.
- Hayashi, H. and Oinuma, K. (1964) Aluminian chlorite from Kamikita mine, Japan: *Clay Sci.* **2**, 22–30.
- Honda, S. (1975) Dioctahedral chlorite, closely associated with kaolinite from the Kamikita mine (Kuroko deposit): *Rep. Res. Inst. Underground Res. Mining College Akita Univ.* **43**, 1–8 (in Japanese).
- Kimbara, K. and Nagata, H. (1974) Clay minerals in the core samples of the mineralized zone of Niida—With special reference to the mineralogical properties of sudoite and tosudite: *Jap. Assoc. Mineral., Petrol. Econ. Geol.* **69**, 239–254 (in Japanese).
- Kramm, U. (1980) Sudoite in low-grade metamorphic manganese-rich assemblages: *N. Jahrb. Mineral., Abh.* **138**, 1–13.
- Lacroix, A. (1912) Sur une nouvelle espèce minérale (mannandonite) des pegmatites de Madagascar: *Bull. Soc. Fr. Minéral.* **35**, 223–226.
- Lacroix, A. (1922) *Minéralogie de Madagascar. I. Géologie, Descriptive Minéralogie*: Librairie Maritime Coloniale, Paris, 481–482.
- Lazarenko, E. K. (1940) Donbassite—A new group of minerals from the Donetz Basin: *Dokl. Akad. Nauk S.S.S.R.* **28**, 519–521 (in Russian).
- Lin, C.-Y. and Bailey, S. W. (1985) Structural data for sudoite: *Clays & Clay Minerals* **33**, 410–414.
- Lister, J. S. (1966) The crystal structure of two chlorites: Ph.D. thesis, Univ. Wisconsin, Madison, Wisconsin, 77 pp.
- Lister, J. S. and Bailey, S. W. (1967) Chlorite polytypism: IV. Regular 2-layer structures: *Amer. Mineral.* **52**, 1614–1631.
- Mathieson, A. McL. and Walker, G. F. (1954) Crystal structure of Mg-vermiculite: *Amer. Mineral.* **39**, 231–255.
- Müller, G. (1961) Vorläufige Mitteilung über ein neues dioktaedrisches Phyllosilikat der Chlorit-Gruppe: *N. Jahrb. Minerals Monatsh.*, 112–120.
- Müller, G. (1963) Zur Kenntnis di-oktaedrischer Vierschicht-Phyllosilikate (Sudoit-Reihe der Sudoit-Chlorit-Gruppe): in *Proc. Internat. Clay Conf., Stockholm 1963, Vol. 1*, I. Th. Rosenqvist and P. Graff-Petersen, eds., Pergamon Press, Oxford, 121–130.
- Peacor, D. R., Rouse, R. C., and Bailey, S. W. (1988) Crystal structure of franklinfurnaceite: A tri-dioctahedral zincosilicate intermediate between chlorite and mica: *Amer. Mineral.* **73**, 876–887.
- Ranoroosa, N. (1986) Etude minéralogique et microthermométrique des pegmatites de champ de la Sahatany, Madagascar: Ph.D. thesis, Univ. Paul-Sabatier, Toulouse, France, 223 pp.
- Rozinova, E. L. and Dubik, O. Yu. (1983) Dioctahedral chlorites: *Mineralog. Zh.* **5**, no. 3, 14–31 (in Russian).
- Schultz, L. G. (1963) Clay minerals in Triassic rocks of the Colorado Plateau: *U.S. Geol. Survey Bull.* **1147-C**, 1–69.
- Shirozu, H. and Bailey, S. W. (1965) Chlorite polytypism: III. Crystal structure of an orthohexagonal iron chlorite: *Amer. Mineral.* **50**, 868–885.
- Shirozu, H. and Bailey, S. W. (1966) Crystal structure of a two-layer Mg-vermiculite: *Amer. Mineral.* **51**, 1124–1143.
- Shirozu, H. and Higashi, S. (1976) Structural investigations of sudoite and regularly interstratified sericite/sudoite: *Mineral. J. (Japan)* **8**, 158–170.
- Stanton, R. L. (1984) The direct derivation of cordierite from a clay-chlorite precursor: Evidence from the Geco mine, Manitowadge, Ontario: *Econ. Geol.* **79**, 1245–1264.
- Strunz, H. (1957) Bor und Beryllium in Phyllosilikaten: *Rend. Soc. Min. Italy* **13**, p. 372.
- Sudo, T. and Sato, M. (1966) Dioctahedral chlorite: in *Proc. Internat. Clay Conf., Jerusalem 1966, Vol. 1*, L. Heller and A. Weiss, eds. Israel Prog. Sci. Transl., Jerusalem, 33–39.
- Thompson, J. G. (1984) <sup>29</sup>Si and <sup>27</sup>Al nuclear magnetic resonance spectroscopy of 2:1 clay minerals: *Clay Miner.* **19**, 220–236.
- Tsukahara, N. (1964) Dioctahedral chlorite from the Furutobe mine, Akita Prefecture, Japan: *Clay Sci.* **2**, 56–75.
- Tsuzuki, Y. and Honda, S. (1977) Three examples of Mg-Al chlorite from Kuroko deposits: *J. Mineral. Soc. Japan* **13**, 85–93 (in Japanese).

Vlasov, V. V. and Drits, V. A. (1967) Study of endogenetic dioctahedral chlorite: in *X-ray Analysis of Mineral Products Vol. 6*, Nedra Press, Moscow, 114–121 (in Russian).

Vrublevskaja, Z. V., Delitsin, I. S., Zvyagin, B. B., and Soboleva, S. V. (1975) Cookeite with a perfect regular struc-

ture, formed by bauxite alteration: *Amer. Mineral.* **60**, 1041–1046.

(Received 21 June 1988; accepted 29 October 1988; Ms. 1802)